**Full Title: Boysenberry and apple juice concentrate reduces acute lung**

**inflammation through increased alternatively activated macrophage activity in**

**an acute mouse model of allergic airways disease**

Odette M. Shaw[1*], Roger D Hurst[2], Janine Cooney[3], Gregory M. Sawyer[1], Hannah

Dinnan[1], Sheridan Martell[1]

[1]Nutrition & Health Group, Food Innovation Portfolio, The New Zealand Institute for

Plant and Food Research Limited, Palmerston North, New Zealand

[2] Food Innovation Portfolio, The New Zealand Institute for Plant and Food Research

Limited, Palmerston North, New Zealand

[3]Biological Chemistry & Bioactives Group, Food Innovation Portfolio, The New

Zealand Institute for Plant and Food Research Limited, Hamilton, New Zealand

***Correspondence:**

Odette Shaw

odette.shaw@plantandfood.co.nz

**Short title:** M2 macrophage reduce acute allergic airways inflammation

**Abstract:**

Bioactive compounds such as anthocyanins, proanthocyanins and other polyphenols are found in a wide variety of fruits and vegetables, and consumption of these have been associated with reduced lung inflammation and improved lung function in asthma and other lung diseases. We investigated whether a combination of Boysenberry and apple juice, found in BerriQi® Boysenberry and apple juice concentrate, could reduce the allergic airways inflammation associated with asthma. We characterised the polyphenolic components in BerriQi® Boysenberry and apple juice concentrate and identified the main compounds as cyanidin glycosides, ellagitannins, and chlorogenic acid. We found that consumption of 2.5 mg/kg of total anthocyanins from the BerriQi® Boysenberry and apple juice concentrate significantly reduced eosinophil infiltration following acute ovalbumin (OVA) exposure in a mouse model of allergic airways inflammation. We found that BerriQi® Boysenberry and apple juice concentrate consumption increased M2 (CD206+) macrophages and the production of the M2-associated cytokines CXCL10 and CCL4 within the lung. These results suggest that consumption of BerriQi® Boysenberry and apple juice concentrate promotes a shift towards an anti-inflammatory environment within the lung leading to reduced immune cell infiltration and tissue damage.

**Keywords: allergic airways inflammation, Boysenberry, apple, anthocyanins, ellagitannins, alternatively activated macrophages.**

## 1. Introduction

Asthma is a heterogeneous, chronic, inflammatory lung disease characterized by reversible airways obstruction, bronchospasm and infiltration of immune cells (1-3). It is estimated that 150 million people are affected by asthma worldwide, with a 5–15% prevalence in children (4) and there is evidence that early life exposure to air pollution caused by vehicle exhaust, environmental dust and industrial processes, increases the severity of asthma in children (5-7). The respiratory symptoms such as cough, and wheeze are worsened by exposure to pollution (8). Proinflammatory cytokine production in response to allergens by immune cells are further increased with concomitant pollution exposure (9-13). Eosinophils, in particular, produce reactive oxygen species, and cytokines, leading to epithelial damage and contribute to mucosal inflammation and the recruitment of other proinflammatory immune cells (14-17). These repeated acute inflammatory responses lead to tissue damage and remodelling, contributing to airway hyperresponsiveness, mucus cell hyperplasia, fixed airway flow obstruction, and loss of lung function over time (18-21).

Large-scale epidemiological studies have found that increased fruit and vegetable consumption correlates with reduced asthma symptoms (22-25). These dietary-related improvements in lung function benefits are also seen in people living in polluted environments (26-28). Fruits and vegetables contain numerous bioactive compounds, including anthocyanins and procyanidins, which have been shown to attenuate lung inflammation in cell and animal models of allergy and asthma (29-34). Human population studies have identified that dietary intake of foods high in polyphenols (28) such as apples, pears (35), carrots, tomatoes (25) and citrus is inversely correlated with the frequency and severity of reported asthma symptoms, especially wheezing and coughing (22, 25, 35, 36). Previously, we have identified

that Boysenberry consumption led to decreased chronic lung inflammation, and improved lung tissue repair in an animal model of chronic allergic lung inflammation (33). Boysenberries contain high concentrations of anthocyanins and ellagitannins (37, 38). We have also found that procyanidin-rich extracts from apple suppressed IL-4 mediated cytokine production in cell culture models of lung epithelial allergic inflammation (32, 39).

There is increasing interest in understanding the mechanisms of action that specific plant bioactives have in the human body. This is partially to better understand the benefits of consuming specific fruits and vegetables, and partially to add value to specific foods through validated health claims. There is also interest in determining if combining specific plants containing different polyphenols can augment the health benefits above those seen with the individual plant. Use of animal models, where dietary intake can be tightly controlled are useful for both demonstrating/revealing the efficacy for identified compounds, and determining the biological mechanisms of action. The aim of this study was to determine whether the combination of Boysenberries and apple, as found in BerriQi® Boysenberry and apple juice concentrate at a dose of 2.5 mg/kg total anthocyanins (TAC) could reduce allergic airways inflammation in response to acute ovalbumin exposure in a mouse model system. We also sought to determine the mechanisms involved in any ameliorating effect.

**2. Methods**

**2.1 Mice and Materials**

C57BL/6J male mice were group housed on 12h light/dark cycle in a conventional animal facility at The New Zealand Institute for Plant and Food Research Limited

(Palmerston North, New Zealand). Mice were fed Prodiet RMH1800 standard chow

for rodents (Lab Diet, St Louis, MO, USA) and filtered water ad libitum throughout

the study, all attempts to minimise suffering were made. All experimental procedures

were approved by the AgResearch Grasslands Animal Ethics Committee (AE

approvals #14839, #14731 and #14016) and carried out in accordance with the

Animal Welfare Act (1999). A proprietary Boysenberry and apple juice concentrate

(BerriQi®) was supplied by Anagenix Ltd (Auckland, New Zealand). Legendplex™

13-plex Th cytokine, proinflammatory cytokine and proinflammatory chemokine

panels, Zombie NIR™ fixable viability dye, and anti-mouse CD3 (clone 17A2), CD4

(clone GK1.5), CD8a (clone 53-6.7), CD80 (clone 16-10A1), CD86 (clone GL-1),

CD11c (clone N418), CD45 (clone 30-F11), CD206 (clone C068C2), CD14 (clone

Sa14-2), Ly6C (clone HK1.4), Gr-1 (clone RB6-8C5), I-A/I-E (MHC class II; clone

M5/114.15.2), and F4/80 (clone BM8) were purchased from Biolegend (San Diego,

CA, USA). Anti-mouse SiglecF (clone E50-2440) and CD11b (clone M1/70) were

from BD Biosciences (San Jose, CA, USA). Ovalbumin (OVA), and Alum were

purchased from Sigma (Auckland, New Zealand). iScript Advanced cDNA kit was

from Bio-Rad Laboratories (Hercules, CA, USA). Unless otherwise stated, all cell

culture media, supplements, Taqman probes and buffers were purchased from Life

Technologies NZ (Auckland, NZ).

**2.2 BerriQi® Formulation and Chemical Composition Analysis**

Boysenberries (*Rubus ursinus* var loganbaccus cv Boysenberry), harvested between

January-March 2017 from the Nelson region of NZ, were processed to a juice

concentrate by Boysenberries NZ (Nelson, NZ). Apples (*Malus domestica* –mixed

varieties) were harvested between April-June 2017 from the Hawkes Bay region of

New Zealand, and juice concentrates prepared by Profruit Ltd. (Hastings, NZ). The

freshly harvested Boysenberries and apples were graded, milled and pressed into

juice, de-seeded by centrifugation, pasteurised, depectinised by enzymatic

treatment, clarified by centrifugation, and concentrated by evaporation by their

respective manufacturers to yield fruit juice concentrates, each with an 8-fold

concentration factor.

Shott NZ Ltd. (Wellington, NZ) then blended the juice concentrates (proprietary

proportions), and added potassium sorbate preservative to a final concentration of

0.12% (w/w), yielding a dark red, non-cloudy syrup. The final BerriQi® Boysenberry

and apple juice concentrate  product measured pH 3.39 (by pH meter) and was

68.3° Brix (by refractometry), with a specific gravity of 1.35 mg/mL comprising 68%

solids measured by both gravimetric and drying methods.

The polyphenol content of the BerriQi® Boysenberry and apple juice concentrate

was determined by liquid chromatography-mass spectrometry (LC-MS) using an

LTQ linear ion trap mass spectrometer fitted with an ESI interface (ThermoFisher

Scientific, San Jose, CA, USA) coupled to an Ultimate 3000 UHPLC and PDA

detector (Dionex, Sunnyvale, CA, USA). Anthocyanins were separated on a

Poroshell 120 SB-C18,  2.1x150 mm, 2.7 µm, analytical LC column (Agilent,

Torrance, CA, USA), maintained at 70°C. The solvents were (A) 5:3:92

acetonitrile:formic acid:water v/v/v and (B) acetonitrile + 0.1% formic acid (flow rate,

200 µL/min). The initial mobile phase, 100% A, was held for 2 min before being

ramped linearly to 88% A at 14 min, returning to 5% A at 15 min and held for 4 min

before resetting to the original conditions. The sample injection volume was 10 µL.

The MS data were acquired in the positive mode. Anthocyanin concentrations are

reported as cyanidin-3-$O$-glucoside equivalents.

Other phenolic compound separation was achieved using a Hypersil GOLD aQ 1.9μ

C18 175Å (Thermo Scientific, Waltham, MA, USA), 150 × 2.1 mm column

maintained at 45°C. The solvents were (A) water + 0.1% formic acid and (B)

acetonitrile + 0.1% formic acid (flow rate, 200 μL/min). The initial mobile phase, 95%

A/5% B, was ramped linearly to 85% A at 10 min, held for 3.75 min, then ramped

linearly to 75% A at 18 min, 67.2% A at 25 min, 50% A at 28 min, 3% A at 29 min and

held for 4 min before resetting to the original conditions. The sample injection volume

was 4 μL. The MS data were acquired in the negative mode.

**2.3 OVA-Induced Airway Inflammation Model**

Allergic airway disease was induced as previously described (34, 40). For the

Boysenberry and apple interventions, mice were randomized into receiving either

water (vehicle control) or 2.5 mg/kg TAC in the BerriQi® Boysenberry and apple

juice concentrate as previously described (34). Briefly, mice were fasted for 4 h

before being orally gavaged with water (control) or at a dose of 2.5 mg/kg body

weight TAC in the BerriQi® Boysenberry and apple juice concentrate made up to a

total volume of 200 μL in water 1h before OVA challenge and again 2 days post-

challenge. Mice were sacrificed 4 days following intranasal ovalbumin challenge and

immune parameters were analyzed.

**2.4 Immune Parameter Analysis**

Bronchoalveolar lavage fluid (BALF) and lung tissues were collected as previously

described and immune cells were phenotyped by flow cytometry (40). Lung tissue

supernatant for cytokine analysis was prepared as previously described (34). Briefly,

the left lung lobe was minced into 500 μL phenol red-free complete RPMI media and

incubated at 37˚C for 30 min before being filtered through a 40 μm mesh and

centrifuged to remove cellular material. The resulting supernatant was analysed for cytokine concentration using Legendplex bead-based multiplex immunoassays as per the manufacturer's instruction. Both cell phenotyping and the cytokine multiplex assays were analyzed using a BD FACSverse (BD Biosciences, San Jose, CA, USA). H&E and AB-PAS histological staining were performed by Massey IVABS histology unit.

**2.5 Real-time qPCR analysis**

Mouse lung tissue was collected 4 days following OVA challenge and snap frozen in liquid nitrogen. The lung samples were crushed into powder using a mortar and pestle with liquid nitrogen to keep the samples frozen. The RNA was extracted from the powder using a TRIZOL total RNA extraction protocol. RNA was quantified using an LVis plate in a POLARstar Omega plate reader (BMG) and the quality of the ribosomal RNA bands confirmed by agarose gel electrophoresis (data not shown). Five µg of RNA from each sample was used as the template for cDNA synthesis using the iScript™ cDNA Synthesis Kit. Taqman® Gene Expression Assays were purchased for each gene of interest. Two housekeeping genes, GAPDH and β-actin, were used as controls to determine the amount of relative gene expression. Taqman® Gene Expression Master Mix was used to PCR amplify the genes in a Bio-Rad™ CFX384™ Real-Time PCR Detection System. Three lung samples per treatment group were prepared and amplified in quadruplicate with the housekeeping genes amplified on the same 384-well plate.

**2.6 Statistical Analysis**

Data were analyzed using one-way analysis of variance (ANOVA) with a Tukey's

post hoc test and graphed in SigmaPlot 12.5 (Systat Software Inc., San Jose, CA,

USA).

**3. Results**

**3.1 Chemical Composition of the Boysenberry and Apple Juice Concentrate**

The results of the LC-MS analysis showed that cyanidin glycosides, ellagitannins,

and chlorogenic acid were the major components in BerriQi® Boysenberry and apple

juice concentrate (Table 1, supplementary figures 1, 2). Minor components included

phloretin 2-O-glucoside, and a mix of phenolic acids, flavonol glycosides, flavanol

monomers and procyanidins. The major classes of phenolic compounds were

anthocyanins (1969 µg/mL) and hydrolysable tannins (946 µg/mL), accounting for

56% and 27%, respectively, of the total phenolics quantified. The most abundant

tannins were ellagic acid (449 µg/mL) and sanguiin H6 (213 µg/mL).

**3.2 Effect of Boysenberry and Apple juice concentrate Intervention on**

**Ovalbumin-Induced Allergic Airways Inflammation**

Acute intranasal OVA exposure resulted in an infiltration of immune cells into the

lung (Figure 1A) and increased mucous production (Figure 1B). Consumption of 2.5

mg/kg TAC BerriQi® Boysenberry and apple juice concentrate reduced the

infiltration of immune cells and decreased OVA-induced mucous production (Figure

1A-B). We quantified the type and number of immune cells infiltrating into the lung,

and found that acute OVA exposure significantly increased (P<0.001) infiltrating

eosinophils (CD45+/CD11b+/SiglecF+), neutrophils (CD45+Ly6C+Gr-1+) and T-cells

(CD45+/CD3+/CD4+ or CD45+/CD3+/CD8a+), compared with the lung of naïve

animals (Figure 1C-F). Compared with animals only exposed to OVA, those that also

consumed 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice concentrate showed a significant decrease (P<0.001) in the number of infiltrating eosinophils, neutrophils and T cells in the lung (Figure 1C-F). We saw no change in the number of CD4+ or CD8+ T cells in the mediastinal (lung-draining) lymph node for any of the treatment groups (Figure 2A-B).

There was a trend towards an increased percentage of CD206+/CD14- macrophages in the lungs of mice that consumed 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice concentrate (Figure 3A). We measured the gene expression of Arg1, Ym-1 and Fizz1 in lung tissue and found that 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice concentrate consumption led to a significant fold- increase in Arg1 (P<0.05) and Fizz1 (P<0.01) gene expression compared to naïve mice (Table 2). We found no significant fold-change in Nos2 or Ym-1 gene expression between any of the treatment groups (Table 2).

Consumption of 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice concentrate led to increased levels of the cytokines IL-17A, CXCL10, and CCL4 (Figure 3B-D)  4 days following OVA challenge, but did not affect the IL-17F concentration (Figure 3E). We saw no effect on the concentrations of the classical Th1/2 cytokines IFNγ, IL-5, IL-9 or IL-10 in either the BerriQi® Boysenberry and apple juice concentrate treated or the OVA alone mice compared to naïve controls (Figure 2C-F).

**4. Discussion**

We evaluated the effects of dietary supplementation with 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice concentrate, on the immune responses in a mouse model of acute allergic airways inflammation. Our results show that consumption of 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice concentrate reduced

granulocyte and local T cell infiltration into the lung after OVA challenge, but did not

alter T cell activation within the lung draining lymph node or the levels of classical

Th-2 and Th-1 cytokines in the lung at four days following OVA challenge. Our

current results indicated that BerriQi® Boysenberry and apple juice concentrate had

little impact on the Th-2/Th-1 mediated allergic response of mice, but rather targeted

innate proinflammatory immune pathways. This is consistent with our previously

reported finding in a mouse model of chronic allergic airways inflammation using 10

mg/kg TAC Boysenberry juice concentrate (33). Chemical composition analysis

showed that the BerriQi® Boysenberry and apple juice concentrate formulation

contained high concentrations of cyanidin glycosides, ellagitannins, and chlorogenic

acid. These compounds have been previously shown to reduce inflammatory

signalling in vitro (38, 41, 42), and in vivo animal models of inflammation (34, 43-46).

Our current results suggest that consumption of 2.5 mg/kg TAC BerriQi®

Boysenberry and apple juice concentrate, which also contains high levels of

ellagitannins and chlorogenic acid, could have broader lung health benefits beyond

allergic asthma disease by promoting the resolution of inflammation caused by

innate immune cell overactivation. Current asthma therapies also suppress both the

adaptive and innate immune responses without affecting the aberrant sensitivity to

the allergen and this can lead to adverse events.

Consumption of BerriQi® Boysenberry and apple juice concentrate had less of an

effect on monocyte/macrophage infiltration into the lung than on granulocyte

infiltration, and there was increased percentage of CD206+ monocytes. This could

represent a shift to an M2 anti-inflammatory phenotype. We then measured the

changes in gene expression for Arg1, Ym-1 and Fizz1, the classic genes for

identifying alternatively activated macrophages (47, 48). Arg1 and Fizz1 gene

expression was significantly increased with BerriQi® Boysenberry and apple juice

concentrate consumption. Consistent with a shift to a more anti-inflammatory

macrophage phenotype, we detected increased levels of CXCL10 and CCL4

cytokines, which are produced by M2 macrophages, in the lungs of mice that

consumed BerriQi® Boysenberry and apple juice concentrate. These results suggest

that the consumption of Boysenberry and apple juice concentrate led to a switch to

the M2 phenotype in OVA-challenged mice. This could be one of the mechanisms by

which BerriQi® Boysenberry and apple juice concentrate consumption contributed to

the resolution of inflammation. Previously, we reported that 10 mg/kg TAC

Boysenberry juice concentrate can increase the abundance of alternatively activated

(M2) macrophages, which promote tissue repair in a chronic model of airways

inflammation (33). Our current results suggest that BerriQi® Boysenberry and apple

juice concentrate, had a similar effect in this mouse model of acute inflammation, in

particular the increase in Arg1 gene expression is similar to our previously reported

study showing increased arginase protein expression by alternatively activated

macrophages (33). Further, research looking at an animal model Th2-mediated

inflammation has identified M2 macrophage derived Fizz1 as a key limiting factor for

Th2-mediated pulmonary inflammation (49).

The mice that consumed 2.5 mg/kg TAC BerriQi® Boysenberry and apple juice

concentrate showed increased levels of the cytokines IL-17A, CXCL10, and CCL4,

but the levels of IL-17F were not affected. High IL-17 and IL-17F levels have been

implicated in asthma pathogenesis, however, there is also evidence that IL-17A (50)

can increase the abundance of MMP-9, an important tissue remodelling protein in

asthma (33) as well as inducing apoptosis of neutrophils and eosinophils (50).

CXCL10 and CCL4 are chemokines that attract monocytes/macrophages, and

CXCL10 may also inhibit the infiltration of eosinophils in response to allergic airways inflammation (51). The combination of increased IL-17A-mediated granulocyte apoptosis and CXCL10-mediated inhibition of granulocyte infiltration could explain how the consumption of the BerriQi® Boysenberry and apple juice concentrate resulted in decreased allergic airways inflammation, in particular the reduced number of eosinophils and neutrophils.

This switch to a M2 macrophage phenotype may be through the Boysenberry and apple polyphenols identified in the BerriQi® Boysenberry and apple juice concentrate directly inhibiting proinflammatory pathways, or through an indirect shortening of the proinflammatory phase. Other studies have shown that increased dietary fibre shortened the duration of the proinflammatory phase, leading to reduced tissue damage in a chronic mouse model of allergic asthma (52). Based on the estimated fibre content in BerriQi® Boysenberry and apple juice concentrate (Supplementary table 1), it is unlikely that the dietary fibre component played a significant role in the immune modulation we observed. However, the combination of anthocyanins with other polyphenols identified in the BerriQi® Boysenberry and apple juice concentrate could have a similar effect on inflammation either by shortening the proinflammatory time course, or promoting the production of anti-inflammatory proteins. Ellagitannins have been shown in cell culture and animal models of chronic inflammatory diseases to reduce proinflammatory prostaglandins (53), cytokines (45, 54), and other proteins (42, 55, 56). Anthocyanins have also been shown to inhibit proinflammatory proteins (57, 58), and activate anti-inflammatory pathways in models of inflammation (59-63). It is possible that the combination of the different polyphenols in the BerriQi® Boysenberry and apple juice concentrate act on a number of different immune pathways to regulate the immune responses to OVA.

We found that mice that consumed BerriQi® Boysenberry and apple juice

concentrate had reduced immune cell infiltration in response to acute OVA challenge

and this could be as a result of a shift towards an anti-inflammatory environment

within the lung. These results highlight the potential of anthocyanin-rich Boysenberry

and apple dietary supplementation to modulate innate immune pathways during

acute allergic lung inflammation. Further work is needed to determine if these

pathways are also altered in other lung inflammatory conditions, such as air pollution

exposure. Clinical studies are needed to show if these findings are translatable to

human health.

**Conflict of Interest**

All authors were employed by The New Zealand Institute of Plant & Food Research

Limited, a New Zealand Crown Research Institute wholly owned by the New Zealand

Government for the purposes of research into sustainable production, elite breeding,

food and health science of horticultural, arable, and seafood products.

OMS and RDH report that they are named on patents related to the formulation of

BerriQi® Boysenberry and apple juice concentrate, but have not received any

financial compensation, nor will receive any personal royalty payments as a result of

this.  None of the other authors declare any other conflicts of interest. Under the

terms of the Innovation Cell™ Collaboration Agreement Plant & Food Research and

Anagenix have a royalty sharing agreement for any royalties that result from the sale

of BerriQi® Boysenberry and apple juice concentrate product.

**Author Contributions**

OMS designed, performed, analyzed and interpreted the in vivo studies, and wrote

and edited the manuscript; JC performed, analyzed and interpreted the chemcial

composition experiments and GMS performed, analyzed and interpreted in vivo

studies, and both contributed to the writing and editing of the manuscript; HD and

SM performed the in vivo studies and helped edit the manuscript; RDH designed and

directed the overall research programme and helped edit the manuscript.

**Funding**

This work was funded by The New Zealand Institute of Plant & Food Research

Limited as a contribution to the Innovation Cell™ Collaboration Agreement executed

between Anagenix Limited and The New Zealand Institute for Plant and Food

Research Limited on 5 September 2016 (PFR reference #33609).

The authors declare that this study received funding from The New Zealand Institute of Plant & Food Research Limited. The funder was not involved in the study design, collection, analysis, and interpretation of data, the writing of this article or the decision to submit it for publication.

**Acknowledgements**

The authors acknowledge the efforts of TC Chadderton, and Brendan Vercoe in securing internal funding for this work. The authors would like to thank Andrew Carroll for assisting with data analysis and Jenny Smith for her invaluable help preparing the manuscript.

**5. References**

1.	Agache I, Akdis CA. Endotypes of allergic diseases and asthma: An important step in building blocks for the future of precision medicine. Allergol Int. 2016;65(3):243-52.

2.	Agrawal DK, Bharadwaj A. Allergic airway inflammation. Curr Allergy Asthma Rep. 2005;5(2):142-8.

3.	Barnes PJ. Pathophysiology of asthma. Br J Clin Pharmacol. 1996;42(1):3-10.

4.	WHO. Prevention of allergy and allergic asthma: based on the WHO/WAO Meeting on the Prevention of Allergy and Allergic Asthma, Geneva, 8-9 January 2002. Management DoND, Arthritis CRDa, Organization WA, Organization. WH, editors. Geneva: World Health Organization; 2003.

5.	Hsu HH, Chiu YH, Coull BA, Kloog I, Schwartz J, Lee A, et al. Prenatal Particulate Air Pollution and Asthma Onset in Urban Children. Identifying Sensitive Windows and Sex Differences. Am J Respir Crit Care Med. 2015;192(9):1052-9.

6.      Miller RL, Peden DB. Environmental effects on immune responses in patients

with atopy and asthma. J Allergy Clin Immunol. 2014;134(5):1001-8.

7.      Jung DY, Leem JH, Kim HC, Kim JH, Hwang SS, Lee JY, et al. Effect of

Traffic-Related Air Pollution on Allergic Disease: Results of the Children's Health and

Environmental Research. Allergy Asthma Immunol Res. 2015;7(4):359-66.

8.      Hoek G, Pattenden S, Willers S, Antova T, Fabianova E, Braun-Fahrlander C,

et al. PM10, and children's respiratory symptoms and lung function in the PATY

study. Eur Respir J. 2012;40(3):538-47.

9.      Acciani TH, Brandt EB, Khurana Hershey GK, Le Cras TD. Diesel exhaust

particle exposure increases severity of allergic asthma in young mice. Clin Exp

Allergy. 2013;43(12):1406-18.

10.     Brandt EB, Biagini Myers JM, Acciani TH, Ryan PH, Sivaprasad U, Ruff B, et

al. Exposure to allergen and diesel exhaust particles potentiates secondary allergen-

specific memory responses, promoting asthma susceptibility. J Allergy Clin Immunol.

2015;136(2):295-303 e7.

11.     Carlsten C, Blomberg A, Pui M, Sandstrom T, Wong SW, Alexis N, et al.

Diesel exhaust augments allergen-induced lower airway inflammation in allergic

individuals: a controlled human exposure study. Thorax. 2016;71(1):35-44.

12.     Kim J, Natarajan S, Vaickus LJ, Bouchard JC, Beal D, Cruikshank WW, et al.

Diesel exhaust particulates exacerbate asthma-like inflammation by increasing CXC

chemokines. Am J Pathol. 2011;179(6):2730-9.

13.     van Voorhis M, Knopp S, Julliard W, Fechner JH, Zhang X, Schauer JJ, et al.

Exposure to atmospheric particulate matter enhances Th17 polarization through the

aryl hydrocarbon receptor. PLoS One. 2013;8(12):e82545.

14.    Trivedi SG, Lloyd CM. Eosinophils in the pathogenesis of allergic airways

disease. Cell Mol Life Sci. 2007;64(10):1269-89.

15.    Amin K, Janson C, Bystrom J. Role of Eosinophil Granulocytes in Allergic

Airway Inflammation Endotypes. Scand J Immunol. 2016;84(2):75-85.

16.    Bossley CJ, Fleming L, Gupta A, Regamey N, Frith J, Oates T, et al. Pediatric

severe asthma is characterized by eosinophilia and remodeling without T(H)2

cytokines. J Allergy Clin Immunol. 2012;129(4):974-82 e13.

17.    Brown JR, Kleimberg J, Marini M, Sun G, Bellini A, Mattoli S. Kinetics of

eotaxin expression and its relationship to eosinophil accumulation and activation in

bronchial biopsies and bronchoalveolar lavage (BAL) of asthmatic patients after

allergen inhalation. Clin Exp Immunol. 1998;114(2):137-46.

18.    Bergeron C, Al-Ramli W, Hamid Q. Remodeling in asthma. Proc Am Thorac

Soc. 2009;6(3):301-5.

19.    Ahdieh M, Vandenbos T, Youakim A. Lung epithelial barrier function and

wound healing are decreased by IL-4 and IL-13 and enhanced by IFN-gamma. Am J

Physiol Cell Physiol. 2001;281(6):C2029-38.

20.    Al-Muhsen S, Johnson JR, Hamid Q. Remodeling in asthma. J Allergy Clin

Immunol. 2011;128(3):451-62.

21.    Brightling CE, Gupta S, Gonem S, Siddiqui S. Lung damage and airway

remodelling in severe asthma. Clin Exp Allergy. 2012;42(5):638-49.

22.    McKeever TM, Britton J. Diet and asthma. Am J Respir Crit Care Med.

2004;170(7):725-9.

23.    Nurmatov U, Devereux G, Sheikh A. Nutrients and foods for the primary

prevention of asthma and allergy: systematic review and meta-analysis. J Allergy

Clin Immunol. 2011;127(3):724-33 e1-30.

24.    Okoko BJ, Burney PG, Newson RB, Potts JF, Shaheen SO. Childhood

asthma and fruit consumption. Eur Respir J. 2007;29(6):1161-8.

25.    Rosenlund H, Kull I, Pershagen G, Wolk A, Wickman M, Bergstrom A. Fruit

and vegetable consumption in relation to allergy: disease-related modification of

consumption? J Allergy Clin Immunol. 2011;127(5):1219-25.

26.    Stevens EL, Rosser F, Forno E, Peden D, Celedon JC. Can the effects of

outdoor    air    pollution    on    asthma    be    mitigated?    J    Allergy    Clin    Immunol.

2019;143(6):2016-8 e1.

27.    Burbank AJ, Duran CG, Pan Y, Burns P, Jones S, Jiang Q, et al. Gamma

tocopherol-enriched    supplement    reduces    sputum    eosinophilia    and    endotoxin-

induced    sputum    neutrophilia    in    volunteers    with    asthma.    J    Allergy    Clin    Immunol.

2018;141(4):1231-8 e1.

28.    Pounis G, Arcari A, Costanzo S, Di Castelnuovo A, Bonaccio M, Persichillo M,

et al. Favorable    association    of    polyphenol-rich    diets    with    lung    function:    Cross-

sectional findings from the Moli-sani study. Respir Med. 2018;136:48-57.

29.    Park SJ, Shin WH, Seo JW, Kim EJ. Anthocyanins inhibit airway inflammation

and    hyperresponsiveness    in    a    murine    asthma    model.    Food    Chem    Toxicol.

2007;45(8):1459-67.

30.    Coleman SL, Shaw OM. Progress in the understanding of the pathology of

allergic asthma and the potential of fruit proanthocyanidins as modulators of airway

inflammation. Food Funct. 2017;8(12):4315-24.

31.    Coleman S, Kruger M, Sawyer G, Hurst R. Procyanidin A2 Modulates IL-4-

Induced    CCL26    Production    in    Human    Alveolar    Epithelial    Cells.    Int    J    Mol    Sci.

2016;17(11):1888.

32.     Sawyer GM, Stevenson DE, McGhie TK, Hurst RD. Suppression of CCL26

and CCL11 generation in human alveolar epithelial cells by apple extracts containing

procyanidins. J Funct Foods. 2017;31:141-51.

33.     Shaw OM, Hurst RD, Harper JL. Boysenberry ingestion supports fibrolytic

macrophages with the capacity to ameliorate chronic lung remodeling. Am J Physiol

Lung Cell Mol Physiol. 2016;311(3):L628-38.

34.     Shaw OM, Nyanhanda T, McGhie TK, Harper JL, Hurst RD. Blackcurrant

anthocyanins modulate CCL11 secretion and suppress allergic airway inflammation.

Mol Nutr Food Res. 2017;61(9):1600868.

35.     Garcia-Larsen V, Thawer N, Charles D, Cassidy A, van Zele T, Thilsing T, et

al. Dietary Intake of Flavonoids and Ventilatory Function in European Adults: A

GA(2)LEN Study. Nutrients. 2018;10(1).

36.     Garcia V, Arts IC, Sterne JA, Thompson RL, Shaheen SO. Dietary intake of

flavonoids and asthma in adults. Eur Respir J. 2005;26(3):449-52.

37.     Cooney JM, Jensen DJ, McGhie TK. LC-MS identification of anthocyanins in

boysenberry extract and anthocyanin metabolites in human urine following dosing. J

Sci Food Agr. 2004;84(3):237-45.

38.     McGhie TK, Martin H, Lunken RC. The combination of analytical-scale HPLC

separation with a TR-FRET assay to investigate JAK2 inhibitory compounds in a

Boysenberry drink. Food Funct. 2012;3(11):1170-5.

39.     Coleman SL, Hurst RD, Sawyer GM, Kruger MC. The in vitro evaluation of

isolated procyanidins as modulators of cytokine-induced eotaxin production in

human alveolar epithelial cells. J Berry Res. 2016;6(2):115-24.

40.     Shaw OM, Harper JL. An efficient single prime protocol for the induction of

antigen-induced airways inflammation. J Immunol Methods. 2013;395(1-2):79-82.

41. Cassidy A, Rogers G, Peterson JJ, Dwyer JT, Lin H, Jacques PF. Higher

dietary anthocyanin and flavonol intakes are associated with anti-inflammatory

effects in a population of US adults. Am J Clin Nutr. 2015;102(1):172-81.

42. El-Shitany NA, El-Bastawissy EA, El-desoky K. Ellagic acid protects against

carrageenan-induced acute inflammation through inhibition of nuclear factor kappa

B, inducible cyclooxygenase and proinflammatory cytokines and enhancement of

interleukin-10 via an antioxidant mechanism. Int Immunopharmacol. 2014;19(2):290-

9.

43. Auclair S, Silberberg M, Gueux E, Morand C, Mazur A, Milenkovic D, et al.

Apple polyphenols and fibers attenuate atherosclerosis in apolipoprotein E-deficient

mice. J Agric Food Chem. 2008;56(14):5558-63.

44. Denis MC, Roy D, Yeganeh PR, Desjardins Y, Varin T, Haddad N, et al. Apple

peel polyphenols: a key player in the prevention and treatment of experimental

inflammatory bowel disease. Clin Sci (Lond). 2016;130(23):2217-37.

45. Guan S, Zheng YY, Yu X, Li WL, Han B, Lu J. Ellagic acid protects against

LPS-induced acute lung injury through inhibition of nuclear factor kappa B,

proinflammatory cytokines and enhancement of interleukin-10. Food Agr Immunol.

2017;28(6):1347-61.

46. Impellizzeri D, Talero E, Siracusa R, Alcaide A, Cordaro M, Maria Zubelia J,

et al. Protective effect of polyphenols in an inflammatory process associated with

experimental pulmonary fibrosis in mice. Br J Nutr. 2015;114(6):853-65.

47. Chung S, Lee TJ, Reader BF, Kim JY, Lee YG, Park GY, et al. FoxO1

regulates allergic asthmatic inflammation through regulating polarization of the

macrophage inflammatory phenotype. Oncotarget. 2016;7(14):17532-46.

48.    Kurowska-Stolarska M, Stolarski B, Kewin P, Murphy G, Corrigan CJ, Ying S,

et al. IL-33 amplifies the polarization of alternatively activated macrophages that

contribute to airway inflammation. J Immunol. 2009;183(10):6469-77.

49.    Nair MG, Du Y, Perrigoue JG, Zaph C, Taylor JJ, Goldschmidt M, et al.

Alternatively activated macrophage-derived RELM-{alpha} is a negative regulator of

type 2 inflammation in the lung. J Exp Med. 2009;206(4):937-52.

50.    Linden A, Dahlen B. Interleukin-17 cytokine signalling in patients with asthma.

Eur Respir J. 2014;44(5):1319-31.

51.    Su YC, Rolph MS, Hansbro NG, Mackay CR, Sewell WA. Granulocyte-

macrophage colony-stimulating factor is required for bronchial eosinophilia in a

murine model of allergic airway inflammation. J Immunol. 2008;180(4):2600-7.

52.    Trompette A, Gollwitzer ES, Yadava K, Sichelstiel AK, Sprenger N, Ngom-Bru

C, et al. Gut microbiota metabolism of dietary fiber influences allergic airway disease

and hematopoiesis. Nat Med. 2014;20(2):159-66.

53.    Karlsson S, Nanberg E, Fjaeraa C, Wijkander J. Ellagic acid inhibits

lipopolysaccharide-induced expression of enzymes involved in the synthesis of

prostaglandin E2 in human monocytes. Br J Nutr. 2010;103(8):1102-9.

54.    Cornelio Favarin D, Martins Teixeira M, Lemos de Andrade E, de Freitas

Alves C, Lazo Chica JE, Arterio Sorgi C, et al. Anti-inflammatory effects of ellagic

acid on acute lung injury induced by acid in mice. Mediators Inflamm.

2013;2013:164202.

55.    Marin M, Maria Giner R, Rios JL, Recio MC. Intestinal anti-inflammatory

activity of ellagic acid in the acute and chronic dextrane sulfate sodium models of

mice colitis. J Ethnopharmacol. 2013;150(3):925-34.

56. Saba, Khan S, Parvez S, Chaudhari B, Ahmad F, Anjum S, et al. Ellagic acid

attenuates bleomycin and cyclophosphamide-induced pulmonary toxicity in Wistar

rats. Food Chem Toxicol. 2013;58:210-9.

57. Esposito D, Chen A, Grace MH, Komarnytsky S, Lila MA. Inhibitory effects of

wild blueberry anthocyanins and other flavonoids on biomarkers of acute and chronic

inflammation in vitro. J Agric Food Chem. 2014;62(29):7022-8.

58. Fu Y, Zhou E, Wei Z, Wang W, Wang T, Yang Z, et al. Cyanidin-3-O-beta-

glucoside ameliorates lipopolysaccharide-induced acute lung injury by reducing

TLR4 recruitment into lipid rafts. Biochem Pharmacol. 2014;90(2):126-34.

59. Chen YF, Shibu MA, Fan MJ, Chen MC, Viswanadha VP, Lin YL, et al. Purple

rice anthocyanin extract protects cardiac function in STZ-induced diabetes rat hearts

by inhibiting cardiac hypertrophy and fibrosis. J Nutr Biochem. 2016;31:98-105.

60. Edirisinghe I, Banaszewski K, Cappozzo J, McCarthy D, Burton-Freeman BM.

Effect of black currant anthocyanins on the activation of endothelial nitric oxide

synthase (eNOS) in vitro in human endothelial cells. J Agric Food Chem.

2011;59(16):8616-24.

61. Khanna S, Roy S, Bagchi D, Bagchi M, Sen CK. Upregulation of oxidant-

induced VEGF expression in cultured keratinocytes by a grape seed

proanthocyanidin extract. Free Radic Biol Med. 2001;31(1):38-42.

62. Koh ES, Lim JH, Kim MY, Chung S, Shin SJ, Choi BS, et al. Anthocyanin-rich

Seoritae extract ameliorates renal lipotoxicity via activation of AMP-activated protein

kinase in diabetic mice. J Transl Med. 2015;13(1):203.

63. Liu Y, Tan D, Tong C, Zhang Y, Xu Y, Liu X, et al. Blueberry anthocyanins

ameliorate radiation-induced lung injury through the protein kinase RNA-activated

pathway. Chem Biol Interact. 2015;242:363-71.

**Table 1. Phenolic compounds detected in BerriQi® (µg/mL).** $M^+$ and $(M-H)^-$ ions

are the pseudomolecular ions used for identification of compounds by liquid

chromatography-mass spectrometry (LC-MS). All identifications confirmed by

MS/MS$^n$ experiments. Peak numbers refer to chromatograms shown in

supplementary data. # detected as [M+formate]$^-$ adduct

| Peak | $M^+$ | $(M-H)^-$ | Compound | µg/mL |
|---|---|---|---|---|
| | | | **Anthocyanins** | |
| 1 | 611 | | Cyanidin 3-*O*-sophoroside | 883 |
| 2 | 449 | | Cyanidin 3-*O*-glucoside | 571 |
| 3 | 481 | | Cyanidin 3-*O*-sambubioside | 24 |
| 4 | 757 | | Cyanidin 3-*O*-(2-glucosylrutinoside) | 411 |
| 5 | 595 | | Cyanidin 3-*O*-rutinoside | 62 |
| 6 | 727 | | Cyanidin 3-*O*-xylosylrutinoside | 18 |
| | | | **Phenolic acids** | |
| 7 | | 169 | Gallic acid | 140 |
| 8 | | 153 | Protocatechuric acid | 35 |
| 10 | | 353 | Chlorogenic acid | 69 |
| 11 | | 179 | Caffeic acid | 7 |
| 15 | | 337 | 4-*p*-Coumaroylquinic acid | 26 |
| 17 | | 337 | 5-*p*-Coumaroylquinic acid | 3 |
| | | | **Flavan-3-ols and procyanidins** | |
| 9 | | 335# | Catechin | 3 |
| 12 | | 577 | Procyanidin B2 | 6 |
| 13 | | 335# | *Epi*-catechin | 21 |
| 25 | | 579 | unknown procyanidin isomer | 3 |
| | | | **Hydrolysable tannins** | |
| 14 | | 1567 | Sanguiin H10 isomer 1 | 9 |
| 16 | | 469 | Sanguisorbic acid dilactone | 120 |
| 19 | | 2036.5 | Galloyl-SH6 | 66 |
| 20 | | 1567 | Sanguiin H10 isomer 2 | 61 |
| 21 | | 2501 | Lambertian C (minus ellagic acid) | 11 |
| 22 | | 2803 | Lambertian C | 17 |
| 24 | | 1869 | Sanguiin H6 | 213 |
| 26 | | 301 | Ellagic acid | 449 |
| | | | **Flavonols** | |
| 27 | | 609 | Quercetin 3-*O*-rutinoside | 6 |
| 28 | | 463 | Quercetin 3-*O*-galactoside | 17 |

| 29 | 477 | Quercetin 3-*O*-glucuronide | 36 |
|----|-----|------------------------------|----|
| 30 | 463 | Quercetin 3-*O*-glucoside | 13 |
| 31 | 433 | Quercetin 3-*O*-pentoside 1 | 9 |
| 32 | 433 | Quercetin 3-*O*-pentoside 3 | 6 |
| 33 | 433 | Quercetin 3-*O*-pentoside 2 | 13 |
| 35 | 447 | Quercetin 3-*O*-rhamnoside | 15 |
| 37 | 301 | Quercetin | 19 |
| | | **Chalcones** | |
| 34 | 567 | Phloretin 2-*O*-xylo-glucoside | 9 |
| 36 | 481[#] | Phloretin 2-*O*-glucoside | 62 |
| | | **Unknowns** | |
| 18 | 563[#] | unknown | 71 |
| 23 | 639 | unknown | 22 |
| | | **Total** | **1557** |

**Table 2: BerriQi® Boysenberry and apple juice concentrate increases**

**alternatively activated macrophage gene expression in the lung.** Mice were

primed with ovalbumin (OVA)/Alum i.p. and then challenged 7 days later with OVA

i.n. (Day 0). Mice were orally gavaged with 2.5 mg/kg total anthocyanins (TAC) in the

BerriQi® Boysenberry and apple juice concentrate (BerriQi) 1h before OVA

challenge and again 2 days post-challenge. Mean fold-change (SEM) in gene

expression was measured by real-time qPCR in lung tissue 4 days post-OVA

challenge. *$P<0.05$, **$P<0.01$ compared with naïve (one-way ANOVA with Tukey's

Post Hoc test) for 4 experimental replicates with n=3 per treatment groups.

| Target Gene | Naïve | OVA | OVA + BerriQi |
|-------------|-------|-----|---------------|
| **Arg1** | 1 (0.5) | 9.4 (5.3) | 26.0 (7.9) * |
| **Ym-1** | 1 (0.4) | 4.4 (1.9) | 5.5 (1.0) |
| **Fizz1** | 1 (0.5) | 15.4 (0.9) ** | 19.6 (3.3) ** |
| **Nos2** | 1 (0.3) | 1.5 (0.8) | 1.7 (0.5) |

**Figure 1: BerriQi® Boysenberry and apple juice concentrate suppresses ovalbumin-induced airway inflammation, and immune cell infiltration.** Mice were primed with ovalbumin (OVA)/Alum i.p and then challenged 7 days later with OVA i.n (Day 0). Mice were orally gavaged with 2.5 mg/kg total anthocyanins (TAC) in the BerriQi® Boysenberry and apple juice concentrate (BerriQi) 1h before OVA challenge and again 2 days post-challenge. (A) Haematoxylin and eosin stained lung tissue from naïve, OVA-challenged and OVA-challenged mice treated with BerriQi® Boysenberry and apple juice concentrate. Magnification 10x (top) and 20x (bottom) Asterisk =cell infiltration. (B) Alcian-blue Periodic acid-Schiff stained lung tissue from naïve, OVA-challenged and OVA-challenged mice treated with BerriQi® Boysenberry and apple juice concentrate. Magnification 10x (top) and 20x (bottom). Arrow=mucous producing goblet cells. (C) Total eosinophil, (D) Total neutrophil, (E) CD4+ T cells and CD8+ T cells in bronchioalveolar lavage fluid (BALF) were determined 4 days post-OVA challenge. Data presented as mean ± SEM $P<0.001$ compared with naïve and OVA challenge + BerriQi® Boysenberry and apple juice concentrate (one-way ANOVA with Tukey's Post Hoc test) for two experimental replicates with n=10 per treatment groups.

**Figure 2: BerriQi® Boysenberry and apple juice concentrate does not alter classical Th-1/Th-2 cells and cytokines.** Mice were primed with ovalbumin (OVA)/Alum i.p and then challenged 7 days later with OVA i.n (Day 0). Mice were orally gavaged with 2.5 mg/kg total anthocyanins (TAC) in the BerriQi® Boysenberry and apple juice concentrate (BerriQi) 1h before OVA challenge and again 2 days post-challenge. Mediastinal lymph node (MSLN) (A) CD4+ and (B) CD8+ T cells number; and lung tissue production of (C) IL-5, (D) IL-9, (E) IL-10 and (F) IFNγ were

determined 4 days post-OVA challenge by Legendplex. Data presented as mean ±

SEM for two experimental replicates with n=10 per treatment groups.

**Figure 3: BerriQi® Boysenberry and apple juice concentrate suppresses**

**ovalbumin-induced airway inflammation through increased IL-17A, CXCL10**

**and CCL4 concentration.** Mice were primed with ovalbumin (OVA)/Alum i.p and

then challenged 7 days later with OVA i.n (Day 0). Mice were orally gavaged with 2.5

mg/kg total anthocyanins (TAC) in the BerriQi® Boysenberry and apple juice

concentrate (BerriQi) 1h before OVA challenge and again 2 days post-challenge. (A)

Percentage of CD206+ macrophages in bronchioalveolar lavage fluid (BALF) and

lung tissue production of (B) CXCL10, (C) IL-17A, (D) CCL4 and (E) IL-17F was

determined 4 days post-OVA challenge by Legendplex. Data presented as mean ±

SEM, $P<0.05$ compared with OVA challenge + BerriQi® Boysenberry and apple juice

concentrate, $P<0.01$ compared with naïve and OVA challenge (one-way ANOVA with

Tukey's Post Hoc test) for two experimental replicates with n=10 per treatment

groups.

**SUPPLEMENTARY DATA**

**Supplementary Figure 1.** Liquid Chromatography-Mass Spectrometry (LC-MS)

chromatogram for BerriQi® concentrate, showing the anthocyanin UV/VIS profile

measured at 520 nm. Peak numbers refer to compounds listed in Table 1.

**Supplementary Figure 2.** Liquid Chromatography-Mass Spectrometry (LC-MS) chromatograms for BerriQi® concentrate, showing PDA (photodiode array) phenolic profiles measured at 200–600 nm, 280 nm and 360 nm. Peak numbers refer to compounds listed in Table 1. X, denotes sorbic acid, a preservative added to the BerriQi concentrate during formulation. This was present at a concentration of ~2000 µg/mL.