Genome-wide targets identification by CRISPRi-Omics for high-titer production of free fatty acids in *Escherichia coli*

Lixia Fang[1,2], Jie Fan[1,2], Congya Wang[1,2], Yingxiu Cao[1,2] * & Hao Song[1,2] *

[1] Frontier Science Center for Synthetic Biology and Key Laboratory of Systems

Bioengineering (Ministry of Education), Tianjin University, Tianjin 300072, China

[2] Collaborative Innovation Center of Chemical Science and Engineering (Tianjin),

School of Chemical Engineering and Technology, Tianjin University, Tianjin 300072,

China

* Correspondence and requests for materials should be addressed to Y.C.

(caoyingxiu@tju.edu.cn) or H.S. (hsong@tju.edu.cn).

Abstract

To construct a superior microbial cell factory for chemical synthesis, a major

challenge is to fully exploit cell potential via identifying and engineering beneficial

gene targets in the sophisticated metabolic networks. Here, we develop an approach

that integrates CRISPR interference (CRISPRi) to readily modulate genes expression

and omics analyses to identify potential targets in multiple cellular processes,

enabling systematical discovery of beneficial chromosomal gene targets that can be

engineered to optimize free fatty acids (FFAs) production in *Escherichia coli*. We

identify 56 beneficial genes via synergistic CRISPRi-Omics strategy, including 46

novel targets functioning in cell structure and division, and signaling transduction that

efficiently facilitate FFAs production. Upon repressing *ihfA* and overexpressing *aidB*

and *tesA'* in *E. coli*, the recombinant strain LihfA-OaidB results in a FFAs titer of

21.6 g L^{-1} in fed-batch fermentation, which, to our best knowledge, is the maximum

FFAs titer by the recombinant *E. coli* reported to date.

Introduction

Microbial biosynthesis of a desired product is a sophisticated process, usually involving the coordination of multiple metabolic intermediates or reactions subject to complicated interactions[1,2]. On the one hand, each of these intermediates flows to other metabolic processes for biosynthesis of cellular components or metabolites, acting as one of the multi-branched nodes in the topology of metabolic landscapes[3]. On the other hand, each reaction in the biosynthetic pathway deeply rely on many cellular processes, such as genes transcription and translation to enable enzymes function[4-6], or cell proliferation to ensure biomass accumulation[7]. As such, manipulation of other metabolic pathways or cellular processes might lead to an unexpectedly improvement on the biosynthesis of a desired product due to distant effects of genetic modulations or unknown regulatory interactions[8,9]. Thus, to construct a superior microbial cell factory for an enhanced product biosynthesis, it is highly desirable to genome-widely identify beneficial gene targets in the highly complicated intracellular interaction networks that can be reengineered for this purpose. As such, the interwoven linkage mechanism between product biosynthesis and other sophisticated genetic and metabolic networks can be better understood.

Free fatty acids (FFAs) are feedstocks for the manufacture of detergents, lubricants, cosmetics, and pharmaceutical ingredients[10,11]. A number of metabolic engineering approaches were developed to facilitate FFAs production in *E. coli*. Overexpression of key enzymes in the FFAs biosynthetic pathway such as acetyl-coenzyme A (CoA)

carboxylase[12-14] and fatty acyl-ACP thioesterase[12,13,15-17], or deletion of key genes in

the beta-oxidation pathway such as *fadD*[12,13,15-17] or *fadE*[15] gene, were studied. In

particular, optimally combinatorial expression of 15 crucial genes in the FFAs

biosynthetic pathway with *fadD* deletion in *E. coli* BL21 resulted in a titer of 8.6 g L^{-1}

of FFAs[18]. However, the cell potential for FFAs overproduction is still not fully

unleashed via modulating the FFAs directly related metabolic pathways alone[19].

Besides, the cellular rewiring mechanism and the linkage between FFAs biosynthesis

and other cellular processes is still largely unknown, and it remains difficult in

systematically identifying beneficial gene targets using laborious and time-consuming

genome engineering methods, such as homologous recombination-based gene

knockout[20]. These hurdles severely restrict the speed and scope of engineering of

microbial hosts to obtain a superior overproducer.

Here, we employ a systems metabolic engineering strategy for rapid, systematical

and effective identification of chromosomal gene targets that can be engineered for

the enhanced production of FFAs in *E. coli*. This strategy makes use of both the

CRISPR interference (CRISPRi) that readily down-regulates genes expression in a

dose-dependent manner to rapidly identify beneficial knockdown targets, and the

omics analyses of different strains that allows acquisition of comprehensive data

related to mechanisms of cellular metabolism to dig potential targets in the genetic

and metabolic networks. By way of this synergistic CRISPRi–Omics strategy, we

identified 46 new beneficial gene targets that efficiently facilitate FFAs production,

which involves in cell division, signaling transduction and transport, cell structure,

and other seemingly irrelevant cellular processes with FFAs biosynthesis. Upon

repressing *ihfA* and overexpressing *aidB* and *tesA'*, the resulting recombinant *E. coli*

strain LihfA-OaidB enabled a FFAs titer of 21.6 g L^{-1} (with productivity of 0.636 g

$L^{-1} h^{-1}$) in fed-batch fermentation, which, to the best of our knowledge, is the

maximally reported FFAs titer and productivity by the recombinant *E. coli* to date.

Our findings not only allow for enhanced biosynthesis of the desired product, but also

lead to a mechanistic understanding of the underlying linkage between product

biosynthesis and multiple cellular processes.

Results

Construction of a non-toxic and functional CRISPRi system in *E. coli*

BL21(DE3). CRISPRi system functioned through the expression of dCas9 and

customized specific sgRNA[21] (**Supplementary Fig. 1a**). However, high-level dCas9

could cause a severe inhibition of cell growth and abnormal expression of genes in a

strain-specific manner[22,23]. To determine a superior dCas9 expression level in

BL21(DE3), we constructed plasmids with dCas9 being expressed under the control

of the promoter P_{T7}, P_{T5}, P_{Trc} and P_{BAD}, respectively (**Supplementary Fig. 2a**). The

expression strengths of these promoters were $P_{T7} > P_{T5} > P_{Trc} > P_{BAD}$, according to our

previous study[24]. With green fluorescent protein (GFP) as a characterization signal

driven by the P_{T7} promoter, fluorescence-based reporter plasmids were further

constructed (**Supplementary Fig. 2a**) and transformed into BL21(DE3), forming five

strains G0 (P_{T7}-GFP), G1 (P_{T7}-GFP-P_{T7}-dCas9), G2 (P_{T7}-GFP-P_{T5}-dCas9), G3

(P_{T7}-GFP-P_{Trc}-dCas9), and G4 (P_{T7}-GFP-P_{BAD}-dCas9), respectively. We found that

the OD_{600} and fluorescence level of the strains G1 and G2 were greatly decreased

from the strain G0, while that of G3 and G4 were little affected compared with G0

(**Supplementary Fig. 1b**). This suggested that the expression of dCas9 at relatively

low level, such as under the control of the P_{Trc} or P_{BAD} promoter, did not cause overt

toxicity in our system.

Then, we examined the transcriptional repression capability of CRISPRi on gene

expression by co-expressing a specific sgRNA. sgRNA-*gfp* was designed to target on

the non-template strand of GFP at a position ~38 nucleotides downstream of the start

codon   to   expect   a   high   repression   efficiency[25].   The   plasmid   Sg-Hgfp

(**Supplementary Fig. 2b**), harboring sgRNA-*gfp* under the control of the P_R promoter,

was constructed and transformed into the strains G0-G4, forming the strains g0-g4,

respectively. The results showed that fluorescence levels of the strains co-expressing

dCas9 and sgRNA-*gfp* (g1-g4) were all dramatically reduced in comparison to the

corresponding strain expressing only dCas9 (G1-G4) (**Supplementary Fig. 1b**). It

indicated   that   the   CRISPRi   system   was   functional   for   effective   transcriptional

repression. In addition, the strain with dCas9 regulated by the P_{Trc} promoter obtained

the highest repression efficiency (93%) (**Supplementary Fig. 1b**). Thus, we selected

dCas9 controlled by the P_{Trc} promoter to construct the CRISPRi system in

BL21(DE3).

Identification of beneficial gene targets in the pathways related to FFAs

metabolism using CRISPRi system. Glycerol, serving as the substrate, is

catabolized into key intermediates glyceraldehyde-3P, pyruvate, acetyl-CoA,

malonyl-ACP in sequence and then converted into acyl-ACP through the fatty acid

biosynthetic pathway (**Fig. 1a**). However, these key intermediates also involved in

other metabolic pathways, resulting in diverted carbon fluxes away from FFAs. The

expression of truncated fatty acyl-ACP thioesterase TesA' (with leader sequence

deleted) could release free fatty acid intermediates from ACP in the cytosol[26,27].

Nevertheless, the synthetic FFAs are degraded naturally by the beta-oxidation process

(**Fig. 1a**). To channel carbon flux towards FFAs and impede products' degradation,

we chose 108 genes in the chromosome that related with the FFAs metabolism and

regulation as candidate targets for modulation by CRISPRi, and classified them into

five modules (upstream carbon flux diversion, downstream carbon flux diversion,

amino acid metabolism, beta-oxidation, and transcription factors) (**Fig. 1a**).

To facilitate large-scale target identification, we constructed a library of 108

synthetic sgRNAs that repress the expression of the 108 genes in the above five

modules with high silencing efficiency[21]. We introduced plasmids harboring each of

the 108 synthetic sgRNAs individually into the starting strain CF co-expressing TesA'

and dCas9 in BL21(ED3) to implement gene perturbations (**Fig. 1b**). The strain CF

transformed with plasmid Sg-0 that expressed sgRNA0 without target site

complementary sequence was named as the strain Control, which produced FFAs

with a tier of 631 mg L^{-1}. Any CRISPRi-engineered strain that could increase FFAs

titer by more than 20% compared with the strain Control was sorted out. As a result,

30 beneficial targets were identified from the library and the highest production of

FFAs reached 1,232 mg L^{-1}, 95% higher than that of the strain Control (**Fig. 2**).

The identified beneficial genes were almost evenly distributed in the modules of

upstream carbon flux diversion (**Fig. 2a**), downstream carbon flux diversion (**Fig. 2b**),

beta-oxidation (**Fig. 2d**), and transcription factors (**Fig. 2e**). It suggested that reducing

the carbon flux that flows to the by-products phosphatidylglycerol (*gpsA*, *plsY*, *pgpB*

and *pgpC*), D-Lactate (*dld*), acetate (*pta*, *yccX* and *eutE*) and biotin (*bioC*, *bioH* and

bioD), or impeding fatty acids degradation (*fadE*, *fadB*, *atoB* and *yqeF*) contributed to

the enhanced production of FFAs. Reducing the expression of a few transcription

factors (including *fabR*, *fadR*, *asnC*, *fnr*, *cra*, *gadW*, *glcC*, *glpR*, *ihfA*, *ihfB* and *torR*)

also enhanced the production of FFAs (**Fig. 2e**), which might benefit from their global

regulation of genes expression. However, reducing amino acids biosynthetic pathways

was unfavorable for FFAs production, as the engineered strains obtained decreased

titer of FFAs or failed to be constructed (**Fig. 2c**). The possible reason was that these

amino acids were necessary for cell viability, but the reduced amount of amino acids

upon CRISPRi perturbation was insufficient to maintain normal cell physiology. Thus,

in addition to the key enzymes in the FFAs biosynthetic pathway, there exist many

chromosomal genes seemingly unrelated to FFAs that could be beneficial to the

enhanced FFAs production.

Then, we tuned repression efficiency towards each of the 30 beneficial genes to

optimize their individual expression level. sgRNAs were designed to bind the

non-template DNA strand at the middle or terminal region to achieve medium or low

repression efficiency towards each target[21] (**Fig. 3a**). The plasmids harboring each of

the 60 synthetic sgRNAs were thus individually introduced into the starting strain CF,

and the production titer of FFAs was determined, showing that 10 strains (LgpsA,

LpfkA, MyccX, LgabD, MgadA, MatoB, LatoB, LglpR, MihfA and LihfA) enabled

further improvement in FFAs production (**Fig. 3b**).

Among the 30 beneficial gene targets, 20 of them (*gpsA*, *plsY*, *pgpB*, *pgpC*, *dld*,

yccX, *bioC*, *bioH*, *bioD*, *gabD*, *gadA*, *asnC*, *fnr*, *cra*, *gadW*, *glcC*, *glpR*, *ihfA*, *ihfB*,

torR) were firstly engineered for FFAs production. We found three favorable

metabolic pathways that have not been studied or modulated for FFAs production

before, namely the biotin, phosphatidylglycerol, and glutamate biosynthetic pathways.

Even in pathways that have been modulated in previous studies, we also identified

novel beneficial targets, such as *dld* in the lactate biosynthetic pathway, *yccX* and

eutE in the acetate formation pathway. In particular, we found a few genes such as *pta*

in the acetate formation pathway that have been deleted in previous studies[28], which

however showed no promotional effect on FFAs production, while CRISPRi

perturbation of the *pta* expression in this study significantly enhanced FFAs

production. Collectively, these results demonstrate that CRISPRi is of use for

systematical identification of beneficial targets that can be engineered for improved

biosynthesis of a desired product.

Identification of potential targets in the cellular network via proteome and

transcriptome analyses. To explore more potential targets that could facilitate FFAs

production, we further focused on transcription factors in consideration of their

complicated regulation functions on metabolic networks. Among all the targeted

transcription factors, repression of *ihfA* with low efficiency (named as the strain

LihfA) achieved a highest FFAs production, followed by the strain HfadR that

repressed *fadR* with high efficiency (**Figs. 2e** and **3b**). IhfA is the alpha subunit of the

integration host factor (IHF) in *E. coli*, which is capable of affecting transcription on a

genome-wide scale[29]. We found that down-regulation of *ihfA* could enhance FFAs

production, and the FFAs titer was negatively correlated with the repression

efficiency (**Fig. 4a**). Unlike IhfA, FadR controls the expression of several genes

directly involved in fatty acid biosynthesis, degradation, and transport across cell

membranes[30]. It is intriguing that up-regulation of FadR (named as the strain OfadR)

could also increase the FFAs titer as demonstrated in a previous study[31] and this study

(**Fig. 4a**).

Even though the strains with FFAs overproducing phenotype have different

genotypes, these strains might have convergent regulatory responses that facilitate

FFAs biosynthesis. Thus, the three above engineered strains (LihfA, HfadR and

OfadR, with knockdown of *ihfA*, *fadR* and overexpression of *fadR*, respectively) with

high-titer FFAs and the strain Control were applied for proteome and transcriptome

analyses to investigate the underlying mechanism of cellular responses to genetic

modulations. Cells were sampled at the early stationary phase (24 h upon IPTG

induction) (**Supplementary Fig. 3**). Hundreds of genes and proteins were

differentially expressed in each pair of LihfA vs. Control, HfadR vs. Control, and

OfadR vs. Control (**Supplementary Figs. 4** and **5**), suggesting the global modulation

of genes in the engineered strains. To identify convergent rewiring in these high-titer

strains, we plotted the differentially expressed genes (DEGs) of the three pairs in a

Venn diagram (**Fig. 4b**). As a result, 24 down-regulated targets and 17 up-regulated

targets were selected based on their significantly differential expression at the protein

level (a and c, respectively) or at the transcript level (b and d, respectively) in at least

two pairs (**Fig. 4b**) (see details in Supplementary Materials). The differential

expression levels of these 41 genes are summarized in **Fig. 4c**. These potentially

favorable gene targets are newly identified and seemingly irrelevant to FFAs

biosynthesis in a direct fashion.

To test the function of these 41 candidate targets on FFAs production, the

corresponding reverse engineering of each gene was conducted. We applied CRISPRi

to repress the expression of the 24 'down-regulated' genes or utilized the P_{BAD}

promoter to overexpress the 17 'up-regulated' genes (**Fig. 5a**). The plasmids carrying

the cassettes towards each candidate gene were transformed into the starting strain CF.

A FFAs assay of the resulting strains showed that modulation of the 19 genes among

the identified targets enhanced FFAs production by more than 20% in comparison to

the strain Control (**Fig. 5b**). Considering that the selected genes were all differentially

expressed in the strain pair of LihfA vs. Control (**Fig. 4c**), these targets were also

manipulated in the strain LihfA to test their effects on FFAs production (**Fig. 5c**). As

a result, 13 engineered strains gained further increase in FFAs production compared

with the strain LihfA (**Fig. 5d**). In particular, the highest FFAs production was

obtained in the strain LihfA-OaidB, in which the *ihfA* gene was repressed with low

silencing efficiency and the *aidB* gene was overexpressed under the P_{BAD} promoter.

The titer is up to 2,052 mg L^{-1}, which is 57% and 225% higher than that of the strains

LihfA and Control, respectively (**Fig. 5d**). In summary, 26 (*norR*, *nrdE*, *npr*, *yihU*,

glpG, *ftsQ*, *creC*, *sdhB*, *pal*, *rplW*, *tyrU*, *crsC*, *rnpB*, *ssrS*, *csrB*, *ryfA*, *sdsR*, *folk*, *aidB*,

yphF, *deoD*, *nnr*, *rihC*, *hofP*, ECD_01466, and *ulaE*) out of the 41 newly identified

genes are determined as the beneficial targets, which are validated to enhance the

FFAs production by reverse engineering.

Exploration of cellular processes facilitating FFAs production. The newly

identified beneficial targets are not in the metabolic pathways that directly related to

FFAs biosynthesis. To mine out the related regulatory mechanisms for FFAs

production, the functions of the 26 beneficial genes were analyzed according to

bioinformatics analysis of the database (NCBI, KEGG, UniProt and Gene Ontology)

and summarized into eight sets (**Fig. 6, Supplementary Table 1**). Based on the FFAs

titer of the corresponding engineered strains, the weight value (W value) of each set

was calculated to characterize their effects on the enhancement of FFAs production.

The eight sets were arranged in descending order of W value as follows: cell division,

transduction and transport, energy metabolism, cellular structure, non-coding RNAs

(ncRNAs), protein metabolism, nucleic acid metabolism, and cofactor metabolism

(**Fig. 6, Supplementary Table 1**). Surprisingly, the processes such as cell division,

signal transduction, protein metabolism and nucleic acid metabolism seemingly not

associated with FFAs biosynthesis in any previous studies that showed favorable

effect on FFAs production. These results indicate that the manipulation of cellular

processes that are seemingly irrelevant with FFAs biosynthesis can also remarkably

boost FFAs production.

Then, we further dissect the underlying connections between these enriched cellular

processes and FFAs production. Down-regulation of *ftsQ* or *pal* significantly

increased the FFAs concentration, inactivation of which resulted in deficiency in cell

division and formed long multi-septated cell chains[32,33], which might associated with

the increased cell volumes for storing FFAs. Down-regulation of *creC*, a sensor for

the early response to phosphate starvation[34], was favorable for enhanced FFAs

production, which is consistent with the promotive effects of phosphate limitation

cultivations[35]. The functioning of *npr* down-regulation and *yphF* up-regulation may

be associated with the reduction of nitrogen assimilation[36] and the elevation of

carbohydrates' uptake[37], indicating a regulatory link between nitrogen and carbon

utilization for the efficient FFAs production. Down-regulation of ATP-consuming

proteins (such as those encoded by the genes *norR*, *nrdE*, *creC* and *rplW*) indirectly

increased ATP availability, a potential bottleneck for FFAs biosynthesis[19]. Beneficial

effects from down-regulation of membrane proteins (such as those encoded by the

genes *glpG*, *ftsQ*, *creC* and *pal*) might associated with microbial tolerance to FFAs[38].

The functioning of down-regulation of *rplW* (encoding 50S ribosomal subunit protein

L23), *tyrU* (encoding tRNA-Tyr) and *rnpB* (encoding M1 RNA, catalytic component

of tRNA processing enzyme RNase P) was associated with the inhibition of protein

synthesis, which increased partitioning of total carbon to lipids[8]. Up-regulation of *nnr*,

facilitating the repair of NAD(P)H hydrates to NAD(P)H[39], could be an important

complement strategy in enhancing NADPH availability, which plays an important

role in FFAs production[19,40]. In all, these findings provide new insights into linkages

between cell functions and product biosynthesis, guiding avenues for further strain

development.

As a result, the engineered *E. coli* strain LihfA-OaidB with simultaneous repression

of *ihfA* and overexpression of *aidB* and *tesA'* achieved the highest titer of FFAs

production. According to the results of combined omics analyses and reverse

engineering, we concluded that down-regulation of *ihfA* induced comprehensive

adjustments of cellular processes that promoted the FFAs production. Nevertheless,

repression of *ihfA* might be unfavorable for cellular survival at acidic pH as IHF is

required for the induction of acid resistance[41,42]. AidB, one of the DNA damage

repairing protein in *E. coli*, is involved in adaptive response of cells, serving to protect

cells from cytotoxic effects of acidified cytoplasm or alkylating agents[43-45]. Thus, we

speculated that up-regulation of *aidB* in the strain LihfA reconciled the stress from the

acidic environment posed by the produced FFAs, facilitating FFAs biosynthesis in the

strain LihfA-OaidB.

Further enhancing FFAs production by fed-batch fermentation. To further

enhance FFAs production of the recombinant strain LihfA-OaidB in a scaled-up

process, fed-batch fermentations were performed in a 5 L fermenter, using the

dO$_2$-stat feeding strategy (details described in Methods). With arabinose and IPTG

induction, *aidB* (DNA alkylation damage repair protein) was overexpressed by the

P$_{BAD}$ promoter, *tesA'* (truncated acyl-CoA thioesterase) was overexpressed by the P$_{T7}$

promoter and *ihfA* (integration host factor subunit alpha) was repressed by CRISPRi

with low silencing efficiency in the strain LihfA-OaidB. The FFAs production and

cell density of the recombinant strain LihfA-OaidB increased along with the glycerol

consumption (**Fig. 7a**). When cells were cultivated for 34 h after IPTG induction,

LihfA-OaidB produced a titer of 21.6 g L^{-1} FFAs, accompanied by 0.636 g L^{-1} h^{-1}

productivity and 0.146 g FFAs per gram glycerol yield, the highest production titer

and productivity in *E. coli* reported to date[18,46]. Extracellularly secreted FFAs were

clearly visible at the top of the culture medium after centrifugation (**Fig. 7b**).

Moreover, floating dead cells or fatty acid particles[47] were precipitated and stuck on

the fermenter inner wall and sensors over the course of the fermentation process

(**Supplementary Fig. 6**), exhibiting an additional production of FFAs that is however

hard to accurately quantify and not counted in the FFAs titer. Thus, the overall actual

FFAs production was higher than that showed in **Fig. 7a**.

312 Discussion

Genome-scale identification of beneficial gene targets in the chromosome for

improving production of a desired product is challenging due to the complicated

intracellular interactions. We here developed a systems metabolic engineering

approach, which took the advantage of CRISPRi to readily and rapidly down-regulate

gene expression and the omics analyses to provide potential targets in the cellular

interaction network, thus enabling rapid, systematical and effective identification of

beneficial targets that can be engineered to optimize FFAs production in *E. coli*. This

synergistic strategy integrated the 'forward' and 'reverse' metabolic engineering

cycles and their interaction[48] (**Supplementary Fig. 7**). In forward metabolic

engineering, CRISPRi-mediated gene down-regulation functions were pursued in this

work for the identification of beneficial chromosomal targets in the competitive or

regulatory pathways of FFAs biosynthesis that can be repressed in high, intermediate

and low levels, unlike gene knockout, to enhance FFAs production. In reverse

metabolic engineering, proteome and transcriptome analyses of the strains with

differential FFAs production provide comprehensive data related to mechanisms of

cellular metabolism, which can be validated experimentally via CRISPRi system and

plasmid mediated expression system. In doing so, we identified beneficial targets that

even seemingly irrelevant with FFAs biosynthesis, which are generally inaccessible to

hypothesis-driven or random gene targeting, aiding in expanding hosts' potential of

enhanced products synthesis. To our knowledge, this is the first report that

systematically identifies beneficial target genes in the cellular interaction network for

enhanced FFAs biosynthesis.

Many new approaches, including in vitro reconstitution of enzymes in the FFAs

biosynthetic pathway[49,50], engineering the reversal of the beta-oxidation pathway[51],

and construction of malonyl-CoA sensors[52,53], have been studied to enhance FFAs

production in *E. coli*. However, these approaches essentially focused on the

optimization of FFAs metabolic pathways alone, not exploiting the full potential of

interwoven intracellular processes. In our study, gene perturbations on the cellular

interaction network, including the competitive pathways of FFAs biosynthesis,

transcription factors and cellular processes that seemingly irrelevant with FFAs

metabolism, were systematically investigated to unleash the cellular potential and

change the cells towards a state of FFAs overproduction. As a result, the best

engineered strain LihfA-OaidB produced FFAs with a titer of 21.6 g L^{-1} (0.636 g L^{-1}

h^{-1} productivity) under fed-batch conditions, which, to the best of our knowledge, is

the maximally reported FFAs titer and productivity by recombinant *E. coli* to date.

Furthermore, the potential of the identified beneficial targets and our constructed

strains for improving FFAs production could be further integrated and amplified by

combinatorial genetic perturbations[54] and exploiting nongenetic variation[46].

Beneficial targets that were identified by our synergistic CRISPRi-Omics strategy

participate in diverse cellular processes, which support the idea that product synthesis

is a complex process and that global optimization of cellular machinery is required for

optimized product synthesis. Generally, these targets primarily associate with

metabolic processes, other cellular processes or regulatory elements. As to metabolic

processes, we identified novel beneficial targets in the common targeted pathways,

such as *yccX* and *eutE* in acetate formation pathway, ATP-consuming protein *nrdE*

and NAD(P)H-hydrate repair enzyme *nnr*, as well as novel pathways that have not

been engineered for FFAs production such as the biotin biosynthetic pathway (*bioC*,

bioH and *bioD*). Our target screening approach also emphasizes the importance of

other cellular processes or functions in the strain improvement for optimized FFAs

production, such as cell division (*ftsQ* and *pal*) related morphology, signaling

transduction (*creC*) and transport (*npr* and *yphF*) related resources utilization, cellular

structure (*glpG*, *ftsQ* and *pal*) related cellular tolerance, protein metabolism (*rplW*,

tyrU, *rnpB*) related resources partition, and nucleic acid metabolism (*aidB*) related

adaptive response. In addition, we also identified beneficial regulatory elements that

have not been engineered for FFAs production, such as the integration host factor *ihfA*,

or novel ncRNAs *ryfA* and *sdsR*, which modulate FFAs production through regulation

of multiple cellular processes. Thus, our findings provide insights into connections

between FFAs biosynthesis and other cellular processes and expand the underlying

mechanistic understanding of the regulation of FFAs production in *E. coli*.

Strains obtained through the identification and modulation of genes have proven to

be a versatile and robust approach for improved FFAs production, which also offers

avenues for engineering more potential targets for further improvement. Knowledge

acquired in this work toward understanding the FFAs biosynthetic machinery can be

extended to other microorganisms for FFAs production, such as yeast and oleaginous

bacteria. The approach for identifying promising modulation candidates is also

applicable to other products without obvious screening phenotypes.

In summary, the synergistic CRISPRi-Omics strategy enabling rapid, systematical

and effective identification of beneficial targets in the intracellular interaction

network provides novel insights and solutions for further increasing productivity of

microbial cell factories and better understanding of the link between product

biosynthesis and the other processes of cellular machinery.

Methods

Experimental materials. Strains and plasmids used in this study are listed in

Supplementary Table 2 and **Supplementary Table 3**, respectively. *E. coli*

*Trans*1-T1 strain was used for cloning. *E. coli* BL21(DE3) and its derived strains were

used for fermentation. Plasmids pACYCDuet-1, YX210, YX212 and YX213 were

used as the vector for expression of dCas9, GFP or TesA'. Plasmid Sg-S (with two

BsaI restriction sites) was constructed and used as a platform vector for expression of

various sgRNAs. Plasmid Sg-BAD was used as the vector for expression of

chromosome genes. Plasmids, except that for sgRNAs expression, were constructed

by standard enzyme digestion and ligation. Target site complementary sequences of

sgRNAs were designed according to previous studies[55]. Custom-designed spacers

were inserted into plasmid Sg-S by one-step Golden Gate assembly[56,57], allowing

rapid construction of plasmids for expression sgRNAs targeting any genomic locus of

interest. Primers annealed to custom-designed spacers and primers used to amplify

genes from *E. coli* genomic DNA were listed in **Supplementary Tables 4-7** and

Supplementary Table 8, respectively. Codon-optimized genes were listed in

Supplementary Table 9.

Culturing conditions and mediums. To assess FFAs production by different

recombinant *E. coli* strains, fermentation was performed in glass tubes. Each strain

was inoculated from a freshly transformed single colony in LB agar plate to 2 mL LB

medium as seed culture. When cell accumulation reached to stationary phase, 1%

(V/V) of seed culture was re-inoculated to 5 mL modified M9 medium in glass tube.

The tube cultures then were induced with 1 mM IPTG when OD_{600} reached ~1.0 and

allowed to grow for an additional 40 h at 30°C and 220 rpm. When the P_{BAD} promoter

was applied, 10 mM arabinose was utilized to induce gene expression at OD_{600} 0.5–

0.6. All tube experiments were performed at least in triplicate. Flask fermentation of

50 mL culture broth was implemented for proteome and transcriptome analyses with

the identical culturing procedures of tube fermentation. M9 medium used for tube and

flask fermentation was as described by Schirmer et al. [58] with some modifications:

17.1 g L^{-1} $Na_2HPO_4 \cdot 12H_2O$, 3 g L^{-1} KH_2PO_4, 0.5 g L^{-1} NaCl, 2 g L^{-1} NH_4Cl, 2 g L^{-1}

yeast extract, 3% glycerol, 0.25 g L^{-1} $MgSO_4 \cdot 7H_2O$, 11.1 mg L^{-1} $CaCl_2$, 10 mg L^{-1}

thiamine and 0.1% (v/v) Triton-X100. 1 mL L^{-1} metal trace stock solution was also

supplemented, which contained 27 g L^{-1} $FeCl_3 \cdot 6H_2O$, 2 g L^{-1} $ZnCl_2$, 2 g L^{-1}

$Na_2MoO_4 \cdot 2H_2O$, 1.9 g L^{-1} $CuSO_4 \cdot 5H_2O$, 0.5 g L^{-1} H_3BO_3. pH was adjusted to about

7.2 by Tris. If necessary, 100 μg mL^{-1} carbenicillin or 34 μg mL^{-1} chloramphenicol

was supplemented.

For fed-batch fermentation, an overnight LB culture (2 mL) of the freshly

transformed single colony was sub-cultured into 200 mL modified mineral medium (6

g L^{-1} NH_4Cl, 8.5 g L^{-1} KH_2PO_4, 0.5 g L^{-1} citrate, 5 g L^{-1} yeast extract, 15 g L^{-1}

glycerol, 1 g L^{-1} $MgSO_4 \cdot 7H_2O$, 0.07 g L^{-1} $CaCl_2 \cdot 2H_2O$, 4 mL L^{-1} metal trace stock

solution, 100 mg L^{-1} thiamine)[59] with 100 μg mL^{-1} carbenicillin and 34 μg mL^{-1}

chloramphenicol. When OD_{600} reached to 3-4, the cultured cells were re-inoculated

into a 5 L bioreactor (T&J, China) with 1.8 L modified mineral medium as above.

Fermentation temperature was set at 30°C and pH was controlled at 7 by feeding 6 N

ammonium hydroxide via an auto pump. Air flow rate was maintained at around 2 L

min^{-1}. The dissolved oxygen (dO_2) concentration was controlled above 30% by

agitation cascade (300–800 r.p.m). When cell density reached about 6 and 15, 10 mM

arabinose and 1 mM IPTG was added into the fermentation cell culture, respectively.

Feeding medium (2.47 g L^{-1} $MgSO_4 \cdot 7H_2O$, 500 g L^{-1} glycerol and 100 g L^{-1} yeast

extract) was fed to the fermentation culture after the initial 15 g L^{-1} glycerol was

almost depleted and the dissolved oxygen showed a sharp increase. The carbon source

restriction strategy was carried out by dO_2 concentration, given that the dO_2 level

immediately spiked once the carbon source was run out. Once the $dO_2 > 60\%$, the

feeding would automatically start at 5 mL min^{-1} until $dO_2 < 60\%$. Antifoam (Sigma)

was added automatically as needed. Broth samples (~5 ml) were collected at a series

of time points to measure cell density and stored at −20°C for further measurements

of residual glycerol and FFAs production.

GFP Fluorescence assay. Fluorescence signal of GFP was used to characterize

repression efficiency of the CRISPRi system. Culturing procedures and induction

process of the GFP reporting strains were the same as tube fermentation of the FFAs

producing strains. For each tube, 200 μL of sample was diluted into the linear range

of the detector with phosphate buffered saline (PBS) after IPTG induction for 20 h.

Fluorescence intensity (excitation at 485 nm and emission at 520 nm) and cell density

(OD_{600}) were detected using 96-well polystyrene plates (black plate with a clear

bottom) (Corning Incorporated 3603, USA) and a microplate reader (SpectraMax M2,

Molecular Devices, USA). The relative fluorescence intensity was first normalized

using OD_{600} and then subtracted that of blank *E. coli* BL21 (DE3). Experiments were

performed in triplicate.

Metabolite extraction and analysis. FFAs titers in whole-cell culture (only FFAs

was measured in this study) were quantified following previously published

methods[60]. Specifically, 0.5 ml of cell culture (or an appropriate volume of cell

culture diluted to 0.5 ml) was acidified with 50 μL of concentrated HCl, spiked with

60 μg of heptadecanoic acid as internal standard. The cell culture was extracted twice

with 0.5 ml of ethyl acetate. The extracted FFAs were then determined using a

Thermo Scientific TRACE 1300 gas chromatograph (GC) equipped with a

TG-WaxMS A column (30 m × 0.32 mm × 0.25 μm; Thermo Scientific) and a Flame

Ionization Detector (FID) operating under constant flow rate of the carrier gas

(nitrogen) at 1 mL min^{-1}. The following temperature program was used: hold at 50°C

for 1 min, then heat to 245°C at 30°C min^{-1} and hold at this temperature for 22.5 min.

Individual fatty acid species were qualified by authentic homologous standards and

quantified by comparing the peak areas with that of the internal standard using the

Chromeleon 7.1 software. Total concentrations of free fatty acids were calculated as

the sum of C_{12} to C_{18} (saturated and monounsaturated).

Glycerol concentration was determined by high-performance liquid

chromatography (HPLC) following Waters standard protocols. Briefly, filtered

culture supernatants were analyzed by a Waters HPLC system including a Waters

e2695 separation module, a Waters 2414 refractive index detector (RID) and an

Aminex HPX-87H column (Bio-Rad). The separation was performed through elution

with 5 mM H_2SO_4 at a flow rate of 0.6 mL min^{-1} at 65°C for 30 min.

Proteome analysis. Cells were harvested by centrifugation at 5000 rpm for 5 min at

24 h after IPTG induction and flash frozen with liquid nitrogen. For protein

preparation, cells were resuspended in 600 μL lysis buffer (0.05 M Tris, pH 8, 1%

SDS, and 8 M Urea) and disrupted by ultrasonication for 5 min (cycles with 2s work

and 3s pause). After centrifugation at 13000 rpm and 4°C for 20 min, the supernatant

was subjected to protein precipitation by adding pre-cooled acetone with 10 mM DTT.

The total precipitate was collected, dried and then re-dissolved in 600 μL Urea buffer

(0.05 M Tris, pH 8, and 8 M Urea). The protein samples were quantitated by the

Bradford method and qualitated by SDS-PAGE. A total of 100 μg of each sample was

trypsin-digested overnight and labeled using 4-plex iTRAQ reagents (AB SCIEX) as

manual described. The tryptic peptides corresponding to each sample were labeled

with its unique iTRAQ tags and proteomics analysis was conducted using a Triple

TOF 6600 system (AB SCIEX). The data were analyzed by Novogene.

Transcriptome analysis. Cells were harvested after 24 h of IPTG induction by quick

centrifugation at 10000 g for 1 min and immediately frozen in liquid nitrogen. Total

RNA was extracted using the RNAprep pure Cell/Bacteria Kit (Tiangen) following

lysozyme treatment. RNA degradation and contamination were monitored on 1 %

agarose gels. RNA concentration was measured by Qubit$^®$ RNA Assay Kit in Qubit$^®$

2.0 Flurometer (Life Technologies, CA, USA) and RNA integrity was assessed on a

2100 Bioanalyzer (Agilent Technologies, CA, USA). rRNA is removed using a

Ribo-zero kit that leaves the mRNA. A total of 3 μg RNA was used as an input per

sample. Sequencing libraries were generated using a NEBNext$^®$UltraTM RNA

Library Prep Kit for Illumina$^®$ (NEB, USA) following the manufacturer's instructions,

and index codes were added to attribute sequences to each sample. Clustering of the

index-coded samples was performed on a cBot Cluster Generation System using the

TruSeq PE Cluster Kit v3-cBot-HS (Illumina) according to the manufacturer's

instructions. After cluster generation, the library preparations were sequenced on an

Illumina Hiseq 4000 platform and paired-end reads were generated. After these steps,

the data were analyzed by Novogene.

Data availability

A reporting summary for this article is available as a Supplementary Information file.

The data that support the findings of this study are available from the corresponding

author upon request. The source data underlying Figs. 2a-e, 3b, 4a-c, 5b, 5d, 6, and 7a,

Supplementary Figs. 1b, 3, 4, and 5 and Supplementary Table 1 are provided as a

Source Data file.

References

1. Keasling, J. D. Manufacturing molecules through metabolic engineering. *Science* **330**, 1355-1358 (2010).

2. Nielsen, J. & Keasling, J. D. Engineering Cellular Metabolism. *Cell* **164**, 1185-1197 (2016).

3. Nielsen, J. It is all about metabolic fluxes. *Journal of Bacteriology* **185**, 7031-7035 (2003).

4. Li, G. W., Burkhardt, D., Gross, C. & Weissman, J. S. Quantifying Absolute Protein Synthesis Rates Reveals Principles Underlying Allocation of Cellular Resources. *Cell* **157**, 624-635 (2014).

5. Cambray, G., Guimaraes, J. C. & Arkin, A. P. Evaluation of 244,000 synthetic sequences reveals design principles to optimize translation in Escherichia coli. *Nature Biotechnology* **36**, 1005-+ (2018).

6. Gill, R. T., Valdes, J. J. & Bentley, W. E. A Comparative Study of Global Stress Gene Regulation in Response to Overexpression of Recombinant Proteins in Escherichia coli. *Metabolic Engineering* **2**, 178-189 (2000).

7. Boer, V. M., Crutchfield, C. A., Bradley, P. H., Botstein, D. & Rabinowitz, J. D. Growth-limiting Intracellular Metabolites in Yeast Growing under Diverse Nutrient Limitations. *Molecular Biology of the Cell* **21**, 198-211 (2010).

8. Ajjawi, I. et al. Lipid production in Nannochloropsis gaditana is doubled by decreasing expression of a single transcriptional regulator. *Nat Biotechnol* **35**, 647-652 (2017).

9. Huang, M., Bao, J., Hallstrom, B. M., Petranovic, D. & Nielsen, J. Efficient protein production by yeast requires global tuning of metabolism. *Nat Commun* **8**, 1131 (2017).

10. Tee, T. W., Chowdhury, A., Maranas, C. D. & Shanks, J. V. Systems metabolic engineering design: fatty acid production as an emerging case study. *Biotechnol Bioeng* **111**, 849-857 (2014).

11. Janssen, H. J. & Steinbuchel, A. Fatty acid synthesis in Escherichia coli and its applications towards the production of fatty acid based biofuels. *Biotechnol Biofuels* **7**, 7 (2014).

12. Lu, X., Vora, H. & Khosla, C. Overproduction of free fatty acids in E. coli: implications for biodiesel production. *Metab Eng* **10**, 333-339 (2008).

13. Lennen, R. M., Braden, D. J., West, R. A., Dumesic, J. A. & Pfleger, B. F. A process for microbial hydrocarbon synthesis: Overproduction of fatty acids in Escherichia coli and catalytic conversion to alkanes. *Biotechnol Bioeng* **106**, 193-202 (2010).

14. Lee, S., Jeon, E., Yun, H. S. & Lee, J. Improvement of Fatty Acid Biosynthesis by Engineered Recombinant Escherichia coli. *Biotechnology and Bioprocess Engineering* **16**, 706-713 (2011).

15. Steen, E. J. et al. Microbial production of fatty-acid-derived fuels and
chemicals from plant biomass. *Nature* **463**, 559-562 (2010).

16. Choi, Y. J. & Lee, S. Y. Microbial production of short-chain alkanes. *Nature*
502, 571-574 (2013).

17. Zhang, X. J., Li, M., Agrawal, A. & San, K. Y. Efficient free fatty acid
production in Escherichia coli using plant acyl-ACP thioesterases. *Metabolic*
Engineering **13**, 713-722 (2011).

18. Xu, P. et al. Modular optimization of multi-gene pathways for fatty acids
production in E. coli. *Nat Commun* **4**, 1409 (2013).

19. He, L. et al. Central metabolic responses to the overproduction of fatty acids
in Escherichia coli based on 13C-metabolic flux analysis. *Biotechnol Bioeng*
111, 575-585 (2014).

20. Datsenko, K. A. & Wanner, B. L. One-step inactivation of chromosomal genes
in Escherichia coli K-12 using PCR products. *Proceedings of the National*
Academy of Sciences of the United States of America **97**, 6640-6645 (2000).

21. Qi, L. S. et al. Repurposing CRISPR as an RNA-guided platform for
sequence-specific control of gene expression. *Cell* **152**, 1173-1183 (2013).

22. Cho, S. et al. High-Level dCas9 Expression Induces Abnormal Cell
Morphology in Escherichia coli. *ACS Synth Biol* **7**, 1085-1094 (2018).

23. Cui, L. et al. A CRISPRi screen in E. coli reveals sequence-specific toxicity of
dCas9. *Nat Commun* **9**, 1912 (2018).

24. Cao, Y. X. et al. Biosynthesis of odd-chain fatty alcohols in Escherichia coli.
Metab Eng **29**, 113-123 (2015).

25. Cao, Y., Li, X., Li, F. & Song, H. CRISPRi-sRNA:
Transcriptional-Translational Regulation of Extracellular Electron Transfer in
Shewanella oneidensis. *ACS Synth Biol* **6**, 1679-1690 (2017).

26. Jiang, P. & Cronan, J. E., Jr. Inhibition of fatty acid synthesis in Escherichia
coli in the absence of phospholipid synthesis and release of inhibition by
thioesterase action. *J Bacteriol* **176**, 2814-2821 (1994).

27. Cho, H. & Cronan, J. E., Jr. Defective export of a periplasmic enzyme disrupts
regulation of fatty acid synthesis. *J Biol Chem* **270**, 4216-4219 (1995).

28. Li, M., Zhang, X., Agrawal, A. & San, K. Y. Effect of acetate formation
pathway and long chain fatty acid CoA-ligase on the free fatty acid production
in E. coli expressing acy-ACP thioesterase from Ricinus communis. *Metab*
Eng **14**, 380-387 (2012).

29. Prieto, A. I. et al. Genomic analysis of DNA binding and gene regulation by
homologous nucleoid-associated proteins IHF and HU in Escherichia coli K12.
Nucleic Acids Research **40**, 3524-3537 (2012).

30. Fujita, Y., Matsuoka, H. & Hirooka, K. Regulation of fatty acid metabolism in
bacteria. *Mol Microbiol* **66**, 829-839 (2007).

31. Zhang, F. et al. Enhancing fatty acid production by the expression of the
regulatory transcription factor FadR. *Metab Eng* **14**, 653-660 (2012).

32.   van den Ent, F. et al. Structural and mutational analysis of the cell division
598   protein FtsQ. *Mol Microbiol* **68**, 110-123 (2008).

599   33.   Gerding, M. A., Ogata, Y., Pecora, N. D., Niki, H. & de Boer, P. A. The
600   trans-envelope Tol-Pal complex is part of the cell division machinery and
601   required for proper outer-membrane invagination during cell constriction in E.
602   coli. *Mol Microbiol* **63**, 1008-1025 (2007).

603   34.   Baek, J. H., Kang, Y. J. & Lee, S. Y. Transcript and protein level analyses of
604   the interactions among PhoB, PhoR, PhoU and CreC in response to phosphate
605   starvation in Escherichia coli. *FEMS Microbiol Lett* **277**, 254-259 (2007).

606   35.   Youngquist, J. T., Rose, J. P. & Pfleger, B. F. Free fatty acid production in
607   Escherichia coli under phosphate-limited conditions. *Appl Microbiol*
608   *Biotechnol* **97**, 5149-5159 (2013).

609   36.   Pfluger-Grau, K. & Gorke, B. Regulatory roles of the bacterial
610   nitrogen-related phosphotransferase system. *Trends Microbiol* **18**, 205-214
611   (2010).

612   37.   Schneider, E. ABC transporters catalyzing carbohydrate uptake. *Res Microbiol*
613   **152**, 303-310 (2001).

614   38.   Rodriguez-Moya, M. & Gonzalez, R. Proteomic analysis of the response of
615   Escherichia coli to short-chain fatty acids. *J Proteomics* **122**, 86-99 (2015).

616   39.   Marbaix, A. Y. et al. Extremely Conserved ATP- or ADP-dependent
617   Enzymatic System for Nicotinamide Nucleotide Repair. *Journal of Biological*
618   *Chemistry* **286**, 41246-41252 (2011).

619   40.   Li, W., Wu, H., Li, M. & San, K. Y. Effect of NADPH availability on free
620   fatty acid production in Escherichia coli. *Biotechnol Bioeng* **115**, 444-452
621   (2018).

622   41.   Bi, H. & Zhang, C. Integration host factor is required for the induction of acid
623   resistance in Escherichia coli. *Curr Microbiol* **69**, 218-224 (2014).

624   42.   Aquino, P. et al. Coordinated regulation of acid resistance in Escherichia coli.
625   *BMC Syst Biol* **11**, 1 (2017).

626   43.   Smirnova, G. V., Oktyabrsky, O. N., Moshonkina, E. V. & Zakirova, N. V.
627   Induction of the alkylation-inducible aidB gene of Escherichia coli by
628   cytoplasmic acidification and N-ethylmaleimide. *Mutat Res* **314**, 51-56 (1994).

629   44.   Bowles, T., Metz, A. H., O'Quin, J., Wawrzak, Z. & Eichman, B. F. Structure
630   and DNA binding of alkylation response protein AidB. *Proc Natl Acad Sci U S*
631   *A* **105**, 15299-15304 (2008).

632   45.   Rippa, V. et al. Preferential DNA damage prevention by the E. coli AidB gene:
633   A new mechanism for the protection of specific genes. *DNA Repair (Amst)* **10**,
634   934-941 (2011).

635   46.   Xiao, Y., Bowen, C. H., Liu, D. & Zhang, F. Exploiting nongenetic cell-to-cell
636   variation for enhanced biosynthesis. *Nat Chem Biol* **12**, 339-344 (2016).

637   47.   Blazeck, J. et al. Harnessing Yarrowia lipolytica lipogenesis to create a
638   platform for lipid and biofuel production. *Nat Commun* **5**, 3131 (2014).

48. Oud, B., van Maris, A. J., Daran, J. M. & Pronk, J. T. Genome-wide analytical
approaches for reverse metabolic engineering of industrially relevant
phenotypes in yeast. *FEMS Yeast Res* **12**, 183-196 (2012).
49. Yu, X., Liu, T., Zhu, F. & Khosla, C. In vitro reconstitution and steady-state
analysis of the fatty acid synthase from Escherichia coli. *Proc Natl Acad Sci U*
S A **108**, 18643-18648 (2011).
50. Liu, Z. et al. In Vitro Reconstitution and Optimization of the Entire Pathway
to Convert Glucose into Fatty Acid. *ACS Synth Biol* **6**, 701-709 (2017).
51. Clomburg, J. M. et al. Integrated engineering of beta-oxidation reversal and
omega-oxidation pathways for the synthesis of medium chain
omega-functionalized carboxylic acids. *Metab Eng* **28**, 202-212 (2015).
52. Xu, P., Li, L., Zhang, F., Stephanopoulos, G. & Koffas, M. Improving fatty
acids production by engineering dynamic pathway regulation and metabolic
control. *Proc Natl Acad Sci U S A* **111**, 11299-11304 (2014).
53. Liu, D., Xiao, Y., Evans, B. S. & Zhang, F. Negative feedback regulation of
fatty acid production based on a malonyl-CoA sensor-actuator. *ACS Synth Biol*
4, 132-140 (2015).
54. Reis, A. C. et al. Simultaneous repression of multiple bacterial genes using
nonrepetitive extra-long sgRNA arrays. *Nat Biotechnol* (2019).
55. Larson, M. H. et al. CRISPR interference (CRISPRi) for sequence-specific
control of gene expression. *Nat Protoc* **8**, 2180-2196 (2013).
56. Bikard, D. et al. Programmable repression and activation of bacterial gene
expression using an engineered CRISPR-Cas system. *Nucleic Acids Res* **41**,
7429-7437 (2013).
57. Engler, C., Kandzia, R. & Marillonnet, S. A one pot, one step, precision
cloning method with high throughput capability. *PLoS One* **3**, e3647 (2008).
58. Schirmer, A., Rude, M. A., Li, X., Popova, E. & del Cardayre, S. B. Microbial
biosynthesis of alkanes. *Science* **329**, 559-562 (2010).
59. Cao, Y. X. et al. Heterologous biosynthesis and manipulation of alkanes in
Escherichia coli. *Metab Eng* **38**, 19-28 (2016).
60. Zhang, F., Carothers, J. M. & Keasling, J. D. Design of a dynamic
sensor-regulator system for production of chemicals and fuels derived from
fatty acids. *Nat Biotechnol* **30**, 354-359 (2012).
672

Acknowledgements

674 This work was supported by the National Key Research and Development Program of

675 China (2018YFA0901300), the National Natural Science Foundation of China (NSFC

676 21621004), and the Young Elite Scientists Sponsorship Program by Tianjin

677 (TJSQNTJ-2018-16).

679 Author contributions

680 L.F., Y.C. and H.S. conceived and designed the experiments. L.F., J.F., and C.W.

681 performed the plasmids construction experiments. L.F. and J.F. performed the

682 fed-batch fermentation experiments. L.F. and Y.C. analyzed the data. L.F. and Y.C.

683 and H.S. wrote the manuscript. H.S. supervised all aspects of the study.

685 Additional information

686 **Competing interests:** The authors declare no competing financial interests.

Figures

Fig. 1 The CRISPRi approach for identification of beneficial knockdown gene targets

from cellular pathways associated with the FFAs metabolism. **a** Schematic of the

metabolic or regulatory pathways related to FFAs biosynthesis in *E. coli*, which were

classified into five modules (upstream carbon flux diversion, downstream carbon flux

diversion, amino acid metabolism, beta-oxidation and transcription factors) and

represented in blue, orange, purple, green and golden background, respectively. Genes

in the above five modules are shown in italics and marked with blue, orange, purple,

green and golden dot, respectively. *2854* represents gene numbered ECD_02854 in

BL21(DE3). The fatty acid biosynthetic pathway is shown in red line and key

metabolites glycerone-P, glyceraldehyde-3P, glycerate-3P, pyruvate, acetyl-CoA and

malonyl-ACP are represented as number 1 to 6, respectively. **b** Plasmid constructs for

the gene expression of *tesA'*, *dCas9* and synthetic sgRNAs.

Fig. 2 Effect of genetic perturbation by CRISPRi on FFAs production. FFAs

production through repression of each gene in upstream carbon flux diversion module

(**a**), downstream carbon flux diversion module (**b**), amino acid metabolism module (**c**),

beta-oxidation module (**d**), and transcription factors module (**e**) by the CRISPRi

system. All target genes were repressed with high efficiency. * represents the strains

with more than 20% increase in FFAs production than that of the strain Control (631

mg L^{-1}). Error bars indicate the standard deviations of three replicates. Source data are

provided as a Source Data file

Fig. 3 Effect of tuning repression efficiency towards target genes on FFAs production.

a Design of sgRNA with high, medium or low repression efficiency towards target

gene. **b** FFAs production under medium or low repression efficiency compared with

that of the high repression efficiency. The repressed genes were screened out from

those genes marked with star in Fig. 2. Error bars indicate the standard deviations of

three replicates. Source data are provided as a Source Data file

Fig. 4 Exploration of potential target genes by proteome and transcriptome analyses.

a Effect of different *ihfA* or *fadR* expression levels on FFAs production. * represents

the strains with high production of FFAs, and these strains were applied to the next

omics analyses. **b** A Venn diagram of differentially expressed genes (DEGs).

Common DEGs in the high-titer strains at the protein level (a, c) or at the transcript

level (b, d) were identified, compared with the strain Control. Green: down-regulated;

red: up-regulated. **c** Expression levels of the selected common DEGs in the strains

LihfA (1), HfadR (2), and OfadR (3) compared with the strain Control. *1466*

represents the gene numbered ECD_01466 in BL21(DE3). Error bars indicate the

standard deviations of three replicates. Source data are provided as a Source Data file

Fig. 5 Reverse engineering of the selected DEGs in the strains CF and LihfA.

Schematic of cassettes used to repress or overexpress the selected DEGs in the strain

CF (**a**), or in the strain LihfA (**c**). The repression was achieved by CRISPRi and the

overexpression was enabled by the P_{BAD} promoter. TesA', dCas9 and sgRNAs were

controlled by the P_{T7}, P_{Trc} and P_R promoter, respectively. *y*: down-regulated gene; *z*:

up-regulated gene. Effect of genetic perturbation of the selected DEGs on FFAs

production in the strain CF (**b**) or in the strain LihfA (**d**). * represents the strain with

FFAs titer increased >20% compared with the reference strain in each bar chart.

Green frame or column: repression; red frame or column: overexpression. Error bars

indicate the standard deviations of three replicates. Source data are provided as a

Source Data file

Fig. 6 Function network plot of the beneficial genes derived from omics analyses. The

size of the node represents the number of genes within the function set. The node's

color represents W value of the function set. The thickness of the edge indicates the

number of genes overlapping two function sets. W value is calculated by averaging

the FFAs ratio of Hx (Ox) / Control or LiHx (LiOx) / LihfA, x referring to each gene

belongs to the function set. Genes and the FFAs ratio are shown in Supplementary

Table 1. Source data are provided as a Source Data file

Fig. 7 Fed-batch production of FFAs in a 5 L bioreactor. **a** Growth, glucose

consumption and FFAs production of the recombinant *E. coli* strain LihfA-OaidB

during fed-batch fermentation. **b** Solid FFAs layer after centrifugation. After

fermentation, 1 ml of sample was picked up and centrifuged at 5000 g for 10 min.

Error bars indicate the standard deviations of three replicates. Source data are

provided as a Source Data file