Variance in within-pair reproductive success drives the opportunity for sexual selection

annually and over the lifetimes of males in a multi-brooded songbird

Ryan R. Germain[1,2,3*], Michael T. Hallworth[4,5], Sara A. Kaiser[1], T. Scott Sillett[4], and Michael S.
Webster[1,2]

[1] Cornell Lab of Ornithology, Cornell University, Ithaca, NY, USA

[2] Department of Neurobiology and Behavior, Cornell University, Ithaca, NY, USA

[3] Current address: Department of Biology & GLOBE Institute, University of Copenhagen,

Copenhagen, Denmark

[4] Migratory Bird Center, Smithsonian Conservation Biology Institute, National Zoological Park,

Washington, DC, USA

[5] Current address: Northeast Climate Adaptation Science Center, University of Massachusetts

Amherst, Amherst, MA, USA

[*]Corresponding author: ryan.r.germain@gmail.com

Running Title: Within-pair success and sexual selection

Keywords: double brooding, female fecundity, within-pair reproduction

Abstract

In pair-bonding species, male reproductive success consists of 'within-pair' offspring produced

with their socially-paired mate(s), and 'extra-pair' offspring produced with additional females

throughout the population. Both reproductive pathways offer distinct opportunities for sexual

selection to operate in wild populations, as each are composed of sub-components of mate

attraction, female fecundity, and paternity allocation. Identifying key sources of variance and

covariance among these components is a crucial step towards understanding the reproductive

strategies that males use to maximize fitness both annually and over their lifetimes. We use 16

years of complete reproductive data from a population of black-throated blue warblers

(*Setophaga caerulescens*) to decompose variance in male annual and lifetime reproductive

success and thereby identify if the opportunity for sexual selection acts consistently over the

lifetimes of individual males and what reproductive strategies likely favor maximum lifetime

fitness. The majority of variance in male reproduction was attributable to within-pair sources of

variance, but the specific effects of individual components of variance differed annually versus

over the lifetimes of individual males. Positive lifetime covariance between within-pair and

extra-pair success indicates that males able to maximize within-pair success likely achieve higher

overall lifetime fitness via both the within-pair and extra-pair reproductive pathways.

Introduction

Sexual selection represents the evolutionary processes that result from differences in mating or

fertilization success within a population (Andersson 1994). The opportunity, or potential, for

sexual selection to operate is proportional to the population-wide variance in mating and/or

reproductive success; if variance is low (e.g. all individuals have similar numbers of mates and

offspring) then sexual selection will be weak, whereas if variance in reproductive output is high

(e.g. few individuals monopolize mating opportunities and produce many offspring), selection

can potentially be strong (Crow 1958, 1991; Emlen and Oring 1977; Wade and Arnold 1980;

Shuster 2009). However, an individual's total reproductive success (T_{RS}) is itself composed of

several component sources, each of which may have different population-wide variances and

hence represent distinct avenues through which sexual selection may operate (Webster et al.

1995; Whittingham and Dunn 2004; Freeman-Gallant et al. 2005; Dolan et al. 2007; Lawler

2007; Lebigre et al. 2012). Determining where the greatest sources of variance and covariance

exist among these components can help identify the mechanisms that lead to the greatest

opportunities for sexual selection to operate, as well the reproductive strategies that may enable

individuals to maximize fitness annually and over their lifetimes.

Sexual selection tends to operate more strongly among males than among females in

many animal populations (Bateman 1948; Wade 1979; Arnold and Duvall 1994; Shuster 2009;

but see Clutton-Brock 2007). Females are often the choosier sex because they are energetically

limited in the number of broods and/or offspring they can produce (i.e. fecundity) and show

relatively little among-individual variation in reproductive success compared to males (e.g.

Hoogland and Foltz 1982; Freeman-Gallant et al. 2005; Lebigre et al. 2012). In contrast, male

reproductive success is often limited by the number of potential mates they can attract and by the

number of offspring that they sire throughout the population, leading to greater variance in T_{RS} as

some males attract numerous females and/or sire many offspring while others have low mating

or siring success (e.g. Freeman-Gallant et al. 2005; Webster et al. 2007; Lebigre et al. 2012).

Male T_{RS} stems from two pathways in socially monogamous and other species that form pair-

bonds: 'within-pair' offspring produced by their social mate(s) and additional, 'extra-pair'

offspring produced throughout the population by females socially paired to other males

(Birkhead and Møller 1995; Webster et al. 1995; Whittingham et al. 2006). Both pathways are

themselves comprised of multiplicative components of mate attraction (i.e. pre-copulatory) and

paternity (i.e. post-copulatory) success, as well as by the fecundity of a male's within-pair and

extra-pair mates (Figure 1). Each constituent component of within-pair and extra-pair

reproductive success (W_{RS}, E_{RS}) may have different genetic, among-individual, and

environmental sources of variation (Webster et al. 2007; Losdat et al. 2015; Evans and Garcia-

Gonzalez 2016) and hence represent separate opportunities for sexual selection to operate.

Several studies have sought to identify sources of variance among mate attraction and

paternity success (e.g. Vedder et al. 2011; Losdat et al. 2015; Evans and Garcia-Gonzalez 2016),

but less emphasis has been placed on the importance of female fecundity in driving variation in

male T_{RS}. In particular, the ability of some females in a population to produce multiple broods

has the potential to drastically increase overall variance in male W_{RS} by doubling the number of

within-pair offspring they can potentially sire (e.g. Nagy and Holmes 2005; Kaiser et al. 2017).

Alternatively, if extra-pair routes are of primary importance for driving the opportunity for

sexual selection in such populations, multi-brooding should not affect variance in W_{RS}, but rather

allow some males to capitalize on additional opportunities to gain extra-pair success with multi-

brooding females throughout the population, leading to greater variance in E_{RS} than in W_{RS} (e.g.

Webster et al. 2007; Lebigre et al. 2012; Losdat et al. 2015). Thus, quantifying the relative

influence of sub-component sources of within-pair and extra-pair female fecundity (i.e. number

of broods, number of offspring per brood) on male T_{RS} has the potential to identify independent

opportunities for sexual selection to operate and thus potential targets of selection in wild

populations.

Much work has focused on the role that E_{RS} can play in driving the opportunity for sexual

selection in socially monogamous species. The advent of molecular paternity assignment has

revealed previously cryptic sources of variance in individual reproductive success (Coltman et al.

1999; Griffith et al. 2002; Uller and Olsson 2008) that often increases the overall opportunity for

sexual selection to operate (e.g. Yezerinac et al. 1995; Dolan et al. 2007; Webster et al. 2007;

Balenger et al. 2009; Vedder et al. 2011; Lebigre et al. 2013). However, some authors suggest

that many of the purported effects of E_{RS} on the opportunity for selection may be inflated by

incomplete sampling of extra-pair sires in the population (Whittingham and Dunn 2004;

Freeman-Gallant et al. 2005; Albrecht et al. 2007), or may vary with annual variation in social-

environmental factors such as population density (Møller and Birkhead 1993; Taff et al. 2013).

Others have noted that W_{RS} is the major source of variance in male T_{RS} in populations with both

frequent and relatively rare extra-pair mating (Whittingham and Dunn 2004; Albrecht et al.

2007), suggesting that variation in E_{RS} may not necessarily be the driving force behind the

opportunity for sexual selection to operate. However, the majority of studies investigating the

opportunity for sexual selection are conducted over relatively short timescales and are thus

limited to decomposing variance in annual reproductive success only. Given within-individual

heterogeneity in annual reproduction due to delayed maturation of sexual signals (e.g. Germain

et al. 2012; Hawkins et al. 2012), social status-dependent access to mates (e.g. Le Boeuf and

Peterson 1969; DuVal 2012), and terminal investment due to senescence (Pianka and Parker

1975; Clutton-Brock 1984; Martin and Festa-Bianchet 2011; Froy et al. 2013; Tarwater and

Arcese 2017; Piper et al. 2018), individual strategies to maximize annual or lifetime reproductive

success may differ, depending on socio-environmental context or individual phenotype. If

drivers of sexual selection acted consistently over the lifetimes of individuals, we would observe

little variation among years in the effects of individual components of W_{RS} and E_{RS} on total

variance in annual reproductive success (T_{ARS}). Similarly, individual components of W_{RS} and E_{RS}

would exhibit consistent effects on T_{ARS} and total variance in lifetime (T_{LRS}) reproductive

success, as well as realized mean annual and lifetime reproductive success in the population

($\overline{ARS}$ and $\overline{LRS}$, respectively). If, however, individual males trade off shorter- versus longer-term

reproductive success, the effects of individual components of variance in W_{RS} and E_{RS} would

exhibit opposing patterns between annual and lifetime reproductive success. Thus, evaluating

how components of reproduction that affect overall variance in T_{ARS} and T_{LRS} may differ, and

quantifying how such components actually affect realized $\overline{ARS}$ and $\overline{LRS}$, are crucial to our

understanding of how the opportunity for sexual selection operates in wild populations.

For males of many species, particularly those that engage in bi-parental care,

reproductive gains through extra-pair mating may be offset by losses in within-pair paternity,

resulting in a negative covariance (and hence a reproductive trade-off) between W_{RS} and E_{RS}. For

instance, seeking extra-pair copulations may come at the expense of losing within-pair paternity

through reduced mate-guarding (Westneat and Stewart 2003; Kokko and Morrell 2005; Dias et

al. 2009; Harts et al. 2016; Reitsma et al. 2017). In contrast, some studies find positive overall

covariance between W_{RS} and E_{RS}, indicating that males with a higher propensity for acquiring

extra-pair mates and/or siring extra-pair young are also more likely to have higher within-pair

success (Yezerinac et al. 1995; Albrecht et al. 2007; Ferree and Dickinson 2014; Reid et al.

2014). We currently have a poor understanding of where potential sources of such trade-offs

exist in wild populations, in part due to the data requirements needed to accurately quantify

covariance among individual components of W_{RS} and E_{RS}. Co-variances have the potential to

dramatically shape the evolutionary dynamics of reproductive systems (Lebigre et al. 2013; Reid

et al. 2014). Evaluating sources of covariance among individual components of W_{RS} and E_{RS} is

therefore a key facet of identifying potential sources of reproductive trade-offs and thus

quantifying the overall opportunity for sexual selection to operate.

Accurately identifying which individual components of (co)variance drive the

opportunity for sexual selection to operate, as well as realized reproductive success in the

population, annually and over the lifetime of individuals requires comprehensive breeding data

which encompasses pairing, female fecundity, and genetic paternity for a large sample of

individual males over their lifetimes. Here, we use 16 years of detailed breeding observations

and genetic paternity data from a longitudinal study of black-throated blue warblers (*Setophaga*

caerulescens) to identify the contributions of components of W_{RS} and E_{RS}, as well as their

covariances, to the opportunity for sexual selection in this socially monogamous, multi-brooded

songbird. We decompose T_{ARS} and T_{LRS} to their component sources of (co)variance to identify

which sources represent the greatest overall opportunity for sexual selection to operate in this

system, both annually and over the lifetimes of individual males. If within-pair reproductive

success consistently drives the opportunity for sexual selection in this multi-brooded species, we

predict that components of W_{RS} will a) explain more of the total variance in both annual and

lifetime reproductive success, b) exhibit relatively stable effects on variance in reproductive

success over time, and c) be positively correlated with higher realized annual and lifetime

reproductive success in the population. Secondly, we test for the presence of reproductive trade-

offs between components of W_{RS} and E_{RS}. If strong sources of negative covariance exist among

components of W_{RS} and E_{RS}, it would suggest that selection favors males trading off one

reproductive pathway for the other. If, however, there is non-negative covariance between W_{RS}

and E_{RS}, it would suggest that males which employ reproductive strategies that achieve higher

success in one pathway experience higher overall fitness on average, and that selection favors

males which capitalize on multiple facets of reproductive success.

Material and Methods

Population monitoring

We monitored an individually marked population of black-throated blue warblers over 16 years

(1999-2015) at the Hubbard Brook Experimental Forest (3,160ha) in New Hampshire, USA

(43˚56′N, 71˚45′W). The breeding ecology of this long-term study population is described in

detail elsewhere (e.g. Rodenhouse et al. 2003; Sillett et al. 2004; Holmes 2011; Kaiser et al.

2017). In brief, black-throated blue warblers are migratory, territorial songbirds found in

relatively high densities at our study site. Arrival and the onset of breeding typically occurs in

early May (Holmes et al. 2017), and males defend 1-4 ha territories until breeding ends in

August (Sillett et al. 2004). Each season, adult males and females are captured in mist nets,

marked with a unique combination of three colored leg bands and one aluminium U.S.

Geological Survey (USGS) leg band, and blood sampled (~70 µl) from the brachial vein. All

nesting attempts by socially-paired mates (i.e. pairs that jointly defend territories and provision

offspring) are monitored on three study plots (35-85 ha). Nestlings are banded with one USGS

leg band and blood sampled (~30 µl) on day 6 after hatching, and then monitored daily until

fledge date (approx. 9 days after hatching, Holmes et al. 2017).

Genetic paternity assignment and calculation of annual/lifetime reproductive success

Detailed methods regarding genetic paternity assignment for this system are described by

Kaiser et al. (2017). Briefly, we genotyped 4097 offspring and >95% of all adults in the study

area (including candidate males adjacent to plot boundaries) at six highly polymorphic

microsatellite loci using Genemapper v.4.1 (Applied Biosystems). We conducted parentage

analyses for each year and study plot separately using CERVUS v3.0 (Kalinowski et al. 2007),

which uses a maximum likelihood-based approach to infer parentage.

We define a male's annual reproductive success as the number of fledglings that he sired

in a given year. Within-pair success represents the number of sired offspring fledged from his

social female(s) nests on his territory, while extra-pair success represents all other fledged

offspring that he sired within the population that year. Lifetime reproductive success is defined

as a male's summed annual reproductive success over his lifetime as a breeder in the population.

Genetic paternity data were not available before 1999 and after 2016 at the time of analysis;

therefore we excluded n = 11 males that were breeders before 1999 and n = 21 males that

survived beyond 2016 from analyses of lifetime reproductive success to avoid underestimating

this metric for males that may have produced offspring outside of the study period.

Statistical analyses

All analyses were performed in R 3.5.1 (R Development Core Team 2018). Our initial

dataset consisted of 1108 observations of male annual reproductive success from 1999-2015. For

all analyses, we excluded observations from male territories that were part of brood manipulation

(n = 50) or food supplementation (n = 79) experiments. To ensure that our analyses included the

most accurate information possible on extra-pair reproductive success, we also excluded males

that had a higher likelihood of siring unsampled extra-pair offspring outside of the study area (n

= 210 males with territories located within 210m [the equivalent of ~3 territory lengths] of the

plot boundaries, Kaiser et al. 2017). Results remained qualitatively similar between the reduced

and full dataset (e.g. variance decomposition results changed by <1% for total annual

reproductive success).

We first decomposed population-wide variance in T_{ARS} and T_{LRS} over the entire study

period to identify the components of variance in annual and lifetime reproductive success that

present the greatest opportunity for sexual selection to operate. Variance in total reproductive

success (T_{RS}, here representing either total annual [T_{ARS}] or lifetime [T_{LRS}] reproductive success)

is the sum of variances of a male's within-pair reproductive success (W_{RS}) and extra-pair

reproductive success (E_{RS}) success, as well as 2× their associated covariance:

$$\mathrm{Var}(T_{RS}) = \mathrm{Var}(W_{RS}) + \mathrm{Var}(E_{RS}) + 2\,\mathrm{Cov}(W_{RS,}\,E_{RS}).$$

Following Webster et al. (1995), each of these components of variance in T_{RS} can be further

partitioned into separate subcomponents of pairing success (i.e. number of mates, M), female

fecundity, and paternity (P) allocation (and their associated covariances), which act

multiplicatively to equal either W_{RS} or E_{RS} for an individual male (Figure 1). However, our

variance decomposition differs from Webster et al. (1995) in that here we further partition

female fecundity into two parts: the number of broods (B) produced per female, given that some

females in our population will double-brood under favorable environmental conditions (Nagy

and Holmes 2005a; Kaiser et al. 2015), and female fecundity per brood (i.e. the number of

offspring produced per brood, F). For ease of interpretation, we present each component of

variance or covariance as the population wide (co)variance of the component itself (e.g. M_w

represents variance in within-pair pairing success, and $\mathrm{Cov}(M_w, M_e)$ represents the covariance

between within-pair and extra-pair pairing success), but see Supporting Information S1 for full

descriptions of how each variance component was calculated. Males that had zero success in one

component of W_{RS} or E_{RS} were not included in analyses of downstream components of variance

decomposition to avoid conflating variance estimates (following Webster et al. 1995; Freeman-

Gallant et al. 2005). Note that this method of variance decomposition also estimates remainder

terms, where the total remainder (D_T) is equal to $D_w + D_e + D_{we}$. These remainder terms capture

multivariate skewness expressed in the higher order moments of the distributions of each

subcomponent of reproductive success, such that total variance in T_{ARS} and T_{LRS} is not a simple

sum of the component variances and covariances (Bohrnstedt and Goldberger 1969; Webster et

al. 1995; Dolan et al. 2007; Lawler 2007).

We identified components of (co)variance in both T_{ARS} and T_{LRS} that constituted a

substantial percentage (arbitrary cut-off of $\geq 10\%$) of the total variance across the full study

period. For T_{ARS}, we further conducted time-series analyses to evaluate whether these key

components in annual reproductive success consistently provided the greatest opportunity for

sexual selection in each year of study (i.e. under varying socio-environmental conditions such as

population density or population age-structure). To do so, we replicated our variance

decomposition (above) for each individual year in the study period, extracted the percent of total

variance in T_{ARS} attributed to each key component, as well as overall (co)variance in W_{RS} and

E_{RS}, and both visually inspected time series plots for presence of lags and conducted

Kwiatkowki-Phillips-Schmidt-Shin (KPSS) tests to determine if each time series was trend

stationary (i.e. lack of overall positive/negative trend over the study period).

Lastly, we determined the relative effect of each key component of W_{RS} and E_{RS} on mean

annual and lifetime reproductive success ($\overline{ARS}$ and $\overline{LRS}$) in the population using generalized

mixed effects models and a model averaging approach. For $\overline{ARS}$, we constructed a global mixed

model (Poisson distribution) which included male identity and year as random effects, and M_w,

B_w, P_w, M_e, B_e as fixed effects (see Table 1 for justification) as well as the interaction between B_w

and B_e, to represent the covariance between these two components that explain a substantial

portion of variance in T_{ARS} (Table 1). All fixed effects were standardized to mean = 0, SD = 1 to

reduce any influence of measurement scale on model results (White and Burnham 1999). We ran

all possible combinations of these fixed effects (n = 40 models total) and selected a subset with a

difference in Akaike Information Criterion (ΔAIC) $\leq$ 7 from the top model. We chose a cutoff of

ΔAIC $\leq$ 7 to ensure that estimates of the relative influences of each predictor on $\overline{ARS}$ were as

conservative and inclusive as possible (Burnham and Anderson 2002). We then recalculated

relative AIC weights from this subset of models, assessed the relative support for each predictor

by summing the AIC weight of each model in which it was included. Averaged parameter

estimates for each predictor were weighted by the cumulative AIC weight of each model in

which it was included to create one representative estimate of the relative effects of each

component on $\overline{ARS}$ (Burnham and Anderson 2002; Germain and Arcese 2014; Germain et al.

2015, 2018). Statistical significance for each fixed effect was assessed by whether 95%

confidence intervals (CIs) overlapped zero.

We repeated this approach for $\overline{LRS}$, constructing a global mixed model (Poisson

distribution) which included male identity and male longevity (years breeding in the population)

as random effects, and M_w, B_w, F_w, P_w, M_e, B_e, F_e, and P_e as fixed effects, as well as the

interactions between $M_w \times F_w$, $M_e \times F_e$, $M_w \times M_e$, $M_w \times F_e$, $M_w \times P_e$, $B_w \times B_e$, $B_w \times P_e$, $F_w \times M_e$,

and $F_e \times F_w$ to represent covariances among these key components of variance in T_{LRS} (see Table

2 for justification). We again ran all possible combinations of these fixed effects (n = 4458

models), selected those within $\Delta AIC \leq 7$ of the top model, and constructed an average model

based on relative AIC weights (as above).

Results

Male black-throated blue warblers exhibited considerable variation in both within-pair (mean =

1.7 young sired ± 1.9SD, range = 0–9) and extra-pair (mean = 1.1 young sired ± 1.5SD, range =

0–9) annual reproductive success, leading to substantial variation in the total number of offspring

produced annually (mean = 2.8 ± 2.3SD, range = 0–14; Figure 2a). Similarly, males varied in the

number of within-pair (mean = 2.76 ± 3.2SD, range = 0–23) and extra-pair (mean = 1.77± 2.4SD,

range = 0–14) offspring produced over their lifetimes. Hence, the Hubbard Brook population had

relatively large variance in lifetime reproductive success (mean = 4.54 $\pm$ 4.7SD, range = 0–28;

Figure 2b) indicating that most males produced few offspring and a small proportion produced

many.

The opportunity for sexual selection in Annual Reproductive Success

286   Decomposition of the total opportunity for sexual selection in annual reproductive

success revealed that the majority of variance (~64%) in T_{ARS} was attributable to variance in

within-pair reproductive success, of which within-pair paternity success (P_w) constituted the

greatest proportion (26.4%, Table 1). Despite this, both the number of within-pair mates (M_w)

and number of broods produced by within-pair mates (B_w) also each accounted for roughly 10%

of T_{ARS}, suggesting multiple distinct opportunities for sexual selection to operate within the W_{RS}

pathway of annual reproductive success (Table 1). In contrast, the ~42% of variance in T_{ARS}

attributable to the extra-pair pathway (E_{RS}) was almost entirely accounted for by the number of

extra-pair mates the male acquired (M_e, 39.6%; Table 1). Overall covariance between W_{RS} and

E_{RS} was slightly negative (-6.2% of total variance in T_{ARS}), with only the positive covariance

between the number of broods by within-pair and extra-pair females (Cov[B_w, B_e]) accounting

for a substantial proportion of variance in T_{ARS} (10.6%, Table 1).

Variance in T_{ARS} varied across the 16 year study, indicating that the overall opportunity

for sexual selection differed among years (Figure 3a). Despite this, variance in T_{ARS} was trend

stationary (KPSS trend = 0.13, p = 0.08), indicating no overall positive/negative trend in the

change in variance over the study period. The percent of T_{ARS} attributable to variance in W_{RS}

likewise varied considerably, but also remained trend stationary (KPSS trend = 0.13, p = 0.07)

and accounted for more total variance than E_{RS} in each year of study (Figure 3b). Among

components of variance in within-pair reproductive success (Figure 3b), both number of within-

pair mates (M_w) and number of within-pair broods (B_w) alternated above and below 10% of

variance in T_{ARS}, and while M_w exhibited no obvious trend (KPSS trend = 0.11, p = 0.1), variance

in B_w bordered on a significant negative trend over the study period (KPSS trend = 0.15, p =

0.05, Figure 3b). Similarly, the proportion of within-pair offspring sired (P_w), which consistently

accounted for the largest percentage of T_{ARS} among within-pair components, also bordered on a

significant negative trend (KPSS trend = 0.14, p = 0.05), due to high within-pair variance during

the early portion of the study. Both E_{RS} and its main component, the number of extra-pair mates

acquired, consistently accounted for ~40% of T_{ARS} and both were trend stationary (E_{RS} – KPSS

trend = 0.05, p = 0.1; M_e – KPSS trend = 0.10, p = 0.1). In contrast, the overall covariance

between W_{RS} and E_{RS} varied from -73% to 28% of total variance in T_{ARS} over the study period,

where negative values indicate a trade-off between within-pair and extra-pair reproductive

success (Figure 3d). Despite this wide range of variance in T_{RS} attributed to $Cov(W_{RS}, E_{RS})$, the

overall covariance between within-pair and extra-pair reproductive success remained trend

stationary (KPSS trend = 0.07, p = 0.1), as did its main constituent component, the covariance

between within-pair and extra-pair number of broods (KPSS test = 0.08, p = 0.1).

Our global model of the effects of each key component on $\overline{ARS}$ was a good fit to the data

(R^2 = 0.43). Our model subset included six models with $\Delta AIC \leq 7$ from the top model; the

cumulative AIC weight of this model subset was <0.98. All five fixed effects and the interaction

between B_w and B_e were included in our final averaged model. Of these, M_w, B_w, P_w, and M_e had

significant, positive effects on $\overline{ARS}$ (Figure 4), indicating that males at Hubbard Brook achieved

the highest annual reproductive success by pairing with multiple double-brooded females,

securing within-pair paternity, and mating with multiple extra-pair females (See Supporting

Information S2 for full parameter estimates from final averaged model and all models in top

subset).

The opportunity for selection in Lifetime Reproductive Success

Decomposition of the opportunity for sexual selection in lifetime reproductive success

revealed that, similar to T_{ARS}, substantially more variance in T_{LRS} was attributable to variance in

within-pair reproductive success than to variance in extra-pair success (46.7% versus 26.7%,

respectively, Table 2). Indeed, every individual component of W_{RS} contributed a substantial

(>10%) proportion of total variance in T_{LRS}, and the covariance between M_w and F_w accounted

for over 50%. Similarly, although M_e again contributed the most of any single component to

variance in E_{RS}, F_e and P_e also accounted for $\geq$10% of total variance. The covariance between M_e

and F_e contributed to almost 100% of variance in T_{LRS}, indicating that the interaction between

these two components is a strong driver of variance in male lifetime reproductive success (Table

2). Several components of within-pair/extra-pair covariances contributed to more than 10% of

the total positive or negative variance in T_{LRS}, suggesting the potential for both synergistic and

trade-off effects between individual components of each pathway. However, overall covariance

between W_{RS} and E_{RS} was positive and accounted for as much total variance in T_{LRS} as E_{RS} itself

(~27%), indicating that there is no overall trade-off between the within-pair and extra-pair

pathways in terms of male lifetime reproductive success in this system. Despite the strong

potential for these listed sources of (co)variance to affect overall population-wide variance in

male T_{LRS}, large, negative sources of (co)variance (D_w, D_e, D_{we}, D_T, Table 2) were unaccounted

for in our decomposition. Although the biological significance of such remainder terms is

difficult to interpret without knowledge of how higher-order moments of the distribution of each

within-pair and extra-pair component affect lifetime reproductive success (e.g. multivariate

skewness: Bohrnstedt and Goldberger 1969; Lawler 2007), their relatively large, negative values

suggest that some amount of realized variation in T_{LRS} is not captured by decomposing variance

to its within-pair and extra-pair components and their associated covariances.

Our global model of the effects of each key component on mean lifetime reproductive

success was an extremely good fit to the data ($R^2 = 0.86$). A total of 147 models were included in

the subset of models within $\Delta AIC \leq 7$ from the top model, which included all fixed effects and

all interaction terms; the cumulative AIC weight of this model subset was <0.99. We detected

significant, positive effects of B_w, F_w, P_w, B_e, F_e, P_e on $\overline{LRS}$, and significant negative effects from

the interactions between M_w x F_w, and M_e x F_e, indicating that males in this population achieve

the highest lifetime reproductive success by pairing with fewer double-brooded females that

produced more offspring per brood and allocated more within-pair paternity to the focal male,

and by mating with fewer double-brooding extra-pair females that likewise produced more

offspring per brood and allocated more paternity towards the focal male. All other terms had CIs

that overlapped zero (Figure 5), and were not considered to have statistically significant effects

on $\overline{LRS}$ (see Supporting Information S3 for full parameter estimates from final averaged model

and all models in top subset).

Discussion

Quantifying the opportunity for sexual selection among different components of

individual reproductive success is a key parameter for determining the evolutionary dynamics of

mating systems within wild populations (Webster et al. 1995; Reid et al. 2014; Losdat et al.

2015). By identifying sources of the opportunity for sexual selection, and determining how the

effects of these sources may change annually or over the lifetimes of individual males, we can

elucidate the reproductive strategies that males employ to maximize reproductive success as well

as determine how such strategies may change over time. In a free living, multi-brooded songbird

population, we found that among-individual variation in within-pair components of reproductive

success consistently accounted for more of the total overall variance in both annual and lifetime

reproductive success than did extra-pair components of variance (Table 1, Table 2, Figure 3).

Several of these within-pair components were likewise associated with higher mean annual

(Figure 4) and lifetime (Figure 5) reproductive success, and generally exhibited non-negative

covariance with extra-pair components, suggesting that allocating resources towards attracting

and defending access to socially-paired mates is likely a key tactic for maximizing annual and

lifetime fitness in species which exhibit environmentally-mediated multiple brooding.

The majority of variance in total annual reproductive success (T_{ARS}) was accounted for by

four individual components of variance and one component of covariance which each explained

≥10% (Table 1). Of these individual components, the number of extra-pair mates that a male

acquired (M_e) accounted for the single largest amount of total variance in T_{ARS}. However, the

number of within-pair mates that a male socially paired with (M_w), as well as the number of

broods these females produced (B_w) and the proportion of within-pair offspring sired by the male

(P_w) accounted for more total variance than M_e and the remaining components of extra-pair

reproduction combined (Table 1). Although the percentage of T_{ARS} accounted for by W_{RS} did vary

somewhat from year-to-year in our study (Figure 3b), the influences of each individual

component of W_{RS} were relatively stable over the longer-term, indicating that individual

behavioral strategies aimed at maximizing annual reproductive success should be under

consistent selection in this population. This finding contrasts with previous work indicating that

the influence of particular variance components on the opportunity for selection can co-vary with

socio-environmental factors like annual breeding density over shorter time periods (e.g. Møller

and Birkhead 1993; Taff et al. 2013; Evans and Garcia-Gonzalez 2016). Total variance in annual

reproductive success was also substantially influenced by one component of covariance, that

between the number of broods produced by a male's socially-paired mate(s) and his extra-pair

mate(s) (Cov[B_w, B_e], Table 1). The overall covariance between W_{RS} and E_{RS} was slightly

negative for T_{ARS} (~ -6%, Table 1) and tended to be negative in most years (Figure 3d); however,

this positive covariance between B_w and B_e was relatively consistent in accounting for ≥10% of

total variance in T_{ARS}, and was never negative over the 16 year study (Figure 3d). This indicates

that while the opportunity for sexual selection to operate on synergistic effects of multiple

brooding among within-pair and extra-pair mates can vary somewhat annually, males never

experienced a trade-off between the number of broods their socially-paired and extra-pair mates

produced. While future studies will need to pinpoint the precise spatial and/or environmental

variables which drive the positive covariance between within-pair and extra-pair double

brooding, our results indicate that this covariance makes a relatively small but important

contribution towards driving the opportunity for sexual selection in this system.

412   In practice, all four key individual components of variance in T_{ARS} (i.e. M_w, B_w, P_w, and

M_e) had significant, positive effects on mean annual reproductive success ($\overline{ARS}$) in black-

throated blue warblers (Figure 4). Previous work from this population likewise indicates that

greater investment in aspects of within-pair reproduction results in a higher reproductive net-gain

for individual males, and that males on high-quality, food-rich territories tend to invest in

behaviors that enhance W_{RS} (e.g. increased mate-guarding and territorial defense) over those that

enhance E_{RS} (e.g. extra-territorial forays; Kaiser et al. 2015, 2017). Food limitation appears to be

the proximate mechanism behind such territorially-driven variation in mating strategies at

Hubbard Brook, as females nesting in food-rich territories are more likely to double brood than

those occupying food-limited habitat (Holmes et al. 1996; Nagy and Holmes 2005a; Kaiser et al.

2015, 2017). Taken together, our results indicate that securing a high-quality breeding territory is

the key strategy for male black-throated blue warblers to maximize annual reproductive success

via the within-pair pathway, and that doing so does not necessarily limit their ability to also

achieve some E_{RS} through mating with multiple extra-pair females. Indeed, doing so may

occasionally result in a higher reproductive net gain via the positive covariance between within-

pair and extra-pair female brooding (Table 1, Figure 3). For instance, males on food-rich

territories with double-brooding females may have neighboring females that likewise double-

brood given greater food resources (Holmes et al. 1996; Nagy and Holmes 2005b), increasing the

availability of extra-pair offspring for the focal male to sire. With warming temperatures, earlier

springs and longer breeding seasons increase the frequency of double-brooding in black-throated

blue warblers and other passerines (Townsend et al. 2013; Both et al. 2019), which may explain

the near-negative trend in B_w over our study (i.e. more double-brooding females will decrease

overall variance in B_w; Figure 3). Increased growing season length with earlier leaf-out date and

later leaf senescence, documented at Hubbard Brook (Richardson et al. 2006) and elsewhere

(Ibáñez Inés et al. 2010; Fridley 2012) could therefore translate to higher realized annual

reproductive success for males able to capitalize on both maximizing W_{RS} among their double-

brooding socially-paired mate(s) and gaining additional E_{RS} with extra-pair females throughout

the population.

Within-pair components of variance also accounted for the majority of variance in total

lifetime reproductive success (T_{LRS}), as we found for T_{ARS}, but the relative influences of within-

pair and extra-pair sources of (co)variance on the opportunity for sexual selection differed

somewhat between annual and lifetime reproduction. In total, W_{RS} components accounted for

~47% of variance in T_{LRS}, whereas components of E_{RS} components accounted for ~27% of

variance (Table 2). Further, more individual components of T_{LRS} accounted for ≥10% of variance

than those of T_{ARS}. Several of these individual components accounted for over half (to almost all)

of the total variance in T_{LRS} before taking negative covariance terms and remainder (D) terms

into account (Table 2). Indeed, a striking feature of our variance decomposition of T_{LRS} is that

several covariance terms, most notably between individual within-pair and extra-pair

components, accounted for substantial proportions of variance, with the overall covariance

between W_{RS} and E_{RS} accounting for as much of the total variance as overall extra-pair variance

(~27%; Table 2). This highlights that potential synergies and tradeoffs among individual

reproductive components may play a much larger role in influencing reproductive success over a

male's lifetime than within a single breeding season. In particular, the largest components of

such covariance were positive and involved the number of within-pair and/or extra-pair mates

and the fecundity per brood of within-pair and/or extra-pair females (i.e. Cov[M_w, M_e], Cov[M_w,

F_e], Cov[F_w, M_e], Cov[F_w, F_e]; Table 2). Substantial, positive within-pair/extra-pair covariances

in the opportunity for sexual selection, as observed here, indicate that there are large potential

lifetime fitness benefits for males able to capitalize on mating with extra-pair females while still

investing in behaviors that enhance W_{RS} with their socially-paired mate(s). Previous research

from this and other systems indicate that local breeding synchrony likely plays a role in the

ability of males to copulate with extra-pair female neighbours while maintaining their own

within-pair reproductive success (Stutchbury and Neudorf 1998; Chuang et al. 1999; Webster et

al. 2001; Stewart et al. 2010). By mating with synchronously fertile females on nearby adjacent

territories, males of some species can reduce the risk of losing within-pair paternity of their own

offspring via reduced mate guarding while seeking potential extra-pair mates over greater

distances throughout the population (Kempenaers 1997; Weatherhead 1997). Future work

investigating environmental drivers of the opportunity for sexual selection should thus consider

the role of local breeding synchrony as it relates to positive covariances between the W_{RS} and E_{RS}

pathways.

Of the components of (co)variance found to have a substantial influence on the

opportunity for sexual selection in lifetime reproductive success, six had significant, positive

effects on $\overline{LRS}$ in the population (Figure 5). Surprisingly, neither the number of within-pair nor

number of extra-pair mates had any significant effect on $\overline{LRS}$, contrary to our results for $\overline{ARS}$

(Figure 4). Instead, all individual components of variance related to female fecundity (B_w, F_w, B_e,

F_e) and male siring success (P_w, P_e) had significant positive effects on $\overline{LRS}$, indicating that

realized male reproductive success is more strongly influenced by female breeding quality and

paternity allocation than aspects of male mate attraction. Indeed, the interaction between number

of within-pair mates and their fecundity per brood, as well as the interaction between number of

extra-pair mates and fecundity per brood of those mates were found to have significant negative

effects on $\overline{LRS}$ (Figure 5), indicating that male black-throated blue warblers experience a lifetime

trade-off in terms of the number and the quality of their mates through both the W_{RS} and E_{RS}

pathways. These results are consistent with the hypothesis that males with multiple within-pair

and/or extra-pair mates suffer reproductive loses annually compared to monogamous males (e.g.

Dunn and Robertson 1993; Poirier et al. 2004). However, such patterns of negative covariance

and lower reproductive success are not widely reported over the lifetimes of breeding males, in

particular when contrasted with the apparent positive effects of more within-pair and extra-pair

mates within a single breeding season. Taken together, our results suggest that males in this

population may sacrifice short-term reproductive gains (i.e. higher $\overline{ARS}$ via more within-pair

and/or extra-pair mates) for longer-term fitness benefits (represented by higher $\overline{LRS}$).

Overall, our results suggest that variance in within-pair reproductive success clearly

drives the opportunity for sexual selection in this multiple-brooded population. While the

components of (co)variance which most strongly affect both the opportunity for sexual selection

and realized reproductive success differed somewhat between short (i.e. annual) and longer (i.e.

lifetime) timescales, components of W_{RS} consistently accounted for more total variance in T_{ARS}

and T_{LRS}, and had more contributions to higher mean annual and lifetime reproductive success,

than did components of E_{RS}. The potential for annual reproductive tradeoffs between within-pair

and extra-pair components of variance were outweighed by the overall positive covariance

among these reproductive pathways. Thus, males able to maximize W_{RS} likely achieve higher

overall lifetime fitness via both the within-pair and extra-pair reproductive pathways. Minimal

observed effects of annual variation in socio-environmental factors on the influences of

components of W_{RS} indicate that consistent selection pressures may offer some males (e.g. those

on food-rich territories, Kaiser et al. 2015, 2017) the opportunity to capitalize on long-term

reproductive benefits (e.g. double-brooding by their socially-paired female), even if such benefits

result in short term losses in annual reproductive success via reduced effort in seeking additional

mates. In contrast, males on food-poor territories may experience more selective pressure to

allocate resources towards extra-pair reproduction given their lower probability of within-pair

success (Kaiser et al. 2015, 2017), leading to greater variance in components of E_{RS} in this subset

of males. Future studies should seek to determine if and how the opportunity for sexual selection

differs between such high vs. low-quality territories, particularly as changing climate conditions

alter breeding-season length and potentially reduce population-wide variance in key components

of reproduction such as the ability of females to initiate multiple broods. Such studies will not

only provide unique insight into how habitat type can influence sexual selection and the lifetime

fitness of individual males, but also advance our understanding of how changing socio-

environmental conditions can ultimately alter the mating systems of wild populations in seasonal

environments.

Acknowledgements

We thank the many undergraduates and field technicians that contributed towards data

collection, G.J. Colbeck, E.R.A. Cramer, and S. Bartos-Smith for assistance in the laboratory and

with genotyping, and R.T. Holmes, M.P. Ayers, and N.L. Rodenhouse for logistic support. This

research is a contribution of the Hubbard Brook Ecosystem Study, part of the Long-Term

Ecological Research network supported by the National Science Foundation (grants awarded to

Cornell University [0640470], the Smithsonian Institution [0640732] and Wellesley College

[064082300]). R.R.G was supported by a postdoctoral grant from the Natural Sciences and

Engineering Research Council of Canada.

Author contributions

R.R.G. and M.S.W. conceived the study, and R.R.G. conducted statistical analyses and wrote the

manuscript with input from all authors. M.T.H, S.A.K, T.S.S. and M.S.W. collected data, and

S.A.K. conducted parentage assignment.

Literature Cited

Albrecht, T., J. Schnitzer, J. Kreisinger, A. Exnerová, J. Bryja, and P. Munclinger. 2007.

Extrapair paternity and the opportunity for sexual selection in long-distant migratory

passerines. Behav. Ecol. 18:477–486.

Andersson, M. 1994. Sexual selection. Princeton University Press, Princeton, NJ.

Arnold, S. J., and D. Duvall. 1994. Animal mating systems: a synthesis based on selection

theory. Am. Nat. 143:317–348.

Balenger, S. L., L. Scott Johnson, H. L. Mays Jr, and B. S. Masters. 2009. Extra-pair paternity in

the socially monogamous mountain bluebird *Sialia currucoides* and its effect on the

potential for sexual selection. J. Avian Biol. 40:173–180.

Bateman, A. J. 1948. Intra-sexual selection in Drosophila. Heredity 2:349–368.

Birkhead, T. R., and A. P. Møller. 1995. Extra-pair copulation and extra-pair paternity in birds.

Anim. Behav.49:843–848.

Bohrnstedt, G. W., and A. S. Goldberger. 1969. On the exact covariance of products of random

variables. J. Am. Stat. Assoc. 64:1439–1442.

Both, C., R. Ubels, and P.-A. Ravussin. 2019. Life-history innovation to climate change: can

single-brooded migrant birds become multiple breeders? J. Avian Biol. 50:e01951.

Burnham, K., and D. Anderson. 2002. Model Selection and Multi-Model Inference: A Practical

Information-Theoretic Approach. Springer, New York, NY.

Chuang, H. C., M. S. Webster, and R. T. Holmes. 1999. Extrapair Paternity and Local Synchrony

in the Black-Throated Blue Warbler. Auk 116:726–736.

Clutton-Brock, T. 2007. Sexual selection in males and females. Science 318:1882–1885.

Clutton-Brock, T. H. 1984. Reproductive effort and terminal investment in iteroparous animals.

Am. Nat. 123:212–229.

Coltman, D. W., D. R. Bancroft, A. Robertson, J. A. Smith, T. H. Clutton-brock, and J. M.

Pemberton. 1999. Male reproductive success in a promiscuous mammal: behavioural

estimates compared with genetic paternity. Mol. Ecol. 8:1199–1209.

Crow, J. F. 1991. Fitness variation on natural populations. Pp. 91–97 *in* Evolution and Animal

Breeding Systems. C.A.B. International, Wallingford.

Crow, J. F. 1958. Some possibilities for measuring selection intensities in man. Hum. Biol. 30:1–

13.

Dias, R. I., P. Salles, and R. H. Macedo. 2009. Mate guarding and searching for extra-pair

copulations: Decision-making when interests diverge. Ecol. Inform.4:405–412.

Dolan, A. C., M. T. Murphy, L. J. Redmond, K. Sexton, and D. Duffield. 2007. Extrapair

paternity and the opportunity for sexual selection in a socially monogamous passerine.

Behav. Ecol. 18:985–993.

Dunn, P. O., and R. J. Robertson. 1993. Extra-pair paternity in polygynous tree swallows. Anim.

Behav. 45:231–239.

DuVal, E. H. 2012. Variation in annual and lifetime reproductive success of lance-tailed

manakins: alpha experience mitigates effects of senescence on siring success. Proc. R.

Soc. B 279:1551–1559.

Emlen, S. T., and L. W. Oring. 1977. Ecology, Sexual Selection, and the Evolution of Mating

Systems. Science 197:215–223.

Evans, J. P., and F. Garcia-Gonzalez. 2016. The total opportunity for sexual selection and the

integration of pre-and post-mating episodes of sexual selection in a complex world. J.

Evol. Biol. 29:2338–2361.

Ferree, E. D., and J. L. Dickinson. 2014. Male western bluebirds that sire extrapair young are

also successful within-pair mates. Anim. Behav. 90: 11–19

Freeman-Gallant, C. R., N. T. Wheelwright, K. E. Meiklejohn, S. L. States, and S. V. Sollecito.

2005. Little effect of extrapair paternity on the opportunity for sexual selection in

savannah sparrows (*Passerculus sandwichensis*). Evolution 59:422–430.

Fridley, J. D. 2012. Extended leaf phenology and the autumn niche in deciduous forest invasions.

Nature 485:359–362.

Froy, H., R. A. Phillips, A. G. Wood, D. H. Nussey, and S. Lewis. 2013. Age-related variation in

reproductive traits in the wandering albatross: evidence for terminal improvement

following senescence. Ecol. Lett. 16:642–649.

Germain, R. R., and P. Arcese. 2014. Distinguishing individual quality from habitat preference

and quality in a territorial passerine. Ecology 95:436–445.

Germain, R. R., M. W. Reudink, P. P. Marra, P. T. Boag, and L. M. Ratcliffe. 2012. Delayed

maturation of multiple signals in a migratory songbird. Behav. Ecol. Sociobiol. 66:419–

431.

Germain, R. R., R. Schuster, K. E. Delmore, and P. Arcese. 2015. Habitat preference facilitates

successful early breeding in an open-cup nesting songbird. Funct. Ecol. 29:1522–1532.

Germain, R. R., R. Schuster, C. E. Tarwater, W. M. Hochachka, and P. Arcese. 2018. Adult

survival and reproductive rate are linked to habitat preference in territorial, year-round

resident Song Sparrows *Melospiza melodia*. Ibis 160:568–581.

Griffith, S. C., I. P. Owens, and K. A. Thuman. 2002. Extra pair paternity in birds: a review of

interspecific variation and adaptive function. Mol. Ecol. 11:2195–2212.

Harts, A. M. F., I. Booksmythe, and M. D. Jennions. 2016. Mate guarding and frequent

copulation in birds: A meta-analysis of their relationship to paternity and male

phenotype. Evolution 70:2789–2808.

Hawkins, G. L., G. E. Hill, and A. Mercadante. 2012. Delayed plumage maturation and delayed

reproductive investment in birds. Biol. Rev. 87:257–274.

Holmes, R. T. 2011. Avian population and community processes in forest ecosystems: Long-

term research in the Hubbard Brook Experimental Forest. Forest Ecol. Manag. 262:20–

32.

Holmes, R. T., S. A. Kaiser, N. L. Rodenhouse, T. S. Sillett, M. S. Webster. 2017. Black-

throated Blue Warbler (*Setophaga caerulescens*), version 3.0. In: Birds of North America

(P. Pyle, and M. A. Patten, eds). Cornell Lab of Ornithology, Ithaca, NY, USA.

Holmes, R. T., P. P. Marra, and T. W. Sherry. 1996. Habitat-specific demography of breeding

black-throated blue warblers (*Dendroica caerulescens*): implications for population

dynamics. J. Anim. Ecol. 65:183–195.

Hoogland, J. L., and D. W. Foltz. 1982. Variance in male and female reproductive success in a

harem-polygynous mammal, the black-tailed prairie dog (Sciuridae: *Cynomys*

ludovicianus). Behav. Ecol. Sociobiol. 11:155–163.

Ibáñez Inés, Primack Richard B., Miller-Rushing Abraham J., Ellwood Elizabeth, Higuchi

Hiroyoshi, Lee Sang Don, Kobori Hiromi, and Silander John A. 2010. Forecasting

phenology under global warming. Philos. T. Roy. Soc. B. 365:3247–3260.

Kaiser, S. A., B. B. Risk, T. S. Sillett, and M. S. Webster. 2017. Ecological and social factors

constrain spatial and temporal opportunities for mating in a migratory songbird. Am. Nat.

189:283–296.

Kaiser, S. A., T. S. Sillett, B. B. Risk, and M. S. Webster. 2015. Experimental food

supplementation reveals habitat-dependent male reproductive investment in a migratory

bird. Proc. R. Soc. B. 282:20142523.

Kalinowski, S. T., M. L. Taper, and T. C. Marshall. 2007. Revising how the computer program

CERVUS accommodates genotyping error increases success in paternity assignment.

Mol. Ecol. 16:1099–1106.

Kempenaers, B. 1997. Does Reproductive Synchrony Limit Male Opportunities or Enhance

Female Choice for Extra-Pair Paternity? Behaviour 134:551–562.

Kokko, H., and L. J. Morrell. 2005. Mate guarding, male attractiveness, and paternity under

social monogamy. Behav Ecol 16:724–731.

Lawler, R. R. 2007. Fitness and extra-group reproduction in male Verreaux's sifaka: An analysis

of reproductive success from 1989–1999. Am. J. Phys. Anthropol. 132:267–277.

Le Boeuf, B. J., and R. S. Peterson. 1969. Social status and mating activity in elephant seals.

Science 163:91–93.

Lebigre, C., P. Arcese, and J. M. Reid. 2013. Decomposing variation in male reproductive

success: age-specific variances and covariances through extra-pair and within-pair

reproduction. J. Anim. Ecol. 82:872–883.

Lebigre, C., P. Arcese, R. J. Sardell, L. F. Keller, and J. M. Reid. 2012. Extra-pair paternity and

the variance in male fitness in song sparrows (*Melospiza melodia*). Evolution 66:3111–

3129.

Losdat, S., P. Arcese, and J. M. Reid. 2015. Double decomposition: decomposing the variance in

subcomponents of male extra-pair reproductive success. J. Anim. Ecol. 84:1384–1395.

Martin, J. G. A., and M. Festa-Bianchet. 2011. Age-independent and age-dependent decreases in

reproduction of females. Ecol. Lett. 14:576–581.

Møller, A. P., and T. R. Birkhead. 1993. Cuckoldry and sociality: a comparative study of birds.

Am. Nat. 142:118–140.

Nagy, L. R., and R. T. Holmes. 2005a. Food limits annual fecundity of a migratory songbird: an

experimental study. Ecology 86:675–681.

Nagy, L. R., and R. T. Holmes. 2005b. To double-brood or not? Individual variation in the

reproductive effort in Black-Throated Blue Warblers (*Dendroica Caerulescens*). Auk

122:902–914.

Pianka, E. R., and W. S. Parker. 1975. Age-specific reproductive tactics. Am. Nat. 109:453–464.

Piper, W. H., K. M. Brunk, G. L. Jukkala, E. A. Andrews, S. R. Yund, and N. G. Gould. 2018.

Aging male loons make a terminal investment in territory defense. Behav Ecol Sociobiol

72:95.

Poirier, N. E., L. A. Whittingham, and P. O. Dunn. 2004. Males achieve greater reproductive

success through multiple broods than through extrapair mating in house wrens. Anim.

Behav. 67:1109–1116.

R Development Core Team. 2018. R: A Language and Environment for Statistical Computing. R

Foundation for Statistical Computing, Vienna, Austria.

Reid, J., P. Arcese, L. Keller, and S. Losdat. 2014. Female and male genetic effects on offspring

paternity: additive genetic (co)variances in female extra-pair reproduction. Evolution

68:3357–2370.

Reitsma, L. R., J. A. Jukosky, A. J. Kimiatek, M. L. Goodnow, and M. T. Hallworth. 2017.

Extra-pair paternity in a long-distance migratory songbird beyond neighbors' borders and

across male age classes. Can. J. Zool. 96:49–54.

Richardson, A. D., A. S. Bailey, E. G. Denny, C. W. Martin, and J. O'keefe. 2006. Phenology of

a northern hardwood forest canopy. Glob. Change Biol. 12:1174–1188.

Rodenhouse, N., T. Sillett, P. Doran, and R. Holmes. 2003. Multiple density–dependence

mechanisms regulate a migratory bird population during the breeding season. Proc. R.

Soc. B. 270:2105–2110.

Shuster, S. M. 2009. Sexual selection and mating systems. PNAS 106 Suppl 1:10009–16.

Sillett, T. S., N. L. Rodenhouse, and R. T. Holmes. 2004. Experimentally reducing neighbor

density affects reproduction and behavior of a migratory songbird. Ecology 85:2467–

2477.

Stewart, S. L. M., D. F. Westneat, and G. Ritchison. 2010. Extra-pair paternity in eastern

bluebirds: effects of manipulated density and natural patterns of breeding synchrony.

Behav Ecol Sociobiol 64:463–473.

Stutchbury, B. J. M., and D. L. Neudorf. 1998. Female control, breeding synchrony, and the

evolution of extra-pair mating systems. Ornithol. Mono. 103–121.

Taff, C. C., C. R. Freeman-Gallant, P. O. Dunn, and L. A. Whittingham. 2013. Spatial

distribution of nests constrains the strength of sexual selection in a warbler. J. Evol. Biol.

26:1392–1405.

Tarwater, C. E., and P. Arcese. 2017. Age and years to death disparately influence reproductive

allocation in a short-lived bird. Ecology 98:2248–2254.

Townsend, A. K., T. S. Sillett, N. K. Lany, S. A. Kaiser, N. L. Rodenhouse, M. S. Webster, and

R. T. Holmes. 2013. Warm springs, early lay dates, and double brooding in a North

American migratory songbird, the black-throated blue warbler. PLOS ONE 8:e59467.

Uller, T., and M. Olsson. 2008. Multiple paternity in reptiles: patterns and processes. Mol. Ecol.

17:2566–2580.

Vedder, O., J. Komdeur, M. van der Velde, E. Schut, and M. J. L. Magrath. 2011. Polygyny and

extra-pair paternity enhance the opportunity for sexual selection in blue tits. Behav Ecol

Sociobiol 65:741–752.

Wade, M. J. 1979. Sexual selection and variance in reproductive success. Am. Nat. 114:742–

747.

Wade, M. J., and S. J. Arnold. 1980. The intensity of sexual selection in relation to male sexual

behaviour, female choice, and sperm precedence. Anim. Behav. 28:446–461.

Weatherhead, P. J. 1997. Breeding synchrony and extra-pair mating in red-winged blackbirds.

Behav Ecol Sociobiol 40:151–158.

Webster, M. S., H. C. Chuang-Dobbs, and R. T. Holmes. 2001. Microsatellite identification of

extrapair sires in a socially monogamous warbler. Behav Ecol 12:439–446.

Webster, M. S., S. Pruett-Jones, D. F. Westneat, and S. J. Arnold. 1995. Measuring the effects of

pairing success, extra-pair copulations and mate quality on the opportunity for sexual

selection. Evolution 49:1147–1157.

Webster, M. S., K. A. Tarvin, E. M. Tuttle, and S. Pruett-Jones. 2007. Promiscuity drives sexual

selection in a socially monogamous bird. Evolution 61:2205–2211.

Westneat, D. F., and I. R. K. Stewart. 2003. Extra-pair paternity in birds: causes, correlates, and

conflict. Annu. Rev. Ecol. Syst. 34:365–396.

White, G., and K. Burnham. 1999. Program MARK: survival estimation from populations of

marked animals. Bird Study 46:120–139.

Whittingham, L. A., and P. O. Dunn. 2004. Effects of extra-pair and within-pair reproductive

success on the opportunity for selection in birds. Behav. Ecol. 16:138–144.

Whittingham, L. A., P. O. Dunn, and M. K. Stapleton. 2006. Repeatability of extra-pair mating

in tree swallows. Mol. Ecol. 15:841–849.

Yezerinac, S. M., P. J. Weatherhead, and P. T. Boag. 1995. Extra-pair paternity and the

opportunity for sexual selection in a socially monogamous bird (*Dendroica petechia*).

Behav Ecol Sociobiol 37:179–188.

Table 1: Decomposition of variance in total annual reproductive success (T_{ARS}) of black-throated

blue warblers across 16 years of study. For each row, variance terms (see Figure 1 for

definitions) represent the focal component of (co)variance in T_{ARS}, but note that actual

calculation of each term involves population-wide means for additional components (see

Supporting Information S1 and Webster et al. 1995 for further details). D denotes remainder

terms for within-pair, extra-pair, and covariance pathways, where $D_T = D_w + D_e + D_{we}$. Values

depict the absolute variance in each component, as well as their relative variance (i.e. divided by

population mean $T_{ARS}^2 = 8.713$), and the percentage of total variance in T_{ARS} attributable to each

term. Crosses (†) denote variance terms which fall above our cutoff of >10% total variance in

T_{ARS}.

Source	Variance term	Absolute variance	Relative variance	% total variance
Total variance in ARS	T_{ARS}	**5.467**	**0.677**	**100**
Within-pair terms	M_w	0.544	0.067	10.0[†]
	B_w	0.591	0.073	10.8[†]
	F_w	0.426	0.053	7.8
	P_w	1.443	0.179	26.4[†]
	$Cov(M_w, B_w)$	0.018	0.002	0.3
	$Cov(M_w, F_w)$	-0.030	-0.004	-0.5
	$Cov(M_w, P_w)$	-0.035	-0.004	-0.6
	$Cov(B_w, F_w)$	-0.027	-0.004	-0.5
	$Cov(B_w, P_w)$	-0.059	-0.007	-1.1
	$Cov(F_w, P_w)$	0.191	0.024	3.5
Total within-pair variance	W_{RS}	**3.487**	**0.432**	**63.8**
	D_W	**0.425**	**0.053**	**7.7**
Extra-pair terms	M_e	2.166	0.269	39.6[†]
	B_e	0.180	0.022	3.3
	F_e	0.130	0.016	2.4
	P_e	0.482	0.060	8.8
	$Cov(M_e, B_e)$	0.014	0.002	0.3
	$Cov(M_e, F_e)$	0.025	0.003	0.5
	$Cov(M_e, P_e)$	-0.075	-0.009	-1.4
	$Cov(B_e, F_e)$	0.005	0.001	0.1
	$Cov(B_e, P_e)$	-0.242	-0.030	-4.4
	$Cov(F_e, P_e)$	-0.196	-0.024	-3.6
Total extra-pair variance	E_{RS}	**2.319**	**0.288**	**42.4**
	D_E	**-0.17**	**-0.022**	**-4.5**
Within-pair and extra-pair covariances	$Cov(M_w, M_e)$	-0.419	-0.052	-7.7
	$Cov(M_w, B_e)$	0.099	0.012	1.8
	$Cov(M_w, F_e)$	-0.009	-0.001	-0.2
	$Cov(M_w, P_e)$	-0.484	-0.060	-8.9
	$Cov(B_w, M_e)$	0.023	0.003	4.2
	$Cov(B_w, B_e)$	0.579	0.072	10.6[†]
	$Cov(B_w, F_e)$	0.008	0.001	0.2
	$Cov(B_w, P_e)$	-0.389	-0.048	-7.1
	$Cov(F_w, M_e)$	0.121	0.015	2.2
	$Cov(F_w, B_e)$	-0.044	-0.005	-0.8
	$Cov(F_w, F_e)$	0.063	0.008	1.2
	$Cov(F_w, P_e)$	-0.096	-0.012	-1.8
	$Cov(P_w, M_e)$	0.090	0.011	1.6
	$Cov(P_w, B_e)$	0.071	0.009	1.3
	$Cov(P_w, F_e)$	-0.025	-0.003	-0.4

	$Cov(P_w, P_e)$	-0.097	-0.012	-1.8
Total WP and EP covariance	$Cov(W_{RS}, E_{RS})$	**-0.339**	**-0.042**	**-6.2**
	D_{WE}	**0.170**	**0.020**	**-0.6**
Remainder	D_T	**0.425**	**0.051**	**3.9**

Table 2: Decomposition of variance in total lifetime reproductive success (T_{LRS}) of black-

throated blue warblers across 16 years of study. For each row, variance terms (see Figure 1 for

definitions) represent the focal component of (co)variance in T_{LRS}, but note that actual calculation

of each term involves population-wide means for additional components (see Supporting

Information S1 and Webster et al. 1995 for further details). D denotes remainder terms for

within-pair, extra-pair, and covariance pathways, where $D_T = D_w + D_e + D_{we}$. Values depict the

absolute variance in each component, as well as their relative variance (i.e. divided by population

mean $T_{LRS}^2 = 20.57$), and the percentage of total variance in T_{LRS} attributable to each term.

Crosses (†) denote variance terms which fall above our cutoff of >10% total variance in T_{LRS}.

Source	Variance term	Absolute value	Standardized value*	% total variance
Total Variance	T_{LRS}	**21.77**	**1.058**	**100**
Within-pair terms	M_w	5.766	0.280	26.5[†]
	B_w	5.331	0.259	24.5[†]
	F_w	7.672	0.373	35.2[†]
	P_w	6.140	0.299	28.2[†]
	$Cov(M_w, B_w)$	1.718	0.084	7.8
	$Cov(M_w, F_w)$	12.383	0.602	56.9[†]
	$Cov(M_w, P_w)$	0.539	0.026	2.5
	$Cov(B_w, F_w)$	-0.213	-0.010	-1.0
	$Cov(B_w, P_w)$	-0.165	-0.008	-0.8
	$Cov(F_w, P_w)$	0.908	0.044	4.2
Total within-pair variance	W_{RS}	**10.171**	**0.494**	**46.7**
	D_W	**-29.908**	**-1.455**	**-137.3**
Extra-pair terms	M_e	19.627	0.954	90.1[†]
	B_e	1.216	0.059	5.6
	F_e	7.345	0.357	33.7[†]
	P_e	3.336	0.162	15.3[†]
	$Cov(M_e, B_e)$	0.318	0.016	1.5
	$Cov(M_e, F_e)$	21.767	1.047	99.9[†]
	$Cov(M_e, P_e)$	-1.491	-0.072	-6.8
	$Cov(B_e, F_e)$	0.259	0.013	1.2
	$Cov(B_e, P_e)$	-1.752	-0.085	-8.1
	$Cov(F_e, P_e)$	-1.773	-0.086	-8.1
Total extra-pair variance	E_{RS}	**5.806**	**0.282**	**26.7**
	D_E	**-43.046**	**-2.083**	**-197.6**
Within-pair and extra-pair covariances	$Cov(M_w, M_e)$	12.860	0.625	59.1[†]
	$Cov(M_w, B_e)$	0.394	0.019	1.8
	$Cov(M_w, F_e)$	8.264	0.402	37.9[†]
	$Cov(M_w, P_e)$	-3.141	-0.145	-14.4[†]
	$Cov(B_w, M_e)$	0.478	0.023	2.2
	$Cov(B_w, B_e)$	3.648	0.177	16.8[†]
	$Cov(B_w, F_e)$	0.389	0.019	1.8
	$Cov(B_w, P_e)$	-2.627	-0.128	-12.1[†]
	$Cov(F_w, M_e)$	15.835	0.770	72.7[†]
	$Cov(F_w, B_e)$	-0.432	-0.021	-2.0
	$Cov(F_w, F_e)$	10.395	0.505	47.7[†]
	$Cov(F_w, P_e)$	-0.319	-0.015	-1.5
	$Cov(P_w, M_e)$	1.101	0.053	5.1

	$Cov(P_w, B_e)$	-0.166	-0.008	-0.8
	$Cov(P_w, F_e)$	0.379	0.018	1.7
	$Cov(P_w, P_e)$	-0.484	-0.024	-2.2
Total WP and EP covariance	$Cov(W_{RS}, E_{RS})$	**5.792**	**0.278**	**26.6**
	D_{WE}	**-40.782**	**-1.991**	**-187.2**
Remainder	D_T	**-113.0736**	**-5.529**	**-522.1**

Figure 1: Flow chart depicting the components of within-pair (WP) and extra-pair (EP) reproductive success, which combine to determine an individual's total reproductive success. Arrows indicate the effects that environmental and phenotypic traits can have on WP and EP components, and hence on total reproductive success. Italic terms (in parentheses) denote abbreviations used for each component of variance. Note that components of within-pair and extra-pair pathways are multiplicative, whereas WP and EP success themselves are additive components of total annual reproductive success (adapted from Webster et al. 1995).

Figure 2: Total number of within-pair (black bars) and extra-pair (white bars) offspring produced by male black-throated blue warblers a) annually (n = 769 male-years) and b) over their complete breeding lifetime (n = 572 males). In each panel, insert (grey bars) depicts total number of offspring (within-pair and extra-pair) produced by these same males.

Figure 3: Time series plots depicting a) yearly variance in total annual reproductive success (T_{ARS}) across a population of black-throated blue warblers, and yearly decompositions of the percent of total variance in T_{ARS} attributable to b) within-pair reproductive success (W_{RS}), c) extra-pair reproductive success (E_{RS}), and d) the covariance between within-pair and extra-pair reproductive success [$Cov(W_{RS}, E_{RS})$]. For panels b, c, and d, colored dashed, dotted, and dot-dashed lines represent the yearly percentage of total variance in T_{ARS} attributable to constituent components of W_{RS}, E_{RS}, and $Cov(W_{RS}, E_{RS})$ found to represent 'substantial variance' ($\geq 10\%$) across the study period (see Table 1), depicted as solid grey lines. For panel d, dashed gray line represents the transition from positive to negative covariance. Note different y-axis scales for panels a) and d).

Figure 4: Effects plots from averaged generalized mixed-effects models testing the relative influence of each key component of (co)variance on mean annual reproductive success $(\overline{ARS})$. Points represent parameter estimates from final averaged model while whiskers depict 95% CIs. Dotted line represents 0, and parameter are considered significant if CIs do not overlap zero. Full results from averaged model and all models in top subset are provided in Supporting Information S2.

Figure 5: Effects plots from averaged generalized mixed-effects models testing the relative influence of each key component of (co)variance on mean lifetime reproductive success ($\overline{LRS}$). Points represent parameter estimates from final averaged model while whiskers depict 95% CIs. Dotted line represents 0, and parameters are considered significant if CIs do not overlap zero. Full results from averaged model and all models in top subset are provided in Supporting Information S3.