1 **CRISPR-mediated Multiplexed Live Cell Imaging of Nonrepetitive Genomic**
2 **Loci**
3
4

5 Patricia A. Clow[1], Menghan Du[1,2], Nathaniel Jillette[1], Aziz Taghbalout[1], Jacqueline J. Zhu[1,5]*,
6 Albert W. Cheng[1,2,3,4,5]*
7
8
9
[1]The Jackson Laboratory for Genomic Medicine, Farmington, CT 06032, USA

[2]Department of Genetics and Genome Sciences, University of Connecticut Health Center,
Farmington, CT 06030, USA

[3]The Jackson Laboratory Cancer Center, Bar Harbor, ME 04609, USA

[4]Institute for Systems Genomics, University of Connecticut Health Center, Farmington, CT
06030, USA

[5]School of Biological and Health Systems Engineering, Arizona State University, Tempe, AZ
85281, USA

*Correspondence to: albert@cheng.bio or jacquelinejufenzhu@gmail.com

Abstract

Three-dimensional (3D) structures of the genome are dynamic, heterogeneous and functionally

important. Live cell imaging has become the leading method for chromatin dynamics tracking.

However, existing CRISPR- and TALE-based genomic labeling techniques have been hampered

by laborious protocols and are ineffective in labeling non-repetitive sequences. Here, we report a

versatile CRISPR/Casilio-based imaging method that allows for a nonrepetitive genomic locus to

be labeled using one guide RNA. We construct Casilio dual-color probes to visualize the

dynamic interactions of DNA elements in single live cells in the presence or absence of the

cohesin subunit RAD21. Using a three-color palette, we track the dynamic 3D locations of

multiple reference points along a chromatin loop. Casilio imaging reveals intercellular

heterogeneity and interallelic asynchrony in chromatin interaction dynamics, underscoring the

importance of studying genome structures in 4D.

Main

Three-dimensional (3D) organization of the genome has pervasive roles in the regulation of

transcription, DNA replication, and DNA repair[1-3]. Sequencing methods[4, 5] can uncover

genome-wide chromatin interactions in bulk cell populations but are unable to resolve

intercellular heterogeneity or temporal dynamics (the "fourth dimension") which instead require

live cell imaging techniques[6]. Four-dimensional (4D) live cell imaging of genomic loci can be

achieved by inserting into the target loci large tandem arrays of sequences that are bound by

well-characterized DNA binding domains fused to fluorescent proteins providing important

insights into the dynamics of chromatin interaction and transcription[7, 8]. However, these

techniques require tedious genome engineering and can be confounded by the side effects of

disruptive introduction of exogenous sequences at the target loci. Live-cell imaging of the

unmodified genome can be conducted with the use of fluorescent proteins fused to

programmable DNA binding proteins such as zinc finger proteins (ZFPs) and transcription

activator-like effectors (TALEs), but the complicated cloning protocols requiring weeks of

development limit their scalability[9-12]. CRISPR/Cas systems offer versatile DNA binding

proteins (e.g., Cas9) that can be programmed by a target-specific guide RNA (gRNA),

revolutionizing the field of genome biology[13]. Fluorescent proteins or fluorophores recruited

by dCas9 or assembled on the gRNA scaffold offer easily programmable platforms for live cell

genomic imaging[14-20]. However, despite current techniques providing robust labeling of

repetitive sequences using a single gRNA, imaging of nonrepetitive sequences requires

simultaneous delivery of up to tens of gRNA to achieve sufficient signal, precluding broad

adoption.

We previously engineered a modular multitasking CRISPR/Cas-based platform called

Casilio by combining dCas9 and engineered Pumilio/FBF (PUF)-tethered effectors[21] (**Fig. 1**).

An effector domain carrying out a specific function (e.g., transcription activation) is fused to a

unique PUF RNA binding domain that can bind to one or more copies of a specific 8-mer motif

inserted to the 3' end of a gRNA. PUF domains contain peptide subunits that can each be

programmed to recognize an RNA-base by changing amino acids contacting the RNA, thereby

allowing designed PUF domains to bind different 8-mer sequences[22, 23]. Casilio capitalizes on

the versatility of PUF domains to allow simultaneous and orthogonal delivery of effector

functions to distinct genome loci, as well as multimerization of effector molecules, for efficient

epigenetic editing[21, 24].

In this study, we developed Casilio for imaging nonrepetitive DNA sequences. We

reasoned that the immense multimerization capability of Casilio will enable concentration of

fluorescent proteins at a target locus to achieve sufficient signal using reduced number of target

DNA binding sites while the multiplexing capability will allow simultaneous multicolor labeling

of distinct loci in the same cell. We first tested Casilio for imaging nonrepetitive sequences

within the *MUC4* gene on chromosome 3 which was previously labeled by dCas9-EGFP direct

fusion with >30 gRNAs tiled over the region[14]. We designed gMUC4 which binds a

nonrepetitive sequence within *MUC4* and inserted 15 copies of Pumilio Binding Site c

(15xPBSc) to recruit Clover-PUFc. To allow the assessment of the specificity of Casilio, we co-

labeled the *MUC4* region with a Cy5-MUC4 DNA fluorescence *in situ* hybridization (FISH)

probe in U2OS and ARPE-19 cells labelled with Casilio/gMUC4 (**Fig. 2a-c**). Casilio achieves

highly specific labeling of the nonrepetitive locus using only one gRNA with 96% and 100%

specificity in U2OS and ARPE-19 cells, respectively. Additional support for specificity is that

the formation of fluorescent spots is dependent on the presence of dCas9 (**Fig. 2d**), and that the

number of fluorescent spots is consistent with the ploidy of the diploid ARPE-19 (**Fig. 2d**) and

haploid HAP1 (**Fig. 2e**) cells. Spot formation efficiency was around ~68% for ARPE-19 (**Fig.**

2d) and ~54% for HAP1 (**Fig. 2e**). These together demonstrate that robust and specific labeling

of nonrepetitive DNA sequence by Casilio requires only one gRNA.

To test whether Casilio can be applied to study the dynamics of chromatin interactions in

live cells, we selected a chromatin interaction, the *MASP1-BCL6*, with 362 kb genomic distance

between the interaction anchors from a published RAD21 ChIA-PET experiment performed in

ARPE-19 cells (ENCSR110JOO, Michael Snyder lab). We then designed a pair of gRNAs to

bind nonrepetitive sequences next to the anchors (**Fig. 3a**). We used Clover-PUFc to label the 5'

anchor (locus A) and PUF9R-iRFP670 to label the 3' anchor (locus B). Live cell microscopy of

ARPE-19 cells transfected with dual-color Casilio probe pairs revealed dynamic chromatin

interactions at second timescales and showcased a high degree of intercellular heterogeneity and

interallelic asynchrony of chromatin interaction at this loop (**Fig. 3b,c; Supplementary Video 1;**

Supplementary Fig. 1,2a). Such insights cannot be readily obtained via bulk cell sequencing

technologies, supporting the advantages of Casilio live-cell imaging for studying the dynamics of

chromatin interactions at single cell and single allele level. To provide future users a method to

assess the specificity of labeling in live-cell settings, we developed a "competitor" assay. Here, a

"cold" specific competitor gRNA without any PBS sites for fluorescent protein recruitment, but

overlaps the target, is co-introduced into cells to interfere with the binding of Casilio imaging

complexes and the formation of target spots (**Fig. 3d**). A non-specific competitor gRNA that

does not overlap the target (such as GAL4[25]) is used as a control. To denote a competitor

gRNA, we prefix the target with "c" to mean cold or competitor. Specific competitor gRNA cA

overlapping with *MASP1-BCL6* locus A greatly reduced the formation of the Clover spots

compared to non-specific competitor cGAL4 control (**Fig. 3e**). As expected, the iRFP670 spots

formed at the non-interfered location (*MASP1-BCL6* locus B) served as an internal control that

was unaffected by either competitor gRNAs.

To test if Casilio can be applied to study the function of cohesin complexes in chromatin

interactions, we took advantage of an engineered HCT116 cell line with endogenous RAD21

fused to an auxin-inducible degron (AID)[26] that allows for the conditional depletion of RAD21

by auxin treatment. A previous study performed Hi-C assays on this cell line and demonstrated

that the depletion of RAD21 in HCT116 cells resulted in loss of loop domains including a ~500

kb loop domain between the *IER5L* promoter (IER5L-P) and its super-enhancer (IER5L-SE)[27].

We used Clover-PUFc and PUF9R-iRFP670 to label the *IER5L* promoter and super-enhancer,

respectively (**Fig. 4a; Supplementary Fig. 3**), and performed live-cell microscopy on untreated

(-Auxin) (**Fig. 4b; Supplementary Video 2; Supplementary Fig. 2b,4**) and RAD21-depleted

(+Auxin) cells (**Fig. 4c; Supplementary Video 3; Supplementary Fig. 2c,5**). Depletion of

RAD21 led to farther pairwise distances of the Casilio spots in cells treated with auxin compared

to the untreated control, with medians of 2.09 μm and 1.21 μm, respectively (Mann-Whitney

two-sided *p*-value<0.00001) (**Fig. 4d**), consistent with the reduced interaction between the

promoter and super enhancer of *IER5L* upon RAD21 depletion[27]. We also used Casilio to

image a "RAD21-independent" interaction that displayed an opposite behavior[27], in which

interaction frequency increased upon RAD21 depletion (**Fig. 4e**). Consistent with the

observation obtained by sequencing experiments, pairwise distances of Casilio spots decreased

upon auxin-mediated RAD21-depletion, with medians of 1.24 μm and 0.57 μm, respectively

(Mann-Whitney two-sided *p*-value<0.00001), for the untreated and auxin-treated cells (**Fig 4f-h;**

Supplementary Videos 4,5; Supplementary Fig. 2d,e).

These data demonstrate that Casilio can image interaction between cis-regulatory

elements to study the contribution of protein factors, such as cohesin, on dynamic chromatin

interactions underlying gene regulation.

Imaging specific interactions of non-coding elements such as enhancers and promoters

provide new insights into gene regulation, while visualizing a continuous stretch of genomic

region will improve our understanding of structural folding dynamics and illuminate the process

of chromatin loop formation. Given that Casilio can image each nonrepetitive locus with a single

gRNA, we next explored the possibility of simultaneously imaging multiple nonrepetitive loci,

with the ultimate goal of tracking the "live" structure of a continuous genomic region. We call

this technique, where sequential Casilio probes are deployed across a stretch of genomic DNA,

"Programmable Imaging of Structure with Casilio Emitted sequence of Signal" or PISCES.

To increase the number of nonrepetitive loci that can be labelled, we developed a three-

color palette. We used three gRNAs (IER5L P-20xPBS$_{48107}$, M-20xPBSc, and SE-25xPBS9R)

(**Fig. 5a**) for labeling three nonrepetitive locations at the *IER5L* P-SE loop, with a three-color

fluorescent protein palette consisting of Clover-PUF$_{48107}$, iRFP670-PUFc, and PUF9R-mRuby2

(**Fig. 5b,c; Supplementary Videos 6,7; Supplementary Fig. 2f,g,6**). These experiments

demonstrated the use of dual or triple-color fluorescent protein combinations to encode a

sequence of PISCES probes for visualizing multiple reference points along a genomic region.

In this study, we present a CRISPR/Cas-based method for live cell fluorescence imaging

of nonrepetitive genomic loci with low gRNA requirement (one gRNA/locus) and high

spatiotemporal resolutions. We applied Casilio to visualize the dynamic interactions of DNA

elements in native, unmodified chromosomes, in the presence or absence of RAD21, showing

RAD21-dependent and independent loops display opposite behaviors upon RAD21 depletion. By

developing a three fluorescent protein palette, we showed that chromatin loops can be imaged at

multiple reference points over time. These revealed the highly heterogeneous and dynamic

nature of chromatin interactions and structural changes not attainable with bulk cell sequencing

approaches, underscoring the need for studying the 4D nucleome with high spatiotemporal

resolution in live cells. Although gRNA-scaffold methods based on MS2 hairpin have been

attempted before to reduce gRNA requirement for non-repetitive DNA labeling, the addition of

14 or more MS2 hairpins to gRNA greatly reduce its expression[18], whereas in Casilio, addition

of up to 47 copies of PBS did not affect dCas9 binding activity[21]. The high signal of target-

bound Casilio complexes compared to the background unbound complexes can be explained by

the observation that gRNA is unstable unless complexed with dCas9 at the target sites[20]. These

unique properties of Casilio enable the reduction of its gRNA requirement compared with

previously published CRISPR-based approaches. Therefore, Casilio not only significantly

reduces the technical challenge of using live cell imaging to study chromatin interactions in hard-

to-transfect cells, but also it simplifies the design of genome-wide gRNA libraries for imaging.

One caveat to bear in mind when conducting and interpreting live-cell genome imaging

experiments is the potential of the interference of the natural genome folding processes by the

binding of the CRISPR reagents, which might be partially alleviated with careful gRNA designs.

It is thus an important future goal to systematically evaluate probe design and produce a design

pipeline that takes into account the potential interference of the underlying genome folding

processes under investigation. Future development of the Casilio imaging toolkit should include

higher-density PISCES probe-set with an expanded palette of fluorescent proteins, in order to

enable the folding of the genomic region to be delineated at a higher resolution. The multiplexing

and expandable nature of Casilio in adapting new effector functions[21], in combination with the

advance in genomic imaging demonstrated here, will enable "on-the-fly" and "plug-and-play"

perturbation of the (epi)genome (e.g., using Casilio activator, repressor and epigenetic editing

modules[21, 24]) and concurrent read-out of dynamic 3D chromatin interactions, thereby

providing unprecedented flexibility and power for the study of the 4D nucleome (**Fig. 1b**).

Methods

Design of chromatin interaction imaging experiments

Fastq files of ARPE-19 RAD21 ChIA-PET experiment (ENCSR110JOO) were downloaded

from ENCODE database and processed by ChIA-PET2[28]. Interactions were ranked by PET

count. Top-ranked interactions with different PET distances were selected. To avoid interference

of CTCF binding, design regions were selected by shifting external to PET regions. JACKIE[29]

was then used to identify unique 1-copy CRISPR sites in the hg38 genome overlapping design

regions of selected loops and further filtered for specificity by Cas-OFFinder[30]. ChIA-PET

loops are displayed on WashU EpiGenome Browser (https://epigenomegateway.wustl.edu/).

Cloning for Casilio imaging

The lentiviral dCas9 expression plasmid (lenti-dCas9-Blast) was generated by PCR-based

mutagenesis of lentiCas9-Blast plasmid (Addgene #52962). Clover fused with PUF RNA-

binding domain were previously described: pAC1447 (Clover_PUFc) (Addgene #73689).

PUF9R-tethered iRFP670 and mRuby2 were created by a combination of PCR (from IDT

gBlocks) and ligation cloning placing iRFP670[31] or mRuby2[32], respectively, downstream of

PUF9R. PUFc-tethered iRFP670 and PUF$_{48107}$ tethered Clover was created by a combination of

PCR (from IDT gBlocks) and ligation cloning placing iRFP670[31] upstream of PUFc or Clover

upstream of PUF$_{48107}$, respectively. These PUF-fluorescent protein fusions contain nuclear

localization signal (NLS) for their localization in the nucleus.

Guide RNA are under control of the human U6 promoter. gRNA spacer sequences were

cloned into sgRNA-PBS expression vectors pCR8-sgRNA-15xPBSc or pCR8-sgRNA-

15xPBS9R via an oligo-annealing protocol[33].

All plasmids were subjected to restriction diagnostic tests and sequenced to ensure

correct cloning. Plasmids will be available on Addgene.

Target sequences for gRNAs

MUC4: GCCGGTGACAGGAAGAGTGC

<u>RAD21 loops in ARPE-19 and HCT116 cells</u>

MASP1-BCL6 loop Locus A: GGGTAAGAAGCCACTAGGGT

MASP1-BCL6 loop Locus B: GCATAGCCGCATTTGAAAGC

IER5L-P: GACTCCCGCGGGTCACTCGG

IER5L-SE: GCCATGCTCCGATAAGGATA

IER5L-M: GTTTACCCCAAGGGTCGCGG

<u>RAD21-independent loop in HCT116 cells</u>

Clover locus: GCTTGCCGACGATGGACCCAT

iRFP670 locus: GAGGATCGAAACGCTGATGCG

<u>Competitor gRNA design</u>

cGAL4: GAACGACTAGTTAGGCGTGTA

c(*MASP1-BCL6*)A (*MASP1-BCL6* loop Locus A competitor, dSaCas9):

GTGCTCCAGGGTAAGAAGCCAC

c*IER5L*-P: GCTTGACTCCCGCGGGTCACT

Cell culture

Human osteosarcoma U2OS cells (ATCC® HTB-96™) were cultivated in Dulbecco's Modified

Eagle's Medium (DMEM) (Sigma) with 10% fetal bovine serum (Lonza), 4% Glutamax (Gibco),

1% Sodium Pyruvate (Gibco) and 1% penicillin-streptomycin (Gibco). Human retinal pigment

epithelial ARPE-19 cells (ATCC® CRL-2302™) were cultivated in DMEM/F12 (Gibco) with

10% fetal bovine serum (Lonza), and 1% penicillin-streptomycin (Gibco). Human chronic

myeloid leukemia HAP1 cells (Horizon™ C859) were cultivated in Iscove's Modified

Dulbecco's Medium (IMDM) with 20% fetal bovine serum (Lonza) and 1% penicillin-

streptomycin (Gibco). HCT116/RAD21-mAID cells were obtained from Kanemaki Lab[26]. The

cells were cultured in McCoy's 5A medium (ATCC) with 10% fetal bovine serum (Lonza), 2

mM L-glutamine (ATCC), penicillin-streptomycin (ThermoFisher Scientific). Incubator

conditions were humidified 37°C and 5% CO_2. Cell lines expressing constitutive dCas9 were

generated by transducing cells with lentiviruses prepared from a lenti-dCas9-Blast plasmid,

followed by Blasticidin selection.

Transfection

U2OS/dCas9 cells were seeded at density of 100,000 cells/compartment in a 35 mm 4-

compartment CELLview™ cell culture dish (Greiner Bio-One) 24 hours before transfection.

Cells were transfected with 550 ng of sgRNA plasmid containing 15 Pumilio Binding Sites

(15xPBS) and 50 ng of Clover-PUF fusion plasmid, using 1.5 µl Lipofectamine 3000

(Invitrogen). Media was changed at 24 hours post-transfection.

ARPE-19/dCas9 cells were seeded at density of 80,000 cells/compartment in 35 mm 4-

compartment CELLview™ cell culture dish (Greiner Bio-One) 17 hours before transfection.

Cells were transfected with 400 ng of each sgRNA-15xPBS plasmid, 40 ng of Clover-PUFc

fusion plasmid, and 40ng of PUF9R-iRFP670 fusion plasmid using 1.5 µl Lipofectamine LTX

(Invitrogen). For competitor experiments, 400ng of the competitor plasmid was added. Media

was changed at 24 hours post-transfection.

HAP1/dCas9 cells were seeded at density of 150,000 cells/compartment in 35 mm 4-

compartment CELLview™ cell culture dish (Greiner Bio-One) 30 hours before transfection.

Cells were transfected with 400 ng of sgRNA-15xPBS plasmid and 40 ng of Clover-PUFc fusion

plasmid DNA using 3.5 µl Turbofectin (OriGene). Media was changed at 24 hours post-

transfection.

HCT116/RAD21-mAID/dCas9 cells were seeded at density of 60,000 cells/compartment

in 35 mm 4-compartment CELLview™ cell culture dish the day before transfection. For each

well, cells were transfected with 300 ng of each sgRNA plasmid and 40 ng of each fluorescent

protein plasmid using 3.5 µL TurboFectin (OriGene). For competitor experiments, 300 ng of the

competitor plasmid was added. For three-color experiments, cells were transfected with 400ng of

each sgRNA plasmid, 40ng of Clover-PUF_{48107} fusion plasmid, 40ng of iRFP670-PUFc fusion

plasmid, and 400ng of PUF9R-mRuby2 fusion plasmid using 3.5 µL TurboFectin (OriGene).

Media was changed at 24 hours post-transfection. Degradation of the AID-tagged RAD21 was

induced by the addition of 500 µM (final concentration) indole-3-acetic acid (IAA/auxin; Sigma

Aldrich) during media change 24 hours prior to imaging if applicable.

Live-cell confocal microscopy

Imaging was performed at 41-51 hours post-transfection. Prior to imaging, cells were stained

with 0.5-1.0 µg/ml Hoechst 33342 prepared in cell culture media for 15-30 minutes, followed by

two media washes. Images were acquired with the Dragonfly High Speed Confocal Platform 505

(Andor) using an iXon EMCCD camera and a Leica HCX PL APO 40x/1.10 W CORR objective

or a Leica HC PL APO 63x/1.47NA OIL CORR TIRF objective mounted on a Leica DMi8

inverted microscope equipped with a live-cell environmental chamber (Okolab) at humidified

37°C and 5% CO_2. Imaging mode was Confocal 25 μm. Hoechst images were acquired with a

200 mW solid state 405 nm laser and 450/50 nm BP emission filter. Clover images were

acquired with a 150 mW solid state 488 nm laser and 525/50 nm BP emission filter. mRuby2

images were acquired with a 150 mW solid state 561 nm laser and 620/60 nm BP emission filter.

iRFP670 images were acquired with a 140 mW solid state 637 nm laser and 700/75 nm BP

emission filter. Z-series covering the full nucleus was acquired at 0.27 μm step size for 40x

objective or 0.13 μm step size for 63x objective. For time-lapse imaging, the Z-series was

acquired at 0.32 μm step size for 40x objective or 0.16-0.2 μm step size for 63x objective. Two-

color images were acquired every 15 seconds. Three-color images were acquired every 15 or 24

seconds. Images are a maximum intensity projection of Z-series. Images were processed in

Fusion software using ClearView-GPU deconvolution with the Robust (Iterative) algorithm (pre-

sharpening 0, 5 iterations, and de-noising filter size 0.1). Linear adjustments in maximum and

minimum levels were applied equally across the entire image to display only nuclear spot

signals, and not background nuclear fluorescence. Example images of our image processing

pipeline are shown in **Supplementary Fig. 2**.

Image analysis

To measure two targeted genomic loci distance in time-lapse images, Fiji image analysis

software was used[34]. Z-series acquired at 0.32 μm step size for 40x objective, and 0.16-0.2 μm

step size for 63x objective was used. If the nucleus drifted over time, the Correct 3D Drift plugin

was used[35]. For segmenting and tracking spots, the TrackMate plugin was used[36]. Blob

diameter was set at 2.0 μm. For each channel, threshold was set to include two spots with the

maximum intensity in the 3D volume of the nucleus. Simple LAP Tracker used 2 μm linking

max distance, 2 μm gap closing max distance, and 2 gap closing max frame gap. Analysis

produced "Spots in track statistics" file which was used to run python script to calculate 3D

distances between spots and generate 3D tracks. To determine and correct for potential chromatic

aberrations, Invitrogen Tetraspeck 0.1 μm Microspheres (ThermoFisher T7279) were imaged.

On the Dragonfly High Speed Confocal Platform 505 (Andor), Z-series were acquired for both

the Leica HCX PL APO 40x/1.10 W CORR objective and the Leica HC PL APO 63x/1.47NA

OIL CORR TIRF objective. TrackMate was then used to localize the positions of the

microspheres. iRFP670 and mRuby2 chromatic aberrations in each X,Y,Z dimension were

calculated relative to Clover, and corrected by offsetting the shifts in each X,Y,Z dimension as

follows:

Objective	Fluorescent Channel	X (μm)	Y (μm)	Z (μm)
40x	iRFP670	-0.035	-0.007	+0.691
40x	mRuby2	-0.053	-0.055	+0.319
63x	iRFP670	-0.001	-0.012	+0.147
63x	mRuby2	-0.018	-0.035	-0.028

DNA-FISH

Cells were cultured on 12 mm circle coverslip on 24-well plates and transfected with Casilio

components. At 48 hours, cells were fixed with 4% formaldehyde at room temperature for 10

min, briefly washed once with 1x PBS, permeabilized with 0.5% PBST for 20 min followed by

1x PBS wash, then dehydrated by a series of 75%, 85% and 100% ethanol for 2 min each. Let

dry. FISH probe (Cy5-MUC4: /5Cy5/CTTCCTGTCACCGAC, IDT custom DNA oligo) and

DNA hybridization buffer were prewarmed and mixed well, added onto slide, covered with cell-

attached coverslip, sealed with rubber cement. The sealed cells and probe were then heated on a

heat block at 80 °C for 2 min, incubated in a dark humidified chamber at 37 °C overnight. After

hybridization, the cell-attached coverslip was unsealed and washed in sequence with 2x SSC at

room temperature for 2 min, 10% formamide/2x SSC for 5 min at 45 °C, 2x SSC at room

temperature for 5 min, 0.2x SSC at 45 °C for 10 min twice, then counterstained with DAPI for 5

min and mounted with SlowFade™ Diamond Antifade Mountant (Invitrogen) for imaging.

Genome track and Hi-C maps

Genome tracks were visualized by UCSC Genome Browser[37]. Hi-C maps were generated

using FAN-C[38].

Acknowledgments

We thank the Jackson Laboratory Microscopy Service managed by Qianru Yu for microscope

usage, Erick Ratamero (Research IT Applications) for imaging analysis support, Jane Cha (JAX

Creative) for graphic design, and Research Program Development for help with manuscript

editing. This work is partially supported by internal funds provided by the Jackson Laboratory as

well as grants from the National Cancer Institute (P30CA034196), the National Human Genome

Research Institute (R01-HG009900), and the National Science Foundation (CCF-1955712).

Author contributions

J.J.Z. and A.W.C. conceived the study. P.A.C., J.J.Z, and A.W.C. designed the experiments.

P.A.C., M.D., N.J., A.T., J.J.Z., and A.W.C. performed the experiments. P.A.C. and M.D.

analyzed the data. P.A.C., A.T., J.J.Z., and A.W.C. wrote the manuscript.

Competing interests

A patent application has been filed on the invention.

Data availability

All data is available in the main text or the supplementary materials. Plasmids will be available

on Addgene. Raw images will be deposited on open repository.

Code availability

Not applicable.

Fig. 1. Schematic of Casilio. (a) Casilio extends CRISPR gRNA with binding sites for Pumilio

RNA binding protein-tethered effectors to allow multimerization, multiplexing and expansion of

functionalities. **(b)** The potential for Casilio to achieve multimodal operations simultaneously,

for example, demethylating histone (using LSD1 module), acetylating histone (CBP),

methylating DNA (DNMT3A), activating (p65HSF1) different targets, and at the same time

monitoring the localization of multiple locations in live cells.

Fig. 2. Validation of Casilio imaging specificity. (a) Schematic showing locations of Casilio

MUC4 gRNA (gMUC4) and Cy5-MUC4 FISH probes. **(b)** Co-labeling of *MUC4* locus by

Casilio (Green) and DNA-FISH (Cy5-MUC4, Red) in U2OS cells (n=8 nuclei; 28 spots;

Specificity=#Red-Green-overlap/#Green=96%). Scale bar, 5 μm **(c)** Co-labeling of Casilio

(Green) and DNA-FISH (Cy5-MUC4, Red) in ARPE-19 cells (n=8 nuclei; 16 spots;

Specificity=100%). Scale bar, 5 μm **(d)** Percentage of green ARPE-19 nuclei with the indicated

numbers of Casilio/gMUC4 spots in live cells without and with dCas9. Counts are indicated on

361 top of the columns. **(e)** Percentage of green HAP1 nuclei with the indicated numbers of

362 Casilio/gMUC4 spots in live cells. Counts are indicated on top of the columns.

Fig. 3. Casilio live-cell imaging of chromatin interactions and validation. (a) Casilio probes

for visualizing chromatin interactions mediated by cohesin, with locus A (5' anchor) labeled by

gRNA-15xPBSc/Clover-PUFc and locus B (3' anchor) labeled by gRNA-15xPBS9R/PUF9R-

iRFP670. The *MASP1-BCL6* loop with anchors 367 kb apart. **(b)** Representative time-lapse

images of the *MASP1-BCL6* loop anchors, locus A near *MASP1* (Clover, green, stealth

arrowhead) and locus B near *BCL6* (iRFP670, red, triangle arrowhead), in ARPE-19 cells. Time-

lapse image strips show two allelic Pair 1 and Pair 2 at the indicated time points. The whole

nucleus at time zero is shown on the left. See Supplementary Video 1. **(c)** Pairwise 3D distances

of fluorescent foci in (b) over time. See Supplementary Fig. 1 for all distance plots of imaged

nuclei (n=20 nuclei; 33 pairs). **(d)** Schematic of competitor experiment. A "cold" competitor

gRNA without Pumilio binding sites (PBS) for fluorescent protein recruitment binds to sequence

overlapping a test target. **(e)** Competitor experiments for validating *MASP1-BCL6*. Column plot

shows percentages of nuclei with the indicated numbers of Clover (green) spots (counts shown

on top of columns) in ARPE-19 cells co-transfected with Clover-PUFc/A-15xPBSc, PUF9R-

iRFP670/B-15xBPS9R and either a non-competing GAL4 gRNA (black columns) or a gRNA

competing with the corresponding A locus of the loop (patterned columns). Representative

images in the non-competed or locus A-competed samples (Clover, green spots and stealth

arrowheads; iRFP, red spots and triangle arrowheads) are shown to the right. Whole nucleus

scale bars, 5 μm. Time-lapse scale bars, 1 μm.

Fig. 4. Casilio visualization of RAD21-dependent and -independent interactions in the

presence and absence of RAD21. (a) Schematic of Casilio imaging of chromatin interactions

between *IER5L* promoter (*IER5L*-P) and a super enhancer (*IER5L*-SE) in HCT116/RAD21-

mAID cells. Hi-C map shows interactions in this region in HCT116/RAD21-mAID cells derived

from a previous study[27]. Upper right off-diagonal and lower left off-diagonal plots pairwise

interaction between genomic bins in untreated and auxin-treated cells, respectively. Dotted

squares mark the area of interest. **(b,c)** Representative images of *IER5L*-P and *IER5L*-SE labeled

by Clover (green, stealth arrowhead), and iRFP670 (red, triangle arrowhead), respectively, in

nuclei of HCT116/RAD21-mAID cells without (b, see Supplementary Video 2) or with auxin

treatment (c, see Supplementary Video 3). Scale bars, 5 μm. **(d)** Boxplot of pairwise spot

distances in untreated (n=1325 measurements of 33 pairs from 18 nuclei; mean=1.52 μm, x-

marked; median=1.21 μm, line inside box) and auxin-treated (n=1435 measurements of 35 pairs

from 20 nuclei; mean= 2.26 μm; median=2.09 μm) cells. ***Mann-Whitney test two-sided

p<0.00001. **(e)** Hi-C map showing the pairwise interaction between genomic bins encompassing

a RAD21-independent loop in untreated (upper right off-diagonal) and auxin-treated (lower left

off-diagonal) cells. Dotted squares mark the area of interest. **(f,g)** Representative images of the

RAD21-independent interaction labeled by Clover (green, stealth arrowhead), and iRFP670 (red,

triangle arrowhead), respectively, in nuclei of HCT116/RAD21-mAID cells without (f, see

Supplementary Video 4) or with auxin treatment (g, see Supplementary Video 5). Scale bars, 5

μm. **(h)** Boxplot of pairwise spot distances in untreated (n=1046 measurements of 27 pairs from

18 nuclei; mean=1.44 μm, x-marked; median=1.24 μm, line inside box) and auxin-treated

(n=921 measurements of 25 pairs from 20 nuclei; mean= 0.76 μm; median= 0.57 μm) cells.

***Mann-Whitney test two-sided *p*<0.00001.

Fig. 5. Three-color three-point live-cell imaging of chromatin loops. **(a)** Casilio probes for visualizing chromatin interactions of *IER5L* promoter and its super-enhancer, with locus P (promoter) labeled by gRNA-20xPBS$_{48107}$/Clover-PUF$_{48107}$, locus M (mid-point) labeled by gRNA-20xPBSc/iRFP670-PUFc, and locus SE (super-enhancer) labeled by gRNA-25xPBS9R/PUF9R-mRuby2. **(b,c)** Representative nuclei (nuclear boundaries outlined) and time-lapse images of *IER5L*-P-SE loop, with *IER5L* promoter, midpoint and super-enhancer labeled by Clover (green, stealth arrowhead), iRFP670 (red, triangle arrowhead), mRuby2 (magenta, open arrowhead), respectively, in HCT116 cells at the indicated time point. Scale bars 5 μm. See Supplementary Videos 6 and 7. See Supplementary Fig. 6 for distance plots of imaged nuclei (n=18 nuclei; 28 alleles).

References

1. Maass, P.G., A.R. Barutcu, and J.L. Rinn, *Interchromosomal interactions: A genomic love story of kissing chromosomes.* J Cell Biol, 2019. **218**(1): p. 27-38.
2. Mishra, A. and R.D. Hawkins, *Three-dimensional genome architecture and emerging technologies: looping in disease.* Genome Med, 2017. **9**(1): p. 87.
3. Krumm, A. and Z. Duan, *Understanding the 3D genome: Emerging impacts on human disease.* Semin Cell Dev Biol, 2019. **90**: p. 62-77.
4. Fullwood, M.J., et al., *An oestrogen-receptor-alpha-bound human chromatin interactome.* Nature, 2009. **462**(7269): p. 58-64.
5. Lieberman-Aiden, E., et al., *Comprehensive mapping of long-range interactions reveals folding principles of the human genome.* Science, 2009. **326**(5950): p. 289-93.
6. Lakadamyali, M. and M.P. Cosma, *Visualizing the genome in high resolution challenges our textbook understanding.* Nature Methods, 2020.
7. Robinett, C.C., et al., *In vivo localization of DNA sequences and visualization of large-scale chromatin organization using lac operator/repressor recognition.* J Cell Biol, 1996. **135**(6 Pt 2): p. 1685-700.
8. Alexander, J.M., et al., *Live-cell imaging reveals enhancer-dependent Sox2 transcription in the absence of enhancer proximity.* Elife, 2019. **8**.
9. Lindhout, B.I., et al., *Live cell imaging of repetitive DNA sequences via GFP-tagged polydactyl zinc finger proteins.* Nucleic Acids Res, 2007. **35**(16): p. e107.
10. Casas-Delucchi, C.S., et al., *Targeted manipulation of heterochromatin rescues MeCP2 Rett mutants and re-establishes higher order chromatin organization.* Nucleic Acids Res, 2012. **40**(22): p. e176.
11. Ma, H., P. Reyes-Gutierrez, and T. Pederson, *Visualization of repetitive DNA sequences in human chromosomes with transcription activator-like effectors.* Proc Natl Acad Sci U S A, 2013. **110**(52): p. 21048-53.
12. Ren, R., et al., *Visualization of aging-associated chromatin alterations with an engineered TALE system.* Cell Res, 2017. **27**(4): p. 483-504.
13. Sander, J.D. and J.K. Joung, *CRISPR-Cas systems for editing, regulating and targeting genomes.* Nat Biotechnol, 2014. **32**(4): p. 347-55.
14. Chen, B., et al., *Dynamic imaging of genomic loci in living human cells by an optimized CRISPR/Cas system.* Cell, 2013. **155**(7): p. 1479-91.
15. Tanenbaum, M.E., et al., *A protein-tagging system for signal amplification in gene expression and fluorescence imaging.* Cell, 2014. **159**(3): p. 635-46.
16. Ma, H., et al., *Multiplexed labeling of genomic loci with dCas9 and engineered sgRNAs using CRISPRainbow.* Nat Biotechnol, 2016. **34**(5): p. 528-30.
17. Qin, P., et al., *Live cell imaging of low- and non-repetitive chromosome loci using CRISPR-Cas9.* Nat Commun, 2017. **8**: p. 14725.
18. Ma, H., et al., *CRISPR-Sirius: RNA scaffolds for signal amplification in genome imaging.* Nat Methods, 2018. **15**(11): p. 928-931.
19. Maass, P.G., et al., *Spatiotemporal allele organization by allele-specific CRISPR live-cell imaging (SNP-CLING).* Nat Struct Mol Biol, 2018. **25**(2): p. 176-184.
20. Wang, H., et al., *CRISPR-mediated live imaging of genome editing and transcription.* Science, 2019. **365**(6459): p. 1301-1305.

21. Cheng, A.W., et al., *Casilio: a versatile CRISPR-Cas9-Pumilio hybrid for gene*
regulation and genomic labeling. Cell Res, 2016. **26**(2): p. 254-7.
22. Chen, Y. and G. Varani, *Engineering RNA-binding proteins for biology.* Febs j, 2013.
280(16): p. 3734-54.
23. Wang, Y., et al., *Engineering splicing factors with designed specificities.* Nat Methods,
2009. **6**(11): p. 825-30.
24. Taghbalout, A., et al., *Enhanced CRISPR-based DNA demethylation by Casilio-ME-*
mediated RNA-guided coupling of methylcytosine oxidation and DNA repair pathways.
Nat Commun, 2019. **10**(1): p. 4296.
25. Gilbert, L.A., et al., *CRISPR-mediated modular RNA-guided regulation of transcription*
in eukaryotes. Cell, 2013. **154**(2): p. 442-51.
26. Natsume, T., et al., *Rapid Protein Depletion in Human Cells by Auxin-Inducible Degron*
Tagging with Short Homology Donors. Cell Rep, 2016. **15**(1): p. 210-218.
27. Rao, S.S.P., et al., *Cohesin Loss Eliminates All Loop Domains.* Cell, 2017. **171**(2): p.
305-320.e24.
28. Li, G., et al., *ChIA-PET2: a versatile and flexible pipeline for ChIA-PET data analysis.*
Nucleic Acids Res, 2017. **45**(1): p. e4.
29. Zhu, J.J. and A.W. Cheng, *JACKIE: Fast enumeration of single- and multi-copy CRISPR*
binding sites. bioRxiv, 2020: p. 2020.02.27.968933.
30. Bae, S., J. Park, and J.S. Kim, *Cas-OFFinder: a fast and versatile algorithm that*
searches for potential off-target sites of Cas9 RNA-guided endonucleases.
Bioinformatics, 2014. **30**(10): p. 1473-5.
31. Shcherbakova, D.M. and V.V. Verkhusha, *Near-infrared fluorescent proteins for*
multicolor in vivo imaging. Nat Methods, 2013. **10**(8): p. 751-4.
32. Lam, A.J., et al., *Improving FRET dynamic range with bright green and red fluorescent*
proteins. Nat Methods, 2012. **9**(10): p. 1005-12.
33. Cong, L., et al., *Multiplex genome engineering using CRISPR/Cas systems.* Science,
2013. **339**(6121): p. 819-23.
34. Schindelin, J., et al., *Fiji: an open-source platform for biological-image analysis.* Nat
Methods, 2012. **9**(7): p. 676-82.
35. Parslow, A., A. Cardona, and R.J. Bryson-Richardson, *Sample drift correction following*
4D confocal time-lapse imaging. J Vis Exp, 2014(86).
36. Tinevez, J.Y., et al., *TrackMate: An open and extensible platform for single-particle*
tracking. Methods, 2017. **115**: p. 80-90.
37. Kent, W.J., et al., *The human genome browser at UCSC.* Genome Res, 2002. **12**(6): p.
996-1006.
38. Kruse, K., C.B. Hug, and J.M. Vaquerizas, *FAN-C: a feature-rich framework for the*
analysis and visualisation of chromosome conformation capture data. Genome Biol,
2020. **21**(1): p. 303.