Consistent behavioural syndromes across seasons in an invasive freshwater fish

Lukas J[1,2, #], Kalinkat G[1], Miesen FW[3], Landgraf T[4], Krause J[1,2], Bierbach D[1,2]

1. Leibniz-Institute of Freshwater Ecology and Inland Fisheries, Department of Biology and Ecology of Fishes, Müggelseedamm 310, D-12587 Berlin, Germany
2. Humboldt University of Berlin, Faculty of Life Sciences, Invalidenstraße 42, D-10115 Berlin, Germany
3. Zoological Research Museum Alexander Koenig, Adenauerallee 160, D-53113 Bonn, Germany
4. Free University of Berlin, Institute of Computer Science, Dahlem Center for Machine Learning and Robotics, Arnimallee 7, D-14195 Berlin, Germany

\# corresponding author: contact@julianelukas.com

Abstract

The linkage between behavioural types and dispersal tendency has been suggested to be a widespread phenomenon and understanding its mechanisms has become a pressing issue in light of global change and biological invasions. Here, we investigate whether individuals who colonize new habitats exhibit a certain set of behavioural types that differs from those remaining in the source population. We focussed on a feral population of guppies (*Poecilia reticulata*) that is assumed to undergo a yearly (re)colonization process. Guppies are among the most widespread invasive species in the world, but in temperate regions these fish can only survive in thermally altered freshwaters due to their tropical origin. The investigated population has sustained in a thermally-altered stream in Germany for over 50 years, where they find year-round suitable water temperatures around a warm-water influx. However, in warm seasons, peripheral parts of the warm-water flume become thermally accessible and may be colonized. We sampled fish from the source population and from a winter-abandoned downstream site in March, June and August. Fish were tested for boldness, sociability and activity involving open field tests as well as interactions with a biomimetic robot as social partner. Guppies differed consistently among each other in all three traits. Behavioural trait expression in the source population differed across seasons, however, we could not detect differences between source and downstream populations. Instead, all sampled populations showed a remarkably stable behavioural syndrome between boldness and activity despite strong changes in water temperatures across seasons. We conclude that random drift (as opposed to a personality-biased dispersal) is a more likely dispersal mode for guppies, at least in the investigated stream system. In the face of highly fluctuating environments, guppies seem to be extremely effective in keeping their behavioural expressions constant, which could help explain their successful invasion and adaptation to disturbed habitats.

Keywords: animal personality, dispersal, range expansion, invasive species, thermally altered freshwaters, guppy

Introduction

Consistent behavioural differences among individuals (e.g. behavioural types or animal personality; Réale et al. 2007) seem to be a ubiquitous biological feature (Bell et al. 2009, Bierbach et al. 2017) that was proposed to have substantial ecological and evolutionary importance (Sih et al. 2012, Wolf and Weissing 2012). For one, behavioural types are thought to be relevant in determining success (Chapple et al. 2012, Carere and Gherardi 2013, Canestrelli et al. 2015) and impact of biological invasions (Cote et al. 2010a, Fogarty et al. 2011, Juette et al. 2014), which is becoming an ever more pressing issue in the light of global change and ongoing translocations of species among habitats (Seebens et al. 2017, 2018).

It is assumed that individuals, which invade novel habitats, are a non-random subset of their source populations (Shine et al. 2011, Sih et al. 2012, Juette et al. 2014, Canestrelli et al. 2015, Spiegel et al. 2017). These invaders may share certain phenotypic and life-history characteristics that are associated with their behavioural types (reviewed in Juette et al. 2014). Behavioural types of invading individuals may determine whether they are transported (by humans) or move (by themselves) into new environments (first invasion stage; after Lockwood et al. 2011), whether they can survive and establish a non-native population (second stage), and whether they may spread to new habitats thereafter (third stage). Behavioural types shown to be relevant in this regard are boldness (Cote et al. 2011, Myles-Gonzalez et al. 2015), aggression (Duckworth and Badyaev 2007, Hudina et al. 2015), activity (Brown and Irving 2014, Thorlacius et al. 2015) as well as sociability (Cote et al. 2010b, 2011, Rasmussen and Belk 2012). Further, it was predicted that individuals carrying correlated suits (i.e., behavioral syndromes, Sih et al. 2012) of bold, aggressive, active and asocial types may predominantly be found at the invasion front (Duckworth and Badyaev 2007, Cote et al. 2010b, 2011, Juette et al. 2014). However, such spatially and/or temporally assorted compositions of behavioural types among populations along invasion gradients have only rarely been investigated under natural conditions in the wild.

The difficulty to study invasion phenomena in the wild is their random and unpredictable occurrence (Lockwood et al. 2011), which makes it nearly impossible to follow active invasion

fronts. In the rare cases where range expansions could be tracked in real time (e.g., western

bluebirds: Duckworth and Badyaev 2007; Afican jewelfish: Lopez et al. 2012; round goby:

Thorlacius et al. 2015; signal crayfish: Hudina et al. 2015; cane toad: Gruber et al. 2017), the

invasion origin is often unknown or unclear due to multiple introduction events. Thus, we need to

identify study systems that allow us to investigate dispersal across sensible temporal and spatial

scales. Thermally altered freshwater systems (TAFs) in the temperate zones provide fruitful

conditions to study this phenomenon. Here, natural and/or anthropogenic warm water influxes

create unique temperature refuges that allow non-native species (especially those of tropical

origin) to establish populations (Langford 1990, Jourdan et al. 2014). While ambient water

temperatures drop in winter (e.g., less than +1°C in the river Rhine; Jourdan et al. 2014), areas

close to constant warm water influx will remain suitable for non-native tropical species to survive.

Once peripheral areas become thermally suitable again in warmer seasons, the winter-abandoned

areas are assumed to be (re)colonized. Such a scenario enables us to investigate the role of

behavioural types during *in situ* dispersal of non-native species from a single, well-determined

source into new (i.e. winter-abandoned) habitats.

Under a personality-biased dispersal hypothesis, where individuals with a certain set of

personality traits are assumed to leave their source population to colonize new habitats, we

predicted that *(i)* the source population's average personality traits should change over the course

of the year due to the drain of the emigrating individuals, *(ii)* newly founded populations should

differ in their average personality traits compared to their source populations, and *(iii)* behavioural

variation should decrease both in the source as well as in the newly founded populations due to

the non-random migration patterns. In addition, *(iv)* population-wide correlations among personality

traits (behavioural syndromes) may emerge or disappear because of assumed non-random

migration patterns.

In addition to these predictions arising from non-random migration patterns, behavioural

types may change due to the influence of seasonal fluctuations in abiotic (e.g., temperature;

Spiegel et al. 2015) and/or biotic (e.g., resource availability and predator pressure; Barbosa et al.

2018) factors common in temperate climates. If so, one would predict that population averages,

behavioural variation and syndromes should change over the course of the year (similar to *i* and *iii*

and *iv*) but in both established and newly founded populations to a similar extent (opposing

prediction *ii*), especially when those populations face similar external conditions.

In Germany, tropical fishes have established populations in several artificially heated

stream systems (Lukas et al. 2017a), the most investigated being the Gillbach near Cologne (see

Fig. 1a; Jourdan et al. 2014, Lukas et al. 2017b, Kempkes et al. 2018). To estimate the seasonal

fluctuations within the sampled range of the stream, we continuously recorded water temperatures

with submerged data-loggers. We sampled feral guppies (*Poecilia reticulata*) from the Gillbach

during different seasons (temporal scale: spring, early summer, late summer) and, if possible, at

different sites (spatial scale: core area of warm water influx versus more peripheral downstream

area; see Fig. 1a) and repeatedly assessed behavioural types in terms of boldness, sociability and

activity in a laboratory setting.

Material& Methods

Study site & Sampling

The Gillbach (Rhine catchment, Germany) has been under the influence of thermal pollution and anthropogenic modifications for over six decades. Its headwaters are exclusively fed by the warm water discharge of the brown coal power plant Niederaußem (Fig. 1a). This creates a temperature regime, which is highly dependent on the volume and temperature of the plant's water discharge. Generally exhibiting above ambient water temperatures (Fig. 1b), the Gillbach harbours several established populations of non-native fish, invertebrate and plant species, many of which are of tropical or subtropical origin (see Supplementary material Appendix 1, Table A2 for a list of native and non-native fish species found in the Gillbach).

Here, feral guppies (*Poecilia reticulata*) are highly abundant and as successful global invaders (Deacon et al. 2011), they are well suited to investigate the expression of behavioural traits across time and space. Samplings were conducted at two sites (~300m transects, see Fig. 1a) in March, June and August 2016. To target guppies, we collected specimen by seine and dip netting (2.5 mm mesh width) until we reached sufficient numbers for behavioural testing (i.e., 35-45 adult individuals). Captured fish were transferred into well-aerated coolers and kept separately by sampling site.

Water temperatures at sites 1 and 2 were monitored using HOBO data loggers (Onset Computer Corporation, Bourne, MA, USA) for a period of 16 months (Fig. 1b). For the period of sampling and behavioural testing (March – August 2016), the Gillbach experienced a thermal gradient of ~2°C/km (mean ± SD, site 1: 25.53°C ± 3.21°C; site 2: 23.49°C ± 3.4°C; Fig. 1b). We expect this trend to be even more pronounced in colder months, thus decreasing the likelihood of survival for guppies outside the core area during winter (see also Jourdan et al. 2014). Interestingly, the temperature range at both sites was similar (min – max, site 1: 8.38°C – 43.11°C; site 2: 10.46°C – 39.5°C), with two cases of temperature drops below the guppies' lower critical minimum temperature (approx. 12°C according to Jourdan et al. 2014).

Figure 1: The Gillbach stream and its altered temperature regime. a) Warm water discharge from a power plant feeds into the Gillbach (map created using © 2019 Google Earth with data from 2009 GeoBasis-DE/BKG) and generates a thermal gradient above ambient water temperatures that is preserved for several kilometres (site 1 – site 2). **b)** 4-hourly water temperature* with error bars indicating monthly means ± SD for each site (site 1 – Gillbach source: red; site 2 – one kilometre downstream: orange). Dotted lines correspond to the three sampling occasions. * Note that due to a loss of loggers, continuous monitoring was interrupted for certain periods (site 1: no data from 11.09.2015 – 24.12.2015; site 2: no data from 24.12.2015 – 19.03.2016 nor from 29.08.2016 – 31.12.2016).

Behavioural assessment

Fish collected for behavioural testing were transferred to the laboratory at Humboldt University of

Berlin, Germany. Fish were kept in one holding tank (270 litres), but were separated into mesh-

divided compartments by collection time and site ('populations' hereafter). Each compartment (24

cm × 58 cm × 40 cm L×W×H) contained artificial java moss and natural gravel. Females and males

were housed communally. Water was maintained at 26°C and kept under continuous aeration and

filtration. Fish were held under diurnal lighting (12:12h light:dark cycle) and fed commercial flake

food (TetraMin® Tetra GmbH) twice a day. Feeding stopped 12 hours before experiments and

recommenced afterwards. To acclimate the fish to laboratory conditions and reduce potential

short-term effects of the different sampling habitats on behavioural trait expression, no behavioural

testing was conducted for at least 30 days (see Fig. 2a for detailed experimental design). One

week prior to testing, adults of suitable size (SL>15 mm) were marked using visible implant

elastomer tags (VIE; Northwest Marine Technology Ltd, Shaw Island, WA, USA) to allow for

individual identification throughout the experiment. To be mindful of capture biases towards a

specific behavioural type, we used all fish within a compartment for testing.

We tested a total of 159 guppies (95 females, 64 males; for population-specific sample

sizes see Fig. 2a). Test fish were introduced to an open field test arena (88 x 88 cm, 75 mm water

depth) filled with aged tap water kept at 26°C. To reduce disturbing stimuli, the arena was

enclosed in black, opaque plastic and illuminated with artificial light reproducing the daylight

spectrum. Behavioural observations were recorded via a camera (Ximea 4K USB 3.1 camera)

mounted above the arena. Position information was extracted using EthoVision 10 XT software

(Noldus Information Technology, Wageningen, Netherlands).

Figure 2: Behavioural testing with Robofish setup. a) Test fish were collected three times and, when

possible, at two sites (site 1 and 2, see Fig. 1a). After an initial 30-day acclimation period, fish were tested a

total of three times (trial I – III, sample sizes indicate tested individuals, for total catch numbers see

Supplementary material Appendix 1). **b-d)** During trials, the Robofish replica was moved remotely through

the test tank (c). Once the focal individual had emerged from its shelter, the biomimetic Robofish initiated its

predefined path (b) and the fish's swimming trajectory was recorded (d).

(1) Boldness: Boldness was measured as latency to emerge from a shelter with a low latency time

indicating high levels of boldness. Fish were introduced into a grey opaque shelter box (Ø 100

mm; top left of each image in Fig. 2b and d) and allowed to acclimate. After one minute, a

sponge was removed from the entrance (40mm x 25mm), allowing the fish to emerge and

explore the test arena. A fish was scored as having emerged when its full body was visible

outside of the shelter. If fish did not emerge after five minutes, the lid of the shelter box was

removed. Fish that failed to emerge within eight minutes were given the maximum score of 480

seconds and the box was removed entirely.

(2) Sociability: A three-dimensional fish replica was stationed 1 cm away from the shelter entrance.

The replica, which resembled a large-sized guppy female (SL=30mm; left fish in Fig. 2c), could

be moved by a robot via a magnetic base (see Landgraf et al. 2016, Bierbach et al. 2018 for

details). Upon the fish's emergence (or the removal of the shelter box), the biomimetic robot

started to move through the arena in a standardized sequence (Fig. 2b) imitating a conspecific.

We calculated the mean distance the fish kept to the biomimetic robot as a proxy of sociability

with lower values indicating higher levels of sociability. Consequently, fish that followed the

robot closer were assumed to be more sociable (see Bierbach et al. 2018).

(3) Activity: After completion of the Robofish's predefined path, the replica was removed from the

arena by hand and once the fish resumed normal swimming behaviour, it was allowed to

explore the arena. As a measure of activity, we calculated the mean velocity with which the fish

explored the open arena during 3 minutes.

After the first behavioural assay, we recorded the body length of each individual. Fish were

transferred back to their respective compartment and allowed 72 hours resting before being

retested following the same protocol. Fish were tested a total of three times in randomized order.

All reported experiments comply with the current German law approved by LaGeSo (G0117/16 to

Dr. D. Bierbach).

Statistical analysis

Datasets with sample sizes less than 10 were excluded from all further analyses (i.e. site 2 in

March and June; see Fig. 2a).

To test for differences in average trait expression over a temporal and/or spatial scale, we

used linear mixed models (LMM, *GENLINMIXED* procedure; SPSS v. 25.0, IBM Corp., Armonk,

NY, USA) with each behavioural trait (boldness, sociability, activity) as a separate response

variable. 'Boldness' and 'sociability' were log10-transformed to fulfil normality assumptions. All

models included sample population as the fixed effect of interest (four levels: March – site 1, June

– site 1, August – site 1, August – site 2). Behaviour in poeciliids is often size- and/or sex-

dependent (Brown and Braithwaite 2004, Harris et al. 2010), thus we included standard length (SL)

as a covariate and 'sex' as an additional fixed factor. To account for the repeated testing of

individuals, we further included 'trial' as a fixed factor and 'fish ID' as a random factor. Post-hoc

LSD tests were employed in case fixed factors were found to have a significant effect. For the

covariate 'body size', we also performed Pearson's correlations to better evaluate the relationship.

(Co)Variance structure was nested within 'population', which enabled us to calculate the

total behavioural variance as well as within- and among-individual variance components in each

population. Based on this, we calculated behavioural repeatability for each of the four populations.

We assumed that repeatability and/or variance estimates differed among populations when 95%

confidence intervals (CIs) were not overlapping (see Nakagawa and Schielzeth 2010).

To evaluate the existence of behavioural syndromes (i.e. correlation among behavioural

traits) within our populations, we used linear regression models to test for significant correlation

between each trait pair. In order to minimize the effects of mean differences in behavioural traits on

trait correlation, we mean-centred each behavioural trait within 'trial' and 'population' (z-

transformation). Each model included one trait as the dependent variable and another as

covariate. We further compared whether trait correlation differed among populations (i.e.,

difference in slopes termed 'covariate nested within population'). Visualizations were done with

GraphPad Prism (version 8.01, GraphPad Software Inc., La Jolla, CA, USA).

Results

Differences in average trait expression among populations

Expression of sociability and activity differed between populations, while boldness did not ('population' effect in Table 1; Fig. 3a).

On a temporal scale, fish sampled at site 1 in August showed higher levels of sociability (EMMs ± SD, August – site 1: 1.18 ± 0.03) compared to March and June populations (March – site 1: 1.3 ± 0.03, June – site 1: 1.27 ± 0.03; post-hoc, August site 1 vs. March – site 1: $p = 0.003$, August site 1 vs. June – site 1: $p = 0.025$). Similarly, fish sampled at site 1 in August were more active (August – site 1: 7.81 ± 0.33) than populations sampled in March and June (March – site 1: 6.38 ± 0.33, June – site 1: 6.03 ± 0.27; post-hoc, August site 1 vs. March – site 1: $p = 0.002$, August site 1 vs. June – site 1: $p < 0.001$).

On a spatial scale, we found no evidence that fish sampled in August at site 1 and site 2 differed in boldness (1.4 ± 0.08; post-hoc, August site 1 vs. August site 2: $p = 0.76$), sociability (1.18 ± 0.04; post-hoc, August site 1 vs. August site 2: $p = 0.92$), nor activity (7.74 ± 0.31; post-hoc, August site 1 vs. August site 2: $p = 0.89$).

Boldness and activity showed a size-dependency (covariate 'body size'; Table 1, Fig. 3b) with smaller individuals being bolder (Pearson's $r = 0.32$; $p < 0.001$) and more active (Pearson's $r = -0.35$; $p < 0.001$) than larger ones. We found no such relation between body size and sociability (Pearson's $r = 0.002$; $p = 0.98$). While sex is correlated with size in the sexually dimorphic guppy, sex did not affect any of the measured behavioural traits (non-sig. 'sex' effect; Table 1). We further observed a habituation to the repeated testing during sociability and activity assays (sig. 'trial' effect; Table 1). Fish were considerably less social (trial 1: 1.03 ± 0.21, trial 2: 1.27 ± 0.21, trial 3: 1.4 ± 0.21) and less active (trial 1: 8.09 ± 0.21, trial 2: 6.93 ± 0.21, trial 3: 5.96 ± 0.21) with each consecutive measurement.

246 **Table 1: Results from linear mixed models**. For each behavioural trait, the LMM included 'population',

247 'trial' and 'sex' as fixed effects and 'body size' (SL in mm) as covariate. Fish ID was used as random factor.

248 Significant effects ($p < 0.05$) are indicated in bold.

Effect	Boldness ($\log_{10}$)		Sociability ($\log_{10}$)		Activity	
	$F_{df1,df2}$	p	$F_{df1,df2}$	p	$F_{df1,df2}$	p
population	$F_{3,469} = 0.57$	0.634	$F_{3,469} = 3.8$	**0.01**	$F_{3,469} = 8.44$	**< 0.001**
trial	$F_{2,469} = 1.72$	0.18	$F_{2,469} = 135.9$	**< 0.001**	$F_{2,469} = 42.32$	**< 0.001**
sex	$F_{1,469} = 0.06$	0.81	$F_{1,469} = 2.28$	0.132	$F_{1,469} = 0.24$	0.622
body size (SL)	$F_{1,469} = 8.43$	**0.004**	$F_{1,469} = 0.02$	0.884	$F_{1,469} = 3.58$	0.059

Figure 3: Average personality traits across temporal and spatial scale. *(a)* Median trait scores (and

IQRs) for boldness* (i.e., time to emerge from shelter), activity (i.e., velocity during open field exploration)

and sociability* (i.e., distance to biomimetic Robofish). Different letters depict significant differences based

on post-hoc LSD tests after global linear mixed models ($p < 0.05$). *(b)* Mean traits scores (± SD) in relation to

body size (standard length; n=159). To visualize the relationships, we show linear regressions lines (red)

calculated from individual means (see main text for significance levels and Pearson's r). * Note that low

scores of boldness and sociability indicate high trait expression.

Among- and within-individual variation in behavioural traits

Fish within each sampled population differed consistently in their expression of boldness,

sociability and activity across trials (sig. repeatability estimates; Table 2). Only fish sampled in

August at site 2 had a significantly higher repeatability estimate for sociability than fish from the

March sampling at site 1 (non-overlapping CIs; Table 2). There was no significant difference

among populations in estimated overall behavioural variances in any of the examined traits (see

'total variation' in Table 2; for within- and among-individual variance estimates see Supplementary

material Appendix 1, Table A2).

Table 2: Repeatability estimates and total behavioural variance estimates. *p*-values are derived from

Wald's *z* statistics (see Supplementary material Appendix 2, Table A3 for within- and among-individual

variation estimates). Significant differences among populations can be assumed when 95%CI do not

overlap.

Popula- tion	Boldness (log$_{10}$)			Sociability (log$_{10}$)			Activity		
	Repeat- ability [95% CI]	*p*	Total Variation [95% CI]	Repeat- ability [95% CI]	*p*	Total Variation [95% CI]	Repeat- ability [95% CI]	*p*	Total Variation [95% CI]
March site 1	0.49 [0.40–0.57]	**0.004**	0.24 [0.15–0.4]	0.26 [0.15–0.4]	0.05	0.06 [0.04–0.1]	0.27 [0.17–0.41]	**0.042**	6.86 [4.3–11.87]
June site 1	0.40 [0.31–0.5]	**0.005**	0.31 [0.2–0.5]	0.32 [0.23–0.43]	**0.011**	0.07 [0.05–0.11]	0.30 [0.21–0.42]	**0.015**	5.75 [3.74-9.33]
August site 1	0.54 [0.47–0.6]	**0.001**	0.34 [0.22–0.54]	0.34 [0.25–0.45]	**0.009**	0.06 [0.04–0.09]	0.38 [0.29–0.47]	**0.005**	7.15 [4.61-11.48]
August site 2	0.53 [0.46–0.6]	**0.001**	0.3 [0.19–0.48]	0.49 [0.42–0.57]	**0.001**	0.08 [0.05–0.13]	0.36 [0.27–0.46]	**0.007**	5.92 [3.81-9.57]

Behavioural syndrome structure

Boldness and activity were negatively correlated in all four sampling populations (sig. correlation

between boldness and activity: $F_{1,157}$ = 52.23; *p* < 0.001; no sig. difference in slopes among

populations: $F_{3,151}$ = 0.45; *p* = 0.715). Fish with high average activity levels also left the shelter

quicker, confirming that activity and boldness are part of a larger behavioural syndrome (Fig. 4).

There was no evidence for correlations between sociability and boldness ($F_{1,157}$ = 0.026; *p* =

0.872; no sig. difference in slopes among populations: $F_{3,151}$ = 1.8; *p* = 0.149), nor between

sociability and activity ($F_{1,157}$ = 0.062; *p* = 0.803; no sig. difference in slopes among populations:

$F_{3,151} = 0.816$; $p = 0.487$). Thus, sociability measured as distance kept to the moving Robofish does

not seem to be part of a behavioural syndrome with neither boldness nor activity.

Figure 4: Correlations between behavioural traits across temporal and spatial scale. Mean trait

scores* (±SD) of individual test fish are shown for each population. Linear regressions (red line and

Pearson's *r*) visualize the relationships between traits (see main text for detailed analysis of trait correlation;

n = 159). * Note that for boldness and sociability lower values indicate a high trait score.

Discussion

Our results document the annual (re)colonization process of an artificially heated stream by a non-native fish in temperate Germany. As guppies dispersed into winter-abandoned areas in warmer months, we expected this process to be driven by dispersal-enhancing behavioural traits. Contrary to expectations, we found no evidence that the source and the newly-founded population differed in terms of boldness, sociability and activity. Likewise, we did not detect a decrease in behavioural variation (except sociability in March and August) indicative of a "thinning out" of disperser's behavioural types. Instead, we found a behavioural syndrome among activity and boldness that was consistently detectable throughout the year and across sampled populations. However, average personality traits in the source population changed over the course of 6 months with August-sampled individuals being more active and more sociable than those sampled earlier in the year, while boldness scores remained unchanged. To sum up, we found no strong evidence for a personality-biased dispersal in this feral population of *Poecilia reticulata*. Instead, the detected seasonal differences in average trait expression (sociability, activity) are likely due to changed environmental factors such as temperature, predator abundance and/or resource availability.

The observed changes in behavioural traits across seasons but not between sites were likely caused by changes in environmental conditions experienced by both populations rather than by a drain of individuals carrying certain behavioural types from source to downstream populations. This assumption is congruent with the Gillbach's temperature profile that clearly shows strong seasonal variations within but not across sites. The observed seasonality in temperature in combination with a variation in day length, food quality/quantity and predation are likely to be important modifiers of population-wide behaviours (Biro et al. 2010, Eccard and Herde 2013, Spiegel et al. 2015, Uchida et al. 2016, Dhellemmes et al. 2020). There is further evidence that prior acclimation could, in fact, alter offspring phenotypes (Beitinger et al. 2000, Seebacher et al. 2014). This might explain the increased activity exhibited by fish in August, as higher developmental temperatures lead to more active phenotypes (Seebacher et al. 2014). Out of the studies finding personality-biased compositions along a dispersal front, only two noted context-

dependency: spatial sorting of behavioural types did not emerge under predation (Cote et al. 2013)

and was lost as the population aged (Thorlacius et al. 2015). Our findings now highlight

heterogeneity across seasons, however, teasing apart the effects of abiotic and biotic fluctuations

in resources and predation will require an experimental approach.

Besides seasonal differences in average behavioural traits, we found stable behavioural

syndromes among sampled populations that involved a strong positive correlation between

boldness and activity. This is consistent with several studies, where this boldness-activity

syndrome has been documented in various taxa (van Oers et al. 2004, Pintor et al. 2008, Wilson

and Godin 2009, Cote et al. 2010b, Eccard and Herde 2013, Muraco et al. 2014), including

guppies (Brown and Irving 2014). Natural selection through predation favours the evolution of

behavioural correlations (Dingemanse et al. 2007, Harris et al. 2010, Dhellemmes et al. 2020). A

positive boldness-activity correlation might thus represent an optimal trait combination in the fact of

predation pressure experienced by both populations. However, the lack of differences in the level

of (individual) behavioural variation among populations might be a result of a random downstream

drift and a low overall behavioural plasticity in this population. This can be a consequence of the

assumed bottleneck origin of this feral population that most likely stems from an unknown number

of released domesticated guppies (see Gertzen et al. 2008 for estimates of propagule pressure for

popular aquarium species). Investigations involving other populations of the guppy and more

distantly related species are highly recommended, especially as several environmental factors are

predicted to impact behavioural variation, especially in fish (e.g., Harris et al. 2010, Laskowski et

al. 2016, Barbosa et al. 2018). Alternatively, guppies and especially those from the Gillbach that

experience highly fluctuating temperature regimes may have the ability to adjust their behavioural

expressions within very short times, e.g., within the 30-days acclimatisation period in the laboratory

used here. It is known that guppies have a remarkable capacity for thermal adaptation (Chung

2001, Jourdan et al. 2014) and have successfully established populations in environments with

very different thermal regimes compared to their native habitats (Deacon et al. 2011). Testing of

individuals after different acclimation periods could be a possibility to investigate this issue further.

Theoretical models (Cote et al. 2010a, Spiegel et al. 2017) as well as several studies

across taxa (Duckworth and Badyaev 2007, Pintor et al. 2009, Cote et al. 2010b, Myles-Gonzalez

et al. 2015, Coates et al. 2019) provide evidence for the prominent role of individual differences in

behaviour during dispersal and invasion processes (Bowler and Benton 2005, Chapple et al. 2012,

Canestrelli et al. 2015). Nevertheless, the picture might be more complicated: our current study

exemplifies that population-wide behavioural analysis can fail to provide evidence for

behaviourally-mediated migration patterns and future analysis may focus on tracking individuals

along their travel routes across seasons and locations. Furthermore, analysis may incorporate the

invasion history, environmental factors or the species-specific genetic backgrounds which all can

affect behavioural variation in wild populations (Sih and Bell 2008, Bergmüller and Taborsky 2010,

Stamps and Groothuis 2010).

References

Barbosa, M. et al. 2018. Individual variation in reproductive behaviour is linked to temporal heterogeneity in
predation risk. - Proceedings of the Royal Society B: Biological Sciences 285: 20171499.

Beitinger, T. L. et al. 2000. Temperature Tolerances of North American Freshwater Fishes Exposed to
Dynamic Changes in Temperature. - Environ Biol Fish 58: 40.

Bell, A. M. et al. 2009. The repeatability of behaviour: a meta-analysis. - Anim Behav 77: 771–783.

Bergmüller, R. and Taborsky, M. 2010. Animal personality due to social niche specialisation.

Bierbach, D. et al. 2017. Behavioural individuality in clonal fish arises despite near-identical rearing
conditions. - Nat Comm 8: 15361.

Bierbach, D. et al. 2018. Using a robotic fish to investigate individual differences in social responsiveness in
the guppy. - Roy Soc Open Sci 5: 181026.

Biro, P. A. et al. 2010. Small within-day increases in temperature affects boldness and alters personality in
coral reef fish. - Proc R Soc B: 71–77.

Bowler, D. E. and Benton, T. G. 2005. Causes and consequences of animal dispersal strategies: relating
individual behaviour to spatial dynamics. - Biol Rev Camb Philos Soc 80: 205–225.

Brown, C. and Braithwaite, V. A. 2004. Size matters: A test of boldness in eight populations of the poeciliid
Brachyraphis episcopi. - Anim Behav 68: 1325–1329.

Brown, C. and Irving, E. 2014. Individual personality traits influence group exploration in a feral guppy
population. - Behav Ecol 25: 95–101.

Canestrelli, D. et al. 2015. Bolder Takes All? The Behavioral Dimension of Biogeography. - Trends Ecol Evol
31: 35–43.

Carere, C. and Gherardi, F. 2013. Animal personalities matter for biological invasions. - Elsevier Ltd.

Chapple, D. G. et al. 2012. Can behavioral and personality traits influence the success of unintentional
species introductions? - Trends Ecol Evol 27: 57–64.

Chung, K. S. 2001. Critical thermal maxima and acclimation rate of the tropical guppy *Poecilla reticulata*. -
Hydrobiologia 462: 5.

Coates, W. D. et al. 2019. Dispersal decisions and personality in a freshwater fish. - Anim Behav 157: 209–
218.

Cote, J. et al. 2010a. Personality-dependent dispersal: characterization, ontogeny and consequences for
spatially structured populations. - Phil Trans R Soc B 365: 4065–4076.

Cote, J. et al. 2010b. Personality traits and dispersal tendency in the invasive mosquitofish (*Gambusia
affinis*). - Proc R Soc B 277: 1571–1579.

Cote, J. et al. 2011. Personality-dependent dispersal in the invasive mosquitofish: group composition
matters. - Proc R Soc B 278: 1670–1678.

Cote, J. et al. 2013. Personality-dependent dispersal cancelled under predation risk. - Proc R Soc B 280:
20132349.

Deacon, A. E. et al. 2011. How Reproductive Ecology Contributes to the Spread of a Globally Invasive Fish.
- PLOS ONE 6: e24416.

Dhellemmes, F. et al. 2020. Comparing behavioural syndromes across time and ecological conditions in a free-ranging predator. - Animal Behaviour 162: 23–33.

Dingemanse, N. J. et al. 2007. Behavioural syndromes differ predictably between 12 populations of three-spined stickleback. - J Anim Ecol 76: 1128–1138.

Duckworth, R. A. and Badyaev, A. V. 2007. Coupling of dispersal and aggression facilitates the rapid range expansion of a passerine bird. - PNAS 104: 15017–15022.

Eccard, J. A. and Herde, A. 2013. Seasonal variation in the behaviour of a short-lived rodent. - BMC Ecol 13: 43.

Fogarty, S. et al. 2011. Social personality polymorphism and the spread of invasive species: a model. - Am Nat 177: 273–287.

Gertzen, E. et al. 2008. Quantifying invasion pathways: fish introductions from the aquarium trade. - Can J Fish Aquat Sci 65: 1265–1273.

Gruber, J. et al. 2017. Geographic divergence in dispersal-related behaviour in cane toads from range-front versus range-core populations in Australia. - Behav Ecol Sociobiol 71: 38.

Harris, S. et al. 2010. Picking personalities apart: estimating the influence of predation, sex and body size on boldness in the guppy *Poecilia reticulata*. - Oikos 119: 1711–1718.

Hudina, S. et al. 2015. Differences in aggressive behaviour along the expanding range of an invasive crayfish: an important component of invasion dynamics. - Biol Invasions 17: 3101–3112.

Jourdan, J. et al. 2014. On the natural history of an introduced population of guppies (*Poecilia reticulata* Peters, 1859) in Germany. - BioInvasions Rec in press.

Juette, T. et al. 2014. Animal personality and the ecological impacts of freshwater non-native species. - Curr Zool 60: 417–427.

Kempkes, M. et al. 2018. Tropische Neozoen in heimischen Fließgewässern. - VerlagsKG Wolf, Magdeburg.

Landgraf, T. et al. 2016. RoboFish: increased acceptance of interactive robotic fish with realistic eyes and natural motion patterns by live Trinidadian guppies. - Bioinspir Biomim 11: 15001.

Langford, T. E. 1990. Ecological Effects of Thermal Discharges. - Elsevier Applied Science Publishers Ltd.

Laskowski, K. L. et al. 2016. Behaviour in a standardized assay, but not metabolic or growth rate, predicts behavioural variation in an adult aquatic top predator Esox lucius in the wild. - J Fish Biol 88: 1544–1563.

Lockwood, J. L. et al. 2011. Non-natives: plusses of invasion ecology. - Nature 475: 36–36.

Lopez, D. P. et al. 2012. Nonnative African jewelfish are more fit but not bolder at the invasion front: a trait comparison across an Everglades range expansion. - Biol Invasions 14: 2159–2174.

Lukas, J. et al. 2017a. Feral guppies in Germany – a critical evaluation of a citizen science approach as biomonitoring tool. - Bull Fish Biol 17: 13–27.

Lukas, J. A. Y. et al. 2017b. On the occurrence of three non-native cichlid species including the first record of a feral population of *Pelmatolapia* (Tilapia) *mariae* (Boulenger, 1899) in Europe. - Roy Soc Open Sci 4: 170160.

Muraco, J. J. et al. 2014. Does male behavioral type correlate with species recognition and stress? - Behav Ecol 25: 200–205.

Myles-Gonzalez, E. et al. 2015. To boldly go where no goby has gone before: Boldness, dispersal tendency,
and metabolism at the invasion front. - Behav Ecol 26: 1083–1090.

Nakagawa, S. and Schielzeth, H. 2010. Repeatability for Gaussian and non-Gaussian data: a practical guide
for biologists. - Biol Rev Camb Philos Soc 85: 935–956.

Pintor, L. M. et al. 2008. Differences in aggression, activity and boldness between native and introduced
populations of an invasive crayfish. - Oikos 117: 1629–1636.

Pintor, L. M. et al. 2009. Behavioral correlations provide a mechanism for explaining high invader densities
and increased impacts on native prey. - Ecology 90: 581–587.

Rasmussen, J. E. and Belk, M. C. 2012. Dispersal behavior correlates with personality of a North American
fish. - Curr Zool 58: 260–270.

Réale, D. et al. 2007. Integrating animal temperament within ecology and evolution. - Biol Rev Camb Philos
Soc 82: 291–318.

Seebacher, F. et al. 2014. Regulation of thermal acclimation varies between generations of the short-lived
mosquitofish that developed in different environmental conditions. - Funct Ecol 28: 137–148.

Seebens, H. et al. 2017. No saturation in the accumulation of alien species worldwide. - Nat Comm: 1–9.

Seebens, H. et al. 2018. Global rise in emerging alien species results from increased accessibility of new
source pools. - PNAS 115: E2264–E2273.

Shine, R. et al. 2011. An evolutionary process that assembles phenotypes through space rather than through
time. - PNAS 108: 5708–5711.

Sih, A. and Bell, A. M. 2008. Insights for Behavioral Ecology from Behavioral Syndromes. - In: Advances in
the Study of Behavior. Elsevier, pp. 227–281.

Sih, A. et al. 2012. Ecological implications of behavioural syndromes. - Ecol Lett 15: 278–289.

Spiegel, O. et al. 2015. When the going gets tough: behavioural type-dependent space use in the sleepy
lizard changes as the season dries. - Proc R Soc B 282: 20151768.

Spiegel, O. et al. 2017. What's your move? Movement as a link between personality and spatial dynamics in
animal populations. - Ecol Lett 20: 3–18.

Stamps, J. and Groothuis, T. G. G. 2010. The development of animal personality: relevance, concepts and
perspectives. - Biol Rev Camb Philos Soc 85: 301–325.

Thorlacius, M. et al. 2015. Behavioral dependent dispersal in the invasive round goby *Neogobius*
melanostomus depends on population age. - Curr Zool 61: 529–542.

Uchida, K. et al. 2016. Seasonal variation of flight initiation distance in Eurasian red squirrels in urban versus
rural habitat. - J Zool 298: 225–231.

van Oers, K. et al. 2004. Realized heritability and repeatability of risk-taking behaviour in relation to avian
personalities. - Proc R Soc B 271: 65–73.

Wilson, A. D. M. M. and Godin, J.-G. J. G. J. 2009. Boldness and behavioral syndromes in the bluegill
sunfish, *Lepomis macrochirus*. - Behav Ecol 20: 231–237.

Wolf, M. and Weissing, F. J. 2012. Animal personalities: consequences for ecology and evolution. - Trends
Ecol Evol 27: 452–461.

Acknowledgement

The authors would like to thank Dr. Udo Rose – biologist at the Erftverband and member of the Erftfischereigenossenschaft (Bergheim, Germany) – for the permission to do research at the Gillbach. Sampling efforts were supported by several fish enthusiasts: Bernd Neu, Geoffrey P.F. Mazué, Vivek Hari Sridhar, as well as Petr Zajicek and Marcus Ebert from IGB Berlin. This study received funding from the Elsa-Neumann-Scholarship from the state of Berlin, the German Ichthyological Society, the German Research Foundation (BI 1828/2-1, LA 3534/1-1, KA 3029/2-1) and through Germany's Excellence Strategy - EXC 2002/1 "Science of Intelligence" (project number 228 390523135).

Authors' contributions:

JL and DB designed the study; JL, FWM, DB and GK jointly collected the fish; DB, TL and JK developed the Robofish system; JL and DB conducted the behavioural experiments, analysed the data and wrote the manuscript; all authors approved the final draft.