1 **Consistent behavioral syndrome across seasons in an**

2 **invasive freshwater fish**

4 Abstract

Understanding the linkage between behavioral types and dispersal tendency has become a

pressing issue in light of global change and biological invasions. Here, we explore whether

dispersing individuals exhibit behavioral types that differ from those remaining in the source

population. We investigated a feral population of guppies (*Poecilia reticulata*) that undergoes a

yearly range shift cycle. Guppies are among the most widespread invasive species in the world,

but in temperate regions these tropical fish can only survive in winter-warm freshwaters.

Established in a thermally-altered stream in Germany, guppies are confined to a warm-water influx

in winter, but can spread to peripheral parts as these become thermally accessible. We sampled

fish from the source population and a winter-abandoned site in March, June and August. Fish were

tested for boldness, sociability and activity involving open-field tests including interactions with a

robotic social partner. Guppies differed consistently among each other in all three traits. Behavioral

trait expression in the source population differed across seasons, however, we could not detect

differences between source and downstream populations. Instead, all sampled populations

exhibited a remarkably stable behavioral syndrome between boldness and activity despite strong

seasonal changes in water temperature and associated environmental factors. We conclude that

random drift (opposed to personality-biased dispersal) is a more likely dispersal mode for guppies,

at least in the investigated stream. In the face of fluctuating environments, guppies seem to be

extremely effective in keeping behavioral expressions constant, which could help explain their

successful invasion and adaptation to disturbed habitats.

Introduction

Consistent behavioral differences among individuals (e.g. behavioural types or animal personality; Réale et al. 2007) seem to be a ubiquitous biological feature (Bell et al. 2009; Bierbach et al. 2017) that was proposed to have substantial ecological and evolutionary importance (Sih et al. 2012; Wolf and Weissing 2012). For one, behavioural types are thought to be relevant in determining success (Chapple et al. 2012; Carere and Gherardi 2013; Canestrelli et al. 2015) and impact of biological invasions (Cote, Clobert, et al. 2010; Fogarty et al. 2011; Juette et al. 2014), which is becoming an ever more pressing issue in the light of global change and ongoing translocations of species among habitats (Wong and Candolin 2015; Seebens et al. 2017; Seebens et al. 2018).

It is assumed that individuals that invade novel habitats are a non-random subset of their source populations (Shine et al. 2011; Sih et al. 2012; Juette et al. 2014; Canestrelli et al. 2015; Spiegel et al. 2017). These invaders may share certain phenotypic and life-history characteristics that are associated with their behavioral types (Juette et al. 2014). Behavioral types of invading individuals may determine whether they are transported (by humans) or move (by themselves) into new environments (first invasion stage; Lockwood et al. 2011), whether they can survive and establish a non-native population (second stage), and whether they may spread to new habitats thereafter (third stage). Behavioral types shown to be relevant in this regard are boldness (Cote et al. 2011; Myles-Gonzalez et al. 2015), aggression (Duckworth and Badyaev 2007; Hudina et al. 2015), activity (Brown and Irving 2014; Thorlacius et al. 2015) as well as sociability (Cote, Fogarty, et al. 2010; Cote et al. 2011; Rasmussen and Belk 2012). Further, it was predicted that individuals carrying correlated suites (i.e., behavioral syndromes, Sih et al. 2012) of bold, aggressive, active and asocial types may predominantly be found at the invasion front (Duckworth and Badyaev 2007; Cote, Fogarty, et al. 2010; Cote et al. 2011; Juette et al. 2014). However, such spatially and/or temporally assorted compositions of behavioral types among populations have only rarely been investigated along naturally occurring invasion gradients.

The difficulty to study invasion phenomena in the wild is their random and unpredictable occurrence (Lockwood et al. 2011), which makes it nearly impossible to follow active invasion

fronts. In the rare cases where range expansions could be tracked in real time (e.g., western

bluebirds: Duckworth and Badyaev 2007; African jewelfish: Lopez et al. 2012; signal crayfish:

Hudina et al. 2015; round goby: Thorlacius et al. 2015; cane toad: Gruber et al. 2017), the invasion

origin is often unknown or unclear due to multiple introduction events. Thus, we need to identify

study systems that allow us to investigate dispersal across sensible temporal and spatial scales.

Thermally altered freshwater systems (TAFs) in the temperate zones provide fruitful conditions to

study this phenomenon. Here, natural and/or anthropogenic warm water influxes create unique

temperature refuges that allow non-native species (especially those of tropical origin) to establish

populations (Langford 1990; Jourdan et al. 2014). While ambient water temperatures drop in winter

(e.g., less than +1°C in the river Rhine; Jourdan et al. 2014), areas close to constant warm water

influx will remain suitable for non-native tropical species to survive. Once peripheral areas become

thermally suitable again in warmer seasons, certain individuals from the sources may disperse into

these areas. Such a scenario enables us to investigate the role of behavioral types during *in situ*

dispersal of non-native species from a single, well-determined source into new (i.e. winter-

abandoned) habitats.

Under a personality-biased dispersal hypothesis, where individuals with a certain set of

personality traits are assumed to leave the source population to colonize new habitats, we

predicted that *(i)* the source population's average personality traits should change over the course

of the year due to the drain of the emigrating individuals, *(ii)* newly founded populations should

differ in their average personality traits compared to their source populations, and *(iii)* behavioral

variation should decrease both in the source as well as in the newly founded population due to the

non-random migration patterns. In addition, *(iv)* population-wide correlations among personality

traits (behavioral syndromes) may emerge or disappear because of assumed non-random

migration patterns.

In Germany, tropical fishes have established populations in several artificially heated

stream systems (Lukas, Kalinkat, et al. 2017), the most investigated being the Gillbach near

Cologne (see Fig. 1a; Jourdan et al. 2014; Lukas, Jourdan, et al. 2017; Kempkes et al. 2018). To

estimate seasonal fluctuations in temperature-associated abiotic and biotic conditions within the

sampled range of the stream, we continuously recorded water temperatures with submerged data-

loggers. We sampled feral guppies (*Poecilia reticulata*) from the Gillbach during different seasons

(temporal scale: spring, early summer, late summer) and, if possible, at different sites (spatial

scale: core of warm water influx versus more peripheral downstream area; see Fig. 1a) and

repeatedly assessed behavioral types in terms of boldness, sociability and activity in a laboratory

setting.

Material& Methods

Study site & Sampling

The Gillbach (Rhine catchment, Germany) has been under the influence of thermal pollution and anthropogenic modifications for over six decades. Its headwaters are exclusively fed by the warm water discharge of the brown coal power plant Niederaußem (Fig. 1a). This creates a temperature regime that is highly dependent on the volume and temperature of the plant's water discharge. Generally exhibiting above ambient water temperatures (Fig. 1b), the Gillbach harbors several established populations of non-native fish, invertebrate and plant species, many of which are of tropical or subtropical origin (see Supplementary material Appendix 1, Table S1 for a list of native and non-native fish species found in the Gillbach).

Here, feral guppies (*Poecilia reticulata*) are highly abundant and as successful global invaders (Deacon et al. 2011) they are well suited to investigate the expression of behavioral traits across time and space. Sampling was conducted at two sites (~300m transects, see Fig. 1a) in March, June and August 2016. The Gillbach's guppy population density increased from March to June and decreased in late summer (Supplementary material Appendix 2, Table S3). This extreme population decline was likely predation-induced as densities of convict cichlids (*Amatitlania nigrofasciata*) and European chub (*Squalius cephalus*) also increased during this period (Supplementary material Appendix 1, Table S1). Sampling commenced until sufficient numbers for behavioral testing were reached (i.e., ~40 adult individuals). Captured fish were transferred into well-aerated coolers and kept separately by sampling site. To reduce a possible sampling bias of certain behavioral types, we collected specimens by seine and dip netting and actively drove fish into the net as recommended by Biro & Dingemanse (2009). While we acknowledge some size selectivity against juvenile guppies introduced by the 2 mm mesh size, our population sample was representative of adult body size variation (see Fig. 3b), thus avoiding indirect bias of behavioral traits that are correlated with size (Polverino et al. 2016).

Water temperatures at the source (site 1) and one kilometer downstream (site 2) were monitored using HOBO data loggers (Onset Computer Corporation, Bourne, MA, USA) for a period

of 12 months (Fig. 1b). For the period of sampling and behavioral testing (March – August 2016),

the Gillbach experienced a thermal gradient of ~2°C/km (mean ± SD, site 1: 25.53°C ± 3.21°C; site

2: 23.49°C ± 3.4°C; Fig. 1b). However, we expect the temperature difference between sites to be

even more pronounced in colder months (i.e. decline of 3.5°C/km in November (Klotz et al. 2013);

4.7°C/km in February (Jourdan et al. 2014)). With temperatures below the population's critical

minimum temperature (~12°C; Jourdan et al. 2014), survival of guppies outside the core area

during winter is highly unlikely. Under similar conditions in June (~1.5°C/km gradient), a previous

study (Emde et al. 2016) found diet composition of the non-native predatory cichlid *Amatitlania*

nigrofasciata to vastly differ between sites, with cichlids exerting more piscivorous predation

pressure at peripheral sites.

Figure 1: The Gillbach stream and its altered temperature regime. a) Warm water discharge

from a power plant feeds into the Gillbach (map created using © 2019 Google Earth with data from

2009 GeoBasis-DE/BKG) and generates a thermal gradient above ambient water temperatures

that is preserved for several kilometers (site 1 – site 2). **b)** 4-hourly water temperature* with error

bars indicating monthly means ± SD for each site (site 1 – Gillbach source: red; site 2 – one

kilometer downstream: orange). Dotted lines correspond to the three sampling occasions. * Note

that due to a loss of loggers, continuous monitoring at site 2 was interrupted (no data from

01.01.2016 – 19.03.2016 nor from 29.08.2016 – 31.12.2016).

Behavioral assessment

Fish collected for behavioral testing were transferred to the laboratory at Humboldt University of

Berlin, Germany. Fish were kept in one holding tank (270 liters), but were separated into mesh-

divided compartments by collection time and site ('populations' hereafter). Each compartment (24

cm ✕ 58 cm ✕ 40 cm L✕W✕H) contained artificial java moss and natural gravel. Females and males

were housed communally. Water was maintained at 26°C and kept under continuous aeration and

filtration. Fish were held under diurnal lighting (12:12h light:dark cycle) and fed commercial flake

food (TetraMin® Tetra GmbH) twice a day. Feeding stopped 12 hours before experiments and

recommenced afterwards. To acclimate the fish to laboratory conditions and reduce potential

short-term effects of the different sampling habitats on behavioral trait expression, no behavioral

testing was conducted for at least 30 days (see Fig. 2a for detailed experimental design). One

week prior to testing, adults of suitable size (SL>15 mm) were marked using visible implant

elastomer tags (VIE; Northwest Marine Technology Ltd, Shaw Island, WA, USA) to allow for

individual identification throughout the experiment. To be mindful of capture biases towards a

specific behavioral type, we used all fish within a compartment for testing.

Figure 2: Behavioral testing with robotic fish. a) Test fish were collected three times and, when possible, at two sites (site 1 and 2, see Fig. 1a). After an initial 30-day acclimation period, fish were tested a total of three times (trial I – III, sample sizes indicate tested individuals, for total catch numbers see Supplementary material Appendix 2, Table S3). **b-d)** During trials, the fish replica was moved remotely through the test tank (c). Once the focal individual had emerged from its shelter, the biomimetic robot initiated its predefined path (b) and the fish's swimming trajectory was recorded (d).

We tested a total of 159 guppies (95 females, 64 males; for population-specific sample sizes see Fig. 2a). Test fish were introduced to an open field test arena (88 x 88 cm, 75 mm water depth) filled with aged tap water kept at 26°C. To reduce disturbing stimuli, the arena was enclosed in black, opaque plastic and illuminated with artificial light reproducing the daylight spectrum. Behavioral observations were recorded via a camera (Ximea 4K USB 3.1 camera) mounted above the arena. Position information was extracted using EthoVision 10 XT software (Noldus Information Technology, Wageningen, Netherlands).

(1) Boldness: Latency to emerge from a shelter was used as a measure of boldness with a low latency time indicating high levels of boldness. Fish were introduced into a grey opaque shelter

box (Ø 100 mm; top left of each image in Fig. 2b and d) and allowed to acclimate. After one

minute, a sponge was removed from the entrance (40mm x 25mm), allowing the fish to emerge

and explore the test arena. A fish was scored as having emerged when its full body was visible

outside of the shelter. If fish did not emerge after five minutes, the lid of the shelter box was

removed. Fish that failed to emerge within eight minutes were given the maximum score of 480

seconds and the box was removed entirely.

(2) Sociability: A three-dimensional fish replica was stationed 1 cm away from the shelter entrance.

The replica, which resembled a large-sized guppy female (SL=30mm; left fish in Fig. 2c), could

be moved by a robot via a magnetic base (see Landgraf et al. 2016; Bierbach et al. 2018 for

details). Upon the fish's emergence (or the removal of the shelter box), the biomimetic robot

started to move through the arena in a standardized sequence (Fig. 2b) imitating a conspecific.

We calculated the mean distance the fish kept to the biomimetic robot as a proxy of sociability

with lower values indicating higher levels of sociability. Consequently, fish that followed the

robot closer were assumed to be more sociable (see Bierbach et al. 2018).

(3) Activity: After completion of the robotic fish's predefined path, the replica was removed from the

arena by hand and once the fish resumed normal swimming behavior, it was allowed to explore

the arena. As a measure of activity, we calculated the mean velocity with which the fish

explored the open arena during 3 minutes.

To minimize handling and exposure to stress (Animal Behaviour 2020), all three traits were

assessed within a single behavioral assay. After the first behavioral assay, we recorded the body

length of each individual. Fish were transferred back to their respective compartment and allowed

72 hours resting before being retested following the same protocol. To reduce acclimatization

effects, we only assessed short-term repeatability by testing fish a total of three times (randomized

order) within the span of a week. All reported experiments comply with the current German law

approved by LaGeSo (G0117/16 to Dr. D. Bierbach).

Statistical analysis

To test for differences in average trait expression over a temporal and/or spatial scale, we used

linear mixed models (LMM, *GENLINMIXED* procedure; SPSS v. 25.0, IBM Corp., Armonk, NY,

USA) with each behavioral trait (boldness, sociability, activity) as a separate response variable.

'Boldness' and 'sociability' were log10-transformed to fulfil normality assumptions. All models

included sample population as the fixed effect of interest (four levels: March – site 1, June – site 1,

August – site 1, August – site 2). The low sample size collected at site 2 during March and June

(n=7 and n=2, respectively; see Fig. 2a) made these estimates uncertain. After verifying that there

were no substantial differences found when running the analyses with and without the above

population samples, these estimates were excluded from final analyses. Behavior in poeciliids is

often size- and/or sex-dependent (Brown and Braithwaite 2004; Harris et al. 2010), thus we

included standard length (SL) as a covariate and 'sex' as an additional fixed factor. To account for

the repeated testing of individuals, we further included 'trial' as a fixed factor and 'fish ID' as a

random factor. Post-hoc LSD tests were employed in case fixed factors were found to have a

significant effect. For the covariate 'body size', we also performed Pearson's correlations to better

evaluate the relationship.

(Co)Variance structure was nested within 'population', which enabled us to calculate the

total behavioral variance as well as within- and among-individual variance components in each

population. Based on this, we calculated behavioral repeatability for each of the four populations

as a measure of consistency. We assumed that repeatability and/or variance estimates differed

among populations when 95% confidence intervals (CIs) were not overlapping (see Nakagawa and

Schielzeth 2010).

To evaluate the existence of behavioral syndromes (i.e. correlation among behavioral traits)

within our populations, we used linear regression models to test for significant correlation between

each trait pair. In order to minimize the effects of mean differences in behavioral traits on trait

correlation, we mean-centered each behavioral trait within 'trial' and 'population' (z-transformation).

Each model included one trait as the dependent variable and another as covariate. We further

compared whether trait correlation differed among populations (i.e., difference in slopes termed

'covariate nested within population'). Visualizations were done with GraphPad Prism (version 8.01,

GraphPad Software Inc., La Jolla, CA, USA).

Results

Differences in average trait expression among populations

Expression of sociability and activity differed between populations, while boldness did not ('population' effect in Table 1; Fig. 3a). On a temporal scale, fish sampled at site 1 in August showed higher levels of sociability (i.e. reduced distance to robotic fish) compared to March and June populations (*post-hoc*, August site 1 *vs.* March – site 1: $p = 0.003$, *vs.* June – site 1: $p = 0.025$). Similarly, fish sampled at site 1 in August were more active than populations sampled in March and June (*post-hoc*, August site 1 *vs.* March – site 1: $p = 0.002$, *vs.* June – site 1: $p < 0.001$). On a spatial scale, we found no evidence that fish sampled in August at site 1 and site 2 differed in boldness, sociability, nor activity (*post-hoc*, August site 1 *vs.* site 2 – boldness: $p = 0.76$, sociability: $p = 0.92$, activity: $p = 0.89$; Fig. 3a).

Boldness and activity showed a size-dependency (sig. effect of covariate 'body size'; Table 1, Fig. 3b) with smaller individuals being bolder (i.e. reduced emergence time; Pearson's $r = 0.32$; $p < 0.001$) and more active (Pearson's $r = -0.35$; $p < 0.001$) than larger ones. We found no such relation between body size and sociability (Pearson's $r = 0.002$; $p = 0.98$). While sex is correlated with size in the sexually dimorphic guppy, sex did not affect any of the measured behavioral traits (non-sig. 'sex' effect; Table 1). We further observed a habituation to the repeated testing during sociability and activity assays (sig. 'trial' effect; Table 1). Fish were considerably less social and less active with each consecutive measurement.

Table 1: Results from linear mixed models.

Effect	Boldness ($\log_{10}$)		Sociability ($\log_{10}$)		Activity	
	$F_{df1,df2}$	p	$F_{df1,df2}$	p	$F_{df1,df2}$	p
population	$F_{3,469} = 0.57$	0.634	$F_{3,469} = 3.8$	**0.01**	$F_{3,469} = 8.44$	**< 0.001**
trial	$F_{2,469} = 1.72$	0.18	$F_{2,469} = 135.9$	**< 0.001**	$F_{2,469} = 42.32$	**< 0.001**
sex	$F_{1,469} = 0.06$	0.81	$F_{1,469} = 2.28$	0.132	$F_{1,469} = 0.24$	0.622
body size (SL)	$F_{1,469} = 8.43$	**0.004**	$F_{1,469} = 0.02$	0.884	$F_{1,469} = 3.58$	0.059

For each behavioral trait, the LMM included 'population', 'trial' and 'sex' as fixed effects and 'body size' (SL in mm) as covariate with the random factor Fish ID. Significant effects ($p < 0.05$) are indicated in bold.

Figure 3: Average personality traits across temporal and spatial scale. *(a)* Median trait scores (and IQRs) for boldness* (i.e., time to emerge from shelter), activity (i.e., velocity during open field exploration) and sociability* (i.e., distance to biomimetic robot). Different letters depict significant differences based on post-hoc LSD tests after global linear mixed models (*p* < 0.05). *(b)* Mean traits scores (± SD) in relation to body size (standard length; *n*=159). To visualize the relationships, we show linear regressions lines (red) calculated from individual means (see main text for significance levels and Pearson's *r*). * Note that low scores of boldness and sociability indicate high trait expression.

Among- and within-individual variation in behavioral traits

Fish within each sampled population differed consistently in their expression of boldness,

sociability and activity across trials (sig. repeatability estimates; Table 2). Only fish sampled in

August at site 2 had a significantly higher repeatability estimate for sociability than fish from the

March sampling at site 1 (non-overlapping CIs; Table 2). There was no significant difference

among populations in estimated overall behavioral variances in any of the examined traits (see

'total variation' in Table 2; for within- and among-individual variance estimates see Supplementary

material Appendix 3, Table S4).

Table 2: Repeatability estimates and total behavioral variance estimates.

Popula-tion	Boldness (log_{10})			Sociability (log_{10})			Activity		
	Repeat-ability [95% CI]	p	Total Variation [95% CI]	Repeat-ability [95% CI]	p	Total Variation [95% CI]	Repeat-ability [95% CI]	p	Total Variation [95% CI]
March site 1	0.49 [0.40–0.57]	**0.004**	0.24 [0.15–0.4]	0.26 [0.15–0.4]	0.05	0.06 [0.04–0.1]	0.27 [0.17–0.41]	**0.042**	6.86 [4.3–11.87]
June site 1	0.40 [0.31–0.5]	**0.005**	0.31 [0.2–0.5]	0.32 [0.23–0.43]	**0.011**	0.07 [0.05–0.11]	0.30 [0.21–0.42]	**0.015**	5.75 [3.74-9.33]
August site 1	0.54 [0.47–0.6]	**0.001**	0.34 [0.22–0.54]	0.34 [0.25–0.45]	**0.009**	0.06 [0.04–0.09]	0.38 [0.29–0.47]	**0.005**	7.15 [4.61-11.48]
August site 2	0.53 [0.46–0.6]	**0.001**	0.3 [0.19–0.48]	0.49 [0.42–0.57]	**0.001**	0.08 [0.05–0.13]	0.36 [0.27–0.46]	**0.007**	5.92 [3.81-9.57]

p-values are derived from Wald's *z* statistics (see Supplementary material Appendix 3, Table S4 for within-
and among-individual variation estimates). Significant differences among populations (bold) can be assumed
when 95%CI do not overlap.

Behavioral syndrome structure

Boldness and activity were correlated in all four population samples (sig. correlation between

boldness and activity: $F_{1,157}$ = 52.23; $p < 0.001$; no sig. difference in slopes among populations:

$F_{3,151}$ = 0.45; p = 0.715). Fish with high average activity levels also left the shelter quicker,

confirming that activity and boldness are part of a larger behavioral syndrome (Fig. 4).

There was no evidence for correlations between sociability and boldness ($F_{1,157}$ = 0.026; p =

0.872; non-sig. difference in slopes: $F_{3,151}$ = 1.8; p = 0.149), nor between sociability and activity

($F_{1,157}$ = 0.062; p = 0.803; non-sig. difference in slopes: $F_{3,151}$ = 0.816; p = 0.487) among

populations. Thus, sociability measured as distance kept to the moving robotic fish does not seem

to be part of a behavioral syndrome with either boldness nor activity.

Figure 4: Correlations between behavioral traits across temporal and spatial scale. Mean trait scores* (±SD) of individual test fish are shown for each population. Linear regressions (red line and Pearson's *r*) visualize the relationships between traits (see main text for detailed analysis of trait correlation; *n* = 159). * Note that for boldness and sociability lower values indicate a high trait score.

Discussion

Our study follows the annual range shifts of non-native fish in an artificially heated stream in

temperate Germany. As guppies dispersed into winter-abandoned areas in warmer months, we

expected this process to be driven by dispersal-enhancing behavioral traits. We found similar

consistent among-individual variation (e.g., significant behavioral repeatability) in the behaviors

'boldness', 'sociability' and 'activity' within all temporal and spatial samples (except a difference

between sociability in March at site 1 and August at site 2). Contrary to our predictions, we did not

detect a decrease in total behavioral variation of the sampled populations indicative of a "thinning

out" of disperser's behavioral types. Further, we found no evidence that the source and the newly-

founded population differed in terms of average behavioral trait expression, though a seasonal

change in activity and sociability (but not boldness) was detectable in the source population:

August-sampled individuals were more active and more sociable than those sampled earlier in the

year. Instead, we found a behavioral syndrome among activity and boldness that was consistently

detectable throughout the year and across sampled populations. To sum up, we found no strong

evidence for a personality-biased dispersal in this feral guppy population and the detected

seasonal differences in average trait expression (sociability, activity) are likely due to changed

environmental factors such as temperature, predator abundance and/or resource availability.

Differences in behavioral traits across seasons but not between sites as found in our study

could be explained by seasonal variation in environmental conditions rather than by a drain of

individuals carrying certain behavioral types from source to downstream populations. The

Gillbach's water temperatures show strongest differences between sites in the colder months but

relatively small differences during the warmer months. Thus, when winter-abandoned peripheral

sites are colonized again in summer, these habitats do not differ much in temperature from the

source at this point in time. Nevertheless, the Gillbach still shows a strong seasonality in

temperature which, in combination with a variation in day length, food quality/quantity and

predation pressure, may likely cause population-wide changes in behaviors across seasons (e.g.,

Biro et al. 2010; Eccard and Herde 2013; Spiegel et al. 2015; Uchida et al. 2016; Barbosa et al.

2018; Dhellemmes et al. 2020) as seen in the source population. For example, prior temperature

acclimation during sensitive developmental phases can alter the phenotypes of subsequent adult

phases (Beitinger et al. 2000; Seebacher et al. 2014). This might explain the increased activity

exhibited by fish in August, as higher developmental temperatures lead to more active phenotypes

(Seebacher et al. 2014). However, teasing apart the effects of abiotic (e.g. temperature) and biotic

(e.g., predation pressure) fluctuations in environmental conditions will require additional

experimentation.

Besides seasonal differences in average behavioral traits, we found stable

behavioral syndromes among sampled populations that involved a strong positive correlation

between boldness and activity. Such a boldness-activity syndrome has been documented in

various taxa (van Oers et al. 2004; Pintor et al. 2008; Wilson and Godin 2009; Cote, Fogarty, et al.

2010; Eccard and Herde 2013; Muraco et al. 2014), including guppies (Brown and Irving 2014).

Natural selection through predation favors the evolution of behavioral correlations (Dingemanse et

al. 2007; Harris et al. 2010; Dhellemmes et al. 2020). The Gillbach's guppy population underwent

strong seasonal changes in size and the observed population decline in summer, while primary

production is highest in temperate regions, suggests strong predation pressure. A positive

boldness-activity correlation might thus represent an optimal trait combination in the face of

predation experienced by both populations (Smith and Blumstein 2010). However, the lack of

differences in the level of (individual) behavioral variation among populations might be a result of a

random downstream drift and a low overall behavioral plasticity in this population. This can be a

consequence of the assumed bottleneck origin of this feral population that most likely stems from

an unknown number of released domesticated guppies (see Gertzen et al. 2008 for estimates of

propagule pressure for popular aquarium species). Investigations involving other populations of the

guppy and more distantly related species are highly recommended, especially as several

environmental factors are predicted to impact behavioral variation, especially in fish (e.g., Harris et

al. 2010; Laskowski et al. 2016; Barbosa et al. 2018). Alternatively, guppies and especially those

from the Gillbach that experience highly fluctuating temperature regimes may have the ability to

adjust their behavioral expressions within very short times, e.g., within the 30-days acclimatization

period in the laboratory used here. It is known that guppies have a remarkable capacity for

behavioral plasticity (Deacon and Magurran 2016) and thermal adaptation (Chung 2001; Jourdan

et al. 2014), and thus have successfully established populations in environments with very different

(thermal) regimes compared to their native habitats (Deacon et al. 2011). Testing of individuals

after different acclimation periods could be a possibility to investigate this issue further.

Theoretical models (Cote, Clobert, et al. 2010; Spiegel et al. 2017) as well as several

studies across taxa (Duckworth and Badyaev 2007; Pintor et al. 2009; Cote, Fogarty, et al. 2010;

Myles-Gonzalez et al. 2015; Coates et al. 2019) provide evidence for the prominent role of

individual differences in behavior during dispersal and invasion processes (Bowler and Benton

2005; Chapple et al. 2012; Canestrelli et al. 2015). Nevertheless, the picture might be more

complicated and context-dependent: spatial sorting of behavioral types did not emerge under

predation pressure (Cote et al. 2013) and was lost as the population aged (Thorlacius et al. 2015).

Moreover, even if there are significant (but small) effects of behavioural trait variation affecting

relevant ecological phenomena, they might be non-detectable in benign laboratory environments

(Schröder et al. 2016; Moiron et al. 2020). Our current study exemplifies that population-wide

behavioral analysis can fail to provide evidence for behaviorally-mediated migration patterns and

future analysis may focus on tracking individuals along their travel routes across seasons and

locations. On top of this, several other feral guppy populations are known in Germany (Lukas,

Kalinkat, et al. 2017) and across Europe (Kempkes et al. 2018) for which similar investigations are

highly advisable in order to test the generality of our herein presented results. Such analyses

should strive to incorporate the invasion history, environmental factors as well as the population-

specific genetic backgrounds which all can affect behavioral variation in wild populations (Sih and

Bell 2008; Bergmüller and Taborsky 2010; Stamps and Groothuis 2010).

References

Animal Behaviour. 2020. Guidelines for the treatment of animals in behavioural research and teaching. Animal Behaviour. 159:I–XI. doi:10.1016/j.anbehav.2019.11.002.

Barbosa M, Deacon AE, Janeiro MJ, Ramnarine I, Morrissey MB, Magurran AE. 2018. Individual variation in reproductive behaviour is linked to temporal heterogeneity in predation risk. Proceedings of the Royal Society B: Biological Sciences. 285(1870):20171499. doi:10.1098/rspb.2017.1499.

Beitinger TL, Bennett WA, McCauley RW. 2000. Temperature Tolerances of North American Freshwater Fishes Exposed to Dynamic Changes in Temperature. Environ Biol Fish. 58:40.

Bell AM, Hankison SJ, Laskowski KL. 2009. The repeatability of behaviour: a meta-analysis. Anim Behav. 77(4):771–783. doi:10.1016/j.anbehav.2008.12.022.

Bergmüller R, Taborsky M. 2010. Animal personality due to social niche specialisation.

Bierbach D, Landgraf T, Romanczuk P, Lukas J, Nguyen H, Wolf M, Krause J. 2018. Using a robotic fish to investigate individual differences in social responsiveness in the guppy. Roy Soc Open Sci. 5(8):181026. doi:10.1098/rsos.181026.

Bierbach D, Laskowski KL, Wolf M. 2017. Behavioural individuality in clonal fish arises despite near-identical rearing conditions. Nat Comm. 8:15361. doi:10.1038/ncomms15361.

Biro PA, Beckmann C, Stamps JA. 2010. Small within-day increases in temperature affects boldness and alters personality in coral reef fish. Proc R Soc B.(September 2009):71–77. doi:10.1098/rspb.2009.1346.

Biro PA, Dingemanse NJ. 2009. Sampling bias resulting from animal personality. Trends in Ecology & Evolution. 24(2):66–67. doi:10.1016/j.tree.2008.11.001.

Bowler DE, Benton TG. 2005. Causes and consequences of animal dispersal strategies: relating individual behaviour to spatial dynamics. Biol Rev Camb Philos Soc. 80(2):205–225. doi:10.1017/S1464793104006645.

Brown C, Braithwaite VA. 2004. Size matters: A test of boldness in eight populations of the poeciliid *Brachyraphis episcopi*. Anim Behav. 68(6):1325–1329. doi:10.1016/j.anbehav.2004.04.004.

Brown C, Irving E. 2014. Individual personality traits influence group exploration in a feral guppy population. Behav Ecol. 25(1):95–101. doi:10.1093/beheco/art090.

Canestrelli D, Bisconti R, Carere C. 2015. Bolder Takes All? The Behavioral Dimension of Biogeography. Trends Ecol Evol. 31(1):35–43. doi:10.1016/j.tree.2015.11.004.

Carere C, Gherardi F. 2013. Animal personalities matter for biological invasions. Elsevier Ltd. http://dx.doi.org/10.1016/j.tree.2012.10.006.

Chapple DG, Simmonds SM, Wong BBM. 2012. Can behavioral and personality traits influence the success of unintentional species introductions? Trends Ecol Evol. 27(1):57–64. doi:10.1016/j.tree.2011.09.010.

Chung KS. 2001. Critical thermal maxima and acclimation rate of the tropical guppy *Poecilla reticulata*. Hydrobiologia. 462:5.

Coates WD, Hale R, Morrongiello JR. 2019. Dispersal decisions and personality in a freshwater fish. Anim Behav. 157:209–218. doi:10.1016/j.anbehav.2019.07.012.

Cote J, Clobert J, Brodin T, Fogarty S, Sih A. 2010. Personality-dependent dispersal: characterization, ontogeny and consequences for spatially structured populations. Phil Trans R Soc B. 365(1560):4065–4076. doi:10.1098/rstb.2010.0176.

Cote J, Fogarty S, Brodin T, Weinersmith K, Sih A. 2011. Personality-dependent dispersal in the invasive mosquitofish: group composition matters. Proc R Soc B. 278(1712):1670–1678. doi:10.1098/rspb.2010.1892.

Cote J, Fogarty S, Tymen B, Sih A, Brodin T. 2013. Personality-dependent dispersal cancelled under predation risk. Proc R Soc B. 280(1773):20132349. doi:10.1098/rspb.2013.2349.

Cote J, Fogarty S, Weinersmith K, Brodin T, Sih A. 2010. Personality traits and dispersal tendency in the invasive
mosquitofish (*Gambusia affinis*). Proc R Soc B. 277(1687):1571–1579. doi:10.1098/rspb.2009.2128.

Deacon AE, Magurran AE. 2016. How Behaviour Contributes to the Success of an Invasive Poeciliid Fish: The
Trinidadian Guppy (*Poecilia reticulata*) as a Model Species. In: Weis JS, Sol D, editors. Biological Invasions and Animal
Behaviour. Cambridge: Cambridge University Press. p. 266–290. [accessed 2018 Jun 19].
https://www.cambridge.org/core/product/identifier/9781139939492%23CT-bp-15/type/book_part.

Deacon AE, Ramnarine IW, Magurran AE. 2011. How Reproductive Ecology Contributes to the Spread of a Globally
Invasive Fish. PLOS ONE. 6(9):e24416. doi:10.1371/journal.pone.0024416.

Dhellemmes F, Finger J-S, Laskowski KL, Guttridge TL, Krause J. 2020. Comparing behavioural syndromes across time
and ecological conditions in a free-ranging predator. Animal Behaviour. 162:23–33. doi:10.1016/j.anbehav.2020.01.009.

Dingemanse NJ, Wright J, Kazem AJN, Thomas DK, Hickling R, Dawnay N. 2007. Behavioural syndromes differ
predictably between 12 populations of three-spined stickleback. J Anim Ecol. 76(6):1128–1138. doi:10.1111/j.1365-
2656.2007.01284.x.

Duckworth RA, Badyaev AV. 2007. Coupling of dispersal and aggression facilitates the rapid range expansion of a
passerine bird. PNAS. 104(38):15017–15022. doi:10.1073/pnas.0706174104.

Eccard JA, Herde A. 2013. Seasonal variation in the behaviour of a short-lived rodent. BMC Ecol. 13(1):43.
doi:10.1186/1472-6785-13-43.

Emde S, Kochmann J, Kuhn T, Doerge DD, Plath M, Miesen FW, Klimpel S. 2016. Cooling water of power plant creates
"hot spots" for tropical fishes and parasites. Parasitol Res. 115(1):85–98. doi:10.1007/s00436-015-4724-4.

Fogarty S, Cote J, Sih A. 2011. Social personality polymorphism and the spread of invasive species: a model. Am Nat.
177(3):273–287. doi:10.1086/658174.

Gertzen E, Familiar O, Leung B. 2008. Quantifying invasion pathways: fish introductions from the aquarium trade. Can J
Fish Aquat Sci. 65(7):1265–1273. doi:10.1139/F08-056.

Gruber J, Brown G, Whiting MJ, Shine R. 2017. Geographic divergence in dispersal-related behaviour in cane toads
from range-front versus range-core populations in Australia. Behav Ecol Sociobiol. 71(2):38. doi:10.1007/s00265-017-
2266-8.

Harris S, Ramnarine IW, Smith HG, Pettersson LB. 2010. Picking personalities apart: estimating the influence of
predation, sex and body size on boldness in the guppy *Poecilia reticulata*. Oikos. 119(11):1711–1718.
doi:10.1111/j.1600-0706.2010.18028.x.

Hudina S, Žganec K, Hock K. 2015. Differences in aggressive behaviour along the expanding range of an invasive
crayfish: an important component of invasion dynamics. Biol Invasions. 17(11):3101–3112. doi:10.1007/s10530-015-
0936-x.

Jourdan J, Miesen FW, Zimmer C, Gasch K, Herder F, Schleucher E, Plath M, Bierbach D. 2014. On the natural history
of an introduced population of guppies (*Poecilia reticulata* Peters, 1859) in Germany. BioInvasions Rec. 3(3).

Juette T, Cucherousset J, Cote J. 2014. Animal personality and the ecological impacts of freshwater non-native species.
Curr Zool. 60(3):417–427. doi:10.1093/czoolo/60.3.417.

Kempkes M, Lukas J, Bierbach D. 2018. Tropische Neozoen in heimischen Fließgewässern. VerlagsKG Wolf,
Magdeburg (Die Neue-Brehm Bücherei (NBB kompakt)).

Klotz W, Miesen FW, Hüllen S, Herder F. 2013. Two Asian fresh water shrimp species found in a thermally polluted
stream system in North Rhine-Westphalia, Germany. Aquat Invasions. 8(3):333–339. doi:10.3391/ai.2013.8.3.09.

Landgraf T, Bierbach D, Nguyen H, Muggelberg N, Romanczuk P, Krause J. 2016. RoboFish: increased acceptance of
interactive robotic fish with realistic eyes and natural motion patterns by live Trinidadian guppies. Bioinspir Biomim.
11(1):15001. doi:10.1088/1748-3190/11/1/015001.

Langford TE. 1990. Ecological Effects of Thermal Discharges. Essex, UK: Elsevier Applied Science Publishers Ltd.

Laskowski KL, Monk CT, Polverino G, Alós J, Nakayama S, Staaks G, Mehner T, Arlinghaus R. 2016. Behaviour in a
standardized assay, but not metabolic or growth rate, predicts behavioural variation in an adult aquatic top predator *Esox*
lucius in the wild. J Fish Biol. 88(4):1544–1563. doi:10.1111/jfb.12933.

Lockwood JL, Hoopes MF, Marchetti MP. 2011. Non-natives: plusses of invasion ecology. Nature. 475:36–36.
doi:10.1038/475036c.

Lopez DP, Jungman AA, Rehage JS. 2012. Nonnative African jewelfish are more fit but not bolder at the invasion front: a
trait comparison across an Everglades range expansion. Biol Invasions. 14(10):2159–2174. doi:10.1007/s10530-012-
0221-1.

Lukas J, Jourdan J, Kalinkat G, Emde S, Miesen FW, Jüngling H, Cocchiararo B, Bierbach D. 2017. On the occurrence
of three non-native cichlid species including the first record of a feral population of *Pelmatolapia* (Tilapia) *mariae*
(Boulenger, 1899) in Europe. Roy Soc Open Sci. 4(6):170160. doi:10.1098/rsos.170160.

Lukas J, Kalinkat G, Kempkes M, Rose U, Bierbach D. 2017. Feral guppies in Germany – a critical evaluation of a citizen
science approach as biomonitoring tool. Bull Fish Biol. 17(1/2):13–27.

Moiron M, Laskowski KL, Niemelä PT. 2020. Individual differences in behaviour explain variation in survival: a
meta-analysis. Gurevitch J, editor. Ecology Letters. 23(2):399–408. doi:10.1111/ele.13438.

Muraco JJ, Aspbury AS, Gabor CR. 2014. Does male behavioral type correlate with species recognition and stress?
Behav Ecol. 25(1):200–205. doi:10.1093/beheco/art106.

Myles-Gonzalez E, Burness G, Yavno S, Rooke A, Fox MG. 2015. To boldly go where no goby has gone before:
Boldness, dispersal tendency, and metabolism at the invasion front. Behav Ecol. 26(4):1083–1090.
doi:10.1093/beheco/arv050.

Nakagawa S, Schielzeth H. 2010. Repeatability for Gaussian and non-Gaussian data: a practical guide for biologists.
Biol Rev Camb Philos Soc. 85(4):935–956. doi:10.1111/j.1469-185X.2010.00141.x.

van Oers K, Drent PJ, de Goede P, van Noordwijk AJ. 2004. Realized heritability and repeatability of risk-taking
behaviour in relation to avian personalities. Proc R Soc B. 271(1534):65–73. doi:10.1098/rspb.2003.2518.

Pintor LM, Sih A, Bauer ML. 2008. Differences in aggression, activity and boldness between native and introduced
populations of an invasive crayfish. Oikos. 117(11):1629–1636. doi:10.1111/j.1600-0706.2008.16578.x.

Pintor LM, Sih A, Kerby JL. 2009. Behavioral correlations provide a mechanism for explaining high invader densities and
increased impacts on native prey. Ecology. 90(3):581–587. doi:10.1890/08-0552.1.

Polverino G, Bierbach D, Killen SS, Uusi-Heikkilä S, Arlinghaus R. 2016. Body length rather than routine metabolic rate
and body condition correlates with activity and risk-taking in juvenile zebrafish *Danio rerio*. Journal of Fish Biology.
89(5):2251–2267. doi:10.1111/jfb.13100.

Rasmussen JE, Belk MC. 2012. Dispersal behavior correlates with personality of a North American fish. Curr Zool.
58(2):260–270.

Réale D, Reader SM, Sol D, McDougall PT, Dingemanse NJ. 2007. Integrating animal temperament within ecology and
evolution. Biol Rev Camb Philos Soc. 82(2):291–318. doi:10.1111/j.1469-185X.2007.00010.x.

Schröder A, Kalinkat G, Arlinghaus R. 2016. Individual variation in functional response parameters is explained by body
size but not by behavioural types in a poeciliied fish. Oikos.(1). doi:10.1007/s00442-016-3701-7.

Seebacher F, Beaman J, Little AG. 2014. Regulation of thermal acclimation varies between generations of the short-lived
mosquitofish that developed in different environmental conditions. Funct Ecol. 28(1):137–148. doi:10.1111/1365-
2435.12156.

Seebens H, Blackburn TM, Dyer E, Genovesi P, Hulme PE, Jeschke JM, Pagad S, Pyšek P, Winter M, Arianoutsou M, et
al. 2017. No saturation in the accumulation of alien species worldwide. Nat Comm.:1–9. doi:10.1038/ncomms14435.

Seebens H, Blackburn TM, Dyer EE, Genovesi P, Hulme PE, Jeschke JM, Pagad S, Pyšek P, Kleunen M van, Winter M,
et al. 2018. Global rise in emerging alien species results from increased accessibility of new source pools. PNAS.
115(10):E2264–E2273. doi:10.1073/pnas.1719429115.

Shine R, Brown GP, Phillips BL. 2011. An evolutionary process that assembles phenotypes through space rather than
through time. PNAS. 108(14):5708–5711. doi:10.1073/pnas.1018989108.

Sih A, Bell AM. 2008. Insights for Behavioral Ecology from Behavioral Syndromes. In: Advances in the Study of
Behavior.   Vol.   38.   Elsevier.   p.   227–281.   [accessed   2020   Feb   5].
https://linkinghub.elsevier.com/retrieve/pii/S0065345408000053.

Sih A, Cote J, Evans M, Fogarty S, Pruitt J. 2012. Ecological implications of behavioural syndromes. Ecol Lett.
15(3):278–289. doi:10.1111/j.1461-0248.2011.01731.x.

Smith BR, Blumstein DT. 2010. Behavioral types as predictors of survival in Trinidadian guppies (*Poecilia reticulata*).
Behav Ecol. 21(5):919–926. doi:10.1093/beheco/arq084.

Spiegel O, Leu ST, Bull CM, Sih A. 2017. What's your move? Movement as a link between personality and spatial
dynamics in animal populations. Ecol Lett. 20(1):3–18. doi:10.1111/ele.12708.

Spiegel O, Leu ST, Sih A, Godfrey SS, Bull CM. 2015. When the going gets tough: behavioural type-dependent space
use in the sleepy lizard changes as the season dries. Proc R Soc B. 282(1819):20151768. doi:10.1098/rspb.2015.1768.

Stamps J, Groothuis TGG. 2010. The development of animal personality: relevance, concepts and perspectives. Biol
Rev Camb Philos Soc. 85(2):301–325. doi:10.1111/j.1469-185X.2009.00103.x.

Thorlacius M, Hellstrom G, Brodin T. 2015. Behavioral dependent dispersal in the invasive round goby *Neogobius*
melanostomus depends on population age. Curr Zool. 61(3):529–542. doi:10.1093/czoolo/61.3.529.

Uchida K, Suzuki K, Shimamoto T, Yanagawa H, Koizumi I. 2016. Seasonal variation of flight initiation distance in
Eurasian red squirrels in urban versus rural habitat. J Zool. 298(3):225–231. doi:10.1111/jzo.12306.

Wilson ADMM, Godin J-GJGJ. 2009. Boldness and behavioral syndromes in the bluegill sunfish, *Lepomis macrochirus*.
Behav Ecol. 20(2):231–237. doi:10.1093/beheco/arp018.

Wolf M, Weissing FJ. 2012. Animal personalities: consequences for ecology and evolution. Trends Ecol Evol. 27(8):452–
461. doi:10.1016/j.tree.2012.05.001.

Wong BBM, Candolin U. 2015. Behavioral responses to changing environments. Behav Ecol. 26(3):665–673.
doi:10.1093/beheco/aru183.

520