The obligate intracellular bacterium *Orientia tsutsugamushi* differentiates into a

developmentally distinct extracellular state

Sharanjeet Atwal[b*], Jantana Wongsantichon[a*], Suparat Giengkam[a*], Kittirat Saharat[a], Yanin

Jaiyen Pittayasathornthun[a], Suthida Chuenklin[b], Loo Chien Wang[e,f], Taerin Chung[c], Hyun Huh[c],

Sang-Hyuk Lee[c,d], Radoslaw M. Sobota[e,f] and Jeanne Salje[a, b, g#]

Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol

University, Bangkok, Thailand[a]

Public Health Research Institute, Rutgers the State University of New Jersey, Newark, USA[b]

Institute for Quantitative Biomedicine, Rutgers University, Piscataway, New Jersey 08854, USA[c]

Department of Physics and Astronomy, Rutgers University, Piscataway, New

Jersey 08854, USA[d]

Functional Proteomics Laboratory, Institute of Molecular and Cell Biology, Agency for Science,

Technology and Research (A*STAR), 138673, Singapore[e]

SingMass - National Mass Spectrometry Laboratory, Institute of Molecular and Cell Biology,

Agency for Science, Technology and Research (A*STAR), Singapore[f]

Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of

Oxford, Oxford, United Kingdom[g]

* These authors contributed equally

[#] Address correspondence to Jeanne Salje: js2522@rutgers.edu

Abstract

Orientia tsutsugamushi (Ot) is an obligate intracellular bacterium in the family Rickettsiaceae

that causes scrub typhus, a severe mite-borne human disease. Its mechanism of cell exit is

unusual amongst Rickettsiaceae, as Ot buds off the surface of infected cells enveloped in

plasma membrane. Here, we show that Ot bacteria that have budded out of host cells are in a

distinct developmental stage compared with intracellular bacteria. We refer to these two stages

as intracellular and extracellular bacteria (IB and EB, respectively). These two forms differ in

physical properties: IB is elongated, and EB is round. Additionally, IB has higher levels of

peptidoglycan and is physically robust compared with EB. The two bacterial forms differentially

express proteins involved in bacterial physiology and host-pathogen interactions, specifically

those involved in bacterial dormancy and stress response, secreted bacterial effectors, and

outer membrane autotransporter proteins ScaA and ScaC. Whilst both populations are

infectious, entry of IB Ot is sensitive to inhibitors of both clathrin-mediated endocytosis and

macropinocytosis, whereas entry of EB Ot is only sensitive to a macropinocytosis inhibitor. Our

identification and detailed characterization of two developmental forms of Ot significantly

advances our understanding of the intracellular lifecycle of an important human pathogen.

Keywords

Orientia tsutsugamushi, Rickettsiales, obligate intracellular bacteria, intracellular pathogens

Author Summary

Orientia tsutsugamushi (Ot) is a bacterial pathogen that causes scrub typhus, a mite-transmitted

human disease. This illness is traditionally known to be endemic in the Asia-Pacific, but recent

reports of *Orientia*-like organisms from the Middle East, Africa, and Latin America suggest that it

may be globally distributed. Scrub typhus is associated with high mortality if not treated promptly

with appropriate antibiotics. Ot is a highly specialized bacterium that can only replicate within

living cells, either within the mite vector or in mammalian or human hosts. Ot exits infected cells

using a unique mechanism that involves budding off the surface of infected cells. We have

discovered that this unusual aspect of its lifecycle involves the bacteria themselves

differentiating into a distinct growth form. Different growth forms have not been described in

other members of the family Rickettsiaceae, and no other family members have been shown to

bud out of host cells in a manner similar to Ot. We find that the two forms of Ot, which we refer

to as intracellular and extracellular bacteria (IB and EB respectively), differ in physical properties

and protein expression and infect cells through different mechanisms. The identification of

structurally and functionally distinct forms of Ot elucidates a vital aspect of this pathogen's

intracellular life cycle. The two forms are likely to have different antibiotic susceptibilities,

therefore our findings may advance the development of novel interventions aimed at inhibiting

Ot growth in scrub typhus patients.

Main

Orientia tsutsugamushi (Ot) is an obligate intracellular Gram-negative bacterium that causes the

mite-transmitted human disease scrub typhus. This condition is one of the most severe

rickettsial diseases; it is associated with a median 6% mortality in untreated patients[1]. Scrub

typhus is a leading cause of severe febrile illness in the Asia-Pacific, a region containing two-

thirds of the world's population. Recent reports from Latin America, the Middle East, and Africa

suggest that the disease is more globally distributed[2]. Knowledge of the fundamental biology of

this organism lags behind that of pathogens of equivalent prevalence and severity. Historical

underestimations of Ot incidence, its classification as a biosafety level 3 pathogen, its fastidious

growth in cultured cells, and the fact that it remains genetically intractable[3] likely account for the

paucity of studies focused on this pathogen.

The order Rickettsiales contains two major families: Anaplasmataceae and Rickettsiaceae[4].

Members of the Anaplasmataceae, which include *Anaplasma* spp. and *Ehrlichia* spp., reside

within membrane-bound vacuoles inside host cells. The developmental cycle of certain

Anaplasmataceae family members is biphasic and involves switching between a non-infectious

intracellular replicative form (reticulate body) and an infectious extracellular form (elementary

body). While members of the Rickettsiaceae, namely *Rickettsia* spp. and Ot, do not appear to

differentiate into distinct forms during their intracellular lifecycle, Ot is unique amongst the

characterized Rickettsiaceae in budding off the surface of infected host cells in a manner

reminiscent of enveloped viruses[5]. This observation led us to investigate whether budded Ot is

in a distinct developmental stage compared with intracellular counterparts.

Ot has been shown to enter host cells using the zipper-like invasion mechanism, which is

common to many Rickettsiales members[4]. This process is initiated by binding of bacterial

surface adhesins to receptors expressed on mammalian cells. Adhesins such as TSA56, ScaA,

and ScaC[6-8] bind to host cell fibronectin and syndecan-4; these proteins then interact with

integrin α5ß1, leading to activation of intracellular signaling that culminates in actin remodeling

and clathrin-mediated endocytosis[9-11]. It is important to emphasize that these studies were

entirely focused on intracellular Ot, as they were performed on bacteria isolated from host cells

by mechanical lysis of infected cells. However, the re-entry mechanism utilized by bacteria that

have budded out of cells may be influenced by the presence of an enveloping host cell

membrane and/or its differentiation into a distinct state.

Here, we show that Ot exists in distinct intracellular (IB) and extracellular (EB) bacterial

populations. We report the characterization of these two states in terms of their physical

properties, metabolic activity, and gene expression profiles and demonstrate that they enter

host cells using different mechanisms.

Results

IB and EB forms of Ot differ in physical properties and metabolic activity.

Ot is typically described as a coccobacillary bacterium. Its small size and propensity to aggregate make it difficult to obtain clear images of this bacterium using conventional light microscopy. To circumvent this challenge, we labelled Ot with an antibody against the abundant surface protein TSA56 and employed stochastic optical reconstruction microscopy (STORM) for high-resolution imaging. We first investigated the shape of bacteria located inside and outside host cells (Fig. 1A, B and Supp. Fig. 1). Since Ot begins to appear in the supernatant of infected L929 cells at high levels five days post infection (Supp. Fig. 2) we focused on the shape of EB Ot beginning five days post infection, whilst we measured the shape of IB Ot throughout the infection cycle. We found that EBs were approximately round in shape with a diameter of about 1.2 μm, and they were often aggregated into clusters. In contrast, IBs adopted diverse round and elongated shapes and varied in dimensions; some bacteria were 3-4 μm in length. Additionally, IB size varied during the course of infection. Initially, and up to one-day post-infection, IBs were similar in length to EBs. At later stages, *i.e.,* 2-7 days following infection, IBs were characterized by longer lengths. These kinetic changes in shape correlated with Ot growth dynamics whereby replication starts 24-48 hours post infection[12]. These differences in shape provide evidence for the existence of two separate bacterial forms.

We next investigated potential differences in bacterial cell wall composition. We previously showed that Ot builds a low-abundance peptidoglycan-like structure[13]. Thus, we assessed relative peptidoglycan levels in IB and EB Ot forms using a NOD-1 reporter assay (Fig. 1C). We detected significantly more peptidoglycan in IB cells compared with EB cells. This finding may account for the observed differences in shape (Fig. 1A) as peptidoglycan is required to build elongated cells.

In related structural studies, we capitalized on our previous finding that the abundant surface

protein TSA56 is involved in forming disulfide bonds with itself or other proteins, leading to the

formation of aggregates[13]. These complexes can be identified as smears following sample

preparation in the absence of reducing agents and evaluation by immunoblot analyses. In this

study, we found evidence for substantially higher disulfide bond formation in EB Ot than IB Ot,

as determined by the presence of a high molecular weight smear (Fig. 1D). We hypothesize that

this crosslinked structure confers EB Ot with additional structural rigidity in the absence of

peptidoglycan. These experiments were performed in the presence of iodoacetamide, an agent

that covalently modifies cysteine residues and prevents them from forming new bonds, thus

ensuring that the cross-links were not formed during sample preparation and processing.

Our next goal was to assess potential changes in genome copy number as, during rapid growth,

bacteria such as *E. coli* carry multiple genome copies within a single cell. To address this issue,

we determined bacterial cell numbers using fluorescence microscopy and bacterial genome

copies using qPCR. We found no differences: Ot IB and EB harbored just over one estimated

genome copy per cell (Fig. 1E). This measurement is an approximation that suffers limitations

associated with absolute quantification using qPCR and microscopy, but any such errors are

expected to be constant between the IB and EB measurements.

Finally, we asked whether IB and EB Ot forms differed in their metabolic activity. Ot is an

obligate intracellular bacterium with limited central metabolic and biosynthetic capability[14] and a

reliance on host cells for carbon and nitrogen sources and metabolic intermediates. These

observations led us to hypothesize that EB bacteria are metabolically inactive owing to their

extracellular location. To test this, we used a microscopy-based assay whereby growing

bacteria are exposed to a clickable methionine analog L-Homopropargylglycine (HPG) that is

subsequently labeled with a fluorophore[15] (Fig. 1F). Using this methodology, we observed that

95 % of IB were translationally active five days post-infection, compared with 0% of EB (Fig.

1G). It is possible that EB Ot undergo some residual metabolic activity that is not detected by

our assay of protein translation. The fact that not all IB were labelled with HPG indicated that

this result did not simply reflect a difference in bacterial permeability between the IB and EB

forms. These results provide an additional line of evidence to support the existence of two

fundamentally different Ot states.

IB and EB forms of Ot exhibit different gene and protein expression profiles.

We next investigated whether the differences in physical properties and metabolic activity

observed in IB and EB forms of Ot were accounted exclusively by physical location relative to

host cells. Exposure of EB forms to an osmotically different extracellular environment combined

with the lack of nutrient availability might account for our observations. However, a second

possibility is that EB forms arise by differentiation of IB into a biochemically distinct state. We

reasoned that bacteria in the same developmental stage but differing in position relative to host

cells would exhibit almost identical protein profiles. To test this, we performed shotgun mass

spectrometric analyses on isolated IB and EB Ot forms. We found that whilst the correlation

coefficient of proteins detected in IB and EB isolated four days post-infection was strong

(correlation coefficient = 0.843) (Supp. Fig. 3) there were several proteins that accumulated to

different levels in the two populations. A complete list of the relative levels of proteins detected

in IB and EB is provided in Supp. Table 1. Pathway analyses revealed that IB Ot was enriched

in proteins involved in protein synthesis (Fig. 2A). In contrast, EB Ot displayed high expression

of SpoT and RpoH, two proteins involved in the stringent response (Fig. 2B). In order to validate

these results, we generated antibodies against four proteins that were present at different levels

in IB and EB Ot: SpoT, RpoH, RpoD and CtrA. We carried out immunoblot analysis of IB and

EB Ot and found that both SpoT and RpoH were present at higher levels in EB than IB, whilst

RpoD and CtrA were present at higher levels in IB Ot (Fig. 2C), consistent with the proteomics

analysis.

We next investigated whether EB and IB Ot exhibited differences in transcript expression levels.

These experiments required the propagation of large amounts of EB because this form of Ot is

metabolically inactive and thus harbors very low mRNA levels. We selected 11 genes, three of

which (*curA, trxB, sodB*) were expressed at similar levels in our proteomic studies comparing

EB and IB. Additionally, we quantified mRNA expression of four genes encoding proteins

enriched in IB Ot (*yccA, nuoB, dapA, scaA*) and four genes enriched in EB Ot (*obgE, spoT,*

tsa56, rpoH). We isolated RNA from IB and EB bacteria four days post-infection and assessed

mRNA levels by qRT-PCR (Fig. 2D). We found no correlation between mRNA and protein levels

in multiple cases. For example, *rpoH* transcript levels were similar in IB and EB Ot, but protein

levels were higher in EB. These observations provide additional evidence supporting the

existence of molecularly distinct bacterial forms and suggest that transcriptional and post-

transcriptional regulatory mechanisms participate in the differentiation process.

Surface proteins are present at different relative levels on IB and EB forms of Ot.

Our differential proteomics analyses revealed that IB and EB have different levels of specific

surface proteins (Fig. 3A). Surface proteins play a central role in Ot function for two reasons.

First, unlike most Gram-negative bacteria, Ot lacks lipopolysaccharide, which generally serves

as a membrane shield. Second, Ot is a cytosolic bacterium, and its surface is in direct contact

with cytoplasmic host components. Ot harbors genes encoding four widely conserved

autotransporter proteins: ScaA, ScaC, ScaD, and ScaE. Using our proteomics dataset, we

detected ScaA and ScaC, but not ScaD or ScaE. ScaA was slightly enriched in IB relative to EB

Ot (1.5X), while ScaC was enriched in EB (1.87X). Ot also harbors genes encoding three

immunogenic surface proteins that dominate the humoral response in human patients, TSA22,

TSA47, and TSA56. We found that TSA22 was enriched in IB relative to EB Ot (3.2X) while

TSA47 and TSA56 were not present at significantly different levels (1.14X and 1.32X increase in

EB compared with IB, respectively). To verify our observations, we generated polyclonal

antibodies against ScaA and ScaC and determined their relative levels in IB and EB bacteria

using immunoblot analyses (Fig. 3B and Supp. Fig. 4). Our experiments up to this point were

performed in mouse fibroblasts (L929) because Ot grows readily in these cells. Moreover, L929

fibroblasts are suitable for characterizing bacterial properties that are not affected by the cellular

environment. However, for our validation studies, we also investigated bacteria grown in primary

human umbilical vein endothelial cells (HUVEC) as a more physiologically relevant model of Ot

infection in humans. Our immunoblot studies in both L929 and HUVECs confirmed enrichment

of ScaA in IB Ot and higher levels of ScaC in EB compared to IB Ot, in agreement with our

mass spectrometric analyses.

ScaA and ScaC as markers of EB and IB forms of Ot

Our observation that 95% of IB is translationally active five days post-infection, compared with

0% of EB, combined with the fact that these Ot forms display opposite relative levels of ScaA

and ScaC led us to investigate whether these surface proteins can be used as *bonafide* markers

of metabolically active IB and metabolically quiescent EB Ot forms, respectively. We used

immunofluorescence microscopy to identify bacteria expressing ScaA and ScaC and then

quantified the number of translationally active ScaA- and ScaC-positive bacteria by measuring

incorporation of HPG. We found (Fig. 3C and Supp. Fig. 7A) that ScaA-positive bacteria, in

contrast to ScaC-positive Ot, were mostly translationally active. We then quantified the number

of ScaA/ScaC positive bacteria inside and outside cells and confirmed that ScaC and ScaA

levels were higher in EB and IB Ot, respectively (Fig. 3D, E and Supp. Fig. 5, 6, 7B). Together,

these results show that ScaA and ScaC are present at different relative levels depending on

bacterial location relative to host cells: metabolically active IB exhibit higher levels of ScaA

relative to EB Ot. Conversely, levels of ScaC in metabolically inactive EB is high relative to that

of IB forms of Ot.

The observation that EB exhibits high *scaA* transcript levels (Fig. 2D) but low levels of ScaA

protein led us to hypothesize that ScaA is translated into protein soon after entry into host cells,

using pre-formed mRNA. We quantified the number of IB Ot expressing ScaA and ScaC at 1-

and 3-hours post-infection and found a decrease in ScaC levels and an increase in ScaA levels

as the infection proceeded (Fig. 3F and Supp. Fig. 7C). These results show that ScaA

accumulates during the first 3 hours after infection, suggesting that *scaA* transcripts are made in

EBs in anticipation of new infections. These results point to the existence of regulatory

mechanisms that control levels of ScaA and ScaC on the surface of Ot cells, depending on

bacterial location and stage of host infection.

IB and EB Ot populations are infectious but differ in physical stability and mechanisms

of entry into host cells.

Previous studies have shown that specific Anaplasmataceae differentiate into distinct

intracellular and extracellular forms and that only the extracellular form is effective at infecting

host cells[16]. This observation led us to investigate whether the ability to infect host cells differs in

IB compared with EB Ot. Growth curve measurements revealed that both bacterial forms could

infect L929 and HUVEC cells and replicate with similar dynamics (Fig. 4A and Supp. Fig. 8).

However, IB appears to retain infectivity more efficiently than EB following physical and osmotic

challenges. Specifically, when we subjected IB and EB Ot to physical stress using a bead mill

for 1-2 minutes, we found that EB displayed a reduction in growth that was dependent on the

time of exposure to stress; in contrast, IB was unaffected by this treatment (Fig. 4B).

Furthermore, osmotic shock resulted in a significantly greater reduction in viability in EB

compared with IB (Fig. 4C). Together, these results demonstrate that IB is physically more

robust than EB; this property is likely due to differences in peptidoglycan content, as described

above.

To investigate whether EB and IB forms of Ot enter host cells using the same invasion

mechanism, we pre-treated HUVEC cells with the clathrin-mediated endocytosis inhibitor

dynasore or the macropinocytosis inhibitors amiloride and zoniporide. We then infected the cells

with IB or EB Ot and quantified the number of intracellular bacteria 3 hours post-infection using

confocal microscopy (Fig. 4D, E). The entry of IB Ot into the cells was blocked by dynasore,

thus confirming the participation of clathrin-mediated endocytosis in this process. IB entry was

also inhibited by amiloride and zoniporide, suggesting additional involvement of

macropinocytosis in bacterial uptake. In contrast, EB Ot entry into cells was not affected by

dynasore treatment but was completely inhibited by pre-incubation with amiloride or zoniporide.

To confirm that EB Ot did not use clathrin-mediated endocytosis to enter cells, we treated

HUVEC and L929 cells with siRNA against clathrin. Control microscopy experiments

demonstrated that clathrin-targeting siRNA reduced the levels of clathrin foci in uninfected cells

(Supp. Fig. 9). We found that entry of IB form Ot required clathrin, whilst entry of EB form Ot did

not (Fig. 4F, G and Supp. Fig. 10). Together these results demonstrate that IB form Ot uses

clathrin-mediated endocytosis to enter cells whilst EB form Ot does not.

To further demonstrate that IB and EB Ot were entering host cells using distinct mechanisms,

we examined the role of the surface adhesins TSA56 and ScaC on bacterial entry. These are

outer membrane proteins that have been shown to be involved in entry of Ot into host cells, in

experiments carried out using what we now describe as IB Ot stage bacteria[6,7]. We used

antibodies targeting ScaC and TSA56 and determined whether these antibodies would block the

entry of Ot into host cells. Our results with IB Ot replicated published studies and showed that

blocking ScaC and TSA56 using antibodies led to a strong reduction in bacterial entry (Fig. 4H).

With EB Ot, however, only a minimal effect was observed (Fig. 4H). This result supports a

model in which IB Ot enters host cells using adhesin-induced clathrin-mediated endocytosis,

whilst in EB Ot the surface proteins are shielded by host derived membrane and use a different

mechanism of entry.

Discussion

Our studies have unveiled a new feature of the obligate intracellular bacterium Ot. This

organism's unique exit strategy, which involves budding off the surface of infected cells, is

associated with differentiation into a molecularly and functionally distinct developmental state

(Fig. 5). While unusual in the family Rickettsiaceae, this property is reminiscent of the biphasic

lifecycle of multiple species in the sister family Anaplasmataceae[4] and the unrelated obligate

intracellular bacteria *Chlamydia* spp.[17]. However, in contrast to these organisms, Ot resides in

the host cytosol in naked form rather than in a membrane-bound vacuole. To our knowledge,

this is the first description of an intracellular bacterium living freely in the host's cytosol and

having a biphasic lifecycle. Our findings have important implications for our understanding of Ot

biology and unveil opportunities to develop better therapeutic strategies to treat scrub typhus.

The physical properties of prokaryotic cell walls are crucial for survival. These stiff structures

maintain cellular shape, protect the interior, counteract osmotic pressure, and provide structural

strength through peptidoglycan layers. Our studies revealed that EB and IB forms of Ot differ in

peptidoglycan content. Lower levels in EB compared with IB make this form more susceptible to

physical and osmotic stress. There is a long-standing question regarding the role of

peptidoglycan in obligate intracellular bacteria. These organisms reside in an osmotically

protective niche and are under selective pressure to reduce the abundance of immunogenic

pathogen-associated molecular patterns (PAMP)s[18]. In the case of pathogenic *Chlamydia*

species, it has been shown that peptidoglycan is retained only at the septum and is required for

cell division[19]. Higher peptidoglycan levels in IB Ot *vs.* EB Ot point to the retention of

peptidoglycan for growth and division rather than structural integrity maintenance.

The observed differences in physical properties between EB and IB forms are complemented by

measurable differences in the accumulation of certain proteins, suggesting that each form

performs unique functions or responds to specific challenges. Proteomics analyses in EB Ot

revealed increased levels of heat shock sigma factor RpoH and the alarmone (p)ppGpp

synthetase SpoT associated with the stringent response[20]. These results suggest that the EB

form of Ot shares features with the dormant state of other bacteria. Notably, Ot is the only

reported member of the Rickettsiaceae to encode a full-length *spoT* gene[4]. This exclusive

feature is in harmony with the fact that developmental differentiation of Ot is highly unusual in

members of this family. The changes associated with differentiation into the EB state likely

serve one or both of the following purposes: ensuring survival in the extracellular state and

priming the bacteria for reinfection of new cells. Indeed, the observation that *scaA* transcripts

are abundant in the EB form and that ScaA protein is synthesized early upon entry into host

cells supports the idea that EB Ot is primed for infection.

Comparing the protein and RNA transcript levels of selected genes revealed a poor correlation

for many genes. A potential explanation is that RNA levels are low in the metabolically inert EB

form, which does not reflect newly translated proteins. Ot strain UT76 encodes only 8 predicted

transcription factors and two sigma factors, RpoD and RpoH[21]. Thus, it is possible that gene

expression is not strictly regulated at the transcriptional level but that other forms of regulation

are prevalent in Ot. We recently showed high antisense transcription levels in Ot, especially in

genes associated with the rampant integrative conjugative element RAGE, which dominates the

Ot genome[22]. There, we found that high antisense transcription correlates with low protein levels

for a particular gene, pointing to this mechanism as a central regulatory strategy in Ot. Gene

expression in the Ot-related, free-living alphaproteobacterium *Caulobacter crescentus* is highly

regulated by proteolysis[23]. Mechanisms based on differential proteolysis may therefore also play

an essential role as critical determinants of protein levels in Ot.

Our observation that EB Ot enters cells through pinocytosis, whereas IB Ot utilizes endocytosis

and, to a lesser extent, pinocytosis resolves a long-standing gap in the field as it was unclear

how budded bacteria enveloped in plasma membrane enter host cells. Physiologically, IB entry

into host cells may be necessary during trafficking into adjacent cells, as previously described

for spotted fever group *Rickettsia* spp.[24]. Additionally, low levels of cell lysis may release IB Ot

forms that can then infect new host cells. The different mechanisms utilized by EB and IB to

enter cells suggest that strategies to escape from the endolysosomal pathway will differ

between the two Ot forms. It is unclear whether and how the differing transcriptional and

proteomic profiles at the point of infection will affect early host-pathogen interactions. Our

finding that IB and EB's growth dynamics are similar suggests that early differences do not

persist through the infection cycle.

In summary, our finding that Ot differentiates into molecularly and functionally distinct forms

provides a novel framework to study the infection cycle of this important obligate intracellular

pathogen and opens avenues to investigate the physiological significance of the distinct

populations characterized.

Fig. 1: Intracellular and extracellular forms of Ot are physically and metabolically distinct.

A. Representative STORM images of IB (5 d.p.i.) and EB (7 d.p.i.) Ot in L929 cells. Bacteria were labelled with a monoclonal antibody against surface protein TSA56 (magenta). Additional images including DIC images of host cells are provided in Supp. Fig. 1. Scale bar = 5 µm. B. Length quantification of individual IB and EB Ot as imaged by STORM. H = hours, D = days post infection. The lengths of 50 individual IB (defined as colocalising with host cells) and EB form (defined as not colocalising with host cells) Ot were measured. Statistical significance was determined using a one-way ANOVA followed by Dunnett's multiple comparisons. C. Quantification of relative peptidoglycan levels through activation of NOD1 reporter cells, as measured by secreted alkaline phosphatase activity. EB Ot induced less activation than IB, indicating lower levels of peptidoglycan. Identical amounts of IB and EB Ot were used, taken from L929 cells 5 days post infection. The results of six independent replicates, carried out on two different days, are shown and the statistical significance was determined using a Mann-Whitney test. Positive control = meso-diaminopimelic acid, negative control = PBS. D. Immunoblot showing the formation of disulfide cross-linked aggregates of the surface protein TSA56, observed in the absence of the reducing agent beta-mercaptoethanol (BME). This effect was stronger in EB compared with IB Ot, taken five days post infection from L929 cells. The expected molecular weight of TSA56 (56 kDa) is indicated with an arrow, and a dashed arrow depicts higher-order aggregates. E. Genome copy number per cell as measured by qPCR; bacterial cell number was determined by immunofluorescence microscopy. Quantification of 10 individual imaging fields is shown, and statistical significance determined using a Mann-Whitney test. F. Confocal microscopy image of an L929 cell infected with Ot for five days and labelled with the clickable methionine analog homopropargylglycine (HPG, green) and an antibody against surface protein TSA56 (red). DNA (Hoechst) is shown in blue. The host cell outline is showed by a dashed line, and an arrow indicates a metabolically inactive bacterium located outside the host cell. G. Quantification of metabolic activity, as measured by a detectable HPG

signal, in IB and EB Ot. L929 cells were infected with Ot for five days, fixed, and labelled with

TSA56 and HPG. Bacteria were identified as being IB or EB, based their position relative to host

cells, and the percentage of each population that was HPG-positive was quantified. The

percentage positivity for ten randomly imaged field is shown here, and this data is

representative of >3 independent repeats of this experiment. Groups were compared using a

Mann-Whitney test. Statistical significance for all data in this figure was determined using

Graphpad Prism software. * p-value ≤ 0.05, ** p-value ≤ 0.01, *** p-value ≤ 0.001, **** p-value ≤

0.0001, ns = not significant.

Fig. 2: Intracellular and extracellular forms of Ot have different levels of certain proteins.

A. Pathway analysis of proteins present at >2X difference between IB and EB forms of Ot (60

proteins in IB, 47 proteins in EB). Bacteria were grown in L929 cells at an MOI of 410:1 and

harvested four days post-infection. IB and EB Ot were isolated and analysed by shotgun

proteomics. B. SpoT and RpoH are present at higher levels in EB compared with IB Ot as

measured by proteomics described in A. C. Immunoblot analysis showing relative levels of four

Ot proteins SpoT, RpoD, RpoH and CtrA in IB and EB Ot. Loading was normalised by

measuring bacterial genome copy number using qPCR and loading the same number of IB and

EB Ot in each pairwise comparison. Samples taken from the same experiment are used for the

four antibodies in this figure, with the same amount loaded in each immunoblot. Uncropped

immunoblot images showing antibody specificity are given in Supp. Fig. 4. This data confirms

the trends observed in the proteomics analysis for these four proteins. D. RTqPCR analysis

showing relative levels of 11 genes expressed in EB and IB Ot. Bacteria were isolated from

L929 cells four days post-infection at an MOI of 672:1, 300:1, and 560:1. The figure depicts

results from three independent replicates. Gene expression levels were normalised using three

genes: *mipZ, ank12*, and *16S.* Mean and standard deviation values are shown. The ratio of

protein levels in EB/IB, as found in the proteomics experiment in 2A, is given above each gene

for comparison.

Fig. 3: Autotransporter proteins ScaA and ScaC are differentially expressed on IB and EB

forms of Ot.

A. Volcano plot showing differential expression of surface proteins in proteomics analysis of IB

and EB forms of Ot. We observed higher expression of ScaC, TSA56, and TSA47 in EB whilst

IB Ot expressed higher ScaA and TSA22 levels B. Immunoblot analysis of IB and EB Ot from

infected L929 and HUVEC cells confirming differential expression of ScaA, and ScaC between

IB and EB forms. Loading was normalised by measuring bacterial genome copy number using

qPCR and loading the same number of IB and EB Ot in each pairwise comparison. Uncropped

immunoblot images showing antibody specificity are given in Supp. Fig. 4. C. Quantification of

translational activity, as determined by a positive confocal microscopy signal observed with the

clickable methionine probe L-homopropylgylglycine (HPG). HUVEC cells were infected with Ot

for five days, then fixed and labelled with HPG and anti-ScaA or anti-ScaC antibodies.

ScaA/ScaC-positive bacteria were scored as being positive or negative for HPG labelling. D.

Confocal microscopy images of HUVEC cells infected with Ot for five days and labelled for DNA

(Hoechst, blue), translational activity (HPG, green), ScaA (anti-ScaA antibody, magenta) and

ScaC (anti-ScaC antibody, red). Scale bar = 10 µm. Examples of extracellular bacteria labelled

with ScaC are shown with a white arrow. More images as well as equivalent experiments in

L929 cells are shown in Supplementary Figures 5-7. E. Quantification of confocal micrographs

of HUVEC cells infected with Ot and labelled as in 3D. The percentage of IB and EB Ot at five

days post-infection that were positively labelled with anti-ScaA and -ScaC antibodies was

counted. F. Quantification of ScaA- and ScaC-positive IB bacteria inside HUVEC cells one- and

three-hours post-infection, showing an increase in ScaA labelling and a decrease in ScaC

labelling. Throughout this figure quantification was carried out by randomly imaging 30 fields

across three independent experiments. Graphs show individual values and means. Statistical

significance was calculated using an unpaired t test with Graphpad Prism software. * p-value ≤

0.05, ** p-value ≤ 0.01, *** p-value ≤ 0.001, **** p-value ≤ 0.0001.

Fig. 4: Both IB and EB populations are infectious but differ in physical stability and mechanism of entry into cells.

IB and EB Ot were isolated from infected L929 cells five days post-infection and grown in fresh

L929 cells in a 24-well plate. The number of intracellular bacteria was then quantified at 3-, 24-,

96-, and 168-hours post-infection using qPCR against *tsa47*, a single copy Ot gene. Doubling

times of IB and EB Ot were similar. Similar results were found in HUVEC cells (Supp. Fig. 8). B.

IB and EB were exposed to physical stress using a bead mill for one or two min; growth was

quantified as in 4A. The bead mill did not affect the growth rate of IB Ot but reduced that of EB

Ot in an exposure-dependent manner. C. IB and EB were exposed to osmotic shock by

suspending the bacteria in water for five min and then growing them in fresh L929 cells.

Bacterial copy number after seven days was quantified by qPCR. EB was more susceptible to

osmotic shock than IB Ot, as shown by a more significant decrease in bacterial number.

Statistical significance was determined using a Mann-Whitney test. For panels A-C each

datapoint was taken from a separate well from a 24-well plate. Images (D.) and quantification

(E.) of the effect of endocytosis inhibitors on entry of IB and EB Ot into HUVEC cells. IB and EB

were isolated from L929 cells five days post-infection and then seeded onto fresh HUVEC cells

that had been treated with inhibitors of clathrin-mediated endocytosis (100 µM dynasore) or

macropinocytosis (50 µM amiloride or 100 µM zoniporide). Cells were fixed three hours post-

infection. Dynasore inhibits entry of IB but not EB, indicating that clathrin-mediated endocytosis

is the major mode of entry of this population, whilst amiloride and zoniporide inhibited entry of

both IB and EB. This observation points to a role for macropinocytosis, but not clathrin-mediated

endocytosis, in EB's entry into mammalian cells. Red = actin (phalloidin), blue = Hoechst (DNA),

green = Ot (anti-TSA56 antibody). Scale bar = 10 µm. Similar results were found in L929 cells

(Supp. Fig. 10). Images (F) and quantification (G) of the effect of 0.2 µM siRNA against clathrin

on entry of EB and IB Ot into HUVEC cells. Experiments were carried out as in D. and E.

Quantification of E. and G. were carried out as follows: 30 fields were randomly imaged over 3

independent experiments (10 fields per experiment). In each field the percentage of infected

host cells was counted. Individual values and means are shown here. Statistical significance in

(E) was determined by comparing all groups using one-way ANOVA, followed by a Dunnett's

multiple comparison tests with the control within the IB or EB group, and in (G.) by performing a

Mann Whitney test comparing the scramble control and the clathrin siRNA treated condition for

IB and EB separately, using GraphPad Prism software. H. Blocking the bacterial surface

adhesins TSA56 and ScaC with anti-TSA56 and anti-ScaC antibodies strongly inhibits entry of

IB but not EB Ot in HUVEC cells. Bacterial entry in the presence of vehicle control or antibodies

was determined by quantifying the number of intracellular bacteria per host cell 30 minutes post

infection using immunofluorescence microscopy of bacteria labelled with HPG. Statistical

significance was determined by comparing all groups using one-way ANOVA, followed by a

Dunnett's multiple comparison tests with the control within the IB or EB group. **** $p \leq 0.0001$;

ns = not significant. * p-value ≤ 0.05, ** p-value ≤ 0.01, *** p-value ≤ 0.001, **** p-value $\leq$

0.0001. Throughout figure blue = IB and red = EB.

Fig.5: An overview of the infection cycle of Ot, showing differentiation between IB and EB

forms.

IB replicate in a micro-colony adjacent to the nucleus where they are elongated in shape and

translationally active. Bacteria move to the cell's edge and transition to an EB form that is round

and translationally inactive. Created with BioRender.com B. Table summarizing the differences

between IB and EB forms of Ot.

Methods

Cell culture, bacterial propagation and quantification, and reagents

Proteomics and qRT-PCR experiments were performed using the mouse fibroblast cell line

L929 kindly provided by Dr. Blacksell at the Mahidol Oxford Tropical Medicine Research Unit,

Bangkok, Thailand. For all other experiments, we used L929 cells (ATCC CCL-1), Primary

Human Umbilical Vein Endothelial Cells (HUVEC) (ATCC PCS-100-010), monkey kidney

epithelial Vero cells (ATCC CRL-1586) or human dermal microvascular endothelium HMEC-1

cells (ATCC CRL-3243), all purchased from ATCC. We cultured cells in either DMEM (L929) or

Large Vessel Endothelial Supplement (LVES, HUVEC) supplemented with 10% FBS (Gibco

16140071) at 37ºC and a 5% CO_2 atmosphere. The Karp-like clinical isolate strain of *Orientia*

tsutsugamushi UT76[21] was used throughout the study. For routine propagation, bacteria were

grown in 25- or 75-cm^2 culture flasks, as described previously[12]. We extracted DNA using

alkaline lysis buffer[12], and then assessed bacterial quantities by determining genome copy

numbers relative to a standard curve. We used a primer/probe set to amplify the single copy

gene *tsa47* (tsa47F   5'-TCCAGAATTAAATGAGAATTTAGGAC-3',   tsa47R   5'-

TTAGTAATTACATCTCCAGGAGCAA-3',   tsa47probe   5'-

[6FAM]TTCCACATTGTGCTGCAGATCCTTC[TAM]-3')   or   the   16S   gene   (Rick16SF   5'-

GCTACACGCGTGCTACAATGG-3',   Rick16SR   5'-   TGTGTACAAGGCCCGAGAACG-3',

Rick16Sprobe 5'-[6FAM] ATCGCTAGTAATCGCGGATCAGCATGCC [TAM]-3'). For drug

inhibition assays, dynasore (Sigma D7693) and amiloride (Sigma A3085) were added to L929 or

HUVEC cells 1 hour before infection with Ot.

Fluorescence microscopy

Immunofluorescence labelling and confocal microscopy. Following fixation, cells were

permeabilized by incubation in 0.5% Triton X-100 for 30 min on ice followed by blocking in

1XPBS (pH7.4) containing either 1 mg/ml BSA or sera (diluted 1:20) matching the labeled

antibody's host, for 30 min at room temperature. Incubations with primary antibodies took place

for 1 hour at 37°C. These studies were conducted using in-house-generated antibodies (*i.e.*, rat

monoclonal antibody against TSA56 diluted 1:200; rabbit polyclonal antibody against Ot ScaC

diluted 1:200) or commercial products (*i.e.*, MA1-164, a Thermo Scientific monoclonal mouse

antibody against LAMP1 diluted 1:200). The fixed cells were washed twice with PBS-BSA and

were then incubated with appropriate secondary antibodies (1:1000 dilutions) for 30 min at 37°C

in the dark. We used the following secondary antibodies: goat anti-mouse IgG superclonal Alexa

Fluor 647 conjugate (Thermo Fisher A28180), goat anti-rat Alexa Fluor 488 (Thermo Fisher

A11006), and goat anti-rabbit Alexa Fluor 594 (Thermo Fisher A11012). The nuclear stain

Hoechst (1:1000 dilution) was supplemented to incubations with secondary antibodies.

Phalloidin Alexa Fluor 594 (1:200, ThermoFisher Scientific A12381) was used to detect host

actin. The cells were washed with 1X PBS before supplementation of mounting medium [20 mM

Tris (pH 8.0); 0.5% N-propyl-gallate; 90% glycerol]. Imaging was performed using a Zeiss

Observer Z1 LSM700 confocal microscope with an HBO 100 illuminating system equipped with

a 63x/1.4 Plan-APOCHROMAT objective lens (Carl Zeiss, Germany) and 405 nm, 488 nm, and

555 nm laser lines. In some cases, we used a Leica TCS SP8 confocal microscope (Leica

Microsystems, Germany) equipped with a 63x/1.4 Plan-APOCHROMAT oil objective lens with a

1.4-mm working distance and 405 nm, 488 nm, 552 nm, and 638 nm laser lines.

Fluorescence intensity measurements were performed manually using Fiji ImageJ software. 30

(Fig. 3C) or 100 (Fig. 1G); individual bacteria were quantified by measuring the brightest pixel

intensity within the bacterial area. Images from individual experiments were acquired under

identical laser intensity and acquisition conditions. Length and width measurements were

performed manually using Fiji Image J software.

STORM imaging and analysis

We acquired 3D-STORM data using a homebuilt multi-color Total Internal Reflection

Fluorescence (TIRF) microscope[25] that is based on an inverted microscope (Nikon, Ti-E), a high

NA objective (Nikon, CFI-Apo 100X, NA 1.49), a 405 nm photoactivation laser (Coherent, OBIS

405 LX 100mW), a 488 nm excitation laser (MPB, 2RU-VFL-P-20000642-B1R), and an EMCCD

camera (Andor, iXon Ultra-888). Samples immunolabelled with Alexa 488 were mounted on the

microscope, transferred to STORM buffer [OxyFluor (Oxyrase), 50 mM betamercaptoethanol

(Sigma), 2.5 mM cyclooctatetraene (COT) (Sigma)], and illuminated with a strong (488 nm)

laser to record single-molecule fluorescence blinking events at a frame rate of 25 Hz.

Illumination at 405 nm was intercalated with pulsed 488nm excitation to enhance

photoactivation of Alexa 488 without exacerbating background signal. A cylindrical lens

(Thorlabs, LJ1516RM-A) was placed in front of the camera to determine the z-coordinates of

fluorescent molecules through the elongated point spread function (PSF) methods[26]. Each

STORM image was reconstructed from single-molecule localization data obtained by analysing

30,000 frames with custom Matlab codes. Briefly, bright single fluorescence spots were

identified by thresholding methods, and their 3-D coordinates were determined by 2-D elliptical

Gaussian fitting along with a pre-calibrated conversion table between eccentricity and z-

coordinate. Focus drift was actively stabilized with Perfect Focus System (Nikon) while acquiring

the data, whereas xy-drift was post-corrected using a redundant cross-correlation algorithm[27].

Each localization event was represented as a 2-D Gaussian of 20 nm standard deviation in

STORM images. For the analysis of bacterial morphology, we selected cells that showed clear

cell boundary, manually adjusted the z-coordinate to locate the cell midplane, and measured

long- and short- axes length.

Click labelling. Our metabolic click-labelling was based on the Click-iT HPG Alexa Fluor

protein Synthesis Assay Kit (Click-iT HPG Alexa Fluor Protein Synthesis Assay Kit,

ThermoFisher C10428) strategy. To incorporate L-homopropargylglycine (HPG), we incubated

infected cells with minimal medium (Dulbecco's Modified Eagle Medium, DMEM, ThermoFisher

21013) lacking L-methionine and containing 25 µM HPG, for 30 min at 37°C. Labelled bacteria

were washed twice with PBS containing 1mg/ml BSA, fixed with 1% formaldehyde, and

subsequently permeabilized with 0.5% Triton X-100 for 20 min on ice. After further washing with

PBS containing 1 mg/ml BSA, we supplemented the Click-iT reaction cocktail (Click-iT HPG

Alexa Fluor Protein Synthesis Assay Kit, ThermoFisher C10428) and incubated the cells for 30

min at room temperature in a light protected environment. The Azide dye (Alexa Fluor 488) was

used at a final concentration of 5 µM. Following completion of the click reaction, the cells were

subjected to immunofluorescence labelling and imaging as above.

Gene expression analysis by RT-qPCR

L929 cells were infected with *Orientia tsutsugamushi* UT76 using multiple MOIs to a target MOI

of 400:1 (quantitative PCR on freshly isolated Ot showed variation of actual MOIs ranging from

200:1 to 672:1 across 3 independent infections). Very high MOIs were used due to the low

levels of EB bacteria combined with the fact that RNA levels in EB Ot were very low. Infected

cells were harvested four days post infection and were stored in RNAProtect Bacteria Reagent

(Qiagen 76506) at -80°C. Total RNA was extracted using the Qiagen RNeasy Plus kit (Qiagen

74136) according to the manufacturer's instructions. One µg of total RNA was incubated with

DNaseI (Thermo Fisher Scientific, AM2238) for 16 h at 37°C, and the samples were then

reverse-transcribed using the iScript reverse transcription supermix (Biorad 170–8841) and

random primers. Controls lacking reverse transcriptase demonstrated that the RNA sample

used for cDNA synthesis was not contaminated with genomic DNA. The resulting cDNA was

then used as template for qPCR using primers for *spoT, rpoH, obgE, yccA, nuoB, dapA, curA,*

sodB, trxB, tsa56, and scaA. RT-qPCR was performed using SYBR green qPCR mix (Biotools

10.609). Expression levels were normalized relative to the housekeeping genes *16sRNA,*

ank12_1 and mipZ.

Primer name	Primer sequences (5' to 3')
spoT forward	GGCAGACATAAAGCACCTTAC
spoT reverse	GAAATCTTGAAAGCTGCCAGG
rpoH forward	CTGCCGAAGAGGAATATATGC
rpoH reverse	CCCCTCGGAAACTAAATCAAC
obgE forward	GCACAAAATGGTAGAGCTGG
obgE reverse	GAATCACCAATCTCACCTACAG
yccA forward	GCTATTGCAGGAACCGCTAC
yccA reverse	CTGAAAGCGACATGCCTACA
nuoB forward	GGGCTAGAACACGCTCGTTA
nuoB reverse	GCACAGCTACCCATTGATATC
dapA forward	GGTAGAACATTAGTGATTGCTGG
dapA reverse	GCGTCTTTGATTGCTACTATCC
curA forward	AGTAGGGCAACGAGTTGGC
curA reverse	CAACAGCGCTACTAGCAGC
sodB forward	GTAGTGGTTGGGTCTGGTTG
sodB reverse	GTTCAGCAAATCGCCAATTTG
trxB forward	GATATGCACTGGTGCTCAAG
trxB reverse	CACCTCCTACAACAACTACTC
tsa56 forward	TGCCCAACAAGAAGAAGATG
tsa56 reverse	TTTAACTTGGCCGACAATCA
scaA forward	TAGTGAAAGGGGACGATTGC

scaA reverse	ATCCGACAAGTTAGGCATGG
*16s*RNA forward	CGGTGAATACGTTCTCGGG
*16s*RNA reverse	GAGGTAATCCAGCCGCAGG
ank12_1 forward	GATAGCAGAGGTTGTACACCG
ank12_1 reverse	CTGATTTGTTAGGCTGCCTGC
mipZ forward	GTGGTGCAGGTAAGACTACTGTTG
mipZ reverse	GTGAATGCTGACGTGAATCAG

Proteomics sample preparation

We used *Orientia tsutsugamushi* strain UT76 to infect L929 mouse fibroblasts at a 410:1 MOI in

six independent culture flasks. High MOIs were used to ensure sufficient material for proteomics

analysis. Despite the high MOI the L929 cells remained intact and attached to the surface of the

flask at the point of harvesting the samples. We replaced the culture medium at one day

postinfection (dpi) to remove bacteria that failed to enter the host. Intracellular and extracellular

bacteria were isolated at four dpi; two separate flasks were harvested as one replicate, yielding

a total of three replicates. L929 cells were grown in 75-cm^2 plastic flasks at 37°C in a 5% CO_2

atmosphere, using DMEM media (Gibco 11965092) supplemented with 10% FBS (Sigma-

Aldrich F7524). At four dpi, the media were transferred to 50-ml conical tubes for extracellular

bacteria isolation. Culture flasks were rinsed with 15 ml of PBS, and cells were scraped using

two ml of fresh media per flask. The detached cells were transferred to two-ml Safe-lock tubes

(Eppendorf 0030120094) and homogenized for three minutes using a Bullet blender[®]

homogenizer (Next Advance #BBY24M) with speed control setting at 8. Host cell debris was

removed by centrifugation at 300 g for three minutes; we then harvested intracellular bacteria

(IB) by an additional three-minute centrifugation at 20,000 g. Before isolation of extracellular

bacteria (EB), we removed possible detached host cells containing IB through a pre-spin step

(1,000 g, 5 min, 4°C). No visible pellet was observed indicating that there were negligible

amounts of detached host cells potentially contaminating the EB prep with IB bacteria.

Following transfer of supernatants to new conical tubes, we harvested extracellular bacteria by

centrifugation at 3000 g for 20 minutes at 4°C. IBs and EBs were washed with 0.3 M sucrose

three times, resuspended in PBS, and lysed with 1% Triton X-100 for 30 minutes at 4°C.

Bacterial proteins were precipitated by addition of 80% ice-cold acetone and overnight

incubation at -20°C. Protein pellets were washed in acetone twice to remove residual Triton X-

100 by centrifugation at 15,000 g for 10 min. Briefly dried protein extracts were then dissolved in

100 μl of 50% trifluoroethanol in 100 mM Triethylammonium bicarbonate (TEAB) buffer. The

samples were reduced by treatment with 20 mM tris(2-carboxyethyl)phosphine (TCEP) for 20

minutes at 55°C, alkylated in 55 mM chloroacetamide (CAA) at room temperature in the dark,

and digested with endoproteinase LysC (LysC: protein ratio=1:50) overnight at 37°C and,

subsequently, with trypsin (trypsin: protein ratio=1:100), also overnight at 37°C. Average

bacterial copy numbers for IBs and EBs were estimated by quantitative PCR yielding 1.5E+08

and 2.6E+09 per replicate, respectively. The whole protein content in each replicate was used

without normalisation at this step due to the fact that IBs and EBs were isolated from different

environment and therefore have diverse amounts of co-purified host protein. Samples were

acidified with Trifluoroacetic acid (TFA) to 1% final concentration and centrifuged to remove

insoluble materials. Trypsinized samples were desalted using an Oasis® HLB 1cc cartridge

(Waters™). Briefly, following cartridge activation with 500 μl of 100% acetonitrile and

equilibration with 500 μl of 0.5% acetic acid, we loaded samples and then washed the cartridge

with 1 ml MS-grade water. Peptides were eluted with 500 μl of 80% acetonitrile in 0.5% acetic

acid, then transferred to low-binding microtubes, and dried using a SpeedVac. Each peptide

pellet was resuspended with 20 μl of MS-grade water. We estimated peptide content using 5%

of each sample, so called 'Mini-runs'. This was to check whether we need to adjust or normalise

sample loadings in the actual runs.. Briefly, individual aliquots in 1 μl of peptides were

supplemented with 4 µl of 1% (v/v) acetonitrile containing 0.06% (v/v) trifluoroacetic acid and

0.5% (v/v) acetic acid, transferred to a 96-well autosampler plate, and then injected (3 µl) into an

Orbitrap Fusion™ Mass Spectrometer. After we screened out host proteins from the Mini-runs

result, the frequency distribution of Ot protein abundances were plotted. Medians of IB and EB

replicates showed abundance values with maximum difference of 1 log scale, suggesting a

single normalisation process may be performed during data analysis. We decided to fractionate

the rest of samples without the need of normalizing sample loadings. The remaining 95% of

each sample (19 µl in MS-grade water) was mixed with 20 µl of 10 mM ammonium formate (pH

10) and centrifuged at 20,000 g for 3 min to eliminate insoluble components. Each peptide

sample was then fractionated through a step-wise desalting process on C18 stage tips using

10%, 15%, 25%, and 50% acetonitrile in 10 mM ammonium formate (pH 10). Fractionated

samples were washed with 60% acetonitrile containing 0.1% formic acid twice to eliminate

residual ammonium formate by evaporation before tandem MS analysis.

LC-MS/MS

Dried samples were resuspended in 10 µl of 2% (v/v) acetonitrile containing 0.06% (v/v)

trifluoroacetic acid and 0.5% (v/v) acetic acid, and 1 µl was loaded onto an autosampler plate.

Online chromatography was performed in an EASY-nLC 1000 (ThermoScientific) instrument

using a one-column setup and 0.1% formic acid in water / 0.1% formic acid in acetonitrile as

mobile phases. The samples were injected and separated on a reversed-phase C18 analytical

column (EASY-Spray LC Column, 75 µm inner diameter × 50 cm, 2 µm particle size;

ThermoScientific) maintained at 50°C and using a 2-23% (v/v) acetonitrile gradient over 60 min,

followed by an increase to 50% over the next 20 min, and to 90% over 5 min. The final mixture

was maintained on the column for 5 min to elute all remaining peptides. Total run duration for

each sample was 90 min at a constant flow rate of 300 nl/min.

Data were acquired using an Orbitrap Fusion Lumos™ mass spectrometer (ThermoScientific)

using the instrument's data-dependent mode. Samples were ionized using 2.5 kV and 300°C at

the nanospray source; positively-charged precursor MS1 signals were detected using an

Orbitrap analyser set to 60,000 resolution, automatic gain control (AGC) target of 400,000 ions,

and maximum injection time (IT) of 50 ms. Precursors with charges 2-7 and having the highest

ion counts in each MS1 scan were further fragmented using collision-induced dissociation (CID)

at 35% normalized collision energy. MS2 signals were analysed by ion trap at an AGC of 15,000

and maximum IT of 50 ms. Precursors used for MS2 scans were excluded for 90 s to avoid re-

sampling of high abundance peptides. The MS1–MS2 cycles were repeated every 3 s until

completion of the run.

Proteomic data analysis

Proteins were identified using Proteome Discoverer™ (v2.3, Thermo Scientific). Raw mass

spectra were searched against *Orientia tsutsugamushi* primary protein sequences derived from

complete genome data for the UT76 strain. Mouse whole proteome sequences were obtained

from Uniprot and included as background. Carbamidomethylation on Cys was set as a fixed

modification; deamidation of asparagine and glutamine, acetylation on protein N termini, and

methionine oxidation were set as dynamic modifications for the search. Trypsin/P was set as the

digestion enzyme and was allowed up to three missed cleavage sites. Precursors and

fragments were accepted if they had a mass error within 10 ppm and 0.8 Da, respectively.

Peptides were matched to spectra at a false discovery rate (FDR) of 1% against the decoy

database.

For label-free quantitation (LFQ), features were extracted from the raw data files using the

Minora Feature Detector processing workflow node on Proteome Discoverer. Chromatographic

feature-matching was performed with a tolerance of 10 min to align the retention times and

enable quantitation of the precursors. No pre-normalization was performed using the Proteome

Discoverer workflow, and raw results were manually normalized using the Median normalization

method. A protein was classified as detected if at least two peptides were detected in a

minimum of two biological replicates; the mean LFQ across the three replicates was used for

further analyses. There was no normalization during protein treatments and peptide injection as

described earlier. Normalization between samples was performed only in the data analysis

process. Briefly, frequency distributions of Ot protein abundances from six separate runs were

plotted (excluded co-purified host proteins). And the median from each plot was used to

normalize the whole set, prior to data trimming, making all data set scaled to the same median

reference value of 1. Normalized abundance values were then averaged from independent

biological replicates (Supplementary table 1). Statistical significance (t-test) was determined

using GraphPad Prism software 8.4.3. Subsequently, proteins detected in both IBs and EBs for

at least twice were selected for protein abundance comparison, described as Ext/Int ratio.

Proteomic data is available from the jPOST repository with the identifier PXD028218.

Immunoblot analyses

L929 and HUVEC cells were grown in 24-well plates and infected with Ot UT76 at an MOI of

12:1. Non-infectious bacteria were removed by replacing the media 24 hours post-infection. EB

and IB populations were prepared four days post-infection. EB was prepared by transferring the

cell supernatant to an Eppendorf tube, pelleting at 16,873 g, then resuspending in 100 µl PBS

containing 1x protease inhibitor cocktail (ThermoFisher 78425). For TSA56 immunoblots,

samples were prepared in the presence of 1 mM iodoacetamide (Sigma-Aldrich catalog number

I6125). IB was prepared by removing adherent cells using trypsinization, spinning the cell pellet,

and resuspending it in PBS supplemented with protease inhibitors; host cells were lysed using a

bead mill for 1 minute. Host cell debris was removed by centrifugation at 500 g for 5 min; the

supernatant was transferred to a clean tube, spun at 16,873 g for 5 minutes, and resuspended

in 100 µl PBS supplemented with protease inhibitors. Bacterial copy number was quantified by

qPCR using *tsa47* primers, as described above. IB and EB samples were diluted to ensure

equal loading of bacterial cells on gels. Samples were diluted in 5X SDS loading buffer with and

without ß-mercaptoethanol and heated at 95ºC for 5 min. Samples were subjected to

electrophoresis on a 4-20% gradient gel (Bio-Rad 4561094) and then transferred to a

nitrocellulose membrane. Following blocking in PBST containing 5% BSA and 5% skim milk for

30 min, we incubated the blots with primary antibody (1:1,000) in PBST-BSA overnight at 4ºC.

Membranes were washed and incubated with secondary antibody (1:10,000) conjugated to

IRDYE 680RD or IRDYE 800CW (LI-COR Biosciences) in PBST-BSA for 1 hour at room

temperature. Membranes were washed, and immunoreactive proteins were visualized using a

LiCor Odyssey CLX imager.

Antibody name	Epitope used for immunisation	Description of epitope	Species and type
TSA56	KLTPPQPTIMPISIADRDF - KLH	Synthetic peptide from strain Karp conjugated to KLH carrier protein	Rat monoclonal
RpoD	MASLRNKNNKIIICNKKTKKNTNVEELASNFV DESEKLSSKISDSDINLDKDFDYTKLGVCPDK IEEEVALASNATEGEFEEDLDDNIYSTDIVRA YLKDMSN	Recombinant protein M1-N103 from strain Karp	Rabbit polyclonal
CtrA	KGHSEAVIRVGHLSINLDTRSVSMYNKDKQK NMPIHLTNKEYKILELLAIKKGTVLTKEMFLSH LYSGIDEPEIKIIDVFICKLRRKLEKASNGINFIE TVWGRGYIFAEFKSNDNNYSLNNVEEE	Recombinant protein K120-E145 from strain Karp	Rabbit polyclonal
SpoT	ELSPIAISDNNKIDKINQNLPIDFAKCCYPLPG DLIIGLCTKTEIVIHTTYCTIASNYVLESIRLLDF SWTSNAQKMFYTSSVQLSVRNKIGSLASITTI LENNNV	Recombinant protein E556-V661 from strain Karp	Rabbit polyclonal
RpoH	NSSDYKQIADESGVSIAEVSEVNQRLSNSDL SLNTPISYHEDDATELVELVPEHRPNQEIILAN QQDLNLKKEVLQNALQTLTQRELYIIRARKLQ DEPDTLEKLSLHFNVSKERIRQIESKAFEKLQ QYVMNTKHKLLHFV	Recombinant protein N158-V299 from strain Karp	Rabbit polyclonal
ScaC	KLH- KIQLQQLQVTLQNSLFLQNKNNNPA + KLH - IAEVSKLLCMLE	Synthetic peptides from strain Karp conjugated to KLH carrier protein	Rabbit polyclonal
ScaA	NNTTESSNDNQGDTGSGATSNPSNPSNPIE GASGTSPATGASDTTGDSDAAGNKEAAGDD SVTDSGASQSNNTTESSNDNQGDTGSGATS	Recombinant protein N50-G179 from strain Gilliam	Rabbit polyclonal

	NPSNPIEGASGTSPATGASDATGVTETTGGS ATGGEVTDG		

Quantification of genome copy number per cell

The genome copy number per cell was calculated by determining the number of bacterial genome copies in a sample of purified EB and IB Ot, using *tsa47* qPCR, and dividing by the number of bacterial cells in the same sample, as measured by confocal microscopy. Three µl of purified bacteria were spread evenly across a well of a multi-well slide, dried, and labelled for immunofluorescence microscopy using TSA56, as described above. The bacteria in a selected field were manually counted to determine the number of bacterial cells in the sample.

Quantification of relative peptidoglycan levels using NOD1 reporter assay

HEK-Blue hNOD1 cells (Invitrogen, hkb-hnod1, USA) were grown in DMEM containing 4.5 g/l glucose, 10% heat-inactivated FBS, 100 U/ml penicillin, 100 mg/ml streptomycin, 100 mg/ml Normocin, 2 mM L-glutamine, and the selective antibiotics blasticidin (30 µg/ml) and Zeocin (100 µg/ml). The cells were seeded on clear-bottom black 96-well plates (Corning 29444-008) 2 days before treatment. Bacterial aliquots from stocks kept in sucrose phosphate glutamine (SPG) at -80°C were heat-inactivated at 90°C for 30 minutes before supplementation to host cells. These studies were carried out in triplicate. After two days, growth media were replaced with HEK-blue detection media (Invivogen hb-det2) for SEAP assessment. The plates were further incubated for six hours at 37°C in a 5% CO_2 atmosphere and then analyzed spectrophotometrically (Synergy H1, Biotek) at 640 nm. Results were analyzed by GraphPad Prism (GraphPad Software, San Diego California USA), and a t-test was performed to compare bacterial SEAP levels to positive control iE-DAP (γ-D-Glu-mDAP) at 10 µg/ml.

Growth curve measurements and bead mill treatment

To determine the growth rate of IB and EB Ot from frozen stock were first grown in two T25 flasks containing a monolayer of L929 cells. At 6 days post infection, IB and EB were prepared and used to inoculate fresh L929 cells and the subsequent bacterial growth was measured. **EB isolation** – the supernatant portion of the culture consisting of EB was transferred to a 50-mL centrifuge tube (Corning 430828) and subjected to a first centrifugation to remove any host cells at 1,500 g for 5 minutes. The supernatant was then transferred into a new 50-ml tube and centrifuged at 3,220 g for 20 minutes to obtain an EB pellet which was resuspended in the fresh media and ready for the next step. **IB isolation** – fresh media was added directly to the culture flasks containing infected cells and these were disassociated from the adherent surface by using a cell scraper (Corning Falcon, catalog no. 353089). The infected cell suspension was then pipetted into a sterile 2-mL microcentrifuge tube. (Eppendorf 022363352) pre-filled with sterile 0.1-0.15 mm diameter glass beads (Biospec 11079101) before introduced to VWR® Bead Mill Homogenizer (VWR 10158-558) at speed 5 for 1 minutes to break the host cells. Subsequently, the purified IB in the supernatant was separated from host cells debris and glass beads by transferring the supernatant into a new 2-mL microcentrifuge tube and spinning at 300 g for 5 minutes. The bacterial concentration in both EB and IB suspension was quantified by qPCR as described above, and EB and IB Ot was infected onto a fresh monolayer of L929 cells in 12-well plates (Thermo Scientific 150628). Bacteria were added at 2.83E+04 bacterial copies/well which corresponds to an MOI of ~1:10. The number of intracellular bacteria was then measured at 3, 24, 96 and 168 hours post infection by *tsa47* qPCR. **Physical stress induced by bead mill** – IB and EB isolation and infection were performed as described above and then exposed to physical stress with 1 or 2 minutes agitation in the bead mill. Bacteria were then infected onto L929 cells at a concentration of 1.00E+06 copies/well (MOI = 2:1). The quantity of intracellular bacteria was then determined at 3, 24, 96, 120, 144 and 168 hours post infection by qPCR. **Osmotic stress induced by water treatment.** Quantification of the effect of osmotic shock on Ot viability was performed in the same way as for physical stress except

purified bacteria were resuspended in pure water for 5 minutes prior to infecting host cells.

Bacterial copy number after 7 days was quantified by qPCR.

Entry inhibition experiments. IB and EB forms of Ot were isolated from infected host cells as

described above. To assess the impact of different interventions on bacterial entry, IB and EB

form Ot were added to uninfected L929 or HUVEC cells, labelled with HPG and fixed three hours

post infection. The number of bacteria inside host cells was counted manually with at least 100

host cells counted for each condition/replicate. To measure the role of clathrin, 0.2 µM siRNA

(clathrin, catalog number 4390825 or scramble catalog number 4390843, ThermoFisher

Scientific) was added 1 day prior to infection. To measure the effect of perturbing clathrin-

mediated endocytosis or macropinocytosis, inhibitors were added for one hour prior to

infection and kept in the media during the three hours of infection. Dynasore hydrate (inhibits

endocytosis, catalog number D7693, Sigma) was used at 100 µM, amiloride (inhibit

micropinocytosis, 5-(N-Ethyl-N-isopropyl, catalog number A3085, Sigma) at 50 µM and

zoniporide hydrochloride hydrate (inhibit micropinocytosis, catalog number SML0076, Sigma) at

100 µM. In order to determine the role of bacterial surface proteins TSA56 and ScaC, IB and EB

Ot were added to uninfected host cells at the same time as antibodies against TSA56 and ScaC

and the number of intracellular bacteria at three hours post infection quantified by microscopy.

Statistical analyses

All statistical analyses were carried out using GraphPad Prism version 9.2.0. Comparisons

between two groups with >10 datapoints per group were carried out using a students unpaired t

test, whilst comparisons between two groups with ≤10 datapoints per group were carried out

using a Mann-Whitney test. Comparisons between more than two groups were carried out using

one-way ANOVA followed by Dunnetts comparisons. Ns = no significant difference detected, P

> 0.05; * P ≤ 0.05; ** P ≤ 0.01; *** P ≤ 0.001; **** P ≤ 0.0001.

Acknowledgements and Funding

We thank each laboratory member for valuable discussions and support. We are grateful to the

following people for providing helpful feedback and insightful comments related to the project

and the manuscript: Jonathan Dworkin from Columbia University, Dave Dubnau from PHRI, and

Diana Stafforini from Life Science Editors. We also appreciate the support from Patricia

Greenberg and Tracy Andrews at the Rutgers University Biostatistics and Epidemiology

Services (RUBIES).

This work was funded by a Royal Society Dorothy Hodgkin Research Fellowship (JS)

(DH140154), a Newton Fund MRC-NSTDA grant (YJ/JS) (MR/N012380/1), and National

Institute of Health/National Institute of Allergy and Infectious Diseases grants (R21AI144385

and R56AI148645) (JS), and A*STAR core funding and Singapore National Research

Foundation under its NRF-SIS "SingMass" scheme (RMS).

Data availability

Source data are provided with this paper as raw data files. Proteomics data is available on

ProteomeXchange (PXD028218) and jPOST (JSPT001299)[28].

References

References

1   Taylor, A., Paris, D. & Newton, P. A Systematic Review of Mortality from Untreated
Scrub Typhus (Orientia tsutsugamushi). *PLoS Negl Trop Dis* **9**, e0003971,
doi:10.1371/journal.pntd.0003971 (2015).
2   Luce-Fedrow, A. *et al.* A Review of Scrub Typhus (Orientia tsutsugamushi and Related
Organisms): Then, Now, and Tomorrow. *Tropical Medicine and Infectious Disease* **3**, 8,
doi:10.3390/tropicalmed3010008 (2018).
3   McClure, E. *et al.* Engineering of obligate intracellular bacteria: progress, challenges and
paradigms. *Nat Rev Microbiol*, doi:10.1038/nrmicro.2017.59 (2017).
4   Salje, J. Cells within cells: Rickettsiales and the obligate intracellular bacterial lifestyle.
Nat Rev Microbiol, doi:10.1038/s41579-020-00507-2 (2021).
5   Rikihisa, Y. & Ito, S. Localization of electron-dense tracers during entry of Rickettsia
tsutsugamushi into polymorphonuclear leukocytes. *Infect Immun* **30**, 231-243 (1980).
6   Lee, J. *et al.* Fibronectin facilitates the invasion of Orientia tsutsugamushi into host cells
through interaction with a 56-kDa type-specific antigen. *J Infect Dis* **198**, 250-257,
doi:10.1086/589284 (2008).
7   Ha, N., Cho, N., Kim, Y., Choi, M. & Kim, I. An autotransporter protein from Orientia
tsutsugamushi mediates adherence to nonphagocytic host cells. *Infect Immun* **79**, 1718-
1727, doi:10.1128/IAI.01239-10 (2011).
8   Ha, N. *et al.* Immunization with an Autotransporter Protein of Orientia tsutsugamushi
Provides Protective Immunity against Scrub Typhus. *PLoS Negl Trop Dis* **9**, e0003585,
doi:10.1371/journal.pntd.0003585 (2015).
9   Chu, H. *et al.* Exploitation of the endocytic pathway by Orientia tsutsugamushi in
nonprofessional phagocytes. *Infect Immun* **74**, 4246-4253, doi:10.1128/IAI.01620-05
(2006).
10   Kim, H., Choi, M. & Kim, I. Role of Syndecan-4 in the cellular invasion of Orientia
tsutsugamushi. *Microb Pathog* **36**, 219-225, doi:10.1016/j.micpath.2003.12.005 (2004).
11   Cho, B., Cho, N., Seong, S., Choi, M. & Kim, I. Intracellular invasion by Orientia
tsutsugamushi is mediated by integrin signaling and actin cytoskeleton rearrangements.
Infect Immun **78**, 1915-1923, doi:10.1128/IAI.01316-09 (2010).
12   Giengkam, S. *et al.* Improved Quantification, Propagation, Purification and Storage of
the Obligate Intracellular Human Pathogen Orientia tsutsugamushi. *PLoS Negl Trop Dis*
9, e0004009, doi:10.1371/journal.pntd.0004009 (2015).
13   Atwal, S. *et al.* Evidence for a peptidoglycan-like structure in Orientia tsutsugamushi.
Mol Microbiol **105**, 440-452, doi:10.1111/mmi.13709 (2017).
14   Min, C. *et al.* Genome-based construction of the metabolic pathways of Orientia
tsutsugamushi and comparative analysis within the Rickettsiales order. *Comp Funct*
Genomics **2008**, 623145, doi:10.1155/2008/623145 (2008).
15   Atwal, S. *et al.* Clickable methionine as a universal probe for labelling intracellular
bacteria. *J Microbiol Methods* **169**, 105812, doi:10.1016/j.mimet.2019.105812 (2020).

16   Rikihisa, Y. Anaplasma phagocytophilum and Ehrlichia chaffeensis: subversive
manipulators of host cells. *Nat Rev Microbiol* **8**, 328-339, doi:10.1038/nrmicro2318
(2010).
17   Elwell, C., Mirrashidi, K. & Engel, J. Chlamydia cell biology and pathogenesis. *Nat Rev*
Microbiol **14**, 385-400, doi:10.1038/nrmicro.2016.30 (2016).
18   Otten, C., Brilli, M., Vollmer, W., Viollier, P. & Salje, J. Peptidoglycan in obligate
intracellular bacteria. *Mol Microbiol* **107**, 142-163, doi:10.1111/mmi.13880 (2018).
19   Jacquier, N., Frandi, A., Pillonel, T., Viollier, P. & Greub, G. Cell wall precursors are
required to organize the chlamydial division septum. *Nat Commun* **5**, 3578,
doi:10.1038/ncomms4578 (2014).
20   Irving, S. E., Choudhury, N. R. & Corrigan, R. M. The stringent response and physiological
roles of (pp)pGpp in bacteria. *Nat Rev Microbiol*, doi:10.1038/s41579-020-00470-y
(2020).
21   Batty EM, C. S., Blacksell SB, Richards A, Paris D, Bowden R, Chan C, Lachumanan R, Day
N, Donnelly P, Chen SL, Salje J. Long-read whole genome sequencing and comparative
analysis of six strains of the human pathogen Orientia tsutsugamushi. *Plos Negl Trop Dis*
(2018).
22   Mika-Gospodorz, B. *et al.* Dual RNA-seq of Orientia tsutsugamushi informs on host-
pathogen interactions for this neglected intracellular human pathogen. *Nat Commun* **11**,
3363, doi:10.1038/s41467-020-17094-8 (2020).
23   Joshi, K. K. & Chien, P. Regulated Proteolysis in Bacteria: Caulobacter. *Annu Rev Genet*
50, 423-445, doi:10.1146/annurev-genet-120215-035235 (2016).
24   Lamason, R. *et al.* Rickettsia Sca4 Reduces Vinculin-Mediated Intercellular Tension to
Promote Spread. *Cell* **167**, 670-683.e610, doi:10.1016/j.cell.2016.09.023 (2016).
25   Lee, S. Optimal integration of wide field illumination and holographic optical tweezers
for multimodal microscopy with ultimate flexibility and versatility. *Opt Express* **26**, 8049-
8058, doi:10.1364/OE.26.008049 (2018).
26   Huang, B., Wang, W., Bates, M. & Zhuang, X. Three-dimensional super-resolution
imaging by stochastic optical reconstruction microscopy. *Science* **319**, 810-813,
doi:10.1126/science.1153529 (2008).
27   Wang, Y. *et al.* Localization events-based sample drift correction for localization
microscopy with redundant cross-correlation algorithm. *Opt Express* **22**, 15982-15991,
doi:10.1364/OE.22.015982 (2014).
28   Okuda, S. *et al.* jPOSTrepo: an international standard data repository for proteomes.
Nucleic Acids Res **45**, D1107-D1111, doi:10.1093/nar/gkw1080 (2017).
879

A.

B.

C.

D.

DNA Actin Ot

DNA Actin Ot

A.

EB

translationally
active Ot

translationally
inactive Ot

IB

nucleus

B.

IB	EB
more peptidoglycan	less peptidoglycan
elongated and round	round
less fragile	more fragile
TSA56 less cross linked	TSA56 highly cross linked
translationally active	translationally inactive
high ScaA/low ScaC	low ScaA/high ScaC
enters cells by clathrin-mediated endocytosis and macropinocytosis	enters cells by macropinocytosis