The obligate intracellular bacterium *Orientia tsutsugamushi* differentiates into a

developmentally distinct extracellular state

Sharanjeet Atwal[b*], Jantana Wongsantichon[a*], Suparat Giengkam[a*], Kittirat Saharat[a], Yanin

Jaiyen[a], Suthida Chuenklin[b], Taerin Chung[c], Hyun Huh[c], Sang-Hyuk Lee[c,d], Radoslaw M. Sobota[e]

and Jeanne Salje[a, b, f#]

Mahidol-Oxford Tropical Medicine Research Unit, Faculty of Tropical Medicine, Mahidol

University, Bangkok, Thailand[a]

Public Health Research Institute, Rutgers the State University of New Jersey, Newark, USA[b]

Institute for Quantitative Biomedicine, Rutgers University, Piscataway, New Jersey 08854, USA[c]

Department of Physics and Astronomy, Rutgers University, Piscataway, New

Jersey 08854, USA[d]

Institute of Medical Biology, Agency for Science, Technology and Research (A*STAR), 138648,

Singapore[e]

Centre for Tropical Medicine and Global Health, Nuffield Department of Medicine, University of

Oxford, Oxford, United Kingdom[f]

* These authors contributed equally

[#] Address correspondence to Jeanne Salje: js2522@rutgers.edu

Abstract

Orientia tsutsugamushi (Ot) is an obligate intracellular bacterium in the family Rickettsiaceae that

causes scrub typhus, a severe mite-borne human disease. Ot is unusual amongst Rickettsiaceae

in budding out of infected cells in a process resembling that used by enveloped viruses. Here, we

show that Ot bacteria that have budded out of host cells are in a distinct developmental stage

compared with intracellular bacteria. We refer to these two stages as intracellular and extracellular

bacteria (IB and EB, respectively). These two forms differ in physical properties: IB is elongated,

and EB is round. Additionally, IB has higher levels of peptidoglycan and is physically robust

compared with EB. The two bacterial forms differentially express proteins involved in bacterial

physiology and host-pathogen interactions, specifically those involved in bacterial dormancy and

stress response, secreted bacterial effectors, and outer membrane autotransporter proteins ScaA

and ScaC. Whilst both populations are infectious, entry of IB Ot is sensitive to inhibitors of both

clathrin-mediated endocytosis and micropinocytosis, whereas entry of EB Ot is only sensitive to

a micropinocytosis inhibitor. Our identification and detailed characterization of two developmental

forms of Ot significantly advances our understanding of the intracellular lifecycle of an important

human pathogen.

Keywords

Orientia tsutsugamushi, Rickettsiales, obligate intracellular bacteria, intracellular pathogens

Author Summary

Orientia tsutsugamushi (Ot) is a bacterial pathogen that causes scrub typhus, a mite-transmitted human disease. This illness is traditionally known to be endemic in the Asia-Pacific, but recent reports from the Middle East, Africa, and Latin America suggest that it may be globally distributed. Scrub typhus is associated with high mortality if not treated promptly with appropriate antibiotics. Ot is a highly specialized bacterium that can only replicate within living cells, either within the mite vector or in mammalian or human hosts. Ot exits infected cells using a unique mechanism that involves budding off the surface of infected cells. We have discovered that this unusual aspect of its lifecycle involves the bacteria themselves differentiating into a distinct growth form. Different growth forms have not been described in other members of the family Rickettsiaceae, and no other family members have been shown to bud out of host cells in a manner similar to Ot. We find that the two forms of Ot, which we refer to as intracellular and extracellular bacteria (IB and EB respectively), differ in physical properties and protein expression and infect cells through different mechanisms. The identification of structurally and functionally distinct forms of Ot elucidates a vital aspect of this pathogen's intracellular life cycle. Our findings are likely to advance the development of novel interventions aimed at inhibiting Ot growth in scrub typhus patients.

Main

Orientia tsutsugamushi (Ot) is an obligate intracellular Gram-negative bacterium that causes the mite-transmitted human disease scrub typhus. This condition is one of the most severe rickettsial diseases; it is associated with a median 6% mortality in untreated patients[1]. Scrub typhus is a leading cause of severe febrile illness in the Asia-Pacific, a region containing two-thirds of the world's population. Recent reports from Latin America, the Middle East, and Africa suggest that the disease is more globally distributed[2]. Knowledge of the fundamental biology of this organism lags behind that of pathogens of equivalent prevalence and severity. Historical underestimations of Ot incidence, its classification as a biosafety level 3 pathogen, its fastidious growth in cultured cells, and the fact that it remains genetically intractable[3] likely account for the paucity of studies focused on this pathogen.

The order Rickettsiales contains two major families: Anaplasmataceae and Rickettsiaceae[4]. Members of the Anaplasmataceae, which include Anaplasma spp. and Ehrlichia spp., reside within membrane-bound vacuoles inside host cells. The developmental cycle of certain Anaplasmataceae family members is biphasic and involves switching between a non-infectious intracellular replicative form (reticulate body) and an infectious extracellular form (elementary body). While members of the Rickettsiaceae, namely Rickettsia spp. and Ot, do not appear to differentiate into distinct forms during their intracellular lifecycle, Ot buds out of host cells in a manner reminiscent of enveloped viruses[5]. This observation led us to investigate whether budded Ot is in a distinct developmental stage compared with intracellular counterparts.

Ot has been shown to enter host cells using the zipper-like invasion mechanism, which is common to many Rickettsiales members[4]. This process is initiated by binding of bacterial surface adhesins to receptors expressed on mammalian cells. Adhesins such as TSA56, ScaA, and ScaC[6-8] bind to host cell fibronectin and syndecan-4; these proteins then interact with integrin $\alpha5\beta1$, leading to activation of intracellular signaling that culminates in actin remodeling and clathrin-mediated

endocytosis[9-11]. It is important to emphasize that these studies were entirely focused on

intracellular Ot, as they were performed on bacteria isolated from host cells by mechanical lysis

of infected cells. However, the re-entry mechanism utilized by bacteria that have budded out of

cells may be influenced by the presence of an enveloping host cell membrane and/or its

differentiation into a distinct state.

Here, we show that Ot exists in distinct intracellular (IB) and extracellular (EB) bacterial

populations. We report the characterization of these two states in terms of their physical

properties, metabolic activity, and gene expression profiles and demonstrate that they enter host

cells using different mechanisms.

Results

IB and EB forms of Ot differ in physical properties and metabolic activity.

Ot is typically described as a coccobacilli bacterium. Its small size and propensity to aggregate

make it difficult to obtain clear images of this bacterium using conventional light microscopy. To

circumvent this challenge, we labelled Ot with an antibody against the abundant surface protein

TSA56 and employed stochastic optical reconstruction microscopy (STORM) for high-resolution

imaging. We first investigated the shape of bacteria located inside and outside host cells (Fig. 1A,

B and Supp. Fig 1). We found that EBs were approximately round in shape with a diameter of

about 1.2 µm. In contrast, IBs adopted diverse round and elongated shapes and varied in

dimensions; some bacteria were 3-4 µm in length. Additionally, IB size varied during the course

of infection. Initially, and up to one-day post-infection, IBs and EBs were similar in length. At later

stages, *i.e.,* 2-7 days following infection, IBs were characterized by longer lengths. These kinetic

changes in shape correlated with Ot growth dynamics whereby replication starts 24-48 hours

point infection[12]. These differences in shape provide evidence for the existence of two separate

bacterial forms.

We next investigated potential differences in bacterial cell wall composition. We previously

showed that Ot builds a low-abundance peptidoglycan-like structure[13]. Thus, we assessed relative

peptidoglycan levels in IB and EB Ot forms using a NOD-1 reporter assay (Fig. 1C). We detected

significantly more peptidoglycan in IB cells compared with EB cells. This finding may account for

the observed differences in shape (Fig. 1A) as peptidoglycan is required to build elongated cells.

In related structural studies, we capitalized on our previous finding that the abundant surface

protein TSA56 is involved in forming disulfide bonds with itself or other proteins, leading to the

formation of aggregates[13]. These complexes can be identified as smears following sample

preparation in the absence of reducing agents and evaluation by immunoblot analyses. In this

study, we found evidence for substantially higher disulfide bond formation in EB Ot than IB Ot, as

determined by the presence of a high molecular weight smear (Fig. 1D). We hypothesize that this

crosslinked structure confers EB Ot with additional structural rigidity in the absence of

peptidoglycan. These experiments were performed in the presence of iodoacetamide, an agent

that covalently modifies cysteine residues and prevents them from forming new bonds, thus

ensuring that the cross-links were not formed during sample preparation and processing.

Our next goal was to assess potential changes in genome copy number as, during rapid growth,

bacteria such as *E. coli* carry multiple genome copies within a single cell. To address this issue,

we determined bacterial cell numbers using fluorescence microscopy and bacterial genome

copies using qPCR. We found no differences: Ot IB and EB harbored just over one estimated

genome copy per cell (Fig. 1E). This measurement is an approximation that suffers limitations

associated with absolute quantification using qPCR and microscopy, but any such errors are

expected to be constant between the IB and EB measurements.

Finally, we asked whether IB and EB Ot forms differed in their metabolic activity. Ot is an obligate

intracellular bacterium with limited central metabolic and biosynthetic capability[14] and a reliance

on host cells for carbon and nitrogen sources and metabolic intermediates. These observations

led us to hypothesize that EB bacteria are metabolically inactive owing to their extracellular

location. To test this, we used a microscopy-based assay whereby growing bacteria are exposed

to a clickable methionine analog ʟ-Homopropargylglycine (HPG). that is subsequently labeled

with a fluorophore[15] (Fig. 1F). Using this methodology, we observed that 86% of IB were

translationally active five days post-infection, compared with 0% of EB (Fig. 1G). It is possible that

EB Ot undergo some residual metabolic activity that is not detected by our assay of protein

translation. The fact that not all IB were labelled with HPG indicated that this result did not simply

reflect a difference in bacterial permeability between the IB and EB forms. These results provide

an additional line of evidence to support the existence of two fundamentally different Ot states.

IB and EB forms of Ot exhibit different gene and protein expression profiles.

We next investigated whether the differences in physical properties and metabolic activity

observed in IB and EB forms of Ot were accounted exclusively by physical location relative to

host cells. Exposure of EB forms to an osmotically different extracellular environment combined

with the lack of nutrient availability might account for our observations. However, a second

possibility is that EB forms arise by differentiation of IB into a biochemically distinct state. We

reasoned that bacteria in the same developmental stage but differing in position relative to host

cells would exhibit almost identical protein profiles. To test this, we performed shotgun mass

spectrometric analyses on isolated IB and EB Ot forms. We found (Supp. Fig. 2) that IB and EB

isolated four days post-infection exhibited distinct protein profiles (correlation coefficient = 0.843).

A complete list of proteins differentially expressed is provided in Supp. Table 1. Pathway analyses

revealed that IB Ot was enriched in proteins involved in protein synthesis (Fig. 2A). In contrast,

EB Ot displayed high expression of SpoT and RpoH, two proteins involved in dormancy and

quiescence (Fig. 2B). Additionally, we observed striking differential expression of effector

proteins: multiple Ankyrin repeat-containing proteins (Anks) were enriched in IB Ot, whereas

tetratricopeptide repeat-containing proteins (TPRs) were enriched in EB Ot (Fig. 2C).

We next investigated whether EB and IB Ot exhibited differences in transcript expression levels.

These experiments required the propagation of large amounts of EB because this form of Ot is

metabolically inactive and thus harbors very low mRNA levels. We selected 12 genes, three of

which (*curA, trxB, sodB*) were expressed at similar levels in our proteomic studies comparing EB

and IB. Additionally, we quantified mRNA expression of five genes encoding proteins enriched in

IB Ot (*yccA, nuoB, dapA, ank12, scaA*) and four genes enriched in EB Ot (*obgE, spot, tsa56*). We

isolated RNA from IB and EB bacteria four days post-infection and assessed mRNA levels by

qRT-PCR (Fig. 2D). We found no correlation between mRNA and protein levels in multiple cases.

For example, RpoH transcript levels were similar in IB and EB Ot, but protein expression was

higher in EB. These observations provide additional evidence supporting the existence of

molecularly distinct bacterial forms and suggest that transcriptional and post-transcriptional

regulatory mechanisms participate in the differentiation process.

Surface proteins are differentially expressed on IB and EB forms of Ot.

Our differential proteomics analyses revealed that IB and EB differentially express specific surface

proteins (Fig. 3A). Surface proteins play a central role in Ot function for two reasons. First, unlike

most Gram-negative bacteria, Ot lacks lipopolysaccharide, which generally serves as a

membrane shield. Second, Ot is a cytosolic bacterium, and its surface is in direct contact with

cytoplasmic host components. Ot harbors genes encoding four widely conserved autotransporter

proteins: ScaA, ScaC, ScaD, and ScaE. Using our proteomics dataset, we detected ScaA and

ScaC, but not ScaD or ScaE. ScaA was slightly enriched in IB relative to EB Ot (1.5X), while ScaC

was enriched in EB (1.87X). Ot also harbors genes encoding three immunogenic surface proteins

that dominate the humoral response in human patients, TSA22, TSA47, and TSA56. We found

that TSA22 was enriched in IB relative to EB Ot (3.2X) while TSA47 and TSA56 were not

significantly differentially expressed (1.14X and 1.32X increase in EB compared with IB,

respectively). To verify our protein expression observations, we generated polyclonal antibodies

against ScaA and ScaC and determined their expression in IB and EB bacteria using immunoblot

analyses (Fig. 3B). Our experiments up to this point were performed in mouse fibroblasts (L929)

because Ot grows readily in these cells. Moreover, L929 fibroblasts are suitable for characterizing

bacterial properties that are not affected by the cellular environment. However, for our validation

studies, we also investigated bacteria grown in primary human umbilical vein endothelial cells

(HUVEC) as a more physiologically relevant model of Ot infection in humans. Our immunoblot

studies in both L929 and HUVECs confirmed enrichment of ScaA in IB Ot and higher expression

of ScaC in EB compared to IB Ot, in agreement with our mass spectrometric analyses.

ScaA and ScaC as markers of EB and IB forms of Ot

Our observation that 86% of IB is translationally active five days post-infection, compared with

0% of EB, combined with the fact that these Ot forms display opposite ScaA and ScaC expression

led us to investigate whether these surface proteins can be used as *bonafide* markers of

metabolically active IB and metabolically quiescent EB Ot forms, respectively. We used

immunofluorescence microscopy to identify bacteria expressing ScaA and ScaC and then

quantified the number of translationally active ScaA- and ScaC-positive bacteria by measuring

incorporation of $_L$-Homopropargylglycine (HPG). We found (Fig. 3C) that ScaA-positive bacteria,

in contrast to ScaC-positive Ot, were mostly translationally active. We then quantified the number

of ScaA/ScaC positive bacteria inside and outside cells and confirmed that ScaC and ScaA

expression were upregulated in EB and IB Ot, respectively (Fig. 3D, E). Together, these results

show that ScaA and ScaC are differentially expressed depending on bacterial location relative to

host cells: metabolically active IB express higher levels of ScaA relative to EB Ot. Conversely,

expression of ScaC in metabolically inactive EB is high relative to that of IB forms of Ot.

The observation that EB expressed high ScaA transcript levels (Fig. 2E) but low levels of ScaA

protein led us to hypothesize that ScaA is translated into protein soon after entry into host cells,

using pre-formed mRNA. We quantified the number of IB Ot expressing ScaA and ScaC at 1 and

3 hours post-infection and found a decrease in ScaC expression and an increase in ScaA

expression as the infection proceeded (Fig. 3F). These results show that ScaA is expressed

during the first 3 hours after infection, suggesting that *scaA* transcripts are made in EBs in

anticipation of new infections. These results point to the existence of regulatory mechanisms that

control the expression of ScaA and ScaC on the surface of Ot cells, depending on bacterial

location and stage of host infection.

IB and EB Ot populations are infectious but differ in physical stability and mechanisms of entry into host cells.

Previous studies have shown that specific Anaplasmataceae differentiate into distinct intracellular and extracellular forms and that only the extracellular form is effective at infecting host cells[16]. This observation led us to investigate whether the ability to infect host cells differs in IB compared with EB Ot. Growth curve measurements revealed that both bacterial forms could infect cells and replicate with similar dynamics (Fig. 4A). However, IB appears to retain infectivity more efficiently than EB following physical and osmotic challenges. Specifically, when we subjected IB and EB Ot to physical stress using a bead mill for 1-2 minutes, we found that EB displayed a reduction in growth that was dependent on the time of exposure to stress; in contrast, IB was unaffected by this treatment (Fig. 4B). Furthermore, osmotic shock resulted in a significantly greater reduction in viability in EB compared with IB (Fig. 4C). Together, these results demonstrate that IB is physically more robust than EB; this property is likely due to differences in peptidoglycan content, as described above.

To investigate whether EB and IB forms of Ot enter host cells using the same invasion mechanism, we pre-treated HUVEC cells with the clathrin-mediated endocytosis inhibitor dynasore or the micropinocytosis inhibitor amiloride. We then infected the cells with IB or EB Ot and quantified the number of intracellular bacteria 3 hours post-infection using confocal microscopy (Fig. 4D, E). The entry of IB Ot into the cells was blocked by dynasore, thus confirming the participation of clathrin-mediated endocytosis in this process. IB entry was also partially inhibited by amiloride, suggesting additional involvement of micropinocytosis in bacterial uptake. In contrast, EB Ot entry into cells was not affected by dynasore treatment but was completely inhibited by pre-incubation with amiloride. These results demonstrate that IB and EB forms of Ot use distinct mechanisms of entry into host cells. Our findings were recapitulated in L929

246   fibroblasts (Supp. Fig. 6), thus establishing that EB Ot enters cells through macropinocytosis,

247   whereas IB Ot utilizes both clathrin-mediated endocytosis and macropinocytosis.

Discussion

Our studies have unveiled a new feature of the obligate intracellular bacterium Ot. This organism's

unique exit strategy, which involves budding off the surface of infected cells, is associated with

differentiation into a molecularly and functionally distinct developmental state. While unusual in

the family Rickettsiaceae, this property is reminiscent of the biphasic lifecycle of multiple species

in the sister family Anaplasmataceae[4] and the unrelated obligate intracellular bacteria *Chlamydia*

spp.[17] and *Coxiella burnettii*[18]. However, in contrast to these organisms, Ot resides in the host

cytosol in naked form rather than in a membrane-bound vacuole. To our knowledge, this is the

first description of an intracellular bacterium living freely in the host's cytosol and having a biphasic

lifecycle. Our findings have important implications for our understanding of Ot biology and unveil

opportunities to develop better therapeutic strategies to treat scrub typhus.

The physical properties of prokaryotic cell walls are crucial for survival. These stiff structures

maintain cellular shape, protect the interior, counteract osmotic pressure, and provide structural

strength through peptidoglycan layers. Our studies revealed that EB and IB forms of Ot differ in

peptidoglycan content. Lower levels in EB compared with IB make this form more susceptible to

physical and osmotic stress. There is a long-standing question regarding the role of peptidoglycan

in obligate intracellular bacteria. These organisms reside in an osmotically protective niche and

are under selective pressure to reduce the abundance of immunogenic pathogen-associated

molecular patterns (PAMP)s[19]. In the case of pathogenic *Chlamydia* species, it has been shown

that peptidoglycan is retained only at the septum and is required for cell division[20]. Higher

peptidoglycan levels in IB Ot *vs.* EB Ot point to the retention of peptidoglycan for growth and

division rather than structural integrity maintenance.

The observed differences in physical properties between EB and IB forms are complemented with

distinct protein expression profiles suggesting that each form performs unique functions or

responds to specific challenges. Proteomics analyses in EB Ot revealed increased expression of

heat shock sigma factor RpoH and the alarmone (p)ppGpp synthetase SpoT associated with the

stringent response[21]. These results suggest that the EB form of Ot shares features with the

dormant state of other bacteria. Notably, Ot is the only reported member of the Rickettsiaceae to

encode a full-length *spoT* gene[4]. This exclusive feature is in harmony with the fact that

developmental differentiation of Ot is highly unusual in members of this family. The changes

associated with differentiation into the EB state likely serve one or both of the following purposes:

ensuring survival in the extracellular state and priming the bacteria for reinfection of new cells.

Indeed, the observation that *scaA* transcripts are abundant in the EB form and that ScaA protein

is synthesized early upon entry into host cells supports the idea that EB Ot is primed for infection.

Comparing the protein and RNA transcript levels of selected genes revealed a poor correlation

for many genes. A potential explanation is that RNA levels are low in the metabolically inert EB

form, which does not reflect newly translated proteins. Ot strain UT76 encodes only 8 predicted

transcription factors and two sigma factors, RpoD and RpoH[22]. Thus, it is possible that gene

expression is not strictly regulated at the transcriptional level but that other forms of regulation are

prevalent in Ot. We recently showed high antisense transcription levels in Ot, especially in genes

associated with the rampant integrative conjugative element RAGE, which dominates the Ot

genome[23]. There, we found that high antisense transcription correlates with low protein levels for

a particular gene, pointing to this mechanism as a central regulatory strategy in Ot. Gene

expression in the Ot-related, free-living alphaproteobacterium *Caulobacter crescentus* is highly

regulated by proteolysis[24]. Mechanisms based on differential proteolysis may therefore also play

an essential role as critical determinants of protein expression in Ot.

Our observation that EB Ot enters cells through pinocytosis, whereas IB Ot utilizes endocytosis

and, to a lesser extent, pinocytosis resolves a long-standing gap in the field as it was unclear how

budded bacteria enveloped in plasma membrane enter host cells. Physiologically, IB entry into

host cells may be necessary during trafficking into adjacent cells, as previously described for

spotted fever group *Rickettsia* spp.[25]. Additionally, low levels of cell may release IB Ot forms that

can then infect new host cells. The different mechanisms utilized by EB and IB to enter cells

suggest that strategies to escape from the endolysosomal pathway will differ between the two Ot

forms. It is unclear whether and how the differing transcriptional and proteomic profiles at the

point of infection will affect early host-pathogen interactions. Our finding that IB and EB's growth

dynamics are similar suggests that early differences do not persist through the infection cycle.

In summary, our finding that Ot differentiates into molecularly and functionally distinct forms

provides a novel framework to study the infection cycle of this important obligate intracellular

pathogen and opens avenues to investigate the physiological significance of the distinct

populations characterized.

Fig. 1: Intracellular and extracellular forms of Ot are physically and metabolically distinct.

A. STORM images of IB and EB Ot in L929 cells. Bacteria were labelled with a monoclonal

antibody against surface protein TSA56. Additional images are provided in Supp. Fig. 1. B. Length

quantification of individual IB and EB Ot as imaged by STORM. C. Quantification of relative

peptidoglycan levels through activation of NOD1 reporter cells, as measured by secreted alkaline

phosphatase activity. EB Ot induced less activation than IB, indicating lower levels of

peptidoglycan. Identical amounts of IB and EB Ot were used, and biological triplicates are shown.

D. Immunoblot showing the formation of disulfide cross-linked aggregates of the surface protein

TSA56, observed in the absence of the reducing agent beta-mercaptoethanol (BME). This effect

was stronger in EB compared with IB Ot. The expected molecular weight of TSA56 (56 kDa) is

indicated with an arrow, and a dashed arrow depicts higher-order aggregates. E. Genome copy

number per cell as measured by qPCR; bacterial cell number was determined by

immunofluorescence microscopy. Quantification of 10 individual imaging fields is shown. F.

Confocal microscopy image of an L929 cell infected with Ot and labelled with the clickable

methionine analog homopropargylglycine (HPG, green) and an antibody against surface protein

TSA56 (red). The host cell outline is showed by a dashed line, and an arrow indicates a

metabolically inactive bacterium located outside the host cell. G. Quantification of metabolic

activity, as measured by a detectable HPG signal, in IB and EB Ot. L929 cells were infected with

Ot for five days, fixed, and labelled with TSA56 and HPG. One-hundred bacteria inside (IB) and

outside cells (EB) were counted and scored as being HPG-positive or -negative. The percentage

of HPG-positive IB and EB Ot, based on individual values from three independent replicates.

Statistical significance (t-test) was determined using Graphpad Prism software. p-value *** ≤

0.001.

Fig. 2: Intracellular and extracellular forms of Ot have distinct gene expression profiles.

A. Pathway analysis of proteins upregulated >2X between IB and EB forms of Ot (60 proteins in IB, 47 proteins in EB). Bacteria were grown in L929 cells at an MOI of 410:1 and harvested four days post-infection. IB and EB Ot were isolated and analysed by shotgun proteomics. B. SpoT and RpoH are upregulated in EB compared with IB Ot as measured by proteomics described in A. C. Differential expression of Ankyrin-repeat-containing proteins (Anks) and tetratricopeptide repeat-containing proteins (TPR) in EB and IB Ot measured by proteomics, as in A. D. RTqPCR analysis showing relative levels of 12 genes expressed in EB and IB Ot. Bacteria were isolated from L929 cells four days post-infection at an MOI of 672:1, 300:1, and 560:1. The figure depicts results from three independent replicates. Gene expression levels were normalised using three genes: *mipZ*, *ank12*, and *16S.* Mean and standard deviation values are shown. The ratio of protein levels in EB/IB, as found in the proteomics experiment in 2A, is given above each gene for comparison.

Fig. 3: Autotransporter proteins ScaA and ScaC are differentially expressed on IB and EB forms of Ot.

A. Volcano plot showing differential expression of surface proteins in proteomics analysis of IB and EB forms of Ot. We observed higher expression of ScaC, TSA56, and TSA47 in EB whilst IB Ot expressed higher ScaA and TSA22 levels B. Immunoblot analysis of IB and EB Ot from infected L929 and HUVEC cells confirming differential expression of ScaA, and ScaC between IB and EB forms. Loading was normalised by measuring bacterial genome copy number using qPCR and loading the same number of IB and EB Ot in each pairwise comparison. Uncropped immunoblot images showing antibody specificity are given in Supp. Fig. 3. C. Quantification of translational activity, as determined by a positive confocal microscopy signal observed with the clickable methionine probe ʟ-homopropylgylglycine (HPG). L929 cells were infected with Ot for five days, then fixed and labelled with HPG and anti-ScaA or anti-ScaC antibodies. ScaA/ScaC-positive bacteria were scored as being positive or negative for HPG labelling. Mean and SD of

three independent replicates are shown. D. Confocal microscopy images of HUVEC cells infected

with Ot for five days and labelled for DNA (hoescht, blue), translational activity (HPG, green),

ScaA (anti-ScaA antibody, magenta) and ScaC (anti-ScaC antibody, red). Scale bar = 10 μm.

Examples of extracellular bacteria labelled with ScaC are shown with a white arrow. More images

as well as equivalent experiments in L929 cells are shown in Supplementary Figures 4 and 5. E.

Quantification of confocal micrographs of HUVEC cells infected with Ot and labelled as in 3D.

The percentage of IB and EB Ot at five days post-infection that were positively labelled with anti-

ScaA and -ScaC antibodies was counted in three independent experiments. Graph shows mean

and SD, and p value determine by unpaired t test. F. Quantification of ScaA- and ScaC-positive

IB bacteria inside HUVEC cells one and three hours post-infection, showing an increase in ScaA

labelling and a decrease in ScaC labelling. * p-value ≤ 0.05, ** p-value ≤ 0.01, *** p-value ≤ 0.001.

Fig. 4: Both IB and EB populations are infectious but differ in physical stability and

mechanism of entry into cells.

A. IB and EB Ot were isolated from infected L929 cells five days post-infection and grown in fresh

L929 cells in a 24-well plate. The number of intracellular bacteria was then quantified at 3, 24, 96,

and 168 hours post-infection using qPCR against *tsa47,* a single copy Ot gene*.* Doubling times

of IB and EB Ot were similar. B. IB and EB were exposed to physical stress using a bead mill for

one or two min; growth was quantified as in 4A. The bead mill did not affect the growth rate of IB

Ot but reduced that of EB Ot in an exposure-dependent manner. C. IB and EB were exposed to

osmotic shock by suspending the bacteria in water for five min and then growing them in fresh

L929 cells. Bacterial copy number after seven days was quantified by qPCR. EB was more

susceptible to osmotic shock than IB Ot, as shown by a more significant decrease in bacterial

number. D. Quantification of the effect of endocytosis inhibitors on entry of IB and EB Ot. IB and

EB were isolated from L929 cells five days post-infection and then seeded onto fresh HUVEC

cells that had been treated with inhibitors of clathrin-mediated endocytosis (dynasore) or

macropinocytosis (amiloride). Cells were fixed three hours post-infection. Dynasore inhibits entry

of IB but not EB, indicating that clathrin-mediated endocytosis is the major mode of entry of this

population, whilst amiloride strongly inhibited EB and, to a lesser extent, IB. This observation

points to a role for macropinocytosis in EB's entry into mammalian cells. Confocal microscopy

images shown in E. Red = actin (phalloidin), blue = hoescht (DNA), green = Ot (anti-TSA56

antibody). Scale bar = 10 µm. Similar results were found in L929 cells (Supp. Fig. 6).

Fig.5: An overview of the infection cycle of Ot, showing differentiation between IB and EB

forms.

IB replicate in a micro-colony adjacent to the nucleus where they are elongated in shape and

translationally active. Bacteria move to the cell's edge and transition to an EB form that is round

and translationally inactive. Created with BioRender.com B. Table summarizing the differences

between IB and EB forms of Ot.

Methods

Cell culture, bacterial propagation and quantification, and reagents

Proteomics and qRT-PCR experiments were performed using the mouse fibroblast cell line L929

kindly provided by Dr. Blacksell at the Mahidol Oxford Tropical Medicine Research Unit, Bangkok,

Thailand. For all other experiments, we used L929 cells purchased from ATCC (ATCC CCL-1) or

Primary Human Umbilical Vein Endothelial Cells (HUVEC), also from ATCC (ATCC PCS-100-

010). We cultured cells in either DMEM (L929) or Large Vessel Endothelial Supplement (LVES,

HUVEC) supplemented with 10% FBS (Gibco 16140071) at 37ºC and a 5% CO_2 atmosphere.

The Karp-like clinical isolate strain of *Orientia tsutsugamushi* UT76[22] was used throughout the

study. For routine propagation, bacteria were grown in 25- or 75-cm^2 culture flasks, as described

previously[12]. We extracted DNA using alkaline lysis buffer[12], and then assessed bacterial

quantities by determining genome copy numbers relative to a standard curve. We used a

primer/probe set to amplify the single copy gene *tsa47* (tsa47F 5'-

TCCAGAATTAAATGAGAATTTAGGAC-3', tsa47R 5'-TTAGTAATTACATCTCCAGGAGCAA-3',

tsa47probe 5'-[6FAM]TTCCACATTGTGCTGCAGATCCTTC[TAM]-3') or the 16S gene

(Rick16SF 5'- GCTACACGCGTGCTACAATGG-3', Rick16SR 5'-

TGTGTACAAGGCCCGAGAACG-3', Rick16Sprobe 5'-[6FAM]

ATCGCTAGTAATCGCGGATCAGCATGCC [TAM]-3'). For drug inhibition assays, dynasore

(Sigma D7693) and amiloride (Sigma A3085) were added to L929 or HUVEC cells 1 hour before

infection with Ot.

Fluorescence microscopy

Immunofluorescence labelling and confocal microscopy. Following fixation, cells were

permeabilized by incubation in 0.5% Triton X-100 for 30 min on ice followed by blocking in 1XPBS

(pH7.4) containing either 1 mg/ml BSA or sera (diluted 1:20) matching the labeled antibody's host,

for 30 min at room temperature. Incubations with primary antibodies took place for 1 hour at 37ºC.

These studies were conducted using in-house-generated antibodies (*i.e.*, rat monoclonal antibody

against TSA56 diluted 1:200; rabbit polyclonal antibody against Ot ScaC diluted 1:200) or

commercial products (*i.e.*, MA1-164, a Thermo Scientific monoclonal mouse antibody against

LAMP1 diluted 1:200). The fixed cells were washed twice with PBS-BSA and were then incubated

with appropriate secondary antibodies (1:1000 dilutions) for 30 min at 37ºC in the dark. We used

the following secondary antibodies: goat anti-mouse IgG superclonal Alexa Fluor 647 conjugate

(Thermo Fisher A28180), goat anti-rat Alexa Fluor 488 (Thermo Fisher A11006), and goat anti-

rabbit Alexa Fluor 594 (Thermo Fisher A11012). The nuclear stain Hoechst (1:1000 dilution) was

supplemented to incubations with secondary antibodies. Phalloidin Alexa Fluor 594 (1:200,

ThermoFisher Scientific A12381) was used to detect host actin. The cells were washed with 1X

PBS before supplementation of mounting medium [20 mM Tris (pH 8.0); 0.5% N-propyl-gallate;

90% glycerol]. Imaging was performed using a Zeiss Observer Z1 LSM700 confocal microscope

with an HBO 100 illuminating system equipped with a 63x/1.4 Plan-APOCHROMAT objective lens

(Carl Zeiss, Germany) and 405 nm, 488 nm, and 555 nm laser lines. In some cases, we used a

Leica TCS SP8 confocal microscope (Leica Microsystems, Germany) equipped with a 63x/1.4

Plan-APOCHROMAT oil objective lens with a 1.4-mm working distance and 405 nm, 488 nm, 552

nm, and 638 nm laser lines.

Fluorescence intensity measurements were performed manually using Fiji ImageJ software. 30

(Fig. 3C) or 100 (Fig. 1G); individual bacteria were quantified by measuring the brightest pixel

intensity within the bacterial area. Images from individual experiments were acquired under

identical laser intensity and acquisition conditions. Length and width measurements were

performed manually using Fiji Image J software.

STORM imaging and analysis

We acquired 3D-STORM data using a homebuilt multi-color Total Internal Reflection

Fluorescence (TIRF) microscope[26] that is based on an inverted microscope (Nikon, Ti-E), a high

NA objective (Nikon, CFI-Apo 100X, NA 1.49), a 405 nm photoactivation laser (Coherent, OBIS

405 LX 100mW), a 488 nm excitation laser (MPB, 2RU-VFL-P-20000642-B1R), and an EMCCD

camera (Andor, iXon Ultra-888). Samples immunolabelled with Alexa 488 were mounted on the

microscope, transferred to STORM buffer [OxyFluor (Oxyrase), 50 mM BME (Sigma), 2.5 mM

COT (Sigma)], and illuminated with a strong (488 nm) laser to record single-molecule

fluorescence blinking events at a frame rate of 25 Hz. Illumination at 405 nm was intercalated with

pulsed 488nm excitation to enhance photoactivation of Alexa 488 without exacerbating

background signal. A cylindrical lens (Thorlabs, LJ1516RM-A) was placed in front of the camera

to determine the z-coordinates of fluorescent molecules through the elongated point spread

function (PSF) methods[27]. Each STORM image was reconstructed from single-molecule

localization data obtained by analysing 30,000 frames with custom Matlab codes. Briefly, bright

single fluorescence spots were identified by thresholding methods, and their 3-D coordinates were

determined by 2-D elliptical Gaussian fitting along with a pre-calibrated conversion table between

eccentricity and z-coordinate. Focus drift was actively stabilized with Perfect Focus System

(Nikon) while acquiring the data, whereas xy-drift was post-corrected using a redundant cross-

correlation algorithm[28]. Each localization event was represented as a 2-D Gaussian of 20 nm

standard deviation in STORM images. For the analysis of bacterial morphology, we selected cells

that showed clear cell boundary, manually adjusted the z-coordinate to locate the cell midplane,

and measured long- and short- axes length.

Click labelling. Our metabolic click-labelling was based on the Click-iT HPG Alexa Fluor protein

Synthesis Assay Kit (Click-iT HPG Alexa Fluor Protein Synthesis Assay Kit, ThermoFisher

C10428) strategy. To incorporate L-homopropargylglycine (HPG), we incubated infected cells

with minimal medium (Dulbecco's Modified Eagle Medium, DMEM, ThermoFisher 21013) lacking

L-methionine and containing 25 µM HPG, for 30 min at 37ºC. Labelled bacteria were washed

twice with PBS containing 1mg/ml BSA, fixed with 1% formaldehyde, and subsequently

permeabilized with 0.5% Triton X-100 for 20 min on ice. After further washing with PBS containing

1 mg/ml BSA, we supplemented the Click-iT reaction cocktail (Click-iT HPG Alexa Fluor Protein

Synthesis Assay Kit, ThermoFisher C10428) and incubated the cells for 30 min at room

temperature in a light protected environment. The Azide dye (Alexa Fluor 488) was used at a final

concentration of 5 µM. Following completion of the click reaction, the cells were subjected to

immunofluorescence labelling and imaging as above.

Gene expression analysis by RT-qPCR

L929 cells were transfected with *Orientia tsutsugamushi* UT76 using multiple MOIs (*i.e.*, 672:1,

300:1, 560:1, 333:1, and 267:1). Infected cells were harvested four days after transfection and

were stored in RNAProtect Bacteria Reagent (Qiagen 76506) at -80ºC. Total RNA was extracted

using the Qiagen RNeasy Plus kit (Qiagen 74136) according to the manufacturer's instructions.

One µg of total RNA was incubated with DNaseI (Thermo Fisher Scientific, AM2238) for 16 h at

37ºC, and the samples were then reverse-transcribed using the iScript reverse transcription

supermix (Biorad 170–8841) and random primers. Controls lacking reverse transcriptase

demonstrated that the RNA sample used for cDNA synthesis was not contaminated with genomic

DNA. The resulting cDNA was then used as template for qPCR using primers for *spoT, rpoH,*

obgE, yccA, nuoB, dapA, curA, sodB, trxB, tsa56, and scaA. RT-qPCR was performed using

SYBR green qPCR mix (Biotools 10.609). Expression levels were normalized relative to the

housekeeping genes *16sRNA, ank12_1 and mipZ*.

Primer name	Primer sequences (5' to 3')

spoT forward	GGCAGACATAAAGCACCTTAC
spoT reverse	GAAATCTTGAAAGCTGCCAGG
rpoH forward	CTGCCGAAGAGGAATATATGC
rpoH reverse	CCCCTCGGAAACTAAATCAAC
obgE forward	GCACAAAATGGTAGAGCTGG
obgE reverse	GAATCACCAATCTCACCTACAG
yccA forward	GCTATTGCAGGAACCGCTAC
yccA reverse	CTGAAAGCGACATGCCTACA
nuoB forward	GGGCTAGAACACGCTCGTTA
nuoB reverse	GCACAGCTACCCATTGATATC
dapA forward	GGTAGAACATTAGTGATTGCTGG
dapA reverse	GCGTCTTTGATTGCTACTATCC
curA forward	AGTAGGGCAACGAGTTGGC
curA reverse	CAACAGCGCTACTAGCAGC
sodB forward	GTAGTGGTTGGGTCTGGTTG
sodB reverse	GTTCAGCAAATCGCCAATTTG
trxB forward	GATATGCACTGGTGCTCAAG
trxB reverse	CACCTCCTACAACAACTACTC
tsa56 forward	TGCCCAACAAGAAGAAGATG
tsa56 reverse	TTTAACTTGGCCGACAATCA
scaA forward	TAGTGAAAGGGGACGATTGC
scaA reverse	ATCCGACAAGTTAGGCATGG
*16s*RNA forward	CGGTGAATACGTTCTCGGG
*16s*RNA reverse	GAGGTAATCCAGCCGCAGG
ank12_1 forward	GATAGCAGAGGTTGTACACCG

ank12_1 reverse	CTGATTTGTTAGGCTGCCTGC
mipZ forward	GTGGTGCAGGTAAGACTACTGTTG
mipZ reverse	GTGAATGCTGACGTGAATCAG

Proteomics sample preparation

We used *Orientia tsutsugamushi* strain UT76 to infect L929 mouse fibroblasts at a 410:1 MOI in

six independent culture flasks. We replaced the culture medium at one-day post-infection (dpi) to

remove bacteria that failed to enter the host. Intracellular and extracellular bacteria were isolated

at four dpi; two separate flasks were harvested as one replicate, yielding a total of three replicates.

L929 cells were grown in 75-cm^2 plastic flasks at 37°C in a 5% CO_2 atmosphere, using DMEM

media (Gibco 11965092) supplemented with 10% FBS (Sigma-Aldrich F7524). At four dpi, the

media were transferred to 50-ml conical tubes for extracellular bacteria isolation. Culture flasks

were rinsed with 15 ml of PBS, and cells were scraped using two ml of fresh media per flask. The

lysates were transferred to two-ml Safe-lock tubes and homogenized for three minutes using a

Bullet blender$^®$ homogenizer (Next Advance #BBY24M) using speed control setting at 8. Host cell

debris was removed by centrifugation at 300xg for three minutes; we then harvested intracellular

bacteria by an additional three-minute centrifugation at 20,000xg. Before isolation of extracellular

bacteria, we removed detached host cells through a pre-spin step (1,000 xg, 5 min, 4°C).

Following transfer of supernatants to new conical tubes, we harvested extracellular bacteria by

centrifugation at 3000xg for 20 minutes at 4°C. Intracellular and extracellular bacteria were

washed with 0.3 M sucrose three times, resuspended in PBS, and lysed with 1% Triton X-100 for

30 minutes at 4°C. Bacterial proteins were precipitated by addition of 80% ice-cold acetone and

overnight incubation at -20°C. Protein pellets were washed in acetone twice to remove residual

Triton X-100 by centrifugation at 15,000xg for 10 min. Briefly dried protein extracts were then

dissolved in 100 μl of 50% trifluoroethanol in 100 mM TEAB buffer. The samples were reduced

by treatment with 20 mM TCEP for 20 minutes at 55°C, alkylated in 55 mM CAA at room

temperature in the dark, and digested with LysC (LysC: protein ratio=1:50) overnight at 37ºC and,

subsequently, with trypsin (trypsin: protein ratio=1:100), also overnight at 37ºC. Samples were

acidified with TFA to 1% final concentration and centrifuged to remove insoluble materials.

Trypsinized samples were desalted using an Oasis® HLB 1cc cartridge (Waters). Briefly, following

cartridge activation with 500 μl of 100% acetonitrile and equilibration with 500 μl of 0.5% acetic

acid, we loaded samples and then washed the cartridge with 1 ml MS-grade water. Peptides were

eluted with 500 μl of 80% acetonitrile in 0.5% acetic acid, then transferred to low-binding

microtubes, and dried using a SpeedVac. We estimated peptide content using 5% of each

sample. Briefly, individual aliquots in 1 μl of MS-grade water were supplemented with 4 μl of 1%

(v/v) acetonitrile containing 0.06% (v/v) trifluoroacetic acid and 0.5% (v/v) acetic acid, transferred

to a 96-well autosampler plate, and then injected (3 μl) into an Orbitrap Fusion™ Mass

Spectrometer. The remaining 95% of each sample (19 μl in MS-grade water) was mixed with 20

μl of 10 mM ammonium formate (pH 10) and centrifuged at 20,000xg for 3 min to eliminate

insoluble components. Each peptide sample was then fractionated through a step-wise desalting

process on C18 stage tips using 10%, 15%, 25%, and 50% acetonitrile in 10 mM ammonium

formate (pH 10). Fractionated samples were washed with 60% acetonitrile containing 0.1% formic

acid twice to eliminate residual ammonium formate by evaporation before tandem MS analysis.

LC-MS/MS

Dried samples were resuspended in 2% (v/v) acetonitrile containing 0.06% (v/v) trifluoroacetic

acid and 0.5% (v/v) acetic acid, and loaded onto an autosampler plate. Online chromatography

was performed in an EASY-nLC 1000 (ThermoScientific) instrument using a one-column setup

and 0.1% formic acid in water / 0.1% formic acid in acetonitrile as mobile phases. The samples

were injected and separated on a reversed-phase C18 analytical column (EASY-Spray LC

Column, 75 μm inner diameter × 50 cm, 2 μm particle size; ThermoScientific) maintained at 50°C

and using a 2-23% (v/v) acetonitrile gradient over 60 min, followed by an increase to 50% over

the next 20 min, and to 90% over 5 min. The final mixture was maintained on the column for 5

min to elute all remaining peptides. Total run duration for each sample was 90 min at a constant

flow rate of 300 nl/min.

Data were acquired using an Orbitrap Fusion Lumos™ mass spectrometer (ThermoScientific)

using the instrument's data-dependent mode. Samples were ionized using 2.5 kV and 300°C at

the nanospray source; positively-charged precursor MS1 signals were detected using an Orbitrap

analyser set to 60,000 resolution, automatic gain control (AGC) target of 400,000 ions, and

maximum injection time (IT) of 50 ms. Precursors with charges 2-7 and having the highest ion

counts in each MS1 scan were further fragmented using collision-induced dissociation (CID) at

35% normalized collision energy. MS2 signals were analysed by ion trap at an AGC of 15,000

and maximum IT of 50 ms. Precursors used for MS2 scans were excluded for 90 s to avoid re-

sampling of high abundance peptides. The MS1–MS2 cycles were repeated every 3 s until

completion of the run.

Proteomic data analysis

Proteins were identified using Proteome Discoverer™ (v2.3, Thermo Scientific). Raw mass

spectra were searched against *Orientia tsutsugamushi* primary protein sequences derived from

complete genome data for the UT76 strain. Mouse whole proteome sequences were obtained

from Uniprot and included as background. Carbamidomethylation on Cys was set as a fixed

modification; deamidation of asparagine and glutamine, acetylation on protein N termini, and

methionine oxidation were set as dynamic modifications for the search. Trypsin/P was set as the

digestion enzyme and was allowed up to three missed cleavage sites. Precursors and fragments

were accepted if they had a mass error within 10 ppm and 0.8 Da, respectively. Peptides were

matched to spectra at a false discovery rate (FDR) of 1% against the decoy database.

For label-free quantitation (LFQ), features were extracted from the raw data files using the Minora

Feature Detector processing workflow node on Proteome Discoverer. Chromatographic feature-

matching was performed with a tolerance of 10 min to align the retention times and enable

quantitation of the precursors. No pre-normalization was performed using the Proteome

Discoverer workflow, and raw results were manually normalized using the Median normalization

method. A protein was classified as detected if at least two peptides were detected in a minimum

of two biological replicates; the mean LFQ across the three replicates was used for further

analyses.

Immunoblot analyses

L929 and HUVEC cells were grown in 24-well plates and infected with Ot UT76 at an MOI of 12:1.

Non-infectious bacteria were removed by replacing the media 24 hours post-infection. EB and IB

populations were prepared four-seven days post-infection. EB was prepared by transferring the

cell supernatant to an Eppendorf tube, pelleting at 16,873xg, then resuspending in 100 µl PBS

containing 1x protease inhibitor cocktail (ThermoFisher 78425). For TSA56 immunoblots,

samples were prepared in the presence of 1 mM iodoacetamide (Sigma-Aldrich catalog number

I6125). IB was prepared by removing adherent cells using trypsinization, spinning the cell pellet,

and resuspending it in PBS supplemented with protease inhibitors; host cells were lysed using a

bead mill for 1 minute. Host cell debris was removed by centrifugation at 500xg for 5 min; the

supernatant was transferred to a clean tube, spun at 16,873xg for 5 minutes, and resuspended

in 100 µl PBS supplemented with protease inhibitors. Bacterial copy number was quantified by

qPCR using *tsa47* primers, as described above. IB and EB samples were diluted to ensure equal

loading of bacterial cells on gels. Samples were diluted in 5X SDS loading buffer with and without

ß-mercaptoethanol and heated at 95ºC for 5 min. Samples were subjected to electrophoresis on

a 4-20% gradient gel (Bio-Rad 4561094) and then transferred to a nitrocellulose membrane.

Following blocking in PBST containing 5% BSA for 30 min, we incubated the blots with primary

antibody (1:1,000) in PBST-BSA overnight at 4ºC. Membranes were washed and incubated with

secondary antibody (1:10,000) conjugated to IRDYE 680RD or IRDYE 800CW (LI-COR

Biosciences) in PBST-BSA for 1 hour at room temperature. Membranes were washed, and

immunoreactive proteins were visualized using a LiCor Odyssey CLX imager.

Quantification of genome copy number per cell

The genome copy number per cell was calculated by determining the number of bacterial genome

copies in a sample of purified EB and IB Ot, using *tsa47* qPCR, and dividing by the number of

bacterial cells in the same sample, as measured by confocal microscopy. Three µl of purified

bacteria were spread evenly across a well of a multi-well slide, dried, and labelled for

immunofluorescence microscopy using TSA56, as described above. The bacteria in a selected

field were manually counted to determine the number of bacterial cells in the sample.

Quantification of relative peptidoglycan levels using NOD1 reporter assay

HEK-Blue hNOD1 cells (Invitrogen, hkb-hnod1, USA) were grown in DMEM containing 4.5 g/l

glucose, 10% heat-inactivated FBS, 100 U/ml penicillin, 100 mg/ml streptomycin, 100 mg/ml

Normocin, 2 mM L-glutamine, and the selective antibiotics blasticidin (30 µg/ml) and Zeocin (100

µg/ml). The cells were seeded on clear-bottom black 96-well plates (Corning 29444-008) 2 days

before treatment. Bacterial aliquots from stocks kept in sucrose phosphate glutamine (SPG) at -

80°C were heat-inactivated at 90°C for 30 minutes before supplementation to host cells. These

studies were carried out in triplicate. After two days, growth media were replaced with HEK-blue

detection media (Invivogen, hb-det2) for SEAP assessment. The plates were further incubated

for six hours at 37°C in a 5% CO_2 atmosphere and then analyzed spectrophotometrically (Synergy

H1, Biotek) at 640 nm. Results were analyzed by GraphPad Prism (GraphPad Software, San

Diego California USA), and a t-test was performed to compare bacterial SEAP levels to positive

control iE-DAP (γ-D-Glu-mDAP) at 10 µg/ml.

Growth curve measurements and bead mill treatment

To determine the growth rate of IB and EB Ot from frozen stock were first grown in two T25

flasks containing a monolayer of L929 cells. At 6 days post infection, IB and EB were prepared

and used to inoculate fresh L929 cells and the subsequent bacterial growth was

measured. **EB isolation** – the supernatant portion of the culture consisting of EB was

transferred to a 50-mL centrifuge tube (Corning, catalog no. 430828) and subjected to a first

centrifugation to remove any host cells at 1,500 rcf for 5 minutes. The supernatant was then

transferred into a new 50-ml tube and centrifuged at 3,220 rcf for 20 minutes to obtain an EB

pellet which was resuspended in the fresh media and ready for the next step. **IB isolation** –

fresh media was added directly to the culture flasks containing infected cells and these

were disassociated from the adherent surface by using a cell scraper (Corning Falcon,

catalog no. 353089). The infected cell suspension was then pipetted into a sterile 2-mL

microcentrifuge tube. (Eppendorf, catalog no. 022363352) pre-filled with sterile 0.1-0.15 mm

diameter glass beads (Biospec, catalog no. 11079101) before introduced to VWR® Bead Mill

Homogenizer (VWR, catalog no. 10158-558) at speed 5 for 1 minutes to break the host cells.

Subsequently, the purified IB in the supernatant was separated from host cells debris and

glass beads by transferring the supernatant into a new 2-mL microcentrifuge tube and

spinning at 300 rcf for 5 minutes. The bacterial concentration in both EB and IB suspension

was quantified by qPCR as described above, and EB and IB Ot was infected onto a fresh

monolayer of L929 cells in 12-well plates (Thermo Scientific, catalog no. 150628). Bacteria

were added at 2.83E+04 bacterial copies/well which corresponds to an MOI of ~1:10. The

number of intracellular bacteria was then measured at 3, 24, 96 and 168 hours post infection

by *tsa47* qPCR. **Physical stress induced by bead mill** – IB and EB isolation and infection

were performed as described above and then exposed to physical stress with 1 or 2 minutes

agitation in the bead mill. Bacteria were then infected onto L929 cells at a concentration of

1.00E+06 copies/well (MOI = 2:1). The quantity of intracellular bacteria was then determined

at 3, 24, 96, 120, 144 and 168 hours post infection by qPCR. **Osmotic stress induced by**

water treatment. Quantification of the effect of osmotic shock on Ot viability was performed

in the same way as for physical stress except purified bacteria were resuspended in pure

water for 5 minutes prior to infecting host cells. Bacterial copy number after 7 days was

quantified by qPCR.

Acknowledgements and Funding

We thank each laboratory member for valuable discussions and support. We are grateful to the

following people for providing helpful feedback and insightful comments related to the project and

the manuscript: Jonathan Dworkin from Columbia University, Dave Dubnau from PHRI, and Diana

Stafforini from Life Science Editors.

This work was funded by a Royal Society Dorothy Hodgkin Research Fellowship (JS)

(DH140154), a Newton Fund MRC-NSTDA grant (YJ) (MR/N012380/1), and National Institute of

Health/National Institute of Allergy and Infectious Diseases grants (R21AI144385 and

R56AI148645).

References

Taylor, A., Paris, D. & Newton, P. A Systematic Review of Mortality from Untreated Scrub Typhus (Orientia tsutsugamushi). *PLoS Negl Trop Dis* **9**, e0003971, doi:10.1371/journal.pntd.0003971 (2015).

Luce-Fedrow, A. *et al.* A Review of Scrub Typhus (Orientia tsutsugamushi and Related Organisms): Then, Now, and Tomorrow. *Tropical Medicine and Infectious Disease* **3**, 8, doi:10.3390/tropicalmed3010008 (2018).

McClure, E. *et al.* Engineering of obligate intracellular bacteria: progress, challenges and paradigms. *Nat Rev Microbiol*, doi:10.1038/nrmicro.2017.59 (2017).

Salje, J. Cells within cells: Rickettsiales and the obligate intracellular bacterial lifestyle. *Nat Rev Microbiol*, doi:10.1038/s41579-020-00507-2 (2021).

Rikihisa, Y. & Ito, S. Localization of electron-dense tracers during entry of Rickettsia tsutsugamushi into polymorphonuclear leukocytes. *Infect Immun* **30**, 231-243 (1980).

Lee, J. *et al.* Fibronectin facilitates the invasion of Orientia tsutsugamushi into host cells through interaction with a 56-kDa type-specific antigen. *J Infect Dis* **198**, 250-257, doi:10.1086/589284 (2008).

Ha, N., Cho, N., Kim, Y., Choi, M. & Kim, I. An autotransporter protein from Orientia tsutsugamushi mediates adherence to nonphagocytic host cells. *Infect Immun* **79**, 1718-1727, doi:10.1128/IAI.01239-10 (2011).

Ha, N. *et al.* Immunization with an Autotransporter Protein of Orientia tsutsugamushi Provides Protective Immunity against Scrub Typhus. *PLoS Negl Trop Dis* **9**, e0003585, doi:10.1371/journal.pntd.0003585 (2015).

Chu, H. *et al.* Exploitation of the endocytic pathway by Orientia tsutsugamushi in nonprofessional phagocytes. *Infect Immun* **74**, 4246-4253, doi:10.1128/IAI.01620-05 (2006).

Kim, H., Choi, M. & Kim, I. Role of Syndecan-4 in the cellular invasion of Orientia tsutsugamushi. *Microb Pathog* **36**, 219-225, doi:10.1016/j.micpath.2003.12.005 (2004).

Cho, B., Cho, N., Seong, S., Choi, M. & Kim, I. Intracellular invasion by Orientia tsutsugamushi is mediated by integrin signaling and actin cytoskeleton rearrangements. *Infect Immun* **78**, 1915-1923, doi:10.1128/IAI.01316-09 (2010).

Giengkam, S. *et al.* Improved Quantification, Propagation, Purification and Storage of the Obligate Intracellular Human Pathogen Orientia tsutsugamushi. *PLoS Negl Trop Dis* **9**, e0004009, doi:10.1371/journal.pntd.0004009 (2015).

Atwal, S. *et al.* Evidence for a peptidoglycan-like structure in Orientia tsutsugamushi. *Mol Microbiol* **105**, 440-452, doi:10.1111/mmi.13709 (2017).

Min, C. *et al.* Genome-based construction of the metabolic pathways of Orientia tsutsugamushi and comparative analysis within the Rickettsiales order. *Comp Funct Genomics* **2008**, 623145, doi:10.1155/2008/623145 (2008).

Atwal, S. *et al.* Clickable methionine as a universal probe for labelling intracellular bacteria. *J Microbiol Methods* **169**, 105812, doi:10.1016/j.mimet.2019.105812 (2020).

16   Rikihisa, Y. Anaplasma phagocytophilum and Ehrlichia chaffeensis: subversive
manipulators of host cells. *Nat Rev Microbiol* **8**, 328-339, doi:10.1038/nrmicro2318
(2010).
17   Elwell, C., Mirrashidi, K. & Engel, J. Chlamydia cell biology and pathogenesis. *Nat Rev*
Microbiol **14**, 385-400, doi:10.1038/nrmicro.2016.30 (2016).
18   Eldin, C. *et al.* From Q Fever to Coxiella burnetii Infection: a Paradigm Change. *Clin*
Microbiol Rev **30**, 115-190, doi:10.1128/CMR.00045-16 (2017).
19   Otten, C., Brilli, M., Vollmer, W., Viollier, P. & Salje, J. Peptidoglycan in obligate
intracellular bacteria. *Mol Microbiol* **107**, 142-163, doi:10.1111/mmi.13880 (2018).
20   Jacquier, N., Frandi, A., Pillonel, T., Viollier, P. & Greub, G. Cell wall precursors are
required to organize the chlamydial division septum. *Nat Commun* **5**, 3578,
doi:10.1038/ncomms4578 (2014).
21   Irving, S. E., Choudhury, N. R. & Corrigan, R. M. The stringent response and physiological
roles of (pp)pGpp in bacteria. *Nat Rev Microbiol*, doi:10.1038/s41579-020-00470-y
(2020).
22   Batty EM, C. S., Blacksell SB, Richards A, Paris D, Bowden R, Chan C, Lachumanan R, Day
N, Donnelly P, Chen SL, Salje J. Long-read whole genome sequencing and comparative
analysis of six strains of the human pathogen Orientia tsutsugamushi. *Plos Negl Trop Dis*
(2018).
23   Mika-Gospodorz, B. *et al.* Dual RNA-seq of Orientia tsutsugamushi informs on host-
pathogen interactions for this neglected intracellular human pathogen. *Nat Commun* **11**,
3363, doi:10.1038/s41467-020-17094-8 (2020).
24   Joshi, K. K. & Chien, P. Regulated Proteolysis in Bacteria: Caulobacter. *Annu Rev Genet*
50, 423-445, doi:10.1146/annurev-genet-120215-035235 (2016).
25   Lamason, R. *et al.* Rickettsia Sca4 Reduces Vinculin-Mediated Intercellular Tension to
Promote Spread. *Cell* **167**, 670-683.e610, doi:10.1016/j.cell.2016.09.023 (2016).
26   Lee, S. Optimal integration of wide field illumination and holographic optical tweezers
for multimodal microscopy with ultimate flexibility and versatility. *Opt Express* **26**, 8049-
8058, doi:10.1364/OE.26.008049 (2018).
27   Huang, B., Wang, W., Bates, M. & Zhuang, X. Three-dimensional super-resolution
imaging by stochastic optical reconstruction microscopy. *Science* **319**, 810-813,
doi:10.1126/science.1153529 (2008).
28   Wang, Y. *et al.* Localization events-based sample drift correction for localization
microscopy with redundant cross-correlation algorithm. *Opt Express* **22**, 15982-15991,
doi:10.1364/OE.22.015982 (2014).
747

Fig. 1: Intracellular and extracellular forms of Ot are physically and metabolically distinct.
A. STORM images of IB and EB Ot in L929 cells. Bacteria were labelled with a monoclonal antibody against surface protein TSA56. Additional images are provided in Supp. Fig. 1. B. Length quantification of individual IB and EB Ot as imaged by STORM. C. Quantification of relative peptidoglycan levels through activation of NOD1 reporter cells, as measured by secreted alkaline phosphatase activity. EB Ot induced less activation than IB, indicating lower levels of peptidoglycan. Identical amounts of IB and EB Ot were used, and biological triplicates are shown. D. Immunoblot showing the formation of disulfide cross-linked aggregates of the surface protein TSA56, observed in the absence of the reducing agent beta-mercaptoethanol (BME). This effect was stronger in EB compared with IB Ot. The expected molecular weight of TSA56 (56 kDa) is indicated with an arrow, and a dashed arrow depicts higher-order aggregates. E. Genome copy number per cell as measured by qPCR; bacterial cell number was determined by immunofluorescence microscopy. Quantification of 10 individual imaging fields is shown. F. Confocal microscopy image of an L929 cell infected with Ot and labelled with the clickable methionine analog homopropargylglycine (HPG, green) and an antibody against surface protein TSA56 (red). The host cell outline is showed by a dashed line, and an arrow indicates a metabolically inactive bacterium located outside the host cell. G. Quantification of metabolic activity, as measured by a detectable HPG signal, in IB and EB Ot. L929 cells were infected with Ot for five days, fixed, and labelled with TSA56 and HPG. One-hundred bacteria inside (IB) and outside cells (EB) were counted and scored as being HPG-positive or -negative. The percentage of HPG-positive IB and EB Ot, based on individual values from three independent replicates. Statistical significance (t-test) was determined using Graphpad Prism software. p-value *** ≤ 0.001.

Fig. 2: Intracellular and extracellular forms of Ot have distinct gene expression profiles.
A. Pathway analysis of proteins upregulated >2X between IB and EB forms of Ot (60 proteins in IB, 47 proteins in EB). Bacteria were grown in L929 cells at an MOI of 410:1 and harvested four days post-infection. IB and EB Ot were isolated and analysed by shotgun proteomics. B. SpoT and RpoH are upregulated in EB compared with IB Ot as measured by proteomics described in A. C. Differential expression of Ankyrin-repeat-containing proteins (Anks) and tetratricopeptide repeat-containing proteins (TPR) in EB and IB Ot measured by proteomics, as in A. D. RTqPCR analysis showing relative levels of 12 genes expressed in EB and IB Ot. Bacteria were isolated from L929 cells four days post-infection at an MOI of 672:1, 300:1, and 560:1. The figure depicts results from three independent replicates. Gene expression levels were normalised using three genes: *mipZ*, *ank12*, and *16S*. Mean and standard deviation values are shown. The ratio of protein levels in EB/IB, as found in the proteomics experiment in 2A, is given above each gene for comparison.

Fig. 3: Autotransporter proteins ScaA and ScaC are differentially expressed on IB and EB forms of Ot.
A. Volcano plot showing differential expression of surface proteins in proteomics analysis of IB and EB forms of Ot. We observed higher expression of ScaC, TSA56, and TSA47 in EB whilst IB Ot expressed higher ScaA and TSA22 levels B. Immunoblot analysis of IB and EB Ot from infected L929 and HUVEC cells confirming differential expression of ScaA, and ScaC between IB and EB forms. Loading was normalised by measuring bacterial genome copy number using qPCR and loading the same number of IB and EB Ot in each pairwise comparison. Uncropped immunoblot images showing antibody specificity are given in Supp. Fig. 3. C. Quantification of translational activity, as determined by a positive confocal microscopy signal observed with the clickable methionine probe ʟ-homopropylgylglycine (HPG). L929 cells were infected with Ot for five days, then fixed and labelled with HPG and anti-ScaA or anti-ScaC antibodies. ScaA/ScaC-positive bacteria were scored as being positive or negative for HPG labelling. Mean and SD of three independent replicates are shown. D. Confocal microscopy images of HUVEC cells infected with Ot for five days and labelled for DNA (hoescht, blue), translational activity (HPG, green), ScaA (anti-ScaA antibody, magenta) and ScaC (anti-ScaC antibody, red). Scale bar = 10 µm. Examples of extracellular bacteria labelled with ScaC are shown with a white arrow. More images as well as equivalent experiments in L929 cells are shown in Supplementary Figures 4 and 5. E. Quantification of confocal micrographs of HUVEC cells infected with Ot and labelled as in 3D. The percentage of IB and EB Ot at five days post-infection that were positively labelled with anti-ScaA and -ScaC antibodies was counted in three independent experiments. Graph shows mean and SD, and p value determine by unpaired t test. F. Quantification of ScaA- and ScaC-positive IB bacteria inside HUVEC cells one and three hours post-infection, showing an increase in ScaA labelling and a decrease in ScaC labelling. * p-value ≤ 0.05, ** p-value ≤ 0.01, *** p-value ≤ 0.001.

Fig. 4: Both IB and EB populations are infectious but differ in physical stability and mechanism of entry into cells.
A. IB and EB Ot were isolated from infected L929 cells five days post-infection and grown in fresh L929 cells in a 24-well plate. The number of intracellular bacteria was then quantified at 3, 24, 96, and 168 hours post-infection using qPCR against *tsa47,* a single copy Ot gene. Doubling times of IB and EB Ot were similar. B. IB and EB were exposed to physical stress using a bead mill for one or two min; growth was quantified as in 4A. The bead mill did not affect the growth rate of IB Ot but reduced that of EB Ot in an exposure-dependent manner. C. IB and EB were exposed to osmotic shock by suspending the bacteria in water for five min and then growing them in fresh L929 cells. Bacterial copy number after seven days was quantified by qPCR. EB was more susceptible to osmotic shock than IB Ot, as shown by a more significant decrease in bacterial number. D. Quantification of the effect of endocytosis inhibitors on entry of IB and EB Ot. IB and EB were isolated from L929 cells five days post-infection and then seeded onto fresh HUVEC cells that had been treated with inhibitors of clathrin-mediated endocytosis (dynasore) or macropinocytosis (amiloride). Cells were fixed three hours post-infection. Dynasore inhibits entry of IB but not EB, indicating that clathrin-mediated endocytosis is the major mode of entry of this population, whilst amiloride strongly inhibited EB and, to a lesser extent, IB. This observation points to a role for macropinocytosis in EB's entry into mammalian cells. Confocal microscopy images shown in E. Red = actin (phalloidin), blue = hoescht (DNA), green = Ot (anti-TSA56 antibody). Scale bar = 10 μm. Similar results were found in L929 cells (Supp. Fig. 6).

Fig.5: An overview of the infection cycle of Ot, showing differentiation between IB and EB forms.
IB replicate in a micro-colony adjacent to the nucleus where they are elongated in shape and translationally active. Bacteria move to the cell's edge and transition to an EB form that is round and translationally inactive. Created with BioRender.com B. Table summarizing the differences between IB and EB forms of Ot.

IB	EB
more peptidoglycan	less peptidoglycan
elongated and round	round
less fragile	more fragile
TSA56 less cross linked	TSA56 highly cross linked
translationally active	translationally inactive
high expression of Anks	high expression of TPRs
high ScaA/low ScaC	low ScaA/high ScaC
enters cells by endocytosis	enters cells by macropinocytosis