TPX2-dependent spindle positioning dictates division site during asymmetric cell division in moss gametophores

Elena Kozgunova[1,2]*, Mari W. Yoshida[1], Ralf Reski[2,3,4] and Gohta Goshima[1,5]*

[1]Division of Biological Science, Graduate School of Science, Nagoya University, Furo-cho, Chikusa-ku, Nagoya, Aichi 464-8602, Japan
[2]Plant Biotechnology, Faculty of Biology, University of Freiburg, Freiburg 79104, Germany
[3]CIBSS – Centre for Integrative Biological Signalling Studies, University of Freiburg, Freiburg 79104, Germany;
[4]Cluster of Excellence *liv*MatS @ FIT – Freiburg Center for Interactive Materials and Bioinspired Technologies, University of Freiburg, Freiburg 79110, Germany
[5]Sugashima Marine Biological Laboratory, Graduate School of Science, Nagoya University, Sugashima, 429-63, Toba 517-0004, Japan

*Correspondence should be addressed to kozgunova@gmail.com;
goshima@bio.nagoya-u.ac.jp
Phone: +81 52-788-6174

Abstract

Asymmetric cell division (ACD) underlies the development of multicellular organisms. The division site in land plants is predetermined prior to mitosis. The localization of the mitotic spindle is considered uninformative for the division plane, unlike in animal ACD, where the cell division site is defined by active spindle-positioning mechanisms. Here, we isolated a hypomorphic mutant of the microtubule-associated protein TPX2 in moss *Physcomitrium patens* and observed abnormal spindle motility during cell division. This defect compromised the position of the division site and produced inverted daughter cell sizes in the first ACD during gametophore development. The phenotype was rescued by restoring endogenous TPX2 function and, unexpectedly, by depolymerizing actin filaments. Thus, we identified an active spindle-positioning mechanism involving microtubules and actin filaments that set the division site in plants, which is reminiscent of the acentrosomal ACD in animals.

Introduction

Chromosome segregation during mitosis and meiosis is driven by a complex microtubule (MT)-based apparatus known as the spindle. Animal spindles are known to be mobile and their final position corresponds to the future cytokinesis site, which in turn could determine daughter cell fate after asymmetric division. The mechanism by which

the spindle is positioned and spatially controls the assembly of the cytokinetic machinery
has been well studied in animals, and the critical roles of force-generating machinery,
such as dynamic MTs, actin, and motor proteins, have been elucidated (Bergstralh et al.,
2017; Kiyomitsu, 2019). However, in plants, it is believed that the preprophase band
(PPB), a plant-specific MT-actin belt formed prior to mitotic entry, determines the future
cell division site (Buschmann and Müller, 2019; Rasmussen and Bellinger, 2018; Verma,
2001). The mitotic spindle always forms perpendicular to and at the site of the PPB,
perhaps by the action of bridging MTs that connect the spindle to the former PPB site
(Ambrose and Cyr, 2008). The spindle position is considered to be static, unless a strong
force (1600-3350 $\times$ g) is applied through centrifugation, which also causes other
cytoplasmic components to translocate (Arima et al., 2018; Ôta, 1961). The static nature
of spindles is consistent with the fact that plants lack centrosomes, which play key roles
in spindle translocation in animal somatic cells (Bergstralh et al., 2017; Kiyomitsu, 2019).
Multiple proteins co-localized to the PPB are required to establish and maintain the
cortical division zone (CDZ), towards which the cytoskeleton-based cytokinetic
machinery, known as the phragmoplast, expands while recruiting cell plate components
(Müller, 2019; Smertenko et al., 2017). However, the essential role of PPB in determining
the cell division site has been challenged recently by a study that characterized cell
divisions in the roots of *Arabidopsis thaliana trm678* mutants lacking PPB. The findings
showed that PPB is not essential for division plane determination (Schaefer et al., 2017).
In addition, global tissue organization and plant development, although more variable,
were comparable to wild-type plants, implying that ACDs are executable without PPBs.
The moss *Physcomitrium patens* (formerly called *Physcomitrella patens*) is an
attractive model plant for studying PPB-independent division plane determination, as
most cell types naturally lack PPBs, but are capable of oriented cell division and
patterning into complex 3D structures, such as gametophores (leafy shoots) (Kosetsu et
al., 2017; Moody et al., 2018). We have previously shown that the MT structure, called
the gametosome, appears in the cytoplasm transiently at prophase and acts as a
determinant of spindle orientation (Kosetsu et al., 2017). However, gametosomes are
dispensable for spindle MT generation and spindle positioning.
In animal cells, the mitotic spindle is assembled through rapid MT nucleation and
amplification aided by multiple proteins, including γ-tubulin, augmin, and TPX2
(targeting factor for Xklp2) (Petry, 2016). A previous study in *A. thaliana*, using a
combination of knockout and cross-species antibody injection, suggested that TPX2 is an
essential gene (Vos et al., 2008). However, these results were recently questioned when
several viable AtTPX2 t-DNA insertion mutants were obtained (Boruc et al., 2019). In
addition to canonical TPX2, several TPX2-like genes lacking one or more functional
domains have been identified in *A. thaliana*. Among them, TPX2L3 lacks a C-terminal
kinesin-binding motif but is strongly associated with Aurora kinases and is essential for
embryogenesis (Boruc et al., 2019). However, the mechanism by which TPX2 contributes
to spindle formation and MT amplification in plant cells remains unknown.

In this study, we aimed to characterize TPX2 function in the spindle assembly of *P.*
patens, wherein many research tools, including inducible RNAi, endogenous gene
tagging, and highly efficient CRISPR are easily applicable (Leong et al., 2018; Yamada
et al., 2016; Yi and Goshima, 2020). In addition to TPX2's role in MT amplification
during early mitosis, we found an unexpected function of TPX2 in maintaining spindle
position during asymmetric cell division in gametophores.
88
Results
90
P. patens TPX2
We identified five genes homologous to *TPX2* in the *P. patens* genome using a
BLAST search and named them *TPX2-1* to *-5* (Figure 1A). TPX2-1 to -4 proteins showed
high similarity to canonical TPX2 in seed plants (e.g., *A. thaliana* and *Orysa sativa*),
whereas TPX2-5 appears to have lost the N-terminal Aurora-binding motif (Boruc et al.,
2019; Tomaštíková et al., 2015; Vos et al., 2008), but retained the highly conserved C-
terminal domains and, to a certain extent, γ-tubulin activation motifs (Alfaro-Aco et al.,
2017) (Figure 1A, B). During mitosis, endogenous TPX2-1, -2, and -4 proteins fused with
Citrine ("Cit") or mNeonGreen ("NG") or SunTag ("ST") in-frame at the C- or N-
terminus, were enriched at the polar region of the spindle and phragmoplast, suggesting
that they preferentially bind to MT minus ends (Figure 1C, Figure 1 – figure supplement
1, Supplemental Video 1). A similar localization has been reported for *Arabidopsis* TPX2
(Boruc et al., 2019; Vos et al., 2008). TPX2-5 was observed as dim speckles in the spindle
and showed more uniform binding to phragmoplast MTs, suggesting that TPX2-5 might
have an additional function (Figure 1C, Figure 1 – figure supplement 1E-F, Supplemental
Video 1). Unlike TPX2 in animals or seed plants (Boruc et al., 2019; Vos et al., 2008),
none of the TPX2 proteins of *P. patens* were sequestered in the nucleus during interphase
(Figure 1 – figure supplement 2).
109
Generation of hypomorphic mutants of *TPX2*
The similarity in amino acid sequences and intracellular localization suggested that
TPX2-1 to -4 have redundant functions. Therefore, we simultaneously targeted these
genes using a previously established CRISPR/Cas9 protocol (Leong et al., 2018). We
isolated a line, named *TPX2 1-4Δ*, in which frameshifts were introduced to all four *TPX2*
genes in the exons present in all transcript variants identified in the Phytozome database
(Figure 2 – figure supplement 1A). The *TPX2 1-4Δ* line developed protonema (tissue
comprised of frequently dividing tip-growing cells) and gametophores in a similar
manner to the parental "GH" line (Figure 2A, Figure 2 – figure supplement 2). We then
attempted to knock out the *TPX2-5* gene in the *TPX2 1-4Δ* background by means of
homologous recombination. However, we could not isolate a knockout line after multiple
attempts. Given the high efficiency of homologous recombination in *P. patens*, the *TPX2-*
5 gene is likely essential in the *TPX2 1-4Δ* background.

Nonetheless, in our attempt to knockout the *TPX2-5* gene, we isolated three lines
with defective gametophore development after two independent transformations (Figure
2A, Figure 2 – figure supplement 2). In these lines, the original *TPX2-5* gene was replaced
with a hygromycin cassette, as confirmed by PCR and sequencing (Figure 2 – figure
supplement 1B, C). However, DNA was amplified by PCR using *TPX2-5* "internal"
primers, and we confirmed by sequencing that all exons of the *TPX2-5* gene remained in
the genome (Figure 2 – figure supplement 1D). These data suggested that the *TPX2-5*
gene removed from the original locus was re-inserted into another genomic locus,
possibly through micro-homology recombination. We verified that *TPX2-5* relocation
resulted in a compromised expression of *TPX2-5* mRNA by qRT-PCR (Figure 2 – figure
supplement 1E). Hereafter, these hypomorphic lines are referred to as "*TPX2-5 HM1*,"
"*TPX2-5 HM2*" and "*TPX2-5 HM3*."
To test if the lines indeed represent a hypomorphic mutant of *TPX2*, we performed a
rescue experiment in which a frameshift repair construct targeting *TPX2-4* was
transfected into the *TPX2-5 HM1* line (Figure 2 – figure supplement 3A). The selected
lines, in which construct integration was confirmed by PCR, no longer showed a mitotic
delay or dwarf gametophore phenotype (Figure 2 – figure supplement 2B, Figure 2 –
figure supplement 3B-D). We also confirmed that the expression level of *TPX2-4* in the
rescue lines was comparable to that in the control (Figure 2 – figure supplement 3E).
These results strongly suggest that the observed abnormal phenotypes were due to
reduced levels of TPX2 expression.
In addition, we attempted to knockout the *TPX2-5* gene in the wild-type background
(i.e., other *TPX2* genes were intact). Once again, we could not isolate a real knockout line,
but isolated a *TPX2-5* translocation line termed *TPX2-5 M*. However, in the wild-type
background, relocation of *TPX2-5* did not yield any noticeable phenotypes (Figure 2 –
figure supplement 2, Figure 2 – figure supplement 4). These results suggest that TPX2-5
has redundant and distinct functions compared to the other TPX2s.
150
Spindle motility and division plane shift in the *TPX2* mutant gametophore
Next, we aimed to determine whether TPX2 plays a role in the mitosis of
gametophore cells, as dwarf gametophores are the most prominent phenotype of the
TPX2-5 HM1,2,3 lines (Figure 2A, Figure 2 – figure supplement 2B). Dwarf organ
development in plants is often associated with defective cytokinesis (Martinez et al.,
2017). Thus, we first used the lipophilic dye FM4-64 to visualize cell plates in
gametophore initials (stem cells) after the first cell division. We observed that the position
of the cell plate shifted to the basal side of the gametophore initial in multiple cells in all
three *TPX2-5* mutants, dramatically skewing the ratio between apical and basal daughter
cells (Figure 2B, C, Figure 2 – figure supplement 4, Supplemental Video 2). Importantly,
this was due to an overall reduced level of TPX2, as the cell plate positioning was normal
in the aforementioned *TPX2-4* frameshift repair lines and *TPX2-5 M* line (Figure 2B, C,
Figure 2 – figure supplement 4, Supplemental Video 2). Although abnormal cell plate

positioning could explain the dwarf gametophore phenotype, it was unclear what caused
defects in the cell division site in the *TPX2-5 HM1,2,3* lines. Therefore, we next aimed to
perform live-cell imaging of the first cell division in the gametophore initial.
To observe mitotic MTs in gametophore initial cells, we introduced another marker,
mCherry-tubulin, to *TPX2-5 HM1*, *TPX2 1-4Δ,* and control GH lines (note that histone
and tubulin were labeled with the same color). The majority of gametophore initial cells
in the *TPX2-5 HM1* line formed a bipolar spindle (90%; *n* = 19; Figure 3A, B), although
slightly smaller than the spindle in the control or *TPX2 1-4Δ* lines (Figure 3C). Strikingly,
we observed that the bipolar metaphase spindle moved unidirectionally over 5 µm to the
basal side in 73% (n = 14) of *TPX2-5 HM1* cells (Figure 3A, D, Supplemental Video 3).
Consequently, the phragmoplast formed close to the basal edge. We concluded that
defective spindle positioning after NEBD causes abnormal cell plate position in the
gametophore initials of the *TPX2-5 HM1* line.
In six cases, spindle motility or rotation was observed in the second or later divisions,
while the first cell division site appeared normal (Supplemental Video 4). This suggested
that defective ACDs in multiple stages underlie dwarf gametophore formation. Defects
in cell expansion could be another possible explanation for the dwarf gametophore
phenotype, as gametophore cells grow by expansion, controlled by turgor pressure and
cytoskeleton dynamics. To test this possibility, we analyzed cell expansion rates in early
gametophore development (after the first cell division) in the control (GH) and *TPX2-5*
HM1 lines (Figure 3 – figure supplement 1A). We did not detect a statistically significant
difference in the increase in cell size during one cell cycle (Figure 3 – figure supplement
1B). Thus, in the early stages, spindle mispositioning appears to be a primary cause of
defective gametophore development in the *TPX2-5 HM1* line.
188
Spindle motility in the *TPX2* mutant is actin-dependent
In animals, spindle motility relies on MTs and actin. To test the involvement of the
cytoskeleton in spindle motility, we first partially depolymerized MTs in the *TPX2-5 HM*
line using low-dosage oryzalin (200 nM), a MT-destabilizing drug. Upon treatment, the
spindle behaviors were more variable and showed overall differences from the untreated
cells. Most notably, in 5 of 16 cells, we observed that the spindle had shifted towards the
apical side of the cells, which was not observed in untreated cells (Figure 3D). In addition,
the maximum spindle speed was decreased (Figure 3E). In the presence of oryzalin, 9 out
of 11 spindles moved slower than 1 µm/min (maximum speed), whereas only 2 out of 13
did so in control cells (DMSO). These data suggest that MTs contribute to spindle motility
to a certain extent.
We also tested the effect on spindle motility of the MT-stabilizing drug taxol at a
final concentration of 10 µM. The frequency of cells showing spindle motility did not
decrease (18 of 23 cells). However, the motility was slower and the traveling distance
was shorter than that of the control cells (Figure 3E-G). The difference was not
statistically significant, possibly because the spindle motility phenotype showed high

variability in the *TPX2-5 HM1* line. This positive but mild effect might be due to the low
sensitivity of *P. patens* to taxol (Doonan et al., 1988). For example, we never observed
mitotic arrest after taxol treatment, which is expected after high-dose taxol treatment of
animal cells(Jordan et al., 1993).
Next, we investigated the effect of F-actin disruption on spindle motility by adding
latrunculin A, an effective actin inhibitor for *P. patens*, at a final concentration of 5 µM.
Actin inhibition does not affect spindle morphology, orientation, or positioning in the
control line (Kosetsu et al., 2017). Interestingly, 5 µM latrunculin A treatment completely
(20 of 20 cells) suppressed the basal motility of the spindle in the *TPX2-5 HM1* mutant
line (Figure 3A, D, Supplemental Video 3). In addition, we tested several lower
concentrations of latrunculin A and found a concentration-dependent response in spindle
motility. For example, only 2 of 18 gametophore initials still displayed basal spindle
motility when treated with 50 nM latrunculin A (Figure 3E-G). We concluded that actin
is a main driving force of basal spindle motility.
219
Roles of TPX2 in spindle assembly in protonema
ACD occurs in gametophores and in protonema tissue. The protonema consists of
tip-growing chloronemal and caulonemal cells. The apical, tip-growing cells frequently
divide, while subapical cells occasionally form a bulge and divide asymmetrically,
producing a new apical tip cell, in a process known as "branching". We did not observe
spindle motility in the protonema of the *TPX2-5 HM1* line (n = 85, including
caulonemal/chloronemal apical cells and branching cells). The identified differences from
the wild type were the cell plate orientation and apical cell growth speed in caulonemal
cells, which might reflect a slight alteration in MT dynamics in the mutant (Figure 3 –
Supplemental Figure 2). However, it is unlikely that these protonema defects have a
significant impact on gametophore development.
A rare spindle phenotype observed in both protonema (n = 1 of 85) and gametophore
initial cells (n = 2 of 19) was spindle collapse, which led to chromosome missegregation,
cytokinesis failure, and multinucleation (Figure 3B, Supplemental Video 5). In an earlier
study, we showed that a certain percentage of protonema cells of *P. patens* were able to
recover after cytokinesis failure and re-enter the cell cycle, resulting in whole genome
duplication (Kozgunova et al., 2019). In the context of gametophore development, diploid
P. patens has fewer gametophores for unknown reasons (Schween et al., 2005). To
exclude the possibility that *HM* mutants have a high number of diploid protonema cells,
compromising gametophore development, we conducted flow cytometry analysis of
ploidy levels in the protonema cells of control (GH), *TPX2 1-4Δ*, and *TPX2-5 HM1,2,3*
lines. We detected no additional peaks that would indicate higher ploidy in any of these
lines (Figure 3 – figure supplement 3).
To gain insights into the molecular function of moss TPX2 towards MTs, we
performed a detailed analysis of cell division phenotypes in the mutant with the greatest
effect. To this end, we selected an inducible *TPX2-5* RNAi lines in the *TPX2 1-4Δ*

background. Since RNAi induction almost completely inhibited cell growth and
gametophore development, we focused on the division of the protonemal apical stem cells
that appear earlier than gametophores. Using time-lapse imaging of MTs and
chromosomes, we observed severe MT phenotypes during the early stages of mitosis
(Figure 4). During prophase, we detected a reduction in the number of perinuclear MTs
(Figure 4E), which was also observed with a γ-tubulin RNAi (Nakaoka et al., 2012), and
abnormal nuclear shape, which is unique to the *TPX2 1-4Δ* mutant (elongated nucleus
prior to NEBD, Figure 4C). After NEBD, 3 out of 52 cells in the RNAi lines failed to
form bipolar spindles, followed by metaphase arrest and chromosome missegregation,
which was never observed in the control lines (Figure 4D). Other cells formed bipolar
spindles; however, the number of MTs in the prometaphase spindle was greatly reduced
(Figure 4F, blue). This was no longer the case at metaphase, where the number of MTs
was similar to control spindles, indicating the recovery of MT numbers during
prometaphase (Figure 4F, green). Phragmoplast formation and expansion were similar to
those in control cells. This is opposite to the findings in augmin-knockdown cells, in
which the number of metaphase spindle MTs and phragmoplast MTs, but not prophase
MTs, was reduced (Nakaoka et al., 2012). A plausible interpretation is that MTs were
generated and reached control levels through γ-tubulin and augmin-dependent MT
amplification during prolonged prometaphase in the *TPX2* lines (Petry et al., 2013). Thus,
our results suggest that the role of TPX2 in MT amplification is dominant during early
mitosis. As γ-tubulin activation motifs (Alfaro-Aco et al., 2017) are partially conserved
in all moss TPX2 proteins (Figure 1B), we suggest that TPX2 is required for γ-tubulin-
dependent MT nucleation in prophase and prometaphase, while augmin takes over from
prometaphase. Other phenotypes observed upon TPX2 depletion included chromosome
missegregation (29%, n = 15) and spindle misposition/orientation (33%, n = 17, Figure
4D, Supplemental Video 6). Overall, these phenotypes suggested the functional
conservation of moss TPX2 with well-studied animal orthologues, namely assisting in
MT formation through nucleation and/or stabilization.
274
275 **Discussion**
276
In plants, defective division sites have been mostly attributed to defects in PPB
formation (Schaefer et al., 2017; Yoneda et al., 2005), phragmoplast guidance errors
(CDZ deficiency) (Lipka et al., 2014; Müller, 2019), or abnormal positioning of the
nucleus in prophase (Kimata et al., 2019; Yamada and Goshima, 2018). The current study
identified an independent and hitherto unappreciated cause of division site abnormality:
spindle motility after NEBD. Furthermore, we identified a conserved protein, TPX2,
which plays a critical role in spindle positioning in *P. patens*. Interestingly, a recent study
using neural stem cells of the embryonic developing mouse neocortex showed that TPX2
knockdown not only affects spindle MT generation but also spindle orientation,
implicating a conserved TPX2-dependent mechanism of spindle positioning (Vargas-

Hurtado et al., 2019). Spindle-specific positioning defects in meiosis II have also been
observed in *jas* and *ps1* mutants of *Arabidopsis* due to abnormal organelle distribution
(Brownfield et al., 2015). However, meiosis II is a unique system in which two spindles
share a cytoplasm and may be partially fused in the absence of an organelle barrier.
Therefore, there is unlikely to be a mechanistic analogy to moss gametophores. Our data
suggest that defective ACD underlie dwarf gametophores in the *TPX2* mutant. However,
it cannot be ruled out that non-mitotic functions of TPX2 are also required for proper
gametophore development. During gametophore differentiation MT organization
undergoes changes to cortical arrays and PPB-like structure has been observed in mature
gametophores (Doonan et al., 1987; Kosetsu et al., 2017). An interesting future study
would be to investigate whether moss TPX2, which is not sequestered in the nucleus
unlike animal TPX2, regulates interphase MT dynamics and PPB formation in mature
gametophore cells.

In addition to TPX2, this study uncovered the involvement of actin in spindle
positioning. This was also an unexpected observation, as the function of actin in plant cell
division has been mostly attributed to phragmoplast guidance during cytokinesis (Livanos
and Müller, 2019; Rasmussen and Bellinger, 2018). Based on our imaging results, we
suggest that actin filaments are the major driving force in basal spindle motility, while
MTs counteract this force under normal conditions. When the overall levels of TPX2 are
decreased, fewer and/or less stable MTs are insufficient to hold the spindle in place
(Figure 5). This model is supported by the observation that 5 µM latrunculin A completely
abolished basal spindle motility and that a less severe spindle motility phenotype was
observed after taxol treatment, which partially stabilized MTs.

Interestingly, actin is known to play an important role in spindle positioning in animal
cells (Almonacid et al., 2014). Of particular interest are animal oocytes, as they lack
centrosomes. In mouse oocytes, spindle migration and symmetry breaking are driven by
changes in the stability of the actin meshwork, the formation of which depends on actin
nucleators, such as formin-2 and myosin II motor (Almonacid et al., 2014; Duan et al.,
2020). Thus, an analogous mechanism may transmit force to transport spindles in moss.
However, it should be noted that actin function emerged only in the background of a
TPX2 mutation. Thus, sufficient numbers of spindle MTs may predominate in the "tug-
of-war" against actin-dependent forces in wild-type moss cells. TPX2 may be a promising
candidate for investigations to elucidate the PPB-independent mechanism of cell division
site positioning in seed plants.

Materials and methods

P. patens **culture and transformation**

P. patens culture and transformation protocols have been described in detail elsewhere
(Yamada et al., 2016). In brief, BCDAT agar medium (BCDAT stands for stock solutions
<u>B</u>, <u>C</u>, <u>D</u>, and <u>a</u>mmonium <u>t</u>artrate) was used for regular culturing at 25°C under continuous

light. Transformation was performed using a standard polyethylene glycol-mediated
method using protoplasts. Transgenic lines were selected with the corresponding
antibiotics and confirmed by genotyping PCR or sequencing in the case of CRISPR-
generated lines. The GH line, expressing histone H2B-mRFP and GFP-α-tubulin, was
used for transformation in the CRISPR and knockout experiments, while the mCherry-α-
tubulin #52 line was used for Citrine and mNeonGreen endogenous tagging. The
transgenic lines generated in this study are listed in the Supplemental Table 1.
335
Molecular cloning
Plasmids for Citrine or mNeon-Green endogenous tagging were assembled using In-
Fusion (Clontech, Mountain View, CA, USA), in which Citrine or mNeonGreen genes, a
G418 resistance cassette (only C-terminal tagging), and homologous recombination
regions (500-800 bp of the respective genes) were connected. A similar strategy was used
to assemble the knockout plasmid for *TPX2-5*, wherein a hygromycin resistance cassette
was flanked by the 5` and 3`UTR regions of the *TPX2-5* gene. A detailed protocol for
endogenous gene tagging and knockouts in *P. patens* has been previously published
(Yamada et al., 2016). CRISPR gRNAs targeting one of the exons were designed using
the online tool, CRISPOR (http://crispor.tefor.net/), based on target gene specificity (off-
target score) and predicted frameshift efficiency. Individual gRNAs were ligated into
pCasGuide/pUC18 vector pre-digested with BsaI. Next, gRNA sites, together with the
U6 promoter and gRNA scaffold region, were amplified by PCR and assembled into a
single multi-gRNA plasmid, also containing a hygromycin resistance cassette for
transient plant selection. The CRISPR protocol has been described in detail elsewhere
(Leong et al., 2018). RNAi vectors were cloned using the Gateway system (Invitrogen,
Carlsbad, CA, USA), with pGG624 as the destination vector. Two independent, non-
overlapping RNAi constructs were prepared for each gene. The full list of plasmids and
primers used in this study are shown in Supplemental Table 2.
355
Gene expression analysis with quantitative real-time PCR (qRT-PCR)
To determine the expression levels of *TPX2-5* and *TPX2-4* genes, total RNA was
extracted from 7-day-old protonema tissue using the innuPREP Plant RNA kit (Analytik
Jena GmbH, Berlin, Germany) according to the manufacturer's instructions. Total RNA
was treated with DNase I (Thermo Fisher Scientific) at 37°C for 1 h to remove genomic
DNA contamination, and RNA quality was checked by gel electrophoresis. For cDNA
synthesis, we used TaqMan Reverse Transcription Reagents (Thermo Fisher Scientific)
with random hexamers, according to the manufacturer's protocols. Primers were designed
using   the   Universal   Probe   Library   (Roche,
https://lifescience.roche.com/en_de/brands/universal-probe-library.html#assay-design-
center) and selected according to the lowest amount of off-target hits identified in the *P.*
patens transcriptome in Phytozome (www.phytozome.net). The efficiency of each primer
pair was confirmed prior to analysis with a qPCR run using a series of cDNA dilutions

(1:2) and a control without cDNA. Melting curves were analyzed to exclude primer pairs
with off-target amplification. The primer pairs used for qRT-PCR are listed in
Supplemental Table 2. Samples using a cDNA equivalent to 50 ng of total RNA from
corresponding lines were prepared in triplicate or, alternatively, two sets of triplicates for
TPX2-5 in the *TPX2-5 HM1, HM2,* and *HM3* lines, with the SensiMix Kit and SYBR
Green (Bioline, Luckenwalde, Germany). Negative controls without template addition,
as well as reverse transcription controls (reverse transcription reaction carried out without
Multiscribe™ RT), were performed for each primer pair and each line. Amplification was
performed in 40 cycles at a melting temperature of 60°C. qRT-PCR data were analyzed
using LightCycler R 480 software (Roche). The relative expression levels of the gene of
interest (GOI), TPX2-5, and TPX2-4, were compared to those of the control (ctrl), internal
housekeeping genes EF1α (Pp3c2_10310V3.1), and the ribosomal protein L21
(Pp3c13_2360V3.1), and was calculated as $2^{(-\Delta C_T)}$, wherein $\Delta C_T = C_T(\text{GOI}) - C_T(\text{ctrl})$ and
primer efficiency was assumed to be 2. C_T was defined as the cycle number at which each
sample reached an arbitrary threshold (Livak and Schmittgen, 2001).
384

Sample preparation for live-cell imaging

The sample preparation method for live cell imaging has been described in detail in a
previous study (Yamada et al., 2016). In brief, a glass-bottom dish coated with a thin layer
of BCD agar medium (BCD stands for stock solutions <u>B</u>, <u>C</u>, and <u>D</u>. Details can be found
in (Yamada et al., 2016)), was inoculated with moss protonema and cultured under
continuous light at 25°C prior to observation. For gametophore induction, 1 µM of the
synthetic cytokinin, benzylaminopurine, or 1µM 2iP（2-isopentenyladenine）, diluted
in 1 mL of distilled water, was added to the 6 – 7-day-old colony and incubated for 10
min. Next, the remaining liquid was aspirated with a pipette, the dish was sealed, and the
sample was cultured as described above for 20 – 24 h prior to gametophore imaging.
Latrunculin A, taxol or oryzalin were diluted in 1 mL of distilled water to final
concentrations. Prior to drug treatment, most of the agar pad from the glass-bottom dish
was cut and removed to minimize dilution. For RNAi induction, 400 µL of 5 µM β-
estradiol, diluted in distilled water, was added to the pre-cultured protonema 4 d prior to
observation. Although β-estradiol was previously supplemented directly to the agar
medium (Miki et al., 2016), we found that it almost entirely inhibited cell growth in *TPX2-*
5 RNAi lines; hence, the protocol was modified.
402

Microscopy and data analysis

Sample preparation is described above. Localization, RNAi, and most of the gametophore
images were acquired using a Nikon Ti microscope (60 × 1.30-NA lens; Nikon, Tokyo,
Japan) equipped with a CSU-X1 spinning-disk confocal unit (Yokogawa, Tokyo, Japan)
and an electron-multiplying charge-coupled device camera (ImagEM; Hamamatsu,
Hamamatsu, Japan). The microscope was controlled using the NIS-Elements software
(Nikon). Imaging after FM4-64 staining, 10 µM taxol treatment, and low-dosage

latrunculin A treatment (50 nM, 5 nM, and 500 pM) was performed using a ZEISS
inverted microscope (25 × 0.8-NA lens or 63 × 1.4-NA lens, Carl Zeiss Microscopy,
Germany) equipped with a CSU-X1 spinning-disk confocal unit (Yokogawa, Tokyo,
Japan) and CCD camera (Photometrics Prime sCMOS). All imaging was performed at 22
– 24 °C in the dark, except for the first division of the gametophore initial, since
gametophore development requires light (3-min light/2-min dark cycle). For single-leaf
imaging, we dissected gametophores using syringe needles and scissors to isolate single
leaves. Leaves were mounted in a drop of water between two coverslips, and images were
acquired with a Nikon Ti microscope (10 × 0.30-NA lens) in bright-field mode. and image
data were analyzed using ImageJ software (National Institutes of Health, Bethesda, MD,
USA). Prism software was used to plot the graphs and perform statistical analyses
(GraphPad, San Diego, CA, USA). Gametophore and moss colony images were acquired
after 4 weeks of culture using a stereoscopic microscope (SZ2-ILST, Olympus
Corporation, Tokyo, Japan) equipped with a digital camera (Axiocam ICc1; Carl Zeiss
Microscopy, Germany) or a Canon EOS 400 camera, respectively.
425
Cell volume analysis
Cell volume was analyzed in gametophore initial cells stained with FM4-64 dye diluted
in distilled water to a final concentration of $10\,\mu M$. The dye was added to the live-imaging
dish before acquisition, without cutting the agar. Imaging was performed immediately
after FM4-64 application. Images were acquired as a Z-stack ($40\,\mu m$, $0.97\,\mu m$ step) using
a ZEISS inverted microscope (25 × 0.8-NA lens) equipped with a CSU-X1 spinning-disk
confocal unit (Yokogawa, Tokyo, Japan) and a CCD camera (Photometrics Prime
sCMOS). Cell boundary predictions and segmentation were performed using PlantSeg
software (Wolny et al., 2020). Default conditions of PlantSeg were used with the
following modifications: images were rescaled using in-build function (for our images
voxel size was $0.26\,\mu m$ on the X-Y dimensions and $0.97\,\mu m$ on the Z dimension) and cell
minimum size was set at 400 000 voxels. The accuracy of the cell predictions was
confirmed by comparing PlantSeg output files to raw images for each cell. In some cases,
when cell boundaries were not predicted accurately, images were deconvolved using
Huygens Professional version 19.04 (Scientific Volume Imaging, Hilversum,
Netherlands) and again run through PlantSeg with a minimum cell size of 300 000 voxels.
If this did not improve the cell boundary predictions, images were excluded from the
analysis. Cell volume was extracted from PlantSeg output tiff files using a threshold
function and the Voxel counter plugin in ImageJ. 3D projections in Supplemental video
2 were created with Imaris software (Bitplane) version 9.7.0.
446
Flow cytometry analysis of ploidy
Moss protonema was homogenized in 5 mL of distilled water and cultured on BCDAT
agar plates covered with cellophane for 6 days before measurement. Next, small portion

of protonema was chopped with a blade in 4′,6-diamidino-2-phenylindole (DAPI)
solution (0.01 mg/L DAPI, 1.07 g/L $MgCl_2 \cdot 6H_2O$, 5 g/L NaCl, 21.11 g/L TRIS in 1 ml
Triton) and filtered through a 30 µm sieve. Fluorescence of the nuclei was determined
with a PAS cell analyzer (Partec, Munster, Germany) using a 100 W high-pressure
mercury lamp (Schween et al., 2003).
455

Sequence analysis
We used MAFFT ver. 7.043 (https://mafft.cbrc.jp/alignment/software/) to align the amino
acid sequences of the selected full-length proteins, and then manually revised them with
MacClade ver. 4.08 OSX (www.macclade.org) to remove gaps. The Jones-Taylor-
Thornton (JTT) model was used to construct maximum-likelihood trees using MEGA5
software (www.megasoftware.net). Statistical support for internal branches by bootstrap
analysis was obtained using 1,000 replications. The gene sequence information discussed
in this article is available under the following accession numbers in Phytozome
(www.phytozome.net): *At*TPX2 (AT1G03780.3); *Os*TPX2 (LOC_Os07g32390.1);
*Pp*TPX2-1 (Pp3c17_11160V3.1); *Pp*TPX2-2 (Pp3c1_25950V3.1); *Pp*TPX2-3
(Pp3c24_8590V3.2); *Pp*TPX2-4 (Pp3c23_4540V3.1); *Pp*TPX2-5 (Pp3c5_10270V3.1);
*Mp*TPX2-1 (Mapoly0016s0083.1); *Mp*TPX2-2 (Mapoly0105s0040.1) or in UNIPROT
(www.uniprot.org): *Hs*TPX2 (Q9ULW0); *Gg*TPX2 (F1NW64); *Xl*TPX2 (Q6NUF4).
469

Acknowledgments
We thank Momoko Nishina and Yuki Nakaoka for assistance with this project, Sebastian
Hoernstein for help and advice on flow cytometry and RNA extraction, Lennard
Bohlender for help with qRT-PCR, Raymundo Alfaro-Aco for comments on the TPX2
functional motifs, Peishan Yi and Mariana Costa for helpful comments on the manuscript,
and the Life Imaging Center at the University of Freiburg for microscopic use. This work
was funded by JSPS KAKENHI (17H06471), JSPS and DFG under the Joint Research
Projects-LEAD with UKRI (to G.G.), and an Alexander von Humboldt Foundation
Postdoctoral Fellowship (to E.K.). Additional support came from the German Research
Foundation (DFG) under Germany's Excellence Strategy (CIBSS – EXC-2189 – Project
ID 390939984 to R.R.). The authors declare no competing interests.

Author contributions
E.K. and G.G. designed the research project, G.G. and R.R supervised the project, E.K.
and M.W.Y. performed experiments, E.K. analyzed the data, E.K., G.G. and R.R. wrote
and edited the manuscript.
486

References

Alfaro-Aco R, Thawani A, Petry S. 2017. Structural analysis of the role of TPX2 in branching microtubule nucleation. *J Cell Biol* **177**:7–11. doi:10.1083/jcb.200611141

Almonacid M, Terret MÉ, Verlhac MH. 2014. Actin-based spindle positioning: New insights from female gametes. *J Cell Sci* **127**:477–483. doi:10.1242/jcs.142711

Ambrose JC, Cyr R. 2008. Mitotic spindle organization by the preprophase band. *Mol Plant* **1**:950–960. doi:10.1093/mp/ssn054

Arima K, Tamaoki D, Mineyuki Y, Yasuhara H, Nakai T, Shimmen T, Yoshihisa T, Sonobe S. 2018. Displacement of the mitotic apparatuses by centrifugation reveals cortical actin organization during cytokinesis in cultured tobacco BY-2 cells. *J Plant Res* **131**:803–815. doi:10.1007/s10265-018-1047-4

Bergstralh DT, Dawney NS, Johnston DS. 2017. Spindle orientation : a question of complex positioning. *Development* **5**:1137–1145. doi:10.1242/dev.140764

Boruc J, Deng X, Mylle E, Besbrugge N, Durme M Van, Demidov D, Tomaštíková ED, Tan T-RC, Vandorpe M, Eeckhout D, Beeckman T, Nowack M, Jaeger G De, Lin H, Liu B, Damme D Van. 2019. The TPX2-LIKE PROTEIN 3 is the primary activator of α Aurora kinases and is essential for embryogenesis. *Plant Physiol.* doi:10.1101/466276

Brownfield L, Yi J, Jiang H, Minina EA, Twell D, Köhler C. 2015. Organelles maintain spindle position in plant meiosis. *Nat Commun* **6**. doi:10.1038/ncomms7492

Buschmann H, Müller S. 2019. Update on plant cytokinesis: rule and divide. *Curr Opin Plant Biol* **52**:97–105. doi:10.1016/j.pbi.2019.07.003

Doonan JH, Cove DJ, Corke FMK, Lloyd CW. 1987. Pre-prophase band of microtubules, absent from tip-growing moss filaments, arises in leafy shoots during transition to intercalary growth. *Cell Motil Cytoskeleton* **7**:138–153. doi:10.1002/cm.970070206

Doonan JH, Cove DJ, Lloyd CW. 1988. Microtubules and microfilaments in tip growth : evidence that microtubules impose polarity on protonemal growth in Physcomitrella patens. *J Cell Sci* **89**:533–540.

Duan X, Li Y, Yi K, Guo F, Wang HY, Wu PH, Yang J, Mair DB, Morales EA, Kalab P, Wirtz D, Sun SX, Li R. 2020. Dynamic organelle distribution initiates actin-based spindle migration in mouse oocytes. *Nat Commun* **11**:277. doi:10.1038/s41467-019-14068-3

Jordan MANN, Toso RJ, Thrower D, Wilson L. 1993. Mechanism of Mitotic Block and Inhibition of Cell Proliferation by Taxol at Low Concentrations. *Proc Natl Acad Sci* **90**:9552–9556.

Kimata Y, Kato T, Higaki T, Kurihara D, Yamada T, Segami S, Morita MT, Maeshima M, Hasezawa S, Higashiyama T, Tasaka M, Ueda M. 2019. Polar vacuolar

distribution is essential for accurate asymmetric division of Arabidopsis zygotes.
Proc Natl Acad Sci U S A **116**:2338–2343. doi:10.1073/pnas.1814160116

Kiyomitsu T. 2019. The cortical force-generating machinery: how cortical spindle-
pulling forces are generated. *Curr Opin Cell Biol* **60**:1–8.
doi:10.1016/j.ceb.2019.03.001

Kosetsu K, Murata T, Yamada M, Nishina M, Boruc J, Hasebe M, Van Damme D,
Goshima G. 2017. Cytoplasmic MTOCs control spindle orientation for asymmetric
cell division in plants. *Proc Natl Acad Sci* **114**:E8847–E8854.
doi:10.1073/pnas.1713925114

Kozgunova E, Nishina M, Goshima G. 2019. Kinetochore protein depletion underlies
cytokinesis failure and somatic polyploidization in the moss Physcomitrella patens.
Elife **8**:1–16. doi:10.7554/eLife.43652

Leong SY, Yamada M, Yanagisawa N, Goshima G. 2018. SPIRAL2 Stabilises
Endoplasmic Microtubule Minus Ends in the Moss *Physcomitrella patens*. *Cell
Struct Funct* **43**:53–60. doi:10.1247/csf.18001

Lipka E, Gadeyne A, Stöckle D, Zimmermann S, De Jaeger G, Ehrhardt DW, Kirik V,
Van Damme D, Müller S. 2014. The phragmoplast-orienting kinesin-12 class
proteins translate the positional information of the preprophase band to establish
the cortical division zone in Arabidopsis thaliana. *Plant Cell* **26**:2617–2632.
doi:10.1105/tpc.114.124933

Livak KJ, Schmittgen TD. 2001. Analysis of relative gene expression data using real-
time quantitative PCR and the 2-ΔΔCT method. *Methods* **25**:402–408.
doi:10.1006/meth.2001.1262

Livanos P, Müller S. 2019. Division Plane Establishment and Cytokinesis. *Annu Rev
Plant Biol* **70**:239–267. doi:10.1146/annurev-arplant-050718-100444

Martinez P, Luo A, Sylvester A, Rasmussen CG. 2017. Proper division plane
orientation and mitotic progression together allow normal growth of maize. *Proc
Natl Acad Sci U S A* **114**:2759–2764. doi:10.1073/pnas.1619252114

Miki T, Nakaoka Y, Goshima G. 2016. Live Cell Microscopy-Based RNAi Screening in
the Moss Physcomitrella patens. *Methods Mol Biol* **1470**:225–246.
doi:10.1007/978-1-4939-6337-9_18

Moody LA, Kelly S, Rabbinowitsch E, Langdale JA. 2018. Genetic Regulation of the
2D to 3D Growth Transition in the Moss Physcomitrella patens. *Curr Biol* **28**:473-
478.e5. doi:10.1016/j.cub.2017.12.052

Müller S. 2019. Plant cell division — defining and finding the sweet spot for cell plate
insertion. *Curr Opin Cell Biol* **60**:9–18. doi:10.1016/j.ceb.2019.03.006

Nakaoka Y, Miki T, Fujioka R, Uehara R, Tomioka A, Obuse C, Kubo M, Hiwatashi Y,
Goshima G. 2012. An Inducible RNA Interference System in Physcomitrella
patens Reveals a Dominant Role of Augmin in Phragmoplast Microtubule
Generation. *Plant Cell* **24**:1478–1493. doi:10.1105/tpc.112.098509

Ôta T. 1961. The Role of Cytoplasm in Cytokinesis of Plant Cells. *Cytologia (Tokyo)*
26:428–447. doi:10.1508/cytologia.26.428

Petry S. 2016. Mechanisms of Mitotic Spindle Assembly. *Annu Rev Biochem* **85**:659–
683. doi:10.1146/annurev-biochem-060815-014528

Petry S, Groen AC, Ishihara K, Mitchison TJ, Vale RD. 2013. Branching microtubule
nucleation in xenopus egg extracts mediated by augmin and TPX2. *Cell* **152**:768–
777. doi:10.1016/j.cell.2012.12.044

Rasmussen CG, Bellinger M. 2018. An overview of plant division-plane orientation.
New Phytol **219**:505–512. doi:10.1111/nph.15183

Schaefer E, Belcram K, Uyttewaal M, Duroc Y, Goussot M, Legland D, Laruelle E, De
Tauzia-Moreau ML, Pastuglia M, Bouchez D. 2017. The preprophase band of
microtubules controls the robustness of division orientation in plants. *Science*
356:186–189. doi:10.1126/science.aal3016

Schween G, Gorr G, Hohe A, Reski R. 2003. Unique tissue-specific cell cycle in
Physcomitrella. *Plant Biol* **5**:50–58. doi:10.1055/s-2003-37984

Schween G, Schulte J, Reski R. 2005. Effect of Ploidy Level on Growth ,
Differentiation , and Morphology in Physcomitrella patens. *Bryologist* **108**:27–35.

Smertenko A, Assaad F, Baluška F, Bezanilla M, Buschmann H, Drakakaki G, Hauser
MT, Janson M, Mineyuki Y, Moore I, Müller S, Murata T, Otegui MS, Panteris E,
Rasmussen C, Schmit AC, Šamaj J, Samuels L, Staehelin LA, Van Damme D,
Wasteneys G, Žárský V. 2017. Plant Cytokinesis: Terminology for Structures and
Processes. *Trends Cell Biol* **27**:885–894. doi:10.1016/j.tcb.2017.08.008

Tomaštíková E, Demidov D, Jeřábková H, Binarová P, Houben A, Doležel J, Petrovská
B. 2015. TPX2 Protein of Arabidopsis Activates Aurora Kinase 1, But Not Aurora
Kinase 3 In Vitro. *Plant Mol Biol Report* **33**:1988–1995. doi:10.1007/s11105-015-
0890-x

Vargas-Hurtado D, Brault JB, Piolot T, Leconte L, Da Silva N, Pennetier C, Baffet A,
Marthiens V, Basto R. 2019. Differences in Mitotic Spindle Architecture in
Mammalian Neural Stem Cells Influence Mitotic Accuracy during Brain
Development. *Curr Biol* **29**:2993-3005.e9. doi:10.1016/j.cub.2019.07.061

Verma DPS. 2001. Cytokinesis and building the cell wall in plants. *Annu rev plant*
Physiol **52**:751–784.

Vos JW, Pieuchot L, Evrard J-L, Janski N, Bergdoll M, De Ronde D, Pérez LH, Sardon
T, Vernos I, Schmit A-C. 2008. The Plant TPX2 Protein Regulates Prospindle
Assembly before Nuclear Envelope Breakdown. *Plant Cell* **20**:2783–2797.
doi:10.1105/tpc.107.056796

Wolny A, Cerrone L, Vijayan A, Tofanelli R, Barro AV, Louveaux M, Wenzl C,
Strauss S, Wilson-Sánchez D, Lymbouridou R, Steigleder SS, Pape C, Bailoni A,
Duran-Nebreda S, Bassel G, Lohmann JU, Tsiantis M, Hamprecht FA, Schneitz K,
Maizel A, Kreshuk A. 2020. Accurate and versatile 3D segmentation of plant

608   tissues at cellular resolution. *Elife* **9**:1–34. doi:10.7554/eLife.57613

609 Yamada M, Goshima G. 2018. The KCH kinesin drives nuclear transport and
610   cytoskeletal coalescence to promote tip cell growth in Physcomitrella patens. *Plant*
611   *Cell* **30**:1496–1510. doi:10.1105/tpc.18.00038

612 Yamada M, Miki T, Goshima G. 2016. Imaging Mitosis in the Moss Physcomitrella
613   patens. *Methods Mol Biol* **1413**:293–326. doi:10.1007/978-3-319-02904-7

614 Yi P, Goshima G. 2020. Transient cotransformation of CRISPR /Cas9 and
615   oligonucleotide templates enables efficient editing of target loci in Physcomitrella
616   patens. *Plant Biotechnol J* 1–3. doi:10.1111/pbi.13238

617 Yoneda A, Akatsuka M, Hoshino H, Kumagai F, Hasezawa S. 2005. Decision of
618   spindle poles and division plane by double preprophase bands in a BY-2 cell line
619   expressing GFP-tubulin. *Plant Cell Physiol* **46**:531–538. doi:10.1093/pcp/pci055

Figure 1. TPX2 homologues and their localization in *P. patens*

(**A**) Phylogeny analysis revealed two distinct groups of TPX2 proteins in *P. patens*: Pp *TPX2-1* to *-4* (blue), which are more similar to *TPX2* genes from seed plants, and atypical *TPX2-5* (magenta). Asterisks mark predicted proteins, numbers show bootstrap values. Bar, 0.5 amino acid substitutions per site. Note that AtTPX2L3 and AtTPX2L2 could not be added to this tree, since they lack the C-terminal region that is conserved in canonical TPX2 proteins. Hs: *Homo sapiens*, Gg: *Gallus gallus*, Xl: *Xenopus laevis*, At: *Arabidopsis thaliana*, Os: *Oryza sativa*, Pp: *Physcomitrium patens*, Mp: *Marchantia polymorpha*. (**B**) Alignment of TPX2 proteins. Conserved residues are boxed, whereas similar amino acids are hatched. (**C**) Localization of endogenous TPX2-1-Citrine, mNeonGreen-TPX2-4 and TPX2-5-mNeonGreen. More uniform spindle localization was detected for TPX2-5. Asterisks indicate autofluorescent chloroplasts. Bar, 10 μm. The full version of mitotic localization data is presented in Figure 1 - figure supplement 1.

Figure 1 – figure supplement 1. Localization of TPX2 proteins during mitosis
Live-cell imaging was performed in caulonemal apical cells of *P. patens*, expressing mCherry-tubulin and
one of the following TPX2 proteins endogenously tagged with a fluorophore: (A) TPX2-1-Citrine; (B)
SunTag-TPX2-1; (C) TPX2-2-mNeonGreen; (D) mNeonGreen-TPX2-4; (E) TPX2-5-mNeonGreen; (F)
mNeonGreen-TPX2-5. The SunTag-TPX2-1 line also expressed scFv-GCN-sfGFP under a β-estradiol-
inducible promoter. Asterisks indicate the autofluorescent chloroplasts. Bars, 10 µm.

Figure 1 – figure supplement 2. Interphase localization of TPX2

(**A**) Live-cell imaging was performed in caulonemal apical cells of *P. patens*, expressing only mCherry-tubulin (upper panels) or mCherry-tubulin and TPX2-1-Citrine (bottom panels). Cytoplasmic signals increased in the TPX2-1-Citrine line. Bars, 20 µm. (**B**) Fluorescence intensity in the nucleus before and after inhibiting nuclear export with 100 nM Leptomycin B (mean ± SEM) after subtracting the cytoplasmic background. Increase in nuclear signals after nuclear export inhibition suggests that TPX2-1-Cit and TPX2-2-NG are actively shuttled between the nucleus and cytoplasm. Bar, 10 µm.

Figure 2. Abnormal cell division site in the gametophore initial of a *TPX2-5 HM1* mutant
(**A**) Representative photos of gametophores after 4 weeks of culture of GH (control), *TPX2 1-4Δ*, and *TPX2-5 HM1* lines. Brightness/contrast have been linearly adjusted. Extended version is presented in Figure 2 - figure supplement 2B. Bar, 2 mm. (**B**) Gametophore initial at the 2-cell stage stained with FM4-64 dye. Normal and defective cell plate positions are indicated with cyan and red arrowheads, respectively. The full version can be found in Figure 2 - figure supplement 4A. Bar, 10 µm (**C**) The apical/basal cell volume ratio was estimated as the apical cell (pink) volume divided by the basal cell (green) volume, measured during the 2-cell stage (mean ± SEM, ****p=0.0001 by one way Anova with Dunnett's multiple comparison test against GH). n = 22, 15, 18, 10 and 18 for GH, TPX2 1-4Δ, *TPX2-5 HM1*, TPX2-4 *repair* #16, and TPX2-4 *repair* #19. The full version can be found in Figure 2 - figure supplement 4B.

Figure 2 – figure supplement 1. Isolation of hypomorphic *TPX2* mutants
(**A**) Schematic explanation of *P. patens* lines created and used in this study and representative sequencing
data of frameshift mutations in the *TPX2 1-4Δ* line. (**B**) Schematic explanation of *TPX2-5* hypomorphic
line selection and genotyping PCR. (**C**) Results of genotyping PCR and sequencing of the *TPX2-5* locus.
For unknown reasons, first round of in F/in R PCR yielded barely visible bands. Therefore, we performed
a second round of PCR using diluted product of the first PCR as the template. (**D**) Results of amplifying
full coding sequence from start to stop codons and sequencing results of the *TPX2-5* gene. Note the band
shift in the *TPX2-5 HM1* line due to loss of introns. (**E**) qRT-PCR analysis of *TPX2-5* expression
normalized against the expression of the internal housekeeping genes *EF1α* and *L21*. Bars represent mean
values ± SD from n = 3.

Figure 2 – figure supplement 2. Development of P. patens colonies and gametophores after 4 weeks of culture

Representative images of moss colonies **(A)** or gametophores **(B)** for all lines (except RNAi) used in this study after 4 weeks of culture on BCDAT agar plates. The brightness and contrast were linearly adjusted. Some photos are the duplicates of those in Figure 2A. Bars, 10 mm and 2 mm, respectively.

Figure 2 – figure supplement 3. Rescue of the *TPX2-5 HM* phenotypes by frameshift repair of the *TPX2-4* gene

(**A**) Schematic illustration of the frameshift repair experiment. In brief, the N-terminus-coding region of the *TPX2-4* gene in the *TPX2-5 HM1* background was tagged with Cerulean flanked with ~500 bp of the 5′-UTR and exon region (without the frameshift mutation) by homologous recombination. Construct integration was verified by PCR. (**B**) Mitotic duration of protonemal cells calculated from NEBD to anaphase onset in GH (control), *TPX2 1-4Δ*, *TPX2-5 HM,* and two independent *TPX2-4* repair lines (mean ± SEM, ****p=0.0001 by one way Anova with Dunnett's multiple comparison test against GH). (**C**) Mitotic progression in the *TPX2-4* repair #16 line. Note that perinuclear MTs and prometaphase spindle formation were restored. Bar, 10 μm. (**D**) Representative images of gametophore leaf cells in GH, *TPX2 1-4Δ*, *TPX2-5 HM*, and *TPX2-4* repair #16 lines. Bar, 100 μm. (**E**) qRT-PCR analysis of *TPX2-4* expression normalized against the expression of the internal housekeeping genes *EF1α* and *L21*. No significant difference between GH and *TPX2-4 repair* #16 and #19 by one-way Anova test and t-test.

Figure 2 – figure supplement 4. Cell division site and apical/basal cell ratio at the 2-cell stage of gametophore development in different TPX2 backgrounds

(**A**) Gametophore initial at the 2-cell stage stained with FM4-64 dye. Bar, 10 µm (**B**) The apical/basal cell volume ratio was estimated as the apical cell volume divided by the basal cell volume, measured during the 2-cell stage (mean ± SEM, ****p=0.0001 by one way ANOVA with Dunnett's multiple comparison test against GH). n = 22, 15, 18, 18, 17, 10, 18 and 8 for GH, TPX2 1-4Δ, *TPX2-5 HM1*, *TPX2-5 HM2*, *TPX2-5 HM3*, TPX2-4 *repair* #16, TPX2-4 *repair* #19 and *TPX2-5 M*.

Figure 3. Spindle position is actively maintained through the interplay between microtubules and F-actin

(**A**) Live-cell imaging of the first asymmetric division in the gametophore initial revealed a link between the metaphase spindle and phragmoplast positioning. The positions of the nucleus and gametosome (prophase MTOC appeared in the apical cytoplasm) are indicated with yellow circles and red arrowheads, respectively. Cyan lines show the position and orientation of the phragmoplast. Cell borders are outlined

with white lines. Bar, 10 µm. **(B)** The frequency and type of spindle defects in gametophore initial mitosis
observed in GH (control), *TPX2 1-4Δ*, and *TPX2-5 HM* lines. Numbers within the columns indicate number
of cells with corresponding phenotypes. **(C)** Area occupied by the metaphase spindle (spindle size) in
gametophore initials. (mean ± SEM; **p = 0.0029, two-tailed Student's t-test) **(D)** Tracking of the spindle
center position from NEBD to anaphase onset. We assigned the starting position as Y = 0 and different X
positions for each sample group. Note that after 5 µM latrunculin A treatment, spindles never showed
motility towards the basal end of the cell, i.e. negative Y-values. Each line represents spindle movement in
a single cell. More than 12 cells were observed for each sample group in three or more independent
experiments. **(E)** Maximum spindle speed (µm/min), **(F)** average spindle speed (µm/min), and **(G)** distance
travelled by the spindle (µm) in the *TPX2-5 HM1* cell line under various treatments. Total number of cells
observed, including those without spindle motility: n = 20 (DMSO), 18 (50 nM LatA), 12 (5 nM LatA), 13
(500 pM LatA), 23 (10 µM taxol) and 16 (200 nM oryzalin). Only spindle motility towards the basal end
of the cell was analyzed. Bars represent mean ± SEM.
721

Figure 3 - figure supplement 1. Analysis of cell growth in early gametophore development

(A) Representative images of cell expansion in gametophore initials during one cell cycle (from
disappearance of phragmoplast to nuclear envelope breakdown) in GH and TPX2-5 HM1 lines. Cell
borders are indicated with yellow lines. Images were acquired as Z-stack and the best focal plate was
selected for analysis. Bar, 10 µm (B) Quantification of the increase in the cell area during one cell cycle.
Cell area was first measured at the end of the cell division. The value was taken as 100%. The next
measurement was taken at the onset of the subsequent cell division and values were plotted as increase
percentage. Only gametophores with basal spindle motility were analyzed for *TPX2-5 HM1* line. Bars
represent mean ± SEM.

Figure 3 - figure supplement 2. Phragmoplast orientation and tip growth defects in the caulonemal tip cells in the *TPX2-5 HM1* line

(A)Representative images of cell plates in caulonemal cells stained with 10 μM FM4-64 dye. Bar, 50 µm (B) Acute angle of caulonemal cells in GH (control), *TPX2 1-4Δ* and *TPX2_5 HM1* lines, n = 24, 24, and 23, respectively (mean±SEM, **** p = 0.0001 one way ANOVA with Dunnett's multiple comparisons test) (C) Tip growth rate of caulonemal cells in GH, TPX2 1-4Δ and TPX2_5 *HM1* lines, n = 46, 50 and 48, respectively (mean±SEM, * p = 0.0144 one way ANOVA with Dunnett's multiple comparisons test).

745 **Figure 3 - figure supplement 3. Ploidy analysis by flow cytometry**

746 Histograms of nuclei isolated from 6-day-old protonema tissues stained with DAPI. Peak corresponding
747 to G2 (2N) in the control line (GH) is indicated by arrow.

748
Figure 4. TPX2 contributed to microtubule amplification in early mitosis

(A) Representative images of the mitosis of protonemal apical cells in GH (control), *TPX2 1-4Δ*, *TPX2-*
5$^{5'UTR}$ RNAi, and *TPX2-5^{exon}* RNAi lines. Green, GFP-tubulin; Magenta, histoneH2B-mRFP. Bar, 10 µm.
(B) Mitotic duration calculated from NEBD to anaphase onset (mean ± SEM; ****$p < 0.0001$, two-tailed
Student's t-test). **(C)** Nucleus shape prior to NEBD, measured as a ratio of nucleus length to nucleus width
(mean ± SEM; ****p < 0.0001, two-tailed Student's t-test). **(D)** Frequency and type of mitotic defects
observed. Bar, 10 µm. **(E)** Fluorescence intensity of perinuclear MTs (mean ± SEM; **$p = 0.0011$, ***$p =$
0.0002; two-tailed Student's t-test). A.U. stands for Arbitrary Units. **(F)** Fluorescence intensity of MTs in
the prometaphase spindle (4 min after NEBD) and metaphase spindle (2 min before anaphase onset),
measured from a single focal plane, with the cytoplasmic background subtracted. A decrease in
fluorescence intensity was detected in prometaphase, but not in metaphase spindles (mean ± SEM **$p =$
0.0018, ****$p \leq 0.0001$, ***$p = 0.0008$; two-tailed Student's t-test).

Figure 5. Schematic representation of the spindle impact on phragmoplast position and orientation in the gametophore initial cells of *P. patens*

The upper panel shows the role of the microtubule-organizing center (gametosome) in spindle and phagmoplast orientation, based on Kosetsu et al. (2017). The bottom panel summarizes the findings of the present study and the role of the microtubule-associated protein TPX2 in spindle positioning. In the scenario where TPX2 function is reduced, the spindle can be transported to the bottom of the gametophore initial cell, compromising the cell plate position and daughter cell ratios in asymmetric cell division. This spindle transport is an actin-dependent process, as it can be completely inhibited by depolymerizing actin filaments with latranculin A (Lat A). Furthermore, stabilizing microtubules with taxol could partially counteract spindle transport, suggesting that under normal conditions, microtubules are able to fix spindle position against actin force.

775 **Supplemental Video 1. Localization of TPX2 proteins during mitosis**
776 Live-cell imaging was performed in *P. patens* apical caulonemal cells expressing
777 mCherry-tubulin (magenta) and one of the following tagged proteins (green): TPX2-1-
778 Citrine, TPX2-2-mNeonGreen, mNeonGreen-TPX2-4, or TPX2-5-mNeonGreen. Images
779 were acquired every 30 s in a single focal plane. Bar, 10 µm.
780
781 **Supplemental Video 2. 3D projection of gametophore initial cells after the first**
782 **division.**
783 Gametophore initials were stained with 10 µM FM4-64. Cell segmentation was done with
784 PlantSeg. The apical cell of the gametophore initial is shown in pink, while the basal cell
785 in green. Bar, 20 µm.
786
Supplemental Video 3. Spindle motility underlies the erroneous phragmoplast
positioning in *TPX2-5 HM1* gametophore initial cells
Live-cell imaging was performed in *P. patens* gametophore initial cells expressing
mCherry-tubulin and histone H2B-mCherry (chromosomes and MTs are labeled with the
same color). Images were acquired as a z-stack (20 µm in 2.5 µm steps) every 5 min, and
the best focal plane was presented. Bar, 10 µm.
793
Supplemental Video 4. Spindle positioning defects in later gametophore
development in the *TPX2-5 HM1* mutant.
Representative videos of spindle misorientation and/or motility in *TPX2-5 HM1*
gametophore initial cells during second (left cell) or later (right cell) cell divisions. The
spindle axis is indicated by a cyan line. Images were acquired as a z-stack (20 µm, 2.5
µm step) every 5 min, and the best focal plane was presented. Bar, 10 µm.
800
Supplemental Video 5. Spindle-collapse phenotype in the gametophore initial of
***TPX2-5 HM1* mutants**
Representative video of spindle collapse followed by chromosome missegregation,
observed in approximately 10% of *TPX2-5 HM1* gametophore initial cells. Images were
acquired as a z-stack (20 µm, 2.5 µm step) every 5 min, and the best focal plane was
presented. Bar, 10 µm.
807
808 **Supplemental Video 6. Mitotic defects in the *TPX2-5* RNAi lines**
809 Representative images of mitotic defects in the *TPX2-5* RNAi lines. Live-cell imaging
810 was performed in *P. patens* protonemal apical cells expressing GFP-tubulin (green) and
811 histone H2B-mCherry (magenta). Images were acquired at a single focal plane every 2
812 min. Bar, 10 µm.
813