1   **Article type:** Research Article

5   **Title:** Endosymbiotic *Rickettsiella* causes cytoplasmic incompatibility in a spider host

6   Laura C. Rosenwald[1], Michael I. Sitvarin[2], Jennifer A. White[1]*

7   [1] *S-225 Agricultural Science Center N, Department of Entomology, University of Kentucky,*

8   *Lexington, Kentucky.*

9   [2]*100 Piedmont Ave. NE, Department of Biology, Georgia State University, Atlanta, Georgia*

10   *To whom correspondence should be addressed.

11   email: jenawhite@uky.edu

Abstract

Many arthropod hosts are infected with bacterial endosymbionts that manipulate host

reproduction, but few bacterial taxa have been shown to cause such manipulations. Here we

show that a bacterial strain in the genus *Rickettsiella* causes cytoplasmic incompatibility (CI)

between infected and uninfected hosts. We first surveyed the bacterial community of the

agricultural spider *Mermessus fradeorum* (Linyphiidae) using high throughput sequencing and

found that individual spiders can be infected with up to five different strains of maternally-

inherited symbiont from the genera *Wolbachia, Rickettsia,* and *Rickettsiella*. The *Rickettsiella*

strain was pervasive, found in all 23 tested spider matrilines. We used antibiotic curing to

generate uninfected matrilines that we reciprocally crossed with individuals infected only with

Rickettsiella. We found that only 13% of eggs hatched when uninfected females were mated

with *Rickettsiella*-infected males; in contrast, at least 83% of eggs hatched in the other cross

types. This is the first documentation of *Rickettsiella*, or any Gammaproteobacteria, causing CI.

We speculate that induction of CI may be much more widespread among maternally-inherited

bacteria than previously appreciated. Further, our results reinforce the importance of thoroughly

characterizing and assessing the inherited microbiome before attributing observed host

phenotypes to well-characterized symbionts such as *Wolbachia*.

Keywords: Araneae**,** endosymbiont coinfection, cytoplasmic incompatibility, *Rickettsiella*,

Wolbachia, Rickettsia

Introduction

Maternally transmitted endosymbiotic bacteria are common in arthropods, and are

estimated to infect nearly half of all arthropod species [1–4]. These inherited endosymbionts

frequently manipulate host reproduction to promote symbiont transmission and spread in the host

population [5]. The most prevalent and well-studied symbiont-driven manipulation is

cytoplasmic incompatibility [5–8]. Cytoplasmic incompatibility (CI) is a conditional sterility

phenotype in which crosses between infected males and uninfected females result in offspring

mortality, sabotaging the production of uninfected progeny. This phenotype functionally

increases the proportion of infected females in the host population over time [5]. Cytoplasmic

incompatibility influences host population dynamics and gene flow, acting both as a gene drive

mechanism that spreads traits through a population [9] and as a reproductive barrier that can

enforce reproductive isolation and contribute toward speciation [10,11]. In application, CI has

been used to locally repress pest populations as a form of sterile insect technique [12], and also

as a gene drive mechanism to reduce disease transmission by mosquitoes [13,14].

To date, few bacterial taxa have been shown to cause CI. The most widespread and well-

investigated are members of the genus *Wolbachia* (Phylum Proteobacteria, Class

Alphaproteobacteria, Order Rickettsiales), which are estimated to infect at least a third of

arthropod species [1,8,15,16]. Recently, another unnamed strain in the Rickettsiales has also

been suggested to induce CI [17]. Finally, members of the more distantly related *Cardinium*

genus (Phylum Bacteroidetes) also induce CI, although using a mechanism that appears to be

independently evolved from *Wolbachia* [18]. No other bacterial taxa have been documented to

cause CI to date.

In the present study, we provide evidence that CI can be caused by a strain of

Rickettsiella (Phylum Proteobacteria, Class Gammaproteobacteria, Order Legionellales)*, a*

bacterium that is not closely related to either *Wolbachia* or *Cardinium* [19]. We began by

characterizing the microbiome of *Mermessus fradeorum* (Araneae: Linyphiidae), an agricultural

spider that had been previously shown to be infected by multiple *Wolbachia* and *Rickettsia*

(Phylum Proteobacteria, Class Alphaproteobacteria, Order Rickettsiales) symbionts, and to

exhibit CI [20]. Upon discovery of an unexpected *Rickettsiella* symbiont that was pervasive

throughout our sampled population, we proceeded to experimentally document that *Rickettsiella*

causes CI in *M. fradeorum*. This is the first demonstration of CI by a Gammaproteobacteria,

which suggests that the taxonomic distribution of bacteria that cause CI may be wider than

previously appreciated, and could have important implications for the population genetics,

biodiversity, and evolution of arthropod hosts.

Methods

Study System

We initially collected *Mermessus fradeorum* from alfalfa at the University of Kentucky's

Spindletop Research Farm (38.127, -84.508). We placed individual spiders into deli cups (6cm

diam.) with hydrated plaster in the bottom for humidity control, and maintained them in the

laboratory at 22°C 24hr dark conditions. We fed the spiders fruit flies (*Drosophila*

melanogaster) at 2-3 day intervals; spiders collected as juveniles were fed collembola (*Sinella*

curviseta) until maturity. We initiated matrilines from individual field-collected gravid females,

or by mating field-collected females to males within the laboratory.

Microbiome Survey

To evaluate endosymbiont diversity in the population of *M. fradeorum*, we characterized

the microbiome of 23 matrilines that were initiated between August and October 2016. For each

matriline, we evaluated the field-collected female specimen when possible; if the field-collected

female died of natural causes (n = 8), we instead randomly selected an adult female F1 offspring.

We withheld food from each spider for at least five days before preserving the specimen in 95%

ethanol. We surface-sterilized each specimen with a series of washes with 0.5% bleach and PCR

water. We then extracted total DNA from each individual specimen using DNEasy Blood and

Tissue kits (Qiagen, Germantown, MD) according the manufacturer's instructions. Using PCR

with dual-indexed 515F/806R primers [21], we amplified the V4 region of bacterial 16S from

each specimen and visualized an aliquot of the resulting product on a 1% agarose gel stained

with GelRed (Biotium). After verifying the presence of a strong band of the expected product

size, we included a 1µl aliquot of the corresponding product in a multiplexed library for

sequencing. This library also included samples from aphids and parasitoid wasps that served as

controls (see below). After purification using GenCatch PCR Cleanup kit (Epoch Life Sciences,

Missouri City, TX) we sequenced the library on an Illumina Miseq 2500 instrument at the

University of Kentucky's core sequencing facility.

Sequences were demultiplexed, trimmed and quality filtered within BaseSpace (Illumina)

then imported into qiime2 (v2017.11) [22] using a manifest. We conducted additional quality

control using deblur [23], implemented in qiime2 using default parameters and a trim length of

251 bases. To determine the taxonomic placement of each operational taxonomic unit (OTU), we

used a naïve Bayes classifier trained on the V4 region of the Greengenes 13_8 99% OTUs

reference database [24].

The most prevalent OTU in the library, a strain of *Rickettsiella*, constituted 46% of total reads and was associated with every component specimen in the multiplexed sample, albeit at low prevalence for non-*M. fradeorum* samples. We inferred that index swapping may have occured between samples that shared either forward or reverse indices, resulting in reads that were misallocated to the wrong sample [25]. To validate that *Rickettsiella* was genuinely present in all *M. fradeorum* samples, we used *Rickettsiella*-specific primers to diagnostically test for the presence of *Rickettsiella* in the 23 matriline samples, as well as the 8 controls (4 *Aphis craccivora* (aphid) specimens and 4 *Habrobracon hebetor* (parasitoid wasp) specimens). Diagnostic primers and protocols followed Duron et al. [19]; samples that tested negative for *Rickettsiella* were rescreened at least two more times, and additionally tested for extraction quality via diagnostic PCR of a segment of the arthropod COI gene [4].

To assess vertical transmission of *Rickettsiella*, we included specimens from several generations in the microbiome dataset. In total, 148 mother-offspring pairs from the 23 matrilines (representing 83 *Rickettsiella*-infected mothers) were initially included in the dataset. As a quality control measure for the final dataset, we retained only pairs in which both mother and offspring yielded more than 5000 reads (117 pairs, representing 73 different mothers). To be scored positive, the *Rickettsiella* read number for a given microbiome needed to exceed the upper 95% confidence boundary of *Rickettsiella* reads found in the eight control samples (145 reads).

Rickettsiella *Cytoplasmic Incompatibility Assay*

To assess the ability of *Rickettsiella* to cause CI, we first created a population of uninfected spiders using antibiotics. Spiderlings originating from seven infected matrilines were

treated with a combination of tetracycline (0.1%) and ampicillin (0.1%) by a fine mist spray

daily until sub-adulthood [20,26]. Upon maturity, treated spiders were mated to one another, and

a subset of those offspring were diagnostically tested for endosymbionts (*Wolbachia, Rickettsia,*

Rickettsiella) [4,20]. Sibships that were negative for all endosymbionts were used to initiate the

uninfected (cured) population for experimental assays. Uninfected spiders used in experiments

were 1, 2, or 3 generations removed from antibiotic treatment.

To determine if *Rickettsiella* was causing CI in *M. fradeorum*, we reciprocally mated

uninfected spiders with spiders that were infected with only *Rickettsiella* (8 matrilines). We

randomly assigned virgin spiders to mate with either a *Rickettsiella*-infected or uninfected

partner, ensuring that siblings and cousins were not paired with one another. Males and females

were paired for 8 hours, and were observed at approximately 15 minute intervals for the first two

hours and periodically thereafter. If the pair was not observed to mate, they were excluded from

the trial. Females were then allowed up to six weeks to lay three egg masses; females that failed

to lay three egg masses within this time period (n=2) were excluded from the sample set. Egg

masses were checked daily for spiderlings. Once spiderlings were observed, we allowed 24 hours

for emergence, then dissected the egg mass to quantify unhatched eggs. Egg masses hatched

within a mean $\pm$ SE of 14.1 $\pm$ 0.3 days. If no spiderlings were observed within 20 days, the entire

egg mass was scored as unhatched, and dissected to count the number of unhatched eggs.

Throughout the experiment, we found a small number of spiderlings (n=9) that hatched but died

prior to exiting the egg mass. These unemerged spiderlings were excluded from the data set, as

they accounted for only 2% of all eggs.

The infection status of all parents in the experiment was validated using diagnostic PCR

for *Wolbachia, Rickettsia,* and *Rickettsiella* [4,19,20]. Final classification of all matings was

based on these diagnostic results; in several instances spiders from "cured" matrilines were

found to be symbiont infected, presumably due to cryptic low titer infections that rebounded in

subsequent generations. In such cases, matings were reclassified or excluded as necessary,

reflecting the actual infection status of both partners. The final distribution of mating cross types

across four trial dates included 18 pairings in which both females and males were infected with

only *Rickettsiella* (+/+), 14 in which the female was *Rickettsiella*-infected and the male was

uninfected (+/-), 12 in which the female was uninfected and the male was *Rickettsiella*-infected

(-/+), and 8 in which both female and male were uninfected (-/-).

We compared total eggs laid across treatments using ANOVA (JMP) after verifying that

model assumptions of normality and homoscedacity were not violated. We compared egg hatch

across treatments using logistic regression (Arc v. 1.06) with Williams correction [27] to account

for moderate overdispersion. To verify that generations removed from curing (1, 2, or 3

generations) did not influence experimental outcome, we considered a separate model that

incorporated this factor, but found no significant effect (Wald = -0.93, p = 0.35). Generation

from curing was therefore not included in the final model.

Results

Microbiome Survey

The most notable feature of the *M. fradeorum* microbiome was the pervasiveness of a

single strain of *Rickettsiella*, present in all 23 matrilines (figure 1). Mean ± SE reads per sample

was 22,054 ± 2,739, of which 16,172 ± 2,109 (80 ± 6%) were *Rickettsiella*. In most specimens

(20/23), *Rickettsiella* constituted the majority of reads, often representing >95% of reads (figure

1, Electronic supplementary material tables S1-S2). Proportional representation of *Rickettsiella*

was lower in samples that were also infected with *Rickettsia* (7 samples) and/or up to 3 strains of

Wolbachia (8 samples), but still constituted half the reads returned for this subset of samples (50

± 9%). In contrast, aphid and parasitoid wasp control samples that were included in the same

sequencing run had few reads of this *Rickettsiella* strain (115 ± 15 reads/sample, constituting 0.3

± 0.05% of reads/sample). Diagnostic validation confirmed the presence of *Rickettsiella* from all

23 matriline samples, and the absence of *Rickettsiella* from the aphids and parasitoid wasps. We

observed 100% vertical transmission efficiency in the unmanipulated mother-offspring pairs in

the dataset (Electronic supplementary material table S3). Across 117 pairs, all offspring of

Rickettsiella-infected mothers were themselves infected with *Rickettsiella*.

Reproductive Manipulation Assay: Cytoplasmic Incompatibility

When we reciprocally mated spiders bearing only *Rickettsiella* with those from matrilines

that had been cured of all facultative symbionts, we found a large difference in the number of

hatchlings produced. Total egg number was consistent and not significantly different across

treatments (figure 2, Electronic supplementary material table S4; ANOVA: $F_{3,48} = 0.56$, p =

0.644), but the odds that these eggs hatched was significantly lower for uninfected females mated

to *Rickettsiella*-infected males, relative to all three other treatments, ($\chi^2 = 111$, d.f. = 3, p <

0.001). Only 12.5 ± 3.6% of eggs hatched in the incompatible cross between uninfected females

and *Rickettsiella*-infected males, compared to 98.4 ± 1.0% of eggs when both parents were

infected, 89.5 ± 4.5% from *Ricketsiella*-infected females mated with uninfected males, and 83.1

± 12.2% when both parents were uninfected.

Discussion

We found that the spider *Mermessus fradeorum* is infected by up to five strains of

bacterial symbiont, including a strain of *Rickettsiella* that was present in 100% of our field-

collected matrilines. We then generated uninfected matrilines, and used crossing experiments to

document that this *Rickettsiella* strain induced cytoplasmic incompatibility in the spider host.

Previous examples of CI-causing bacteria have been restricted to *Wolbachia* and closely-allied

bacteria in the Alphaproteobacteria, and *Cardinium* in the Bacteroidetes [6,7,17,28]. Ours is the

first demonstration of a Gammaproteobacteria, *Rickettsiella*, causing CI.

Previous studies with this spider species diagnostically documented three endosymbiotic

bacterial strains (two *Wolbachia* and one *Rickettsia*) associated with bacterially-induced

reproductive manipulations, but could not attribute causality to individual bacterial strains [20].

The present microbiome study confirmed the presence of these same three symbiont strains at

similar field frequencies as the previous work, and additionally revealed a third strain of

Wolbachia along with the ubiquitous *Rickettsiella*. We were able to screen archived specimens

from the previous study [20] and found that the CI-inducing matriline from that study was in fact

co-infected with *Rickettsiella* and one strain of *Wolbachia* (L. Rosenwald, unpublished results).

It is therefore likely that the CI observed in this previous study was at least partially attributable

to *Rickettsiella*, although both co-infecting symbionts may have contributed to the CI phenotype

[29]. Focus on taxa that are well-established as reproductive manipulators, such as *Wolbachia*,

may lead researchers to overlook co-infecting bacteria that cause or contribute to manipulated

host reproductive phenotypes, as co-infections are common in arthropods [1,4].

The mechanism by which *Rickettsiella* induces CI is unknown. In *Wolbachia,* CI is

induced by phage-encoded operons in which one gene modifies the sperm of infected males and

the other rescues the modification in the cytoplasm of the egg of infected females [30,31]. In

Cardinium, the mechanism to induce CI appears to be independently evolved from *Wolbachia,*

and is not phage associated [18].   No previously described *Rickettsiella* strains have been

documented to cause CI, and genetic resources for making mechanistic inferences are lacking.

However, strains of *Rickettsiella* infect a wide range of arthropods [19,32], sometimes as a

pathogen [32], sometimes a mutualist [33], and sometimes as a potential reproductive

manipulator [34]. Speculatively, it is possible that horizontal gene transfer among co-infecting

symbionts may be responsible for *Rickettsiella's* ability to induce CI [35,36], particularly given

the high frequency of symbiotic co-infection in spiders in general [1,3,4,37] and *M. fradeorum* in

particular.

Our data also reinforce the perception that spiders can harbor exceptionally diverse

communities of co-infecting symbionts, rivaling the most symbiont-rich inherited communities

among insects [4,38,39]. The discovery of an unexpected reproductive manipulator within such a

community highlights the hazard of focusing only on well-characterized manipulative taxa (e.g.

Wolbachia) when reproductive anomalies are observed in a host population. Our data emphasize

the need for careful characterization of the entire inherited microbiome before attributing

reproductive phenotypes to particular bacterial strains.

Reproductive manipulation by endosymbionts, through mechanisms such as CI, can

influence arthropod behavioral ecology [40], population dynamics [41], and evolutionary

trajectory [42]. As the known roster of symbionts capable of undertaking these manipulations

increases, the potential for understanding and potentially exploiting their effects on the biology

of their hosts increases as well. CI is currently being used as a gene drive mechanism to decrease

disease transmission in mosquitoes [12]; newly discovered CI-inducing symbionts, such as

described in the present work, have similar potential in a wider range of arthropod hosts. In

particular, *Wolbachia* is rare or absent in ticks (Class Arachnida, Order Acari), but strains of

Rickettsiella are routine [19,34,43,44]. It would be worthwhile to evaluate the function of

Rickettsiella and other members of tick microbiomes, similarly to what we have done with

Rickettsiella in the spider *M. fradeorum*, as the potential opportunities for symbiont-based pest

control and vector management are substantial.

Data accessibility

The datasets supporting this article have been uploaded as part of the electronic supplementary

material (Electronic supplementary material, tables S1-S4). High throughput sequencing data is

available in the NCBI Sequence Read Archive under accession number PRJNA604923.

Acknowledgements

We thank M. Curry and A. Styer for their previous work on this system, K. Welch for his *M.*

fradeorum collections, and K. Athey for confirmation and initiation of *M. fradeorum* matrilines.

We additionally thank S. Ferrell, A. Lindsey, K. O'Hearn and G. Moses for laboratory

assistance.

Funding

This work was supported by a grant from the Kentucky Science and Engineering Foundation as

per Grant/Award Agreement 148-502-16-377 with the Kentucky Science and Technology

Corporation, and the National Institute of Food and Agriculture, U.S. Department of Agriculture

(Hatch No. 1004208).

References

1.   Duron O, Bouchon D, Boutin S, Bellamy L, Zhou L, Engelstädter J, Hurst GD. 2008 The
diversity of reproductive parasites among arthropods: *Wolbachia* do not walk alone. *BMC*
Biol. **6**, 1–12. (doi:10.1186/1741-7007-6-27)
2.   Weinert LA, Araujo-Jnr E V., Ahmed MZ, Welch JJ. 2015 The incidence of bacterial
endosymbionts in terrestrial arthropods. *Proc. R. Soc. B Biol. Sci.* **282**, 3–8.
(doi:10.1098/rspb.2015.0249)
3.   Zhang L, Yun Y. 2018 Insights into the bacterial symbiont diversity in spiders. *Ecol. Evol.*
8, 4899–4906. (doi:10.1002/ece3.4051)
4.   White JA, Styer A, Rosenwald LC, Curry MM, Welch KD, Athey KJ, Chapman EG. 2020
Endosymbiotic bacteria are prevalent and diverse in agricultural spiders. *Microb. Ecol.* **79**,
472–481. (doi:10.1007/s00248-019-01411-w)
5.   Engelstädter J, Hurst GDD. 2009 The ecology and evolution of microbes that manipulate
host reproduction. *Annu. Rev. Ecol. Evol. Syst.* **40**, 127–149.
(doi:10.1146/annurev.ecolsys.110308.120206)
6.   O'Neill SL, Giordano R, Colbert AM, Karr TL, Robertson HM. 1992 16S rRNA
phylogenetic analysis of the bacterial endosymbionts associated with cytoplasmic
incompatibility in insects. *Proc. Natl. Acad. Sci.* **89**, 2699–2702.
(doi:10.1073/pnas.89.7.2699)
7.   Hunter MS, Perlman SJ, Kelly SE. 2003 A bacterial symbiont in the Bacteroidetes induces
cytoplasmic incompatibility in the parasitoid wasp *Encarsia pergandiella*. *Proc. R. Soc. B*
Biol. Sci. **270**, 2185–2190. (doi:10.1098/rspb.2003.2475)
8.   Werren JH, Baldo L, Clark ME. 2008 *Wolbachia*: Master manipulators of invertebrate
biology. *Nat. Rev. Microbiol.* **6**, 741–751. (doi:10.1038/nrmicro1969)
9.   Sinkins SP, Gould F. 2006 Gene drive systems for insect disease vectors. *Nat. Rev. Genet.*
7, 427–435. (doi:10.1038/nrg1870)
10.   Bordenstein SR, O'Hara PF, Werren JH. 2001 *Wolbachia*-induced incompatibility
precedes other hybrid incompatibilities in *Nasonia*. *Nature* **409**, 707–710.
(doi:10.1038/35055543)
11.   Gebiola M *et al.* 2016 'Darwin's corollary' and cytoplasmic incompatibility induced by
Cardinium contribute to speciation in *Encarsia* wasps (Hymenoptera : Aphelinidae).
Evolution. **70**, 2447–2458.
12.   Mains JW, Brelsfoard CL, Rose RI, Dobson SL. 2016 Female adult *Aedes albopictus*
suppression by *Wolbachia*-infected male mosquitoes. *Sci. Rep.* **6**, 1–7.
(doi:10.1038/srep33846)
13.   Walker T *et al.* 2011 The wMel *Wolbachia* strain blocks dengue and invades caged *Aedes*
aegypti populations. *Nature* **476**, 450–455. (doi:10.1038/nature10355)
14.   Pan X, Zhou G, Wu J, Bian G, Lu P, Raikhel AS, Xi Z. 2012 *Wolbachia* induces reactive
oxygen species (ROS)-dependent activation of the Toll pathway to control dengue virus in
the mosquito *Aedes aegypti*. *Proc. Natl. Acad. Sci. U. S. A.* **109**, E23–E31.
(doi:10.1073/pnas.1116932108)
15.   Zug R, Hammerstein P. 2012 Still a host of hosts for *Wolbachia*: Analysis of recent data
suggests that 40% of terrestrial arthropod species are infected. *PLoS One* **7**, 7–9.
(doi:10.1371/journal.pone.0038544)
16.   Sazama EJ, Scot P, Wesner JS. 2019 Insect-symbiont interactions bacterial endosymbionts

are common among, but not necessarily within, insect species. *Environ. Entomol.* **48**, 127–
133. (doi:10.5063/F1)

17. Takano S *et al.* 2017 Unique clade of alphaproteobacterial endosymbionts induces
complete cytoplasmic incompatibility in the coconut beetle. *Proc. Natl. Acad. Sci.* **114**,
6110–6115. (doi:10.1073/pnas.1618094114)

18. Penz T, Schmitz-Esser S, Kelly SE, Cass BN, Müller A, Woyke T, Malfatti SA, Hunter
MS, Horn M. 2012 Comparative genomics suggests an independent origin of cytoplasmic
incompatibility in *Cardinium hertigii. PLoS Genet.* **8**, 10.
(doi:10.1371/journal.pgen.1003012)

19. Duron O, Cremaschi J, McCoy KD. 2016 The high diversity and global distribution of the
intracellular bacterium *Rickettsiella* in the polar seabird tick *Ixodes uriae. Microb. Ecol.*
71, 761–770. (doi:10.1007/s00248-015-0702-8)

20. Curry MM, Paliulis L V, Welch KD, Harwood JD, White JA. 2015 Multiple
endosymbiont infections and reproductive manipulations in a linyphiid spider population.
Heredity. **115**, 146–152. (doi:10.1038/hdy.2015.2)

21. Kozich JJ, Westcott SL, Baxter NT, Highlander SK, Schloss PD. 2013 Development of a
dual-index sequencing strategy and curation pipeline for analyzing amplicon sequence
data on the Miseq Illumina sequencing platform. *Appl. Environ. Microbiol.* **79**, 5112–
5120. (doi:10.1128/AEM.01043-13)

22. Caporaso JG *et al.* 2010 QIIME allows analysis of high- throughput community
sequencing data. *Nat. Methods* **7**, 335–336. (doi:10.1038/nmeth0510-335)

23. Bokulich NA, Subramanian S, Faith JJ, Gevers D, Gordon JI, Knight R, Mills DA,
Caporaso JG. 2013 Quality-filtering vastly improves diversity estimates from Illumina
amplicon sequencing. *Nat. Methods* **10**, 57–59. (doi:10.1038/nmeth.2276)

24. DeSantis TZ *et al.* 2006 Greengenes, a chimera-checked 16S rRNA gene database and
workbench compatible with ARB. *Appl. Environ. Microbiol.* **72**, 5069–5072.
(doi:10.1128/AEM.03006-05)

25. Larsson AJM, Stanley G, Sinha R, Weissman IL, Sandberg R. 2018 Computational
correction of index switching in multiplexed sequencing libraries. *Nat. Methods* **15**, 305–
307. (doi:10.1038/nmeth.4666)

26. Vanthournout B, Swaegers J. 2011 Spiders do not escape reproductive manipulations by
Wolbachia. BMC Evol. Biol. **11**. (doi:10.1186/1471-2148-11-15)

27. Williams D. 1982 Extra-binomial variation in logistic linear models. *J. R. Stat. Soc. Ser. C*
(Applied Stat.) **31**, 144–148.

28. Gotoh T, Noda H, Ito S. 2007 *Cardinium* symbionts cause cytoplasmic incompatibility in
spider mites. *Heredity.* **98**, 13–20. (doi:10.1038/sj.hdy.6800881)

29. Zhu LY, Zhang KJ, Zhang YK, Ge C, Gotoh T, Hong XY. 2012 *Wolbachia* strengthens
Cardinium-induced cytoplasmic incompatibility in the spider mite *Tetranychus piercei*
McGregor. *Curr. Microbiol.* **65**, 516–523. (doi:10.1007/s00284-012-0190-8)

30. Le Page DP *et al.* 2017 Prophage WO genes recapitulate and enhance *Wolbachia*-induced
cytoplasmic incompatibility. *Nature* **543**, 243–247. (doi:10.1038/nature21391)

31. Beckmann JF, Ronau JA, Hochstrasser M. 2017 A *Wolbachia* deubiquitylating enzyme
induces cytoplasmic incompatibility. *Nat. Microbiol.* **2**, 1–7.
(doi:10.1038/nmicrobiol.2017.7)

32. Cordaux R, Paces-Fessy M, Raimond M, Michel-Salzat A, Zimmer M, Bouchon D. 2007
Molecular characterization and evolution of arthropod-pathogenic *Rickettsiella* bacteria.

Appl. Environ. Microbiol. **73**, 5045–5047. (doi:10.1128/AEM.00378-07)

33.   Tsuchida T, Koga R, Horikawa M, Tsunoda T, Maoka T, Matsumoto S, Simon J-C,
Fukatsu T. 2010 Symbiotic bacterium modifies aphid body color. *Science.* **330**, 1102–
1105. (doi:10.1126/science.1195463.)

34.   Kurtti TJ, Palmer AT, Oliver JH. 2002 *Rickettsiella*-like bacteria in *Ixodes woodi* (Acari:
Ixodidae). *J. Med. Entomol.* **39**, 534–540. (doi:10.1603/0022-2585-39.3.534)

35.   Ochman H, Lawerence JG, Groisman EA. 2000 Lateral gene transfer and the nature of
bacterial innovation. *Nature* **405**, 299–304.

36.   Ishmael N *et al.* 2009 Extensive genomic diversity of closely related *Wolbachia* strains.
Microbiology **155**, 2211–2222. (doi:10.1099/mic.0.027581-0)

37.   Goodacre SL, Martin OY, Thomas CFG, Hewitt GM. 2006 *Wolbachia* and other
endosymbiont infections in spiders. *Mol. Ecol.* **15**, 517–527. (doi:10.1111/j.1365-
294X.2005.02802.x)

38.   Ferrari J, West JA, Via S, Godfray HCJ. 2011 Population genetic structure and secondary
symbionts in host-associated populations of the pea aphid complex. *Evolution* **66**, 375–
390. (doi:10.5061/dryad.8qb00)

39.   Russell JA *et al.* 2013 Uncovering symbiont-driven genetic diversity across North
American pea aphids. *Mol. Ecol.* **22**, 2045–2059. (doi:10.1111/mec.12211)

40.   Angelella G, Nalam V, Nachappa P, White J, Kaplan I. 2018 Endosymbionts differentially
alter exploratory probing behavior of a nonpersistent plant virus vector. *Microb. Ecol.* **76**,
453–458. (doi:10.1007/s00248-017-1133-5)

41.   Jaenike J. 2009 Coupled population dynamics of endosymbionts within and between
hosts. *Oikos* **118**, 353–362. (doi:10.1111/j.1600-0706.2008.17110.x)

42.   Cordaux R, Bouchon D, Grève P. 2011 The impact of endosymbionts on the evolution of
host sex-determination mechanisms. *Trends Genet.* **27**, 332–341.
(doi:10.1016/j.tig.2011.05.002)

43.   Duron O *et al.* 2017 Evolutionary changes in symbiont community structure in ticks. *Mol.*
Ecol. **26**, 2905–2921. (doi:10.1111/mec.14094)

44.   Tijsse-Klasen E, Braks M, Scholte EJ, Sprong H. 2011 Parasites of vectors - *Ixodiphagus*
hookeri and its *Wolbachia* symbionts in ticks in the Netherlands. *Parasites and Vectors* **4**,
1–7. (doi:10.1186/1756-3305-4-228)

Figures

Figure 1. Proportional composition of 16S bacterial reads from 23 adult female *Mermessus*

fradeorum spiders. All spiders were either field collected or adult offspring of different field

collected mothers. Each shade represents a different bacterial strain; different strains from the

same genus are designated by different numbers in the legend. Bacteria known to be maternally

transmitted are brightly colored (yellow, blue, red), other strains are shades of gray and brown.

Strain sequences, read numbers, and composition of the "Other" category available in Electronic

supplemental material Tables S1 and S2.

Figure 2. Mean ± SE total eggs laid (A) and proportion eggs that hatched (B) in reciprocal

crosses between *Mermessus fradeorum* females and males that varied in *Rickettsiella* infection

status. Column color indicates expected offspring infection status: Yellow columns reflect

offspring of *Rickettsiella*+ mothers, white columns reflect offspring of *Rickettsiella*– mothers.