The neuroplasticity of division of labor: worker polymorphism, compound eye structure

and brain organization in the leafcutter ant *Atta cephalotes*

Sara Arganda[1, 2, 3*]

Andrew P. Hoadley[1*]

Evan S. Razdan[1]

Isabella B. Muratore[1]

James F. A. Traniello[1,4]

[1]Department of Biology, Boston University, Boston, USA

[2]Research Centre on Animal Cognition (CRCA), Centre for Integrative Biology (CBI), Toulouse

University, CNRS, UPS, Toulouse 31062, France

[3]Departamento de Biología y Geología, Física y Química Inorgánica, Área de Biodiversidad y

Conservación, Universidad Rey Juan Carlos, Madrid, Spain

[4]Graduate Program for Neuroscience, Boston University, Boston, MA 02215

* These authors contributed equally to the study

Dedicated to the memory of Eldridge Adams, brilliant behavioral ecologist, colleague, and

friend.

Corresponding Author:

S. Arganda

ORCID: 0000-0002-9875-2379

email: sara.arganda@urjc.es

phone: +34 91 488 85 19

Abstract

Our understanding of how the design of peripheral sensory structures is coupled with neural

processing capacity to adaptively support division of labor is limited. Workers of the remarkably

polymorphic fungus-growing ant *Atta cephalotes* are behaviorally specialized by size: the

smallest workers (minims) tend fungi in dark subterranean chambers while larger workers

perform tasks mainly outside the nest. These strong differences in worksite light conditions are

predicted to influence sensory and processing requirements for vision. We found that eye

structure and visual neuropils have been be selected to maximize task performance according to

light availability. Minim eyes had few ommatidia, large interommatidial angles and eye

parameter values, suggesting selection for visual sensitivity over acuity. Large workers had

larger eyes with disproportionally more and larger ommatidia, and smaller interommatidial

angles and eye parameter values, reflecting peripheral sensory adaptation to ambient rainforest

light. Additionally, optic lobe and mushroom body collar volumes were disproportionately small

in minims, and within the optic lobe, lamina and lobula relative volumes increased with worker

size whereas the medulla decreased. Visual system phenotypes thus correspond to task

specializations in dark or light environments and reflect a functional neuroplasticity

underpinning division of labor in this socially complex agricultural ant.

Keywords (5): task performance, optic lobe, compound eye, ommatidia, social brain

Introduction

Morphology, behavior, and nervous system structure appear to be integrated (Corral-

López et al. 2017; Gordon et al. 2017; Iglesias et al. 2018). For example, body size correlates

with optical sensitivity and resolution in insect vision (Spaethe and Chittka 2003; Rutowski et al.

2009; Palavalli-Nettimi and Narendra 2018; Taylor et al. 2019). Ants are an ideal model system

to examine relationships among behavior, body size, and neuroanatomy because workers have

evolved as task specialists in several clades (Hölldobler and Wilson 1990). Scaling patterns of

brain size and brain compartment substructure among polymorphic workers, moreover, appear to

correspond to foraging ecology and the sensory and cognitive demands of task performance

(Gronenberg 2008; Muscedere and Traniello 2012; Gordon et al. 2017). Although olfactory

inputs are principal information sources in ants (Hölldobler and Wilson 1990; Czaczkes et al.

2015), vision can be significant in foraging ecology and navigation (Knaden and Graham 2016;

Narendra et al. 2017). To home, foragers may use a celestial compass (Wehner 2003; Muller and

Wehner 2006), optic flow (Ronacher and Wehner 1995), visual cues and landmark panoramas

(Graham and Cheng 2009; Müller and Wehner 2010; Schwarz et al. 2011; Huber and Knaden

2015; Freas et al. 2018), polarized light (Zeil et al. 2014), and canopy patterns (Hölldobler 1980;

Beugnon et al. 2005; Rodrigues and Oliveira 2014). Addtionally, visual navigation has been

associated with peripheral receptor structure, and primary and higher-order processing brain

centers (Gronenberg and Hölldobler 1999; Wehner 2003; Ehmer and Gronenberg 2004; Muller

and Wehner 2006; Knaden and Graham 2016), and worker behavioral development may be

associated with light-exposure and cued neuroanatomical reorganization in the visual system

(Stieb et al. 2010, 2012; Yilmaz et al. 2016).

Ant ommatidia are photoreceptive units that may change in number and structure

according to visual needs (Moser et al. 2004; Narendra et al. 2016a). Ommatidia structure affects

visual capacity: larger ommatidia enhance light sensitivity, ommatidia number determine image

resolution, and lower interommatidial angle improves acuity (Land 1997). Reproductive and

worker division of labor in social insects may have selected for differences in compound eye

structure (Schwarz et al. 2011; Streinzer et al. 2013). In some ant species, ommatidia number and

size scale with worker body size (Menzel and Wehner 1970; Bernstein and Finn 1971; Klotz et

al. 1992; Baker and Ma 2006; Schwarz et al. 2011), vary in males and females (Narendra et al.

2016b), and scale differently among polymorphic workers within individual compound eyes and

between colonies (Perl and Niven 2016a, b). In bull ants (Greiner et al. 2007; Narendra et al.

2011) and bees (Jander and Jander 2002; Greiner et al. 2004) photoreceptor diameter and eye

area increase in nocturnal species in comparison to diurnal species, increasing visual sensitivity.

Ommatidia facet diameter is generally smaller in diurnal than nocturnal ants (Narendra et al.

2017), but eye size patterns vary (e.g.(Menzi 1987)).

Visual input from the compound eyes travels to the optic lobes (OL) for primary

processing (Gronenberg and Hölldobler 1999). OL investment reflects visual ecology in social

insects: in subterranean species, workers are eyeless and OLs are absent whereas diurnal solitary

foragers have enormous eyes and their OLs occupy 33% of their brains (Gronenberg and

Hölldobler 1999). In paper wasps, queens remain inside the nest and have smaller OLs than

foraging workers (O'Donnell et al. 2014), and in the weaver ant *Oecophylla smaragdina*, minor

workers nurse brood, rarely leave the nest, and have disproportionally smaller OLs than majors,

which forage and defend territory (Kamhi et al. 2017).

The OL is comprised of three regions: lamina (contrast detection), medulla (color vision

processing and small field motion), and lobula (color vision processing, wide field motion

detection, and shape and panorama construction) (Strausfeld 1989; Gronenberg 2008; Dyer et al.

2011). OL interneurons project to the collar of the mushroom body (MB) calyx for higher-order

processing (Gronenberg 2001; Farris 2016). In ants, males, queens and worker brains show

differential investment in the medulla, lobula and MB collar (Ehmer and Gronenberg 2004),

reflecting different visual ecologies. Peripheral sensory structure should correlate with higher-

order processing ability in task-specialized workers, but this linkage is not well understood.

To investigate visual phenotypes within the worker caste, we investigated variation in the

structure of the compound eyes, OL, and MB collar in morphologically and behaviorally

differentiated workers of the fungus-growing ant *Atta cephalotes*. Worker head widths (HW)

range from 0.6 to 4.5mm; this striking polymorphism is associated with the frequency (Wilson

1980) and efficiency (Wetterer 1991; van Breda and Stradling 1994) of leaf harvesting, fungal

comb maintenance, brood care, hygienic behaviors, and colony defense. The smallest workers

(minims, HW<1.2mm) primarily tend brood and the fungal comb in dark underground chambers

(Wilson 1980) whereas media workers (HW=1.2-3.0mm) harvest plant material, traveling along

foraging trails beneath rainforest canopy, and the largest workers (majors, HW>3.0mm), are

responsible for defense (Powell and Clark 2004; Hölldobler and Wilson 2010). Medias use

vision during orientation along trails (Vilela et al. 1987; Vick 2005). Size-variable workers thus

have different social roles and experience environments strongly differing in ambient light

intensity and visual complexity. It is unlikely that a single eye structure and sensory processing

ability has evolved in all workers. We hypothesized that *A. cephalotes* visual system

organization is associated with the visual ecology of size-related division of labor and has

resulted from selection for adaptive plasticity in ommatidia structure, OL organization and MB

collar investment. Specifically, we predicted that workers engaging in within-nest or outside-nest

activities (in darkness or light, respectively) would vary in compound eye structure and relative

investment in the OL and its constituent parts, and in the MB collar to support the requirements

of vision associated to task performance.

Methods

Laboratory cultures

Queenright *A. cephalotes* colonies were collected in Trinidad (July 2014) and maintained

in a Harris environmental chamber (25°C, 50% relative humidity, 12:12h photoperiod). Artificial

nests were constructed from multiple plastic boxes (11cm×18cm×13cm each) connected by

plastic tubing (ID=2.5cm). Boxes housing fungal combs had dental stone floors with embedded

pebbles to provide air circulation for the fungus. Colonies were fed locally collected leaves free

of chemicals and organic produce on alternate days, supplemented with rolled oats, apple, and

orange mesocarp.

Worker size variation and tissue sampling

We sampled polymorphic workers from three colonies (Ac09, Ac20, and Ac21). *A.*

cephalotes appears to exhibit triphasic allometry, with three worker size classes (subcastes):

minims (HW across the eyes <1.2mm), medias (HW 1.2-3.0mm) and major workers

(HW>3.0mm). Each worker was anesthetized on ice and brains were dissected in ice-cold

HEPES buffered saline. Compound eyes were removed and stored in 70% ethanol for

processing. Because dissection is delicate, we were not always able to preserve the brain and

eyes of the same individual.

Compound eye imaging and structural measurements

Ninety-two intact compound eyes were imaged to create 3D stacks (Fig. 1G) to measure

ommatidia number (ON), average ommatidial diameter (D), and interommatidial angle ($\Delta\phi$).

Eyes were stored in 70% ethanol, washed in 100% ethanol (3×10 min) before mounting. We

measured one eye per worker. Extraneous cuticle was removed to allow eyes to lie flat and were

then mounted in methyl salicylate between coverslips and imaged using a Fluoview 1 confocal

microscope (λ=488nm, step size=3.1μm) with a 20x air objective (NA=0.5, CA=2). Cuticle has

natural fluorescence. Eye data were recorded blind to subcaste by randomly assigning

identification numbers to eyes. To quantify ommatidia number, image stacks were flattened in

ImageJ (Abràmoff et al. 2004) and facets were counted using the Cell Counter plugin. Volume

renderings were viewed in Amira 6.0 to verify counts.

Mean ommatidial diameter was calculated from the average diameter of 5 or 10 randomly

selected ommatidia from each eye. Eye surface area was calculated from the mean ommatidial

diameter (surface area=ON×π×[0.5xD]^2), and ommatidial density (number of ommatidia per

surface area unit) was calculated by dividing the number of ommatidia by eye surface area

(Yilmaz et al. 2014). To quantify interommatidial angle ($\Delta\phi$), image stacks were re-sectioned in

the yz plane to obtain a virtual cross section of the eye. ImageJ was used to estimate local eye

radius R (Schwarz et al. 2011), which with the mean ommatidial diameter (D, in μm) for that

eye, interommatidial angle (in radians) could be estimated as $\Delta\phi = D/R$ (Schwarz et al. 2011).

Eye parameter (P), which indicates the extent of trade-offs between sensitivity and resolution,

was calculated as $\Delta\phi \times D$ (Snyder 1977; Rutowski et al. 2009); lower values of P indicate

enhanced acuity, while compromising sensitivity.

Immunohistochemistry and confocal microscopy

After dissection, brains were placed in 16% Zn-formaldehyde (Ott 2008), fixed overnight

at approximately 18ºC on a shaker, washed in HBS (6×10 min) and then fixed in Dent's Fixative

(80% methanol, 20% DMSO) for minimally 1h. Brains were next washed in 100% methanol and

either stored at -17°C or immediately processed. Brains were washed in 0.1M Tris buffer

(pH=7.4) and blocked in PBSTN (5% neutral goat serum, 0.005% sodium azide in 0.2% PBST)

at 18ºC for 1 hour before incubation for 3 days at room temperature in primary antibody (1:30

SYNORF 1 in PBSTN; monoclonal antibody 3C11obtained from DSHB, University of Iowa, IA,

USA). They were washed (6×10 min) in 0.2% PBST and incubated in secondary antibody (1:100

AlexaFluor 488 goat anti-mouse in PBSTN) for 4 days at room temperature. Brains were then

washed a final time (6×10 min in 0.2% PBST) and dehydrated in an ethanol series (10min/step,

30/50/70/95/100/100% ethanol in 1x PBS), cleared with methyl salicylate, and mounted on

stainless steel slides for imaging.

Sixty-three brains were imaged using an Olympus Fluoview 1 confocal microscope

(λ=488nm, step size=3.1μm) with either a 10x air objective (NA=0.3, CA=1), or a 20x air

objective (NA=0.5, CA=2). Voxel depth was multiplied by a factor of 1.59 to correct for axial

shortening due to mounting in methyl salicylate (Bucher et al. 2000). Brain image stacks were

manually segmented using Amira 6.0 and Amira 2019.2 software to quantify neuropil volumes.

Given their bilateral symmetry, we segmented one hemisphere per brain, chosen randomly. Our

study goal required that only the lamina, medulla and lobula of the OL and MB calyces

(separating lip and collar) were segmented separately (Fig. 2A), the rest of the central brain

regions and the suboesophageal ganglion were segmented as a whole. Brain data collection was

blind to worker HW, although extreme size differences were obvious. Nevertheless, due the

randomized coding of brains, subcaste could not be determined with certainty by the annotator.

We calculated the volume of the brain hemisphere, the absolute and relative volume of OL

(relative to total brain volume), the absolute and relative volume of MB collar (relative to total

brain volume), and the relative volumes of OL subregions (relative to total OL volume).

Statistical evaluation

Statistical evaluations were performed in R (version 3.3.0, Team 2016) using the

'segmented' package to analyze eye and brain metric scaling (V. R. M. Muggeo 2008). To assess

allometries in eye structure and brain volumes in relation to worker size, least-square means

regression was used on log10-transformed values to estimate *a* and *b* in the scaling equation

$y=aM^b$, as *log10(y)=log(a)+b×log(M)*. To test the null hypothesis (H_0) of isometry, a separate

linear model was calculated and tested against different slope values depending on the metric.

The slope for H_0 were *b*=0.0 (linear vs. constant values), *b*=1.0 (linear vs. linear), *b*=2.0 (linear

vs. surface area) and *b*=3.0 (linear vs. volume) (Kaspari and Weiser 1999).

The Davies test was used to determine if there is a statistically significant change in slope

or a 'breakpoint' in a linear relationship (Davies 2002). We observed that the significance of

some changes in slope depended on a single data point; therefore, we accepted the change in

slope only if its significance was always below 0.05 when removing any point from the dataset.

The 'segmented' package was further used to estimate the location of the breakpoint. If the

Davies' test revealed two piecewise linear relationships in a scaling relationship, least-square

means regression was calculated and tested against isometry independently.

To further explore whether increased investment in primary visual neuropil might have

an impact in higher-order visual processing neuropil, we assessed allometry in the ratio of

volumes of the optic lobes and MB collar according to HW. We also calculated a least-square

means regression on log10-transformed values and tested against isometry (*b*=0.0).

Results

Eye Structure

The eyes of media and major workers had significantly more ommatidia than minims

(Fig. 1A) and showed a significant change in the scaling of ommatidia number and worker size

(Davies test, p<0.001) at a HW of 1.38mm (95% CI: 1.20 to 1.58mm). Piecewise linear models

calculated for both slopes were significant (p<0.001, Multiple R^2=0.989) with a slope shift from

2.03 (95% CI: 1.91 to 2.13) to 1.35 (95% CI: 1.25 to 1.45). Piecewise linear models were also

significantly different from isometry ($b=0$; $p<0.001$). Media and major workers also had larger

ommatidial diameter (Fig. 1B). The relationship between ommatidial diameter and worker size

showed no significant breakpoint (Davies test, $p>0.5$), and these variables were significantly

correlated ($F_{(1,90)}=1217$, $p<0.001$, $R^2=0.93$). The slope of the regression line was 0.25 (95% CI:

0.24 to 0.27), which was significantly different from isometry ($b=1.0$; $F_{(1,90)}=1217$, $p<0.001$).

Compound eye size (total eye surface area) was correlated with HW (Fig. 1C). Davies'

test showed a significant change in the scaling of total eye size and worker size ($p<0.001$) at a

HW of 1.44mm (95% CI: 1.20 to 1.72mm). Piecewise linear models calculated for both slopes

were found to be significant ($p<0.001$, Multiple $R^2=0.988$) with a slope shift from 2.52 (95% CI:

2.36 to 2.68) to 1.84 (95% CI: 1.70 to 1.97). Piecewise linear models were significantly different

from isometry ($b=2.0$; $p<0.001$), although the effect size was small.

The density of ommatidia decreased with HW (Fig. 1D). The relationship between the

density of ommatidia and worker size showed no significant breakpoint (Davies test, $p>0.5$ and

these variables showed a significant correlation ($F_{(1,90)}=1217$, $p<0.001$, $R^2=0.93$). The slope was

-0.50 (95% CI: -0.53 to -0.48), also significantly different from isometry ($b=0.0$; $F_{(1,90)}=1217$,

$p<0.001$).

Interommatidial angle decreased as worker size increased (Fig. 1E). Davies' test for a

change in slope showed a significant change in the scaling relationship between interommatidial

angle and worker size ($p<0.001$) at a HW of 1.25mm (95% CI: 1.03 to 1.51mm). Piecewise

linear models were calculated for both slopes and found to be significant ($p<0.001$, Multiple

$R^2=0.84$), with a slope shift from -0.98 (95% CI: -1.21 to -0.76) to -0.33 (95% CI: -0.43 to -

0.23). Piecewise linear models were also significantly different from isometry ($b=0$; $p<0.001$).

Eye parameter decreased with worker size in minims (Fig. 1F). Davies' test showed a

significant change in the scaling relationship between eye parameter and worker size ($p<0.001$)

at a HW of 1.26 mm (95% CI: 1.01 to 1.57 mm). Piecewise linear models were calculated for

both slopes and found to be significant ($p<0.001$, Multiple $R^2=0.588$) with a slope shift from -

0.73 (95% CI: -0.95 to -0.51) to -0.07 (95% CI: -0.20 to 0.05). The first segment of the piecewise

linear models was found significantly different from isometry ($b=0$; $p<0.001$), but the second

segment was not ($b=0$; $p=0.205$).

Brain Structure

Larger workers had significantly larger brains (Fig. 2B). The relationship between brain

volume and worker size showed no significant breakpoint (Davies test, p>0.05) and these

variables showed a significant positive correlation ($F_{(1,61)}$=13.91, p<0.001, R^2=0.18) with a slope

of 0.37 (95% CI: 0.17 to 0.56) significantly different from isometry (b=3.0; $F_{(1,61)}$=13.91,

p<0.001). We found greater OL investment in media and major workers (Fig. 2C). Relative OL

volume and worker size showed a significant positive correlation ($F_{(1,61)}$=271.2, p<0.001,

R^2=0.81) with no significant breakpoint (Davies test, p=1). The slope of the regression line was

0.76 (95% CI: 0.67 to 0.85) and significantly different from isometry (b=0.0; $F_{(1,61)}$=271.2,

p<0.001).

Within the OL, relative investment in lamina increased with worker size (Fig. 2C1).

Relative lamina volume and HW showed a significant positive correlation ($F_{(1,61)}$=37.77,

p<0.001, R^2=0.37) with no significant breakpoint (Davies test, p>0.05). The slope of the

regression line was 0.64 (95% CI: 0.43 to 0.85) and significantly different from isometry (b=0.0;

$F_{(1,61)}$=37.77, p<0.001). Relative investment in the medulla, in contrast, decreased with worker

size (Fig. 2C2). Relative medulla volume and HW showed a significant negative correlation

($F_{(1,61)}$=51.1, p<0.001, R^2 of 0.45) with no significant breakpoint (Davies test, p>0.05) and a

slope of -0.17 (95% CI: -0.21 to -0.12), significantly different from isometry (b=0.0; $F_{(1,61)}$=51.1,

p<0.001). Finally, as for the lamina, relative investment in the lobula increased with worker size

(Fig. 2C3). Relative lobula volume and HW showed a significant positive correlation

($F_{(1,61)}$=13.43, p<0.001, R^2=0.17) and no significant breakpoint (Davies test, p=0.362). The slope

of the regression line was 0.21 (95% CI: 0.10 to 0.32) and significantly different from isometry

(b=0.0; $F_{(1,61)}$=13.43, p<0.001).

Media and major workers also invested relatively more in the MB collar (Fig. 2D).

Relative collar volume and worker size showed a significant positive correlation ($F_{(1,61)}$=13.03,

p<0.001, R^2=0.16) and no significant breakpoint (Davies test, p=0.42). The slope of the

regression line was 0.19 (95% CI: 0.08 to 0.29), which was also significantly different from

isometry (b=0.0; $F_{(1,61)}$=13.03, p<0.001). Despite investing more in the MB collar, larger workers

had a lower collar:OL volume ratio (Fig. 2E). The relationship between this ratio and worker size

showed a significant negative correlation ($F_{(1,61)}$=67.17, p<0.001, R^2=0.52), no significant

breakpoint (Davies test, p=0.69), and a slope of -0.57 (95% CI: -0.71 to -0.43), significantly

different from isometry (b=0.0; $F_{(1,61)}$=67.17, p<0.001).

Discussion

Social insect compound eyes and visual information processing neuropils enable adaptive behavioral performance according to cognitive challenges of navigation and ambient light levels (Jander and Jander 2002; Mares et al. 2005; Kapustjanskij et al. 2007; Warrant 2008; Narendra et al. 2011, 2016a; Streinzer et al. 2013; Yilmaz et al. 2014; Bulova et al. 2016). Eye size and ommatidia number correlate with worker size in ants, including polymorphic species, and may be associated with task performance (Menzel and Wehner 1970; Klotz and Reid 1992; Schwarz et al. 2011). In polymorphic *A. cephalotes* workers, in which task performance is strongly correlated with body size-related division of labor, differences in worksite sensory ecology appear to select for visual system polyphenism.

Division of labor and eye structure in A. cephalotes

A. cephalotes workers perform tasks in the complete darkness of fungal comb chambers and in the filtered light epigaeic environment beneath rainforest canopy. Media workers forage day and night (Cherrett 1968), and use trail pheromones as well. Although olfaction appears to be the dominant sensory modality for foraging in many ants, visual information facilitates trail-following in *Atta laevigata* [69], and other ant species (Beugnon and Fourcassié 1988) alter their use of chemicals or vision depending on light conditions. In *A. cephalotes,* improved forager visual ability may enable flexibility in the use of orientation cues and social signals as ambient light levels change.

Minims tend fungi deep underground, medias harvest leaves from their habitat and labor inside the nest, and majors appear to exclusively perform defense and trail maintenance outside. We hypothesized that eye structure variation among subcastes would reflect adaptation to worksite light availability and visual demands for task performance. We expected the eyes of minims to structurally enable light sensitivity over resolution, whereas larger worker eyes were predicted to favor spatial resolution over sensitivity. It is unclear how minims make use of visual information and what level of spatial resolution and sensitivity is needed to work effectively on the fungal comb. Minims, however, also perform some tasks outside the nest, "hitchhiking" on transported leaves during day and night to defend against fly parasites (Linksvayer et al. 2016). We found that the number and size of ommatidia and eye surface area were significantly smaller in minims (Fig. 1A-C), suggesting less capacity to capture light and less reliance on vision to

perform their tasks. The larger ommatidia of media and major workers (Fig. 1B) indicate greater

sensitive to light. However, ommatidia size increased hypometrically with body size: although

the ommatidia of minims were the smallest, relative ommatidia size was greater in minims than

in medias and majors. This may enable minim worker eyes to collect more light than expected

from their size, suggesting adaptation to darkness and light (Greiner 2006; Yilmaz et al. 2014).

Alternatively, the small size of minim worker eyes and ommatidia may be due to a body size

constraint: eye size is as large as developmentally possible to ensure at least a marginal ability to

capture light, which may be needed for parasitic fly defenses. Minims also showed a higher

density of ommatidia than media and majors (Fig. 1D). Interommatidial angle decreased with

worker size, indicating higher visual acuity in larger workers (Fig. 1E). Eye parameter values

(Fig. 1F) were significantly higher for minims, but for larger workers, values were lower and not

correlated with size. This suggests minim worker eyes are adapted to enhance sensitivity rather

than acuity, whereas larger worker eyes structure have been selected for sensitivity and acuity.

Higher acuity is adaptive outside the nest, as it allows resolving more distal objects. The

significant breakpoints (HW 1.0-1.8mm) found in the linear regressions for ommatidia number,

eye surface area, interommatidial angle and eye parameter (Fig. 1A,C,E,F) suggest structural

changes to accommodate the body size-associated transition between inside and outside nest

division of labor in *A. cephalotes*. Comparisons of eye structure between diurnal, cathemeral,

and nocturnal ant (Greiner et al. 2007; Narendra et al. 2013; Yilmaz et al. 2014; Ogawa et al.

2019) and bee species (Greiner et al. 2004) are generally consistent with our predictions.

Although eye structure determines light sensitivity and visual acuity, other anatomical,

physiological, and behavioral adaptations modify visual abilities: variations in the size of

rhabdomers (Greiner et al. 2004; Gonzalez-Bellido et al. 2011; Narendra et al. 2017),

microsaccadic rhabdomere contractions and microvilli refractory time (Juusola et al. 2016), or

pupillary systems mediated by pigment ommatidial cells (Narendra et al. 2013, 2016a) Such

visual adaptations in *A. cephalotes* polymorphic workers remain to be studied.

Division of labor and visual neuropil size and structure

In ant species characterized by morphological differentiated subcastes, workers are

predicted to vary neurobiologically to support the sensory demands of specialized tasks

(Muscedere and Traniello 2012; Kamhi et al. 2015; Gordon et al. 2019). If metabolic costs

associated with the production and maintenance of brains is high, then selection should favor the

reduction of neuropil size (e.g.(Aiello and Wheeler 1995; Niven and Laughlin 2008)). We found

that brain volume increased with worker size, but larger workers had brains smaller than

expected from their body size (Fig. 2A). We expected *A. cephalotes* workers would invest

differentially in brain compartments due to their body size-related task repertoires, and found

that larger workers had larger eyes and an allometric increase in OL volume (Fig. 2B).

Conversely, some diurnal moths have smaller eyes but larger optic lobes than nocturnal species

(Stöckl et al. 2016), a pattern also found in analogous brain region in teleost fishes (Iglesias et al.

2018). *A. cephalotes* OL were disproportionally larger in larger workers, and consistent with our

prediction, minims showed disproportionately less OL investment. This suggests a task-related

increasing need for primary visual information processing in larger workers.

Our analysis revealed that lamina, medulla and lobula increased with worker size (Fig.

S1), maybe due to higher exposure to light in larger workers active outside the nest (as in

(Yilmaz et al. 2016)). Within the OLs, larger workers possessed disproportionally larger lamina

and lobula, but a disproportionally smaller medulla (Fig. 2C1-C3). These OL subregion

allometries suggest that minims might be better at detecting small-field motion whereas larger

workers might be better at processing contrast, wide-field motion, shape, and panorama

information. This neuroplasticity seems to adaptively support *A. cephalotes* task specialization

inside and outside the nest. We also found a disproportional investment in the MB collar in

larger workers (Fig. 2D). Enlarged MBs in social hymenopterans might be the result of ancestral

neuroanatomical adaptations to process novel visual information [60]. This evolutionary scenario

across phylogenetically diverse ant species appears to be reflected in *A. cephalotes* subcastes that

vary in visual ecologies. Our results suggest that the increased need for visual cognition in larger

workers is greater for primary processing than for higher-order processing. In *Myrmecia* species,

nocturnal workers invested relatively less in OL but relatively more in the MB, including the

collar, than diurnal workers (Sheehan et al. 2019). Our results showed that minims had the

highest collar:optic lobe ratio (Fig. 2E), apparently as an adaptation to performing tasks in

darkness. Collaterally, studies of gene expression differences in whole brains of *A. cephalotes*

subcastes revealed a significant worker size-related increase in the level of a gene associated

with rod cell development, mirroring the higher demand for visual acuity and larger eye

structures in larger workers (Muratore et al., unpublished data). This trend was also true for a

gene associated with growth factor activity, potentially contributing to the allometric OL

enlargement and other brain regions.

Conclusions

We found optical and neural plasticity are associated with the complex agrarian division

of labor of *A. cephalotes* workers. Previous studies describe differences in eye structure (Menzel

and Wehner 1970; Bernstein and Finn 1971; Klotz et al. 1992; Baker and Ma 2006; Schwarz et

al. 2011) or visual neuropil investment (O'Donnell et al. 2018). Our results advance our

understanding of ant visual system functionality by demonstrating caste-related compound eye

and brain plasticity that has evolved in response to worksite light levels. Worker polymorphism

has been shown to be correlated with patriline in the several leafcutting ant species (Hughes et al.

2003; Evison and Hughes 2011), suggesting a potential link between genetic variation and the

neuroanatomical patterns described here. Division of labor underpinning the fungicultural habits

of *A. cephalotes* appears to have played an important selective role in worker visual system

evolution. Worker behavior in this species, however, depends on visual and olfactory

information that likely varies with the cognitive requirements of tasks. The influence of these

factors on the spatial resolving power and sensitivity of eyes and macroscopic and cellular

structure of *A. cephalotes* brains requires further study.

Conflicts of interest/Competing interests

The authors declare that they have no conflict of interest

Author's contributions

APH and ESR prepared brains for imaging and registered images; APH, ESR and SA analyzed

neuropil volumes; ESR initiated and designed the eye metric imaging protocol, registered eye

images, and measured eye structure; APH and SA performed statistical analysis; APH collected

ant colonies; IBM analyzed confocal images and contributed to manuscript content; ESR and

APH prepared the first draft of the manuscript; JFAT, APH, ESR, and SA conceptualized and

designed the study. JFAT secured funding for the study; JFAT and SA wrote the manuscript. All

authors edited and approved the final content of the manuscript.

Acknowledgments

We are very grateful for the assistance of Dr. Darcy Gordon (study design, scaling analysis, and

statistics), Dr. Christopher Starr and Ricardo "Monkey" Pillai (colony collection), Dr. Todd

Blute (confocal training), Dr. Alfonso Pérez-Escudero (data analysis and interpretation), and Dr.

J. Frances Kamhi (insightful comments on the manuscript). This research was supported by

National Science Foundation grant IOS 1354291 to JFAT, a Marie Skłodowska-Curie Individual

Fellowship BrainiAnts-660976 and Ayudas destinadas a la atracción de talento investigador a la

Comunidad de Madrid en centros de I+D to SAC, the Undergraduate Research Opportunities

Program at Boston University to ESR, and a National Science Foundation Graduate Research

Fellowship (DGE-1247312) to APH.

References

Abràmoff MD, Hospitals I, Magalhães PJ, Abràmoff M (2004) Image Processing with ImageJ. Biophotonics Int 11:36–42

Aiello L, Wheeler P (1995) The expensive-tissue hypothesis: the brain and the digestive system in human and primate evolution. Curr Anthropol 36:199–221

Baker GT, Ma PWK (2006) Morphology and number of Ommatidia in the compound eyes of Solenopsis invicta, Solenopsis richteri, and their hybrid (Hymenoptera: Formicidae). Zool Anz 245:121–125. https://doi.org/10.1016/j.jcz.2006.06.001

Bernstein S, Finn C (1971) Ant compound eye: Size-related ommatidium differences within a single wood ant nest. Experientia 27:708–710. https://doi.org/10.1007/BF02136977

Beugnon G, Fourcassié V (1988) How do red wood ants orient during diurnal and nocturnal foraging in a three dimensional system? II. Field experiments. Insectes Soc 35:106–124

Beugnon G, Lachaud JP, Chagné P (2005) Use of long-term stored vector information in the neotropical ant *Gigantiops destructor*. J Insect Behav 18:415–432. https://doi.org/10.1007/s10905-005-3700-8

Bucher D, Scholz M, Stetter M, et al (2000) Correction methods for three-dimensional reconstructions from confocal images: I. Tissue shrinking and axial scaling. J Neurosci Methods 100:135–143. https://doi.org/10.1016/S0165-0270(00)00245-4

Bulova SJ, Purce K, Khodak P, et al (2016) Into the black and back: the ecology of brain investment in Neotropical army ants (Formicidae: Dorylinae). Sci Nat 103:31. https://doi.org/DOI 10.1007/s00114-016-1353-4

Cherrett JM (1968) The foraging behaviour of *Atta cephalotes* L. (Hymenoptera, Formicidae). J Anim Ecol 37:387. https://doi.org/10.2307/2955

Corral-López A, Garate-Olaizola M, Buechel SD, et al (2017) On the role of body size, brain size, and eye size in visual acuity. Behav Ecol Sociobiol 71:179. https://doi.org/10.1007/s00265-017-2408-z

Czaczkes TJ, Grüter C, Ratnieks FLW (2015) Trail Pheromones, An integrative view of their role in social insect colony organization. Annu Rev Entomol 60:581–599. https://doi.org/10.1146/annurev-ento-010814-020627

Davies R (2002) Hypothesis testing when a nuisance parameter is present only under the
alternative: Linear Model Case. Biometrika 89:484–489

Dyer AG, Paulk AC, Reser DH (2011) Colour processing in complex environments: Insights
from the visual system of bees. Proc R Soc B Biol Sci 278:952–959.
https://doi.org/10.1098/rspb.2010.2412

Ehmer B, Gronenberg W (2004) Mushroom body volumes and visual interneurons in ants:
comparison between sexes and castes. J Comp Neurol 469:198–213.
https://doi.org/10.1002/cne.11014

Evison SEF, Hughes WOH (2011) Genetic caste polymorphism and the evolution of polyandry
in *Atta* leaf-cutting ants. Naturwissenschaften 98:643–649. https://doi.org/10.1007/s00114-
011-0810-3

Farris SM (2016) Insect societies and the social brain. Curr Opin Insect Sci 1–8.
https://doi.org/10.1016/j.cois.2016.01.010

Freas CA, Wystrach A, Narendra A, Cheng K (2018) The view from the trees: Nocturnal bull
ants, *Myrmecia midas*, use the surrounding panorama while descending from trees. Front
Psychol 9:1–15. https://doi.org/10.3389/fpsyg.2018.00016

Gonzalez-Bellido PT, Wardill TJ, Juusola M (2011) Compound eyes and retinal information
processing in miniature dipteran species match their specific ecological demands. Proc Natl
Acad Sci U S A 108:4224–4229. https://doi.org/10.1073/pnas.1014438108

Gordon DG, Ilieş I, Traniello JFA (2017) Behavior, brain, and morphology in a complex insect
society: trait integration and social evolution in the exceptionally polymorphic ant *Pheidole*
rhea . Behav Ecol Sociobiol 71:166. https://doi.org/https://doi.org/10.1007/s00265-017-
2396-z

Gordon DG, Zelaya A, Arganda-Carreras I, et al (2019) Division of labor and brain evolution in
insect societies: Neurobiology of extreme specialization in the turtle ant *Cephalotes varians*.
PLoS One 14:1–16. https://doi.org/10.1371/journal.pone.0213618

Graham P, Cheng K (2009) Ants use the panoramic skyline as a visual cue during navigation.
Curr Biol 19:R935–R937. https://doi.org/10.1016/j.cub.2009.08.015

Greiner B (2006) Adaptations for nocturnal vision in insect apposition eyes. Int Rev Cytol
250:1–46. https://doi.org/10.1016/S0074-7696(06)50001-4

Greiner B, Narendra A, Reid SF, et al (2007) Eye structure correlates with distinct foraging-bout
timing in primitive ants. Curr Biol 17:879–880. https://doi.org/10.1016/j.cub.2007.08.015

Greiner B, Ribi WA, Warrant EJ (2004) Retinal and optical adaptations for nocturnal vision in
the halictid bee *Megalopta genalis*. Cell Tissue Res 316:377–390.
https://doi.org/10.1007/s00441-004-0883-9

Gronenberg W (2008) Structure and function of ant (Hymenoptera: Formicidae) brains: strength
in numbers. Myrmecol News 11:25–36

Gronenberg W (2001) Subdivisions of hymenopteran mushroom body calyces by their afferent
supply. J Comp Neurol 435:474–489. https://doi.org/10.1002/cne.1045

Gronenberg W, Hölldobler B (1999) Morphologic representation of visual and antennal
information in the ant brain. J Comp Neurol 412:229–240.
https://doi.org/10.1002/(SICI)1096-9861(19990920)412:2<229::AID-CNE4>3.0.CO;2-E

Hölldobler B (1980) Canopy orientation : a new kind of orientation in ants. Science (80-)
210:86–88

Hölldobler B, Wilson EO (2010) The leafcutter ants: civilization by instinct. WW Norton &
Company

Hölldobler B, Wilson EO (1990) The Ants. Harvard University Press, Cambridge, MA

Huber R, Knaden M (2015) Egocentric and geocentric navigation during extremely long foraging paths of desert ants. J Comp Physiol A Neuroethol Sensory, Neural, Behav Physiol 201:609–616. https://doi.org/10.1007/s00359-015-0998-3

Hughes WOH, Sumner S, Van Borm S, Boomsma JJ (2003) Worker caste polymorphism has a genetic basis in *Acromyrmex* leaf-cutting ants. Proc Natl Acad Sci U S A 100:9394–9397. https://doi.org/10.1073/pnas.1633701100

Iglesias TL, Dornburg A, Warren DL, et al (2018) Eyes Wide Shut: the impact of dim-light vision on neural investment in marine teleosts. J Evol Biol 31:1082–1092. https://doi.org/10.1111/jeb.13299

Jander U, Jander R (2002) Allometry and resolution of bee eyes (Apoidea). Arthropod Struct Dev 30:179–193. https://doi.org/10.1016/S1467-8039(01)00035-4

Juusola M, Dau A, Song Z, et al (2016) Microsaccadic information sampling provides *Drosophila* hyperacute vision. Elife 6:e26117. https://doi.org/10.7554/eLife.26117

Kamhi JF, Nunn K, Robson SKA, Traniello JFA (2015) Polymorphism and division of labour in a socially complex ant: neuromodulation of aggression in the Australian weaver ant, *Oecophylla smaragdina*. Proc Biol Sci 282:20150704-. https://doi.org/10.1098/rspb.2015.0704

Kamhi JF, Sandridge-Gresko A, Walker C, et al (2017) Worker brain development and colony organization in ants: Does division of labor influence neuroplasticity? Dev Neurobiol 77:1072–1085. https://doi.org/10.1002/dneu.22496

Kapustjanskij A, Streinzer M, Paulus HF, Spaethe J (2007) Bigger is better: Implications of body size for flight ability under different light conditions and the evolution of alloethism in bumblebees. Funct Ecol 21:1130–1136. https://doi.org/10.1111/j.1365-2435.2007.01329.x

Kaspari M, Weiser MD (1999) The size–grain hipothesis and interespecific scaling in ants. Fuctional Ecol 13:530–538. https://doi.org/10.1046/j.1365-2435.1999.00343.x

Klotz JH, Reid BL (1992) The use of spatial cues for structural guideline orientation in Tapinoma sessile and Camponotus pennsylvanicus (Hymenoptera: Formicidae). J Insect Behav 5:71–82. https://doi.org/10.1007/BF01049159

Klotz JH, Reid BL, Gordon WC (1992) Variation of ommatidia number as a function of worker size in *Camponotus pennsylvanicus* (DeGeer) (Hymenoptera: Formicidae). Insectes Soc 39:233–236. https://doi.org/10.1007/BF01249297

Knaden M, Graham P (2016) The sensory ecology of ant navigation: from natural environments to neural mechanisms. Annu Rev Entomol 61:63–76. https://doi.org/10.1146/annurev-ento-010715-023703

Land MF (1997) Visual Acuity in Insects. Annu Rev Entomol 42:147–177. https://doi.org/10.1146/annurev.ento.42.1.147

Linksvayer TA, Mccall AC, Jensen RM, et al (2016) The function of hitchhiking behavior in the leaf-cutting ant Atta cephalotes. Biotropica 34:93–100. https://doi.org/10.1111/j.1744-7429.2002.tb00245.x

Mares S, Ash L, Gronenberg W (2005) Brain allometry in bumblebee and honey bee workers. Brain Behav Evol 66:50–61. https://doi.org/10.1159/000085047

Menzel R, Wehner R (1970) Augenstrukturen bei verschieden großen Arbeiterinnen von *Cataglyphis bicolor* Fabr. (Formicidae, Hymenoptera). Z Vgl Physiol 68:446–449. https://doi.org/10.1007/BF00297741

Menzi U (1987) Visual adaptation in nocturnal and diurnal ants. J Comp Physiol A 160:11–21.

https://doi.org/10.1007/BF00613437

Moser JC, Reeve JD, Bento JMS, et al (2004) Eye size and behaviour of day- and night-flying
leafcutting ant alates. J Zool 264:69–75. https://doi.org/10.1017/S0952836904005527

Muller M, Wehner R (2006) The significance of direct sunlight and polarized skylight in the
ant's celestial system of navigation. Proc Natl Acad Sci 103:12575.
https://doi.org/10.1073/pnas.0604430103

Müller M, Wehner R (2010) Path integration provides a scaffold for landmark learning in desert
ants. Curr Biol 20:1368–1371. https://doi.org/10.1016/j.cub.2010.06.035

Muscedere ML, Traniello JFA (2012) Division of labor in the hyperdiverse ant genus *Pheidole* is
associated with distinct subcaste-and age-related patterns of worker brain organization.
PLoS One 7:e31618. https://doi.org/10.1371/journal.pone.0031618

Narendra A, Alkaladi A, Raderschall CA, et al (2013) Compound eye adaptations for diurnal and
nocturnal lifestyle in the intertidal ant, *Polyrhachis sokolova*. PLoS One 8:e76015.
https://doi.org/10.1371/journal.pone.0076015

Narendra A, Greiner B, Ribi WA, Zeil J (2016a) Light and dark adaptation mechanisms in the
compound eyes of *Myrmecia* ants that occupy discrete temporal niches. J Exp Biol
219:2435–2442. https://doi.org/10.1242/jeb.142018

Narendra A, Kamhi JF, Ogawa Y (2017) Moving in dim light: behavioral and visual adaptations
in nocturnal ants. Integr Comp Biol 57:1104–1116. https://doi.org/10.1093/icb/icx096

Narendra A, Ramirez-Esquivel F, Ribi WA (2016b) Compound eye and ocellar structure for
walking and flying modes of locomotion in the Australian ant, *Camponotus consobrinus*.
Sci Rep 6:1–10. https://doi.org/10.1038/srep22331

Narendra A, Reid SF, Greiner B, et al (2011) Caste-specific visual adaptations to distinct daily
activity schedules in Australian *Myrmecia* ants. Proc R Soc B Biol Sci 278:1141–1149.
https://doi.org/10.1098/rspb.2010.1378

Niven JE, Laughlin SB (2008) Energy limitation as a selective pressure on the evolution of
sensory systems. J Exp Biol 211:1792–804. https://doi.org/10.1242/jeb.017574

O'Donnell S, Bulova S, Barrett M, Von Beeren C (2018) Brain investment under colony-level
selection: Soldier specialization in *Eciton* army ants (Formicidae: Dorylinae). BMC Zool
3:1–6. https://doi.org/10.1186/s40850-018-0028-3

O'Donnell S, Clifford MR, Bulova SJ, et al (2014) A test of neuroecological predictions using
paperwasp caste differences in brain structure (Hymenoptera: Vespidae). Behav Ecol
Sociobiol 68:529–536. https://doi.org/10.1007/s00265-013-1667-6

Ogawa Y, Ryan LA, Palavalli-Nettimi R, et al (2019) Spatial resolving power and contrast
sensitivity are adapted for ambient light conditions in Australian *Myrmecia* ants. Front Ecol
Evol 7:1–10. https://doi.org/10.3389/fevo.2019.00018

Ott SR (2008) Confocal microscopy in large insect brains: Zinc-formaldehyde fixation improves
synapsin immunostaining and preservation of morphology in whole-mounts. J Neurosci
Methods 172:220–230. https://doi.org/10.1016/j.jneumeth.2008.04.031

Palavalli-Nettimi R, Narendra A (2018) Miniaturisation decreases visual navigational
competence in ants. J Exp Biol 221:jeb177238. https://doi.org/10.1242/jeb.177238

Perl CD, Niven JE (2016a) Colony-level differences in the scaling rules governing wood ant
compound eye structure. Sci Rep 6:1–8. https://doi.org/10.1038/srep24204

Perl CD, Niven JE (2016b) Differential scaling within an insect compound eye. Biol Lett 12:0–3.
https://doi.org/10.1098/rsbl.2016.0042

Powell S, Clark E (2004) Combat between large derived societies: A subterranean army ant

established as a predator of mature leaf-cutting ant colonies. Insectes Soc 51:342–351.
https://doi.org/10.1007/s00040-004-0752-2

Rodrigues PAP, Oliveira PS (2014) Visual navigation in the Neotropical ant *Odontomachus*
hastatus (Formicidae, Ponerinae), a predominantly nocturnal, canopy-dwelling predator of
the Atlantic rainforest. Behav Processes 109:48–57.
https://doi.org/10.1016/j.beproc.2014.06.007

Ronacher B, Wehner R (1995) Desert ants *Cataglyphis fortis* use self-induced optic flow to
measure distances travelled. J Comp Physiol A 177:21–27.
https://doi.org/10.1007/BF00243395

Rutowski RL, Gislén L, Warrant EJ (2009) Visual acuity and sensitivity increase allometrically
with body size in butterflies. Arthropod Struct Dev 38:91–100.
https://doi.org/10.1016/j.asd.2008.08.003

Schwarz S, Narendra A, Zeil J (2011) The properties of the visual system in the Australian desert
ant *Melophorus bagoti*. Arthropod Struct Dev 40:128–134.
https://doi.org/10.1016/j.asd.2010.10.003

Sheehan ZBV, Kamhi JF, Seid MA, Narendra A (2019) Differential investment in brain regions
for a diurnal and nocturnal lifestyle in Australian *Myrmecia ants*. J Comp Neurol 527:1261–
1277. https://doi.org/10.1002/cne.24617

Snyder AW (1977) Acuity of compound eyes: Physical limitations and design. J Comp Physiol
A 116:161–182. https://doi.org/10.1007/BF00605401

Spaethe J, Chittka L (2003) Interindividual variation of eye optics and single object resolution in
bumblebees. J Exp Biol 206:3447–3453. https://doi.org/10.1242/jeb.00570

Stieb SM, Hellwig A, Wehner R, Rössler W (2012) Visual experience affects both behavioral
and neuronal aspects in the individual life history of the desert ant *Cataglyphis fortis*. Dev
Neurobiol 72:729–742. https://doi.org/10.1002/dneu.20982

Stieb SM, Muenz TS, Wehner R, Rössler W (2010) Visual experience and age affect synaptic
organization in the mushroom bodies of the desert ant *Cataglyphis fortis*. Dev Neurobiol
70:408–423. https://doi.org/10.1002/dneu.20785

Stöckl A, Heinze S, Charalabidis A, et al (2016) Differential investment in visual and olfactory
brain areas reflects behavioural choices in hawk moths. Sci Rep 6:1–10.
https://doi.org/10.1038/srep26041

Strausfeld NJ (1989) Beneath the Compound Eye: Neuroanatomical Analysis and Physiological
Correlates in the Study of Insect Vision. In: Facets of Vision. Springer, Berlin, Heidelberg,
pp 317–359

Streinzer M, Brockmann A, Nagaraja N, Spaethe J (2013) Sex and caste-specific variation in
compound eye morphology of five honeybee species. PLoS One 8:1–9.
https://doi.org/10.1371/journal.pone.0057702

Taylor GJ, Tichit P, Schmidt MD, et al (2019) Bumblebee visual allometry results in locally
improved resolution and globally improved sensitivity. Elife 8:1–32.
https://doi.org/10.7554/eLife.40613

Team RC (2016) R: A Language and Environment for Statistical Computing. R Found. Stat.
Comput.

V. R. M. Muggeo (2008) segmented: An R Package to Fit Regression Models with Broken-Line
Relationships. R News 3:343–4. https://doi.org/10.1159/000323281

van Breda JM, Stradling DJ (1994) Mechanisms affecting load size determination in *Atta*
cephalotes L. (Hymenoptera, Formicidae). Insectes Soc 41:423–435.

https://doi.org/10.1007/BF01240645

Vick K (2005) Use of multiple cues for navigation by the leaf-cutter ant *Atta cephalotes*.
University of Maryland, College Park, MD

Vilela EF, Jaffé K, Howse PE (1987) Orientation in leaf-cutting ants (Formicidae: Attini). Anim
Behav 35:1443–1453. https://doi.org/10.1016/S0003-3472(87)80017-9

Warrant EJ (2008) Seeing in the dark: Vision and visual behaviour in nocturnal bees and wasps.
J Exp Biol 211:1737–1746. https://doi.org/10.1242/jeb.015396

Wehner R (2003) Desert ant navigation: How miniature brains solve complex tasks. J Comp
Physiol A Neuroethol Sensory, Neural, Behav Physiol 189:579–588.
https://doi.org/10.1007/s00359-003-0431-1

Wetterer JK (1991) Allometry and the geometry of leaf-cutting in *Atta cephalotes*. Behav Ecol
Sociobiol 29:347–351. https://doi.org/10.1007/BF00165959

Wilson EO (1980) Caste and division of labor in leaf-cutter ants (Hymenoptera: Formicidae:
Atta): II. The ergonomic optimization of leaf cutting. Behav Ecol Sociobiol 7:57–165

Yilmaz A, Aksoy V, Camlitepe Y, Giurfa M (2014) Eye structure, activity rhythms, and
visually-driven behavior are tuned to visual niche in ants. Front Behav Neurosci 8:1–9.
https://doi.org/10.3389/fnbeh.2014.00205

Yilmaz A, Lindenberg A, Albert S, et al (2016) Age-related and light-induced plasticity in opsin
gene expression and in primary and secondary visual centers of the nectar-feeding ant
Camponotus rufipes. Dev Neurobiol 76:1041–1057. https://doi.org/10.1002/dneu.22374

Zeil J, Ribi WA, Narendra A (2014) Polarisation vision in ants, bees and wasps. In: Polarized
Light and Polarization Vision in Animal Sciences, Second Edition

Figure 1: Compound eye structure in polymorphic *A. cephalotes* workers. A. Log-log plot of the ommatidia number as a function of worker HW showing a significant change of slope at 1.38mm. **B.** Log-log plot of average ommatidial diameter as a function of worker HW. **C.** Log-log plot of the eye surface area (SA) as a function of worker HW showing a significant change of slope at 1.44mm. **D.** Log-log plot of ommatidia density (ommatidial number/eye SA) as a function of worker HW. **E.** Log-log plot of the interommatidial angle (rad) as a function of worker HW showing a significant change of slope at 1.25mm. **F** Log-log plot of the eye parameter as a function of worker HW (significant change of slope at 1.26mm). **G.** Z-projections

of confocal images of eyes from workers with variable HW (scale bar=100μm). A-F: Each pink

point represents a single eye. Solid (significantly different from isometry) or dashed (not

significantly different from isometry) black lines show linear regression or piecewise linear

regressions as appropriate. Purple patches represent 95% confidence intervals of regression lines.

Dashed grey lines are the best-fitting isometric regression models.

Figure 2: Volumes of polymorphic *A. cephalotes* worker brains and brain compartments.

A. 3D reconstruction of the brain hemisphere of an *A. cephalotes* worker (HW ~4mm). **B.** Log-

log plot of hemisphere brain volume as a function of worker HW. **C.** Log-log plot of relative OL

volume as a function of worker HW. **C1.** Log-log plot of relative volume of OL lamina as a

function of worker HW. **C2.** Log-log plot of relative volume of OL medulla as a function of

worker HW.**C3.** Log-log plot of relative volume of OL lobula as a function of worker HW.**D.**

Log-log plot of relative MB collar volume as a function of worker HW. **E.** Log-log plot of MB

collar: OL volume ratio as a function of worker HW. B-E: Legend as in figure 1.

674