Aerobic scope does matter in temperature-size rule, but only under optimal conditions

Running title: Aerobic scope in the temperature-size rule

Aleksandra Walczyńska[*], Anna Maria Labecka, Mateusz Sobczyk

Institute of Environmental Sciences, Jagiellonian University, Gronostajowa 7, 30-387 Krakow,
Poland

*corresponding author

E-mail: aleksandra.walczynska@uj.edu.pl

Phone: +48 126645203

Abstract

We united the theoretical predictions on the factors responsible for the occurrence and
evolutionary significance of the temperature-size rule. We tested the causal connection among
them assuming that (i) the temperature-size rule is the response to temperature-dependent
oxygenic conditions, (ii) body size decrease is a consequence of cell shrinkage in response to
exposure to hypoxia, (iii) this response enables to keep the wide scope for aerobic performance,
and (iv) it prevents the decrease in fitness. We conducted our tests on three clones of the rotifer
Lecane inermis with different thermal preferences. These clones were exposed to three
experimental regimes: mild hypoxia, severe hypoxia driven by a too high temperature, and severe
hypoxia driven by an inadequate oxygen concentration. The results showed that our causative
reasoning was generally correct, but only under mildly hypoxic conditions. In more stressful
environments, rotifers had clone- and condition-specific responses, which in fact were equally
successful in terms of the levels of fitness. Our results join for the first time all factors connecting
the cause and effect in the temperature-size rule. They indicate the importance of the conditions
under which it should be tested. The most important messages from this study were that (i) a
decrease in the body size was one of but not the only option for preventing fitness reduction
under hypoxia, and (ii) such a response to higher temperature enabled the maintenance of wide
aerobic scope in clone-specific, thermally optimal conditions.

Key words: aerobic scope, body size, fitness, rotifers, SDA, temperature-size rule

Introduction

The phenotypic body size decrease with increasing temperature (temperature-size rule, TSR; Atkinson, 1994) has created a major puzzle for evolutionary ecologists (Berrigan and Charnov, 1994) because of its counterintuitive pattern. The reason for this confusion is that, according to the theory of optimal energy allocation, organisms growing in advantageous warm conditions should maximize their size to increase their own fitness, while those growing in harsh cold conditions should not postpone their maturity for long because of the risk of premature death. There have been different hypotheses on why the size response to temperature contradicts this pattern. Among them, the most promising is the hypothesis on the adaptive size response under oxygen deficiency (but see also Audzijonyte et al., 2019). The relative oxygen concentration decreases with increasing temperature (see Wetzel, 2001 for aquatic systems), and then the steeper increase in oxygen demands compared to oxygen supply causes oxygenic stress in ectothermic organisms (Atkinson et al., 2006; Verberk et al., 2011; Woods, 1999). The decrease in cell size has been suggested to increase the efficiency of oxygen transport to the mitochondria (Woods, 1999), and as a consequence, the body size shrinks as well. The adaptive hypothesis proposed was that temperature defines the window for the aerobic scope, namely the range of aerobic metabolism that can be performed under given conditions, and the ability to maintain the flexible efficiency of aerobic respiration prevents a possible decrease in fitness (Poertner, 2010). This mechanism was linked to the body size adjustment to temperature by Atkinson et al. (2006). The hypothesis on the oxygen-driven size-to-temperature response has been positively verified both indirectly (Berner et al., 2007; Harrison et al., 2010; Rollinson and Rowe, 2018a; Rollinson and Rowe, 2018b; Santilli and Rollinson, 2018; Verberk and Atkinson, 2013) and directly (Czarnoleski et al., 2015; Frazier et al., 2001; Hoefnagel and Verberk, 2015). However, results relating this pattern directly to organismal fitness are scarce (Prokosch et al., 2019; Walczyńska et al., 2015a), while such a reference is the only way that enables reliable conclusions on the possible evolutionary meaning of TSR.

In this study, we linked the hypothesized proximate and ultimate mechanisms behind the phenotypically driven size decrease with increasing temperature for the case of the rotifer *Lecane inermis*. We aimed to thoroughly test the theoretical reasoning behind hypoxia driving body size shrinkage to prevent the reduction in fitness, which is presented in the literature. Because the TSR has been empirically confirmed to be condition-sensitive and to be followed only in an

optimal thermal range (Walczyńska et al., 2016), we conducted our tests under optimal and

suboptimal conditions; the latter was further divided between hypoxia driven by stress-inducing

high temperatures or by the stress-inducing low oxygen levels. This distinction between mild *vs.*

harsh conditions enabled inference on the possible thresholds for TSR performance, additionally

referred to fitness.

We examined the aerobic scope, our hypothesized ultimate mechanism, by estimating the specific

dynamic action (SDA). This measure refers to the increasing metabolic expenditures in animals

and is associated with ingestion, digestion, assimilation and absorption of food (McCue, 2006;

Secor, 2009), when the increasing oxygen consumption was mostly associated with the

biochemical transformation of food and the synthesis of the new proteins (Jordan and Steffensen,

2007). In practice, it was estimated as a difference in the respiration rate between the fed and

hungry individuals that cannot be maintained at rest. The hypothesized link between aerobic

scope and SDA is that the larger the oxygen consumption associated with SDA, the wider the

capacity of aerobic metabolism (aerobic scope). This assumption was previously used for the

case of *Daphnia magna* (Chopelet et al., 2008). Our approach was to compare the oxygen

consumption rate in hypoxia for animals previously exposed to normoxic or hypoxic conditions.

We expected that hypoxia would lower the SDA amplitude in hypoxia-naïve animals compared

to hypoxia-experienced animals. This hypothesis was based on Fry (1947), who wrote that

dissolved oxygen acts as a limiting factor for metabolic rate; thus, its reduction decreases aerobic

scope and, as a consequence, the energy budgets of aquatic breathers. Such an arresting effect of

hypoxia on SDA width was previously reported for a cod fish (Jordan and Steffensen, 2007) and

carabid beetles (Gudowska and Bauchinger, 2018). We hypothesized that hypoxia selects for

small cell size, followed by small body size, to maintain high effectiveness of aerobic metabolism

and to maximize fitness (Fig. 1A). We expected this pattern to differ across the hypoxia

treatments, with the more direct response under mild hypoxia. Therefore, our main goal was to

compare all traits examined in rotifers exposed to normoxia and those exposed to hypoxia, across

three experimental treatments.

Apart from conducting the study in optimal or stressful conditions, we also tested for possible

differences in the responses of animals acclimated to different thermal conditions: cold,

intermediate, and warm. Such a distinction addresses an issue of the possible linkage between the

thermal preferences and thermally induced body size adjustment. We used three clones of the

rotifer *L. inermis*, a species that has previously been found to follow the TSR in the laboratory

and in the field (Kiełbasa et al., 2014), to show the adaptive body size response to thermally

driven oxygen conditions based on the cell size adjustment (Walczyńska et al., 2015a) and to

perform simple two-point TSR control within its lifecycle, by the mother and during egg

development (Walczyńska et al., 2015b). To our knowledge, this is the first attempt to test for the

proximate and ultimate mechanisms behind the TSR in such a comprehensive way.

Materials and Methods

Clones studied. The *Lecane inermis* Bryce rotifer species has previously been tested for different

aspects of size response to thermal and oxygenic conditions. Its thermal performance curve was

also examined, and the optimal temperature was estimated to be 30°C (Walczyńska et al., 2016).

This species is very convenient for the studies on the changes in phenotypic response between the

maternal and offspring generations because of its short generation time, estimated previously for

the range between 15 °C and 25 °C as two days (Walczyńska et al., 2015b). We chose three *L.*

inermis clones that were founded from one individual each, which were isolated from activated

sludge from a small wastewater treatment plants in southern Poland. The clones differed in their

acclimation conditions due to exposure to laboratory conditions representing relatively low,

intermediate or high temperatures. Thus, we assumed the existence of differences in thermal

preferences among the clones. Our predictions were confirmed in a previous study conducted for

six clones (Stuczyńska et al., 2020). A general pattern found was the linkage between the pre-

experimental thermal exposure and clonal thermal preference during the study. The three clones

used here refer to clones Cold, Int2 and Warm2 from the previous study and are here named Cold

for the cold preference, Int for intermediate temperature preference and Warm for high heat

preference, respectively. Preceding the experiment, all clones were kept in Petri dishes at 25°C in

the dark in Żywiec spring water (Poland) and fed patented Novo powder (a patent by Pajdak-Stós

et al., 2017).

Experimental setup. The experiment was divided into three parts that constituted different

hypoxic regimes (Fig. 1 B): MH – mild hypoxia (30°C and 80% or 40% O_2) was followed by

SHT – severe hypoxia with temperature above the optimum (38°C and 80% or 40% O_2) and then

by SHO2 – severe hypoxia with very low oxygen level (30°C and 80% or 20% O_2). In each part,

the three rotifer clones were exposed to either normoxia (80% O_2) or hypoxia (40% or 20% O_2).

The choice of 80% normoxia instead of 100% was dictated by the fact that it is much easier to

maintain stable 80% than 100% oxygen concentrations, and in this case, we were assured that the

rotifers were not exposed to fluctuating oxygen conditions that could act as an uncontrolled

factor. The relative 80% oxygen concentration was equivalent to approximately 6 mg/L O_2,

which is very close to the optimal value for this species living in nature (7.5 mg/L O_2) (Berzins

and Pejler, 1989) and much higher than the oxygen concentration frequently set under the

activated sludge conditions (2 mg/L O_2)(Kiełbasa et al., 2014) from which all the studied clones

were established. The value of 2-3 mg/L O_2 was also provided as a standard threshold for

hypoxia (reviewed in Hrycik et al., 2017).

The experimental conditions were set in four temperature-regulated water baths (Memmert,

Germany) connected to a 4-channel OxyReg O_2 regulation system (Loligo Systems, Denmark)

calibrated to 0% and 100% air-saturated water. A stable level of hypoxia was maintained by

adding pure (comestible) nitrogen to the baths. The relative oxygen concentrations of 80, 40 and

20% O_2 were equivalent to approximately 17% (~6.0 mg/L), 8% (~2.8 mg/L) and 4% (~1.5

mg/L) of the atmospheric oxygen, respectively. Two baths were set to normoxia and two to

hypoxia, with the same temperature in all. Six 100 mL cages made of 5 µm mesh enclosed in

slide frames sealed with silicone were placed in each water bath (two cages-replicates per clone

in each bath). This type of cage has been successfully used previously (Walczyńska et al., 2015a).

Each water bath was filled with Żywiec spring water as the medium. The initial densities of

rotifers planted at the beginning of experimental parts were 110, 120 and 250 indiv./mL for MH,

SHT and SHO2 stages, respectively. These numbers were dictated by the clone that was the least

numerous at the moment of counting; our aim was to start with the highest possible number of

rotifers, even across clones. Such initial densities of rotifers were far below the population

carrying capacity because *L. inermis* rotifers may reach densities as high as tens of thousands per

mL (45 000 indiv/mL was reported by Walczyńska et al., 2017). The rotifers in each cage were

fed 200 µL of commercial bioproduct Bio-Trakt®, which has been successfully used in *L.*

inermis mass culture previously (Fiałkowska et al., 2019). During or just after the three days of

each experimental stage, we took samples to estimate the physiological (specific dynamic action

(SDA), which is a proxy for aerobic scope, and nucleus size, which is a proxy for cell size, and

life history traits (body size, population growth rate, egg ratio), as presented in Fig. 1B.

Oxygen consumption. The hypothesis of the decrease in size under hypoxic conditions to

maintain a wide aerobic scope was tested by estimating the SDA. This physiological difference in

respiration rate between fed and hungry animals was previously used as a proxy for aerobic scope

in the case of *Dapnia magna* (Chopelet et al., 2008). We estimated the SDA using

spectrophotometry (Orion^{TM} AquaMate 8000 UV-VIS, Thermo Scientific, USA) with a modified

Winkler titration (Carpenter, 1965; Chopelet et al., 2008; Roland et al., 1999), which has been

utilized as a highly accurate method for determining dissolved oxygen (Helm et al., 2012). In this

method, the absorbance measure is proportional to the amount of oxygen that is dissolved in a

sample as a consequence of the chemical reaction chain. For each experimental stage, we

sampled the rotifers from the normoxia and hypoxia treatments and measured the aerobic scope

of both groups exposed to hypoxic conditions, respective to the given stage. We expected that the

rotifers exposed to experimental hypoxia in the preceding three days should have wider SDAs in

the test under hypoxia than the normoxia-exposed rotifers. Two clones, Cold and Warm, were

examined for SDA because the Int clone did not proliferate well enough to achieve reasonable

sample sizes. On the third day of each experimental stage, we collected 10 mL samples from each

cage and pooled the two replicates within the clone×oxic regime combination. For logistical

reasons, the clones were analysed separately but within the same day for the cases of SHT stage

and SHO2 stage. Because we did not conduct an analysis for both clones after the MH stage, we

repeated this stage at the end of the experiment to collect the missing data on SDA for the Cold

clone.

Each clone was divided into two 50 mL crystallizers with a 5 mL rotifer sample and 15 mL of

hypoxic medium taken directly from the respective water bath representing the experimental

hypoxic conditions. By applying this procedure, we diluted the nutritional particles sampled

along with the rotifers. The rotifers in one crystallizer were then fed with 10 µL of Bio-Trakt®

per 1 mL of sample (treatment: fed). The rotifers in the second crystallizer were not fed

(treatment: hungry). The random division of clone samples into the fed and hungry treatments

caused that the average rotifer body size was similar in both regimes. Therefore, we could assume

that the possible differences in oxygen consumption were not related to size differences.

Both samples were incubated for 2 h at the same temperature the rotifers experienced during the

respective experimental stage. The relatively short exposure to starving conditions was dictated

by the fact that rotifers become starved relatively quickly and that severe starvation has a very

profound effect on metabolism (reviewed by Galkovskaya, 1995).

After this period, from both treatments, we fished the swimming, healthy-looking rotifers in

bunches of 30 and transferred them to the wells of 24-well tissue plates filled with membrane

made of parafilm, with four or five replicates (= rotifer bunches) per treatment. The rotifers from

each well were then quantitatively transferred to individual 2 mL autosampler vials (Agilent

Technologies, USA) with hypoxic medium taken directly from one experimental water bath with

hypoxic conditions. The vials (five replicates per fed and hungry treatments) were filled

according to the following procedure: 1 mL of medium, the rotifer sample from a well, and

another 1 mL of medium until a convex meniscus was achieved. The vials were then closed with

dedicated caps punctured beforehand with a sterile needle to remove excess fluid and air. The

control tubes were prepared following the same procedure but without the animals. The samples

were incubated for another two (38°C) or three (30°C) hours in hypoxic conditions, with the level

of hypoxia set to 40%, 40% and 20% O_2 for the experimental MH, SHT and SHO2 stages,

respectively.

After incubation, the Winkler chemicals were added following the procedure adopted from

Carpenter (1965): 16 µL of $MnCl_2 \cdot 4H_2O$ (3 M), 16 µL of NaI (4 M) in NaOH (8 M), and, after

vigorous shaking to cause precipitation, 16 µL of H_2SO_4 (5 M). The vials were then shaken to

dissolve the precipitate, and 800 µL subsamples were taken directly to spectrophotometric micro-

cuvettes (Brand, Germany). The absorbance was measured immediately at a wavelength of 440

nm within the first 10 minutes to prevent iodine evaporation (Roland et al., 1999). The calibration

curve for absorbance was estimated for 0, 15, 35 and 50% O_2 (in relative values), with 5

replicates per oxygen level. In the final estimation, the extreme values were excluded, and the

average was calculated for the remaining three values per oxygen level. The calibration curve

fitting was estimated at $R^2 = 0.9953$.

The control samples were checked for their reliability for each combination of hypoxia type and

clone separately. In all but two cases, the controls showed similar and reliable patterns. In the

case of the MH treatment with the Warm clone, the control samples showed a pattern indicating

some sort of methodological error (supplementary materials, Fig. S1), and only one value

attained for the first sample was taken into consideration. In the case of the SHT treatment with

the Cold clone, one control sample had a very low value (Fig. S1), which made it unreliable. The

final control value was calculated as a mean from the remaining two samples. SDA was

estimated both as a difference in raw absorbance gained for the fed rotifers and that for hungry

rotifers (absolute SDA) and as their division (factorial SDA). The comparison of these two

approaches for the aerobic scope calculation was suggested by Clark et al. (2013).

The amount of consumed O_2 was calculated from % to mg/L using the equation describing the

relationship between these two variables provided by the OxyReg system. This was possible only

for 30°C, due to the relatively wide range of O_2 (%) covered (from approximately 19 to 95%),

while the much shorter range for 38°C made this estimation less reliable.

Cell size. We measured the nucleus size, a proxy for cell size, according to the method developed

and used before for the same species (Walczyńska et al., 2015a). Such a procedure was dictate by

the fact that cell size was reported to correlate with DNA size (Jalal et al., 2013; Kozlowski et al.,

2003), meaning that nucleus size might be a convenient estimate of cell size (reviewed by

Cavalier-Smith, 2005). The rotifers were sampled on the third day of each experimental part.

Approximately 50 swimming rotifers were fished and placed in 1 mL Eppendorf tubes, and the

samples were fixed with 10% buffered neutral formalin (Avantor Performance Materials Poland

S.A., Poland). The staining procedure was based on nuclear red (Carl Roth, Germany), as

described in Walczyńska et al. (2015a). We measured two perpendicular diameters of 10 nuclei

per rotifer using ImageJ (NIH, USA), and we calculated their product. The nuclei represent

different animal organs. In this context, it is worth noting that rotifers do not possess any

specialized respiratory cells, and they respire through the whole body surface (Galkovskaya,

1995).

Body size. We estimated the strength of the response to hypoxia in comparison to normoxia by

measuring the body size. For this purpose, we sampled the rotifers on the second day of the

experiment and fixed them with Lugol solution. The rotifers were photographed (Eclipse 80i

microscope (Nikon, Japan) equipped with a DS-Ri2 camera assisted by NIS-Elements D software

(Nikon)). The length and width were measured in ImageJ, and their product acted as a proxy for

body area, as previously applied to the rotifers *Brachionus* sp. (Walczyńska and Serra, 2014) and

L. inermis (Walczyńska et al., 2015a).

Fitness measures. Two measures of fitness were compared: population growth rate and egg

ratio, which is the number of eggs laid by an average female. To estimate the former, we sampled

1 mL from each cage (two cages × two water baths gave four replicates per clone×oxic regime

combination), fixed with Lugol solution and counted the rotifers. These procedures were used

twice: one day (count I) and after two days (count II) after the onset of the stage. During count II,

the egg number was noted along with the female number for egg ratio estimation.

Statistical analyses. All analyses were conducted in SAS 9 (SAS, 2013) in PROC MIXED,

method = reml. In the case of SDA, the GLM model included four fixed factors: hypoxia type

(experimental stage), oxic regime of rotifer origin (normoxia *vs*. hypoxia), and clone and food

regime (fed *vs*. hungry), along with all their interactions. The non-significant interactions in the

model were excluded using backward elimination.

Possible differences in the nuclei size and body size were tested separately in GLM models that

included three fixed factors: hypoxia type, clone and oxic regime, and their interactions. The non-

significant interactions were excluded using backward elimination. In the case of body size, the

differences in size among the hypoxia types caused the considerable lack of homogeneity of

variance, which prevented statistical analysis. Therefore, the data were standardized by

recalculating them to the units of standard deviation for each experimental stage (= hypoxia type)

separately.

Both fitness measures were analysed with ANCOVA (PROC MIXED) and were not analysed as

ratios, which has been cited as a more appropriate method (Raubenheimer, 1995; Raubenheimer

and Simpson, 1992). The models included three fixed factors: hypoxia type, clone and oxic

regime, and their interactions. In the case of population growth rate *r*, the dependent variable was

the rotifer number from count II, while that from count I was a covariate. In the case of the egg

ratio, the dependent variable was the number of eggs, while a covariate was the number of rotifer

females in count II. The dependent variables and covariates were natural logarithm-transformed.

In the initial stages, we included all the possible first-level interactions of the covariates with the

main factors to test whether they were appropriately used in the models. In both models, the

covariates were found to significantly affect the dependent variables and to not interact with the

main factors. In the next step, we constructed the models of all the main factors and possible

interactions, plus a covariate. The non-significant interactions were removed using backward

elimination.

Results

Oxygen consumption. The results for the absolute oxygen consumption, which was calculated

from the equation O_2 [mg/L] $= 0.0752 \times O_2$ [%], were compared with other published data for

different rotifer species obtained from the literature (Table 1). Generally, the oxygen

consumption rate for the units provided in Table 1 was approximately 4 times higher in fed

rotifers than in hungry rotifers. This estimated difference was previously shown to be 2-3-fold

(Galkovskaya, 1995), although the units and starvation times for this comparison were unclear.

The oxygen consumption differed among hypoxia types, clones, rotifer origins (normoxia or

hypoxia), and food regimes, as well as for some interactions (Table 2). Most interestingly, the O_2

consumption of the hungry and fed rotifers significantly differed between the clones and the

hypoxia types, according to the significant *hypoxia type×clone×food* interaction (Table 2). In

each case, the fed rotifers consumed more oxygen than the hungry rotifers due to the metabolic

costs associated with SDA. The ceiling of these costs was relatively constant across hypoxia

types in the case of the Warm clone and was elevated in cold rotifers exposed to hypoxia driven

by stress-inducing high temperatures. The SDA-related costs were generally higher in the rotifers

that had experienced hypoxia in the experiment than in hypoxia-naïve rotifers (Fig. 2A). The

negative value for the Warm clone from normoxia in the MH treatment was caused by two

elevated points (see supplementary materials, Fig. S1), but the analysis for outliers showed that

there was no objective reason for their removal from the dataset. The pattern for the SDA was

similar regardless of the measure (whether absolute or factorial; Fig. 2). Generally, the width of

the SDA across the hypoxic treatments was similar for the Cold clone, while in the Warm clone,

the values were much higher under mild hypoxia than under the two severe hypoxia treatments.

In mildly hypoxic conditions, both clones showed wider SDAs when previously exposed to

hypoxia during the experiment than those that were hypoxia-naïve. This pattern was not repeated

in the following severe hypoxia treatment. The cold clone maintained relatively high and similar

SDA, regardless of previous hypoxia experience, while the warm clone displayed a clearly

narrower SDA in the rotifers experiencing hypoxia than in those taken from the normoxia

treatment (Fig. 2).

Cell size. In total, we measured 1386 nuclei in 143 animals. For the experimental SHO2 stage,

we were only able to take measurements for the Warm clone; the samples of other two clones

were lost, most probably because of some unknown error with fixing. The mean nuclei size of

1.99 $\mu m^2 \pm 0.62$ SD was close to the 1.62 μm^2 achieved previously at 32°C for the different *L.*

inermis clones (Walczyńska et al., 2015a). We found a significant effect of the type of hypoxia

and its interaction with the clone and oxygenic conditions on the nuclei size (Table 3), yet there

was no effect of the oxygenic conditions alone. The clones differed in cell size, and their diverse

responses to the oxygenic conditions varied with hypoxia type (significant *hypoxia*

type×clone×oxic regime interaction in Table 3). In general, the Cold clone had the largest nuclei,

followed by the Int clone, and those of the Warm clone were the smallest. The response to

hypoxia was clone-specific; a decrease in cell size under hypoxic conditions was shown by Cold

clone (and less apparently by the Warm clone) in the MH stage and by Int and Warm clones in

the SHT stage (Fig. 3A).

Body size. In total, we measured 1676 individuals. The mean body size of 2679 $\mu m^2 \pm 354$ SD

was similar to the mean of 2411 μm^2 shown for a different *L. inermis* clone across four thermal

regimes (Walczyńska et al., 2015a). We did not find a difference between the experimental

series, which was expected because of the preceding standardization. The body size was affected

by the oxic regime and differed across clones. There was also a significant interaction between

these two effects, and they also interacted significantly with the hypoxia type (Table 4; Fig. 3B).

Fitness measures. The mean population growth rate *r* was 0.64 ± 0.08 SD indiv/day, which was

similar to the value of 0.5 indiv/day that was previously shown at 30°C for a different *L. inermis*

clone (Walczyńska et al., 2016). The mean egg ratio was estimated to be 0.16 ± 0.02 eggs/female.

Both measures of fitness showed similar patterns across experimental regimes. The fitness

differed (or tended to differ, in the case of egg ratio) across hypoxia types and clones (Table 5),

with the lowest values in SHT stage and with the Warm clone showing generally the highest

fitness (Fig. 4). There was a strong lack of difference between rotifers from normoxia or hypoxia

in any measure (Table 5).

Discussion

The most important result from this study was that we generally confirmed the hypotheses that (i) body size decreases in response to hypoxia, (ii) as a consequence of cell size decrease, (iii) preventing the narrowing of aerobic metabolism and (iv) the subsequent decrease in fitness. Strikingly, we observed this causative pattern only for the mildly hypoxic conditions and not for the two severe hypoxia treatments (Fig. 5). In the latter case, the pattern was more complicated and clone-specific. However, in each case, the rotifers originating from the hypoxic conditions displayed a similar level of fitness to those from the normoxic conditions (Fig. 4), which could indicate some alternative mechanisms to the plastic size response that effectively prevented the decrease in fitness under stress-inducing hypoxic conditions. In this study, we confirmed the importance of the 'optimal thermal range' within which the TSR operates (Atkinson et al., 2003; Walczyńska et al., 2016), but we extended this issue by relating the optimality of the conditions to temperature acting in close tandem with oxygen. Additionally, the patterns we obtained concurred with a meta-analysis on 52 species of aquatic invertebrates provided by Galic et al. (2019), who found that (i) the lower the oxygen concentration was, the larger the variation in the measured response traits and that (ii) the respiration was more sensitive than growth to hypoxic conditions. We will now present the results in greater detail.

Proximate mechanisms. Because of the methodological problems, we did not manage to collect the complete dataset for the cell size response. Our partial results show that in the MH treatment, the consequent body size decrease in hypoxia compared to normoxia was associated with a mechanism other than cell shrinkage in the case of one clone. In contrast, the cell and body size differences between normoxic and hypoxic rotifers showed totally opposite patterns in all clones exposed to severe temperature-driven hypoxia. A similar opposing pattern was also found in the case of Warm clone in the SHO2 treatment. The question of whether the variation in body size matches the variation in cell size was classified as fundamental by Leinaas et al. (2016). The suggestions on the differing patterns of this relationship under optimal *vs*. stressful conditions were previously invoked by Walczyńska et al. (2015a) and Walczynska et al. (2018), while the canalization *vs*. plasticity in body *vs*. cell size, which differed across *D. melanogaster*'s organs, was shown by McDonald et al. (2018). In a study in which different life stages of *D. melanogaster* were exposed to hypoxia, the cell size response was found to be stage-specific (Heinrich et al., 2011). All these results showed that the issue of body and cell size adjustment

was not trivial and much can be performed in this field. The novel contribution of this study to

that issue was to show that the body-cell size match was not only due to the condition but was

also clone-dependent and apparently varied with thermal preferences; in a warm-preferring clone,

too high temperature in the SHT treatment caused a decrease in cell size under hypoxia, but this

was not followed by an adequate body size adjustment, while in the cold-preferring clone, the

response was reversed under the same conditions. The intermediate clone showed a similar

pattern to the warm clone in this treatment, but its body size response was opposite (to both

remaining clones) in the SHO2 treatment. Based on these results, we speculated that the specific

cue of plastic response to temperature-dependent hypoxia varied with organismal thermal

preferences.

Ultimate mechanisms. We found that under mild hypoxic conditions (MH treatment), the

rotifers were able to adjust their metabolisms to the oxic conditions in a relatively quick time of

three days (approximately 2 generations), so that the aerobic scope of the rotifers from hypoxic

conditions was on average 30% wider than those of the rotifers from normoxic conditions that

were exposed to acute hypoxia. The rotifers experiencing stressful hypoxic conditions for three

days, either those driven by too high of a temperature (the SHT treatment) or those driven by too

low of an oxygen concentration (SHO2 treatment), displayed similar (Cold clone) or even

narrower (Warm clone) aerobic performances than the hypoxia-naïve rotifers (Fig. 2B, C).

Despite that, all clones performed similarly well (a fitness measure) under normoxia and hypoxia

across the three hypoxia regimes (Fig. 4), though the highest average population growth rate was

under mild hypoxia (approximately 15% higher than in the SHT treatment and approximately 4%

higher than in the SHO2 treatment), followed by the result for the SHO2 treatment

(approximately 1% higher than in the SHT treatment). The decoupling between growth (though

measured at the individual level) and SDA was previously found by Chopelet et al. (2008) in

Daphnia magna clones originating from thermally distinct geographical regions. The authors

concluded that individual growth was not dictated by aerobic capacity measured as SDA. In our

study, the generally worse performance displayed under severe hypoxia driven by too high of a

temperature than under that driven by too low of an oxygen concentration indicated that the cue

for sensing hypoxia was not indifferent for the organisms. In the case of animals that were

tolerant with respect to the oxygenic conditions (Galkovskaya, 1995), the overly elevated

temperatures seemed to be more stressful than too little oxygen. Our rotifers that were exposed to

severe hypoxia apparently switched to alternative physiological mechanisms that we did not examine. However, it was known from previous research that this group of animals was able to switch from the regular low-efficiency and rapid lactate metabolic pathway under hypoxia to a more efficient but slower glucose-succinate metabolic pathway when exposed to severely hypoxic conditions (Esparcia et al., 1992).

Interclonal differences. We summarize the patterns of the different strategies at the interclonal level in Fig. 5:

- Under mild hypoxia, all three clones decreased in body size, and only the Int clone used a different mechanism than cell shrinking. Regardless of the mechanism, all three clones maintained relatively high fitness under normoxia and hypoxia, and in the known cases, this was linked with keeping a wide scope for aerobic metabolism.
- Under severe hypoxia driven by too high of a temperature, each clone showed a different pattern of either decreasing cell size or body size, or, in the case of the Int clone, neither. All three clones maintained relatively high fitness despite the complex and variable body and cell size response. There was also an intriguing difference between the Cold and Warm clones: the former maintained high fitness under hypoxia by widening the aerobic scope but achieved only through body size decrease, without an adequate cell size response, while the latter displayed high performance without the widening the aerobic scope or decrease in body size under hypoxia, but with the clear pattern of cell size response.
- Under severe hypoxia driven by inadequately low oxygen concentrations, two clones decreased in body size, and in the case of the Warm clone, this response was realized without cell shrinking. None of the two clones investigated managed to maintain the wide aerobic scope under hypoxia compared to normoxia, but all three clones maintained relatively high fitness in both oxic regimes.

The reliability of our results was confirmed by the pattern found for the Warm clone, the smallest one, which displayed the highest fitness in the hot experimental conditions. An interesting image resulted from the comparison between the two measures of fitness that we applied. It seemed that in our short study, the clones exposed to stressful conditions invested either in higher fecundity (high egg ratio) or in accelerating development (high population growth rate). In the longer time

scale, the two mechanisms would probably cause similar results in the overall fitness. The

clearest pattern was shown by the Int clone, which had the slowest development (so that we even

did not manage to examine its SDA), but its fecundity was comparable with the Cold clone across

the regimes. The differences in reproductive strategy in response to hypoxia were previously

found across four rotifer species by Kirk et al. (1999).

The role of oxygen deficiency in the TSR. We concur with the conclusions from previous

reports on the diverse response of ectotherms to high temperature combined with hypoxia.

However, we also showed room for TSR in this general perspective; plasticity in size (decrease in

high temperature/hypoxia) occurs within the relatively mild conditions, most likely as the

primary mechanism of maintenance of the wide scope for aerobic metabolism. Above a certain

threshold, which was previously designated by Walczyńska et al. (2016), the response changes;

the plastic response was replaced by physiological mechanisms preventing, or counteracting, any

possible physiological damage. Along with the growing number of empirical studies confirming

the role of oxygen behind the size-to-temperature response, there has also been some criticism.

These doubts are associated with the ambiguous results for higher animals, which have mostly

been fishes (reviewed in Audzijonyte et al., 2019). In the study conducted on *Daphnia magna*

Kielland et al. (2019) confirmed the limiting role of oxygen availability for ectotherms

performance at high temperatures but concluded that plasticity did not counteract the negative

effects of temperature-dependent hypoxia at very high temperatures. In our opinion, the results

we present show that oxygen is the limiting factor (sensu Fry, 1947), inducing the decrease in cell

size and the consequent body size decrease, but this causative relationship should be expected

only under the optimal conditions, which are dependent on many factors, especially organismal

life strategy and thermal preferences. Let us provide an example from everyday life. Imagine that

one wants to buy a house. If there are enough funds for it, one simply buys the house; if there are

not, then there are different potential strategies for gaining the necessary money: taking an

additional job, saving more from the current salary, borrowing from family or friends, or taking a

bank loan. In each case, the result is the same: there is enough money to buy the house. However,

there are always various strategy-dependent costs, which are in each case higher than if there had

been adequate funds at the beginning. The lowest-cost strategy for struggling with mild,

ecologically relevant hypoxia is a plastic response. The possible and differing physiological costs

for organisms struggling with severe oxygen deficiency in their environment await discovery,

although we already know that they are most likely associated with hypoxia sensing at the
mitochondrial level (Sokolova, 2018; Sokolova et al., 2019). A comparison of physiological
responses to mild or severe hypoxia in insects was reviewed by Harrison et al. (2018). The two
approaches, ecological and physiological, should be matched to determine how costly the
response to temperature-dependent oxygen concentrations is, with a distinction between mild and
severe conditions.

Acknowledgments. We are very grateful to Witold Strojny for conducting the
spectrophotometric analyses. The manuscript was edited by American Journal Experts. This work
was supported by the National Science Centre Poland (OPUS 2015/19/B/NZ8/01948) and by the
Jagiellonian University (DS/INoS/757/2019). **The authors declare no conflict of interest.**

Data availability. The data supporting the results will be archived in a public repository of the
Jagiellonian Univeristy and the data DOI will be included at the end of the article. The data will
be made publicly available at the time of publication.

References

Atkinson, D. (1994). Temperature and organism size – a biological law for ectotherms. *Advances*
in Ecological Research **25**, 1-58.
Atkinson, D., Ciotti, B. J. and Montagnes, D. J. (2003). Protists decrease in size linearly with
temperature: ca. 2.5% degrees C(-1). *Proceedings of the Royal Society B* **270**, 2605-11.
Atkinson, D., Morley, S. A. and Hughes, R. N. (2006). From cells to colonies: at what levels of
body organization does the 'temperature-size rule' apply? *Evolution & Development* **8**, 202-214.
Audzijonyte, A., Barneche, D. R., Baudron, A. R., Belmaker, J., Clark, T. D., Marshall, C. T.,
Morrongiello, J. R. and van Rijn, I. (2019). Is oxygen limitation in warming waters a valid mechanism to
explain decreased body sizes in aquatic ectotherms? *Global Ecology and Biogeography* **28**, 64-77.
Belyatskaya, Y. S. (1959). Using the floating microrespirometer for measuring gaseous
interchange in zooplankton (Russian), vol. 7, pp. 315-317: Academy of Sciences, B.S.S.R.
Berner, R. A., VandenBrooks, J. M. and Ward, P. D. (2007). Evolution - Oxygen and evolution.
Science **316**, 557-558.
Berrigan, D. and Charnov, E. L. (1994). Reaction norms for age and size at maturity in response
to temperature – a puzzle for life historians. *Oikos* **70**, 474-478.
Berzins, B. and Pejler, B. (1989). Rotifer occurence in relation to oxygen content *Hydrobiologia*
183, 165-172.
Carpenter, J. H. (1965). The accuracy of the winkler method for dissolved oxygen analysis.
Limnology and Oceanography **10**, 135-140.
Cavalier-Smith, T. (2005). Economy, speed and size matter: Evolutionary forces driving nuclear
genome miniaturization and expansion. *Annals of Botany* **95**, 147-175.
Chopelet, J., Blier, P. U. and Dufresne, F. (2008). Plasticity of growth rate and metabolism in
Daphnia magna populations from different thermal habitats. *Journal of Experimental Zoology A* **309**, 553-
62.

Clark, T. D., Sandblom, E. and Jutfelt, F. (2013). Aerobic scope measurements of fishes in an era
of climate change: respirometry, relevance and recommendations. *Journal of Experimental Biology* **216**,
2771-2782.

Czarnoleski, M., Ejsmont-Karabin, J., Angilletta, M. J., Jr. and Kozlowski, J. (2015). Colder
rotifers grow larger but only in oxygenated waters. *Ecosphere* **6**.

Doohan, M. (1973). An energy budget for adult Brachionus plicatilis Muller (Rotatoria).
Oecologia **13**, 351-362.

Esparcia, A., Serra, M. and Miracle, M. R. (1992). Relationships between oxygen concentration
and patterns of energy metabolism in the rotifer Brachionus plicatilis *Comparative Biochemistry and*
Physiology B-Biochemistry & Molecular Biology **103**, 357-362.

Fiałkowska, E., Klimek, B., Marchlewicz, A., Koverba-Soroka, W., Starzycka, J., Walczyńska, A.
and Pajdak-Stós, A. (2019). Diversity and function of the microbial community under strong selective
pressure of rotifers. *Journal of Basic Microbiology* **59**, 775-783.

Frazier, M. R., Woods, H. A. and Harrison, J. F. (2001). Interactive effects of rearing temperature
and oxygen on the development of Drosophila melanogaster. *Physiological and Biochemical Zoology* **74**,
641-650.

Fry, F. E. J. (1947). Effects of the environment on animal activity. *Publications of the Ontario*
Fisheries Research Laboratory **68**, 1-62.

Galic, N., Hawkins, T. and Forbes, V. E. (2019). Adverse impacts of hypoxia on aquatic
invertebrates: a meta-analysis. *Science of the Total Environment* **652**, 736-743.

Galkovskaya, G. A. (1963). Utilisation of food for growth and conditions for maximum
production of the rotifer Brachionus calyciflorus Pallas (Russian). *Zoologicheskii Zhurnal* **42**, 506-512.

Galkovskaya, G. A. (1995). Oxygen consumption rate in rotifers. *Hydrobiologia* **313**, 147-156.

Gudowska, A. and Bauchinger, U. (2018). Food consumption in ground beetles is limited under
hypoxic conditions in response to ad libitum feeding, but not restricted feeding. *Journal of Insect*
Physiology **107**, 7-13.

Harrison, J. F., Greenlee, K. J. and Verberk, W. (2018). Functional Hypoxia in Insects: Definition,
Assessment, and Consequences for Physiology, Ecology, and Evolution. In *Annual Review of Entomology,*
Vol 63, vol. 63 (ed. M. R. Berenbaum), pp. 303-325.

Harrison, J. F., Kaiser, A. and VandenBrooks, J. M. (2010). Atmospheric oxygen level and the
evolution of insect body size. *Proceedings of the Royal Society B* **277**, 1937-46.

Heinrich, E. C., Farzin, M., Klok, C. J. and Harrison, J. F. (2011). The effect of developmental
stage on the sensitivity of cell and body size to hypoxia in Drosophila melanogaster. *Journal of*
Experimental Biology **214**, 1419-1427.

Helm, I., Jalukse, L. and Leito, I. (2012). A highly accurate method for determination of dissolved
oxygen: Gravimetric Winkler method. *Analytica Chimica Acta* **741**, 21-31.

Hirata, H. and Yamasaki, S. (1987). Effect of feeding on the respiration rate of the rotifer
Brachionus plicatilis. *Hydrobiologia* **147**, 283-288.

Hoefnagel, K. N. and Verberk, W. (2015). Is the temperature-size rule mediated by oxygen in
aquatic ectotherms? *Journal of Thermal Biology* **54**, 56-65.

Hrycik, A. R., Almeida, L. Z. and Hook, T. O. (2017). Sub-lethal effects on fish provide insight into
a biologically-relevant threshold of hypoxia. *Oikos* **126**, 307-317.

Jalal, M., Wojewodzic, M. W., Laane, C. M. M. and Hessen, D. O. (2013). Larger Daphnia at
lower temperature: a role for cell size and genome configuration? *Genome* **56**, 512-520.

Jordan, A. D. and Steffensen, J. F. (2007). Effects of ration size and hypoxia on specific dynamic
action in the cod. *Physiological and Biochemical Zoology* **80**, 178-185.

Kielland, O. N., Bech, C. and Einum, S. (2019). Warm and out of breath: thermal phenotypic
plasticity in oxygen supply. *Functional Ecology* **33**, 2142-2149.

Kiełbasa, A., Walczyńska, A., Fiałkowska, E., Pajdak-Stós, A. and Kozłowski, J. (2014). Seasonal
changes in the body size of two rotifer species living in activated sludge follow the Temperature-Size
Rule. *Ecology and Evolution* **4**, 4678-4689.
Kirk, K. L., Ellis, J. and Taylor, J. (1999). Physiological responses to variable environments:
storage and respiration in starving rotifers. *Freshwater Biology* **42**, 637-644.
Kozlowski, J., Konarzewski, M. and Gawelczyk, A. T. (2003). Cell size as a link between
noncoding DNA and metabolic rate scaling. *Proc Natl Acad Sci U S A* **100**, 14080-5.
Leinaas, H. P., Jalal, M., Gabrielsen, T. M. and Hessen, D. O. (2016). Inter- and intraspecific
variation in body- and genome size in calanoid copepods from temperate and arctic waters. *Ecology and*
Evolution **6**, 5585-5595.
McCue, M. D. (2006). Specific dynamic action: A century of investigation. *Comparative*
Biochemistry and Physiology a-Molecular & Integrative Physiology **144**, 381-394.
McDonald, J. M. C., Ghosh, S. M., Gascoigne, S. J. L. and Shingleton, A. W. (2018). Plasticity
Through Canalization: The Contrasting Effect of Temperature on Trait Size and Growth in Drosophila.
Frontiers in Cell and Developmental Biology **6**.
Pajdak-Stós, A., Fiałkowska, E., Fyda, J., Kocerba-Soroka, W. and Sobczyk, M. (2017). A method
of mass culture of Lecane rotifers.
Poertner, H. O. (2010). Oxygen- and capacity-limitation of thermal tolerance: a matrix for
integrating climate-related stressor effects in marine ecosystems. *Journal of Experimental Biology* **213**,
881-893.
Pourriot, R. and Deluzarches, M. (1970). Sur la consommation d'oxygene par les Rotifers.
Annales de Limnologie **6**, 229-248.
Prokosch, J., Bernitz, Z., Bernitz, H., Erni, B. and Altwegg, R. (2019). Are animals shrinking due to
climate change? Temperature-mediated selection on body mass in mountain wagtails. *Oecologia* **189**,
841-849.
Raubenheimer, D. (1995). Problems with ratio analysis in nutritional indices. *Functional Ecology*
9, 21-29.
Raubenheimer, D. and Simpson, S. J. (1992). Analysis of covariance: an alternative to nutritional
indices. *Entomologia Experimentalis et Applicata* **62**, 221-231.
Roland, F., Caraco, N. F. and Cole, J. J. (1999). Rapid and precise determination of dissolved
oxygen by spectrophotometry: Evaluation of interference from color and turbidity. *Limnology and*
Oceanography **44**, 1148-1154.
Rollinson, N. and Rowe, L. (2018a). Oxygen Limitation at the Larval Stage and the Evolution of
Maternal Investment per Offspring in Aquatic Environments. *American Naturalist* **191**, 604-619.
Rollinson, N. and Rowe, L. (2018b). Temperature-dependent oxygen limitation and the rise of
Bergmann's rule in species with aquatic respiration. *Evolution* **72**, 977-988.
Santilli, J. and Rollinson, N. (2018). Toward a general explanation for latitudinal clines in body
size among chelonians. *Biological Journal of the Linnean Society* **124**, 381-393.
SAS. (2013). SAS/STAT User's Guide: Cary NC, SAS Institute Inc.
Secor, S. M. (2009). Specific dynamic action: a review of the postprandial metabolic response.
Journal of Comparative Physiology B-Biochemical Systemic and Environmental Physiology **179**, 1-56.
Sokolova, I. (2018). Mitochondrial Adaptations to Variable Environments and Their Role in
Animals' Stress Tolerance. *Integrative and Comparative Biology* **58**, 519-531.
Sokolova, I. M., Sokolov, E. P. and Haider, F. (2019). Mitochondrial mechanisms underlying
tolerance to fluctuating oxygen conditions: lessons from hypoxia-tolerant organisms. *Integrative and*
Comparative Biology **59**, 938-952.

Stuczyńska, A., Sobczyk, M., Fiałkowska, E., Kocerba-Soroka, W., Pajdak-Stós, A., Starzycka, J.
and Walczyńska, A. (2020). Clonal thermal preferences affect the strength of the temperature-size rule.
bioRxiv.
Verberk, W. C. E. P. and Atkinson, D. (2013). Why polar gigantism and Palaeozoic gigantism are
not equivalent: effects of oxygen and temperature on the body size of ectotherms. *Functional Ecology*
27, 1275-1285.
Verberk, W. C. E. P., Bilton, D. T., Calosi, P. and Spicer, J. I. (2011). Oxygen supply in aquatic
ectotherms: Partial pressure and solubility together explain biodiversity and size patterns. *Ecology* **92**,
1565-1572.
Vinberg, G. G. (1937). Respiratory intensity in the Rotatoria Anuraea aculeata Ehrbg (Russian).
Bulletin of Experimental Biology and Medicine **4**, 464-466.
Walczynska, A., Labecka, A. M. and Sobczyk, M. (2018). What may a fussy creature reveal about
body/cell size integration under stressful conditions? *Development Genes and Evolution* **228**, 179-188.
Walczyńska, A., Kiełbasa, A. and Sobczyk, M. (2016). 'Optimal thermal range' in ectotherms:
defining criteria for tests of the temperature-size rule. *Journal of Thermal Biology* **60**, 41-48.
Walczyńska, A., Labecka, A. M., Sobczyk, M., Czarnoleski, M. and Kozłowski, J. (2015a). The
Temperature-Size Rule in Lecane inermis (Rotifera) is adaptive and driven by nuclei size adjustment to
temperature. *Journal of Thermal Biology* **54**, 78-85.
Walczyńska, A. and Serra, M. (2014). Inter- and intraspecific relationships between performance
and temperature in a cryptic species complex of the rotifer Brachionus plicatilis. *Hydrobiologia* **734**, 17-
26.
Walczyńska, A., Sobczyk, M., Czarnoleski, M. and Kozłowski, J. (2015b). The temperature-size
rule in a rotifer is determined by the mother and at the egg stage. *Evolutionary Ecology* **29**, 525-536.
Walczyńska, A., Sobczyk, M., Fialkowska, E., Pajdak-Stós, A., Fyda, J. and Wiąckowski, K.
(2017). Interaction between a bacterivorous ciliate Aspidisca cicada and a rotifer Lecane inermis:
Doozers and Fraggles in aquatic flocs. *Microbial Ecology* **75**, 569-581.
Wetzel, R. G. (2001). Limnology. Lake and river ecosystems. San Diego, San Francisco, New York,
Boston, London, Sydney, Tokyo: Elsevier Academic Press.
Woods, H. A. (1999). Egg-mass size cell size: effects of temperature on oxygen distribution.
American Zoologist **39**, 244-252.

643 **Table 1.** Comparison of the absolute oxygen consumption measured under different conditions in
644 different rotifer species.

Rotifer species	O$_2$ cons. [mg×indiv^{-1}×h^{-1}]	O$_2$ cons. [mL×indiv^{-1}×h^{-1}]	Measurements conditions	References
Asplanchna priodonta		0.48×10^{-5} and 0.71×10^{-5}	20°C, well-fed rotifers and after 28 h starvation, respectively	Kirk et al. (1999)
Asplanchna silvestrii		1.65×10^{-5} and 1.54×10^{-5}	20°C, well-fed rotifers and after 57 h starvation, respectively	Kirk et al. (1999)
Brachionus calyciflorus		$0.16\text{-}0.62 \times 10^{-5}$	Large numbers, unknown age distribution, different conditions	Pourriot and Deluzarches (1970), after Doohan (1973)
Brachionus calyciflorus		0.22×10^{-5}	-	Belyatskaya (1959), after Doohan (1973)
Brachionus calyciflorus		0.5×10^{-5}	-	Galkovskaya (1963) after Doohan (1973)
Brachionus calyciflorus		0.31×10^{-5} and 0.13×10^{-5}	20°C, well-fed rotifers and after 53 h starvation, respectively	Kirk et al. (1999)
Brachionus plicatilis		0.27×10^{-5}	female without eggs; 20°C, sea water with no food	Doohan (1973)
Brachionus plicatilis		$0.1\text{-}0.7 \times 10^{-5}$	20‰ salinity, 20°C	Hirata and Yamasaki (1987)
Keratella quadrata	0.08×10^{-5} and 0.11×10^{-5}		14.5-18°C and 20°C, respectively	Vinberg (1937), after Galkovskaya (1995)
Lecane inermis, **two clones**	0.09×10^{-5} and 0.38×10^{-5}	0.07×10^{-5} and 0.29×10^{-5}	30°C, 2 h starved and fed, respectively	this study
Polyarthra dolichoptera		0.09×10^{-5} and 0.027×10^{-5}	2 and 5°C, respectively	Galkovskaya (1995)
Synchaeta pectinata		0.21×10^{-5} and 0.22×10^{-5}	20°C, well-fed rotifers and after 18 h starvation, respectively	Kirk et al. (1999)

Table 2. Final results of the GLM model for oxygen consumption in two *L. inermis* clones.

Effect	F-value (df num, df den)	P-value
hypoxia type	(2, 91) 38.81	< 0.0001
clone	(1, 91) 7.69	0.0067
oxic regime	(1, 91) 100.88	< 0.0001
food regime	(1, 91) 643.58	< 0.0001
h. type × clone	(2, 91) 20.15	< 0.0001
h. type × oxic regime	(2, 91) 0.63	0.5371
clone × oxic regime	(1, 91) 0.81	0.3698
h. type × food regime	(2, 91) 4.81	0.0104
clone × food regime	(1, 91) 2.59	0.1108
oxic regime × food regime	(1, 91) 0.01	0.9034
h. type × clone × food regime	(2, 91) 8.21	0.0005
h. type × oxic regime × food regime	(2, 91) 3.88	0.0240

646   hypoxia type – subsequent experimental stages, standardized within each stage to the SD units

647   oxic regime – the origin of the animals (rotifers could originate from the normoxic or hypoxic experimental

648   conditions, but their respiration was always measured under hypoxia)

649   food regime – fed or hungry rotifers

658   **Table 3.** Final results of the GLM model for the nuclei size (a proxy for cell size) in two *L.*

659   *inermis* clones.

Effect	F-value (df num, df den)	P-value
hypoxia type	(2, 126) 4.39	0.0144
clone	(2, 126) 27.04	< 0.0001
oxic regime	(1, 126) 1.98	0.1615
h. type × clone	(2, 126) 7.98	0.0005
h. type × oxic regime	(2, 126) 7.03	0.0013
clone × oxic regime	(2, 126) 1.29	0.2789
h. type × clone × oxic regime	(4, 126) 14.72	< 0.0001

660   hypoxia type – subsequent experimental stages, standardized within each stage to the SD units

661   oxic regime – the origin of the animals (rotifers could originate from the normoxic or hypoxic experimental

662   conditions, but their respiration was always measured under hypoxia)

675 **Table 4.** Final results of the GLM model for body size standardized to SD units among the

676 hypoxia types.

Effect	F-value (df num, df den)	P-value
hypoxia type	(2, 1662) 1.09	0.3370
clone	(2, 1662) 30.23	< 0.0001
oxic regime	(1, 1662) 4.76	0.0292
h. type × clone	(4, 1662) 18.26	< 0.0001
h. type × oxic regime	(2, 1662) 9.00	0.0001
clone × oxic regime	(2, 1662) 6.37	0.0018

677 hypoxia type – subsequent experimental stages, standardized within each stage to the SD units

678 oxic regime – the origin of the animals (rotifers could originate from the normoxic or hypoxic experimental

679 conditions, but their respiration was always measured under hypoxia)

693   **Table 5.** Final results of the GLM ANCOVA models for the two fitness measures, population

694   growth rate r and egg ratio.

Effect	F-value (df num, df den)	P-value
population growth rate r: rotifers number as a dependent variable		
hypoxia type	(2, 65) 9.13	0.0003
clone	(2, 65) 12.98	< 0.0001
oxic regime	(1, 65) 0.62	0.4340
initial number	(1, 65) 11.30	0.0013
egg ratio: eggs number as a dependent variable		
hypoxia type	(2, 65) 3.12	0.0507
clone	(2, 65) 2.99	0.0572
oxic regime	(1, 65) 0.20	0.6587
final number	(1, 65) 50.81	< 0.0001

695   hypoxia type – subsequent experimental stages, standardized within each stage to the SD units

696   oxic regime – the origin of the animals (rotifers could originate from the normoxic or hypoxic experimental

697   conditions, but their respiration was always measured under hypoxia)

698   the initial and final numbers – the rotifers counted in count I and count II, respectively

Fig. 1. Concept of the study. A – hypothetical effect of hypoxia on the traits estimated: hypoxia causes cell size and body size decrease (proximate mechanisms) to maintain the wide aerobic scope to prevent the possible fitness decline (ultimate mechanisms). B – Experimental setup: the study was divided into three subsequent stages with different hypoxic conditions; five traits were investigated within three days in each stage. SDA – specific dynamic action (a proxy for aerobic scope); MH stage – hypoxia type = mild hypoxia; SHT stage – hypoxia type = severe hypoxia with high temperature; SHO2 stage – hypoxia type = severe hypoxia with low oxygen concentration.

Fig. 2. Results for oxygen consumption (in %), absolute aerobic scope and factorial aerobic
scope for warm and cold clones across three regimes of different hypoxia types. A – oxygen
consumption for rotifers exposed to experimental hypoxia or normoxia, then divided into groups
of fed or hungry and tested for respiration rates under hypoxic conditions (4 or 5 replicates per
group tested). Least square means ± s.e.. B – absolute scope for aerobic metabolism as a
difference between the oxygen consumption of fed and hungry rotifers; absolute SDA. C –
factorial scope for aerobic metabolism as a division of the amount of oxygen consumed by the
fed rotifers to that consumed by the hungry rotifers, factorial SDA.

Fig. 3. Results for cell- and body-size response. The interactions among hypoxia type (MH –

mild hypoxia; SHT – severe hypoxia driven by too high of a temperature; SHO2 – severe

hypoxia driven by too low of an oxygen concentration), oxic regime (N – normoxia; H –

hypoxia) and clone in the GLM analysis for nuclei size (a proxy for cell size) (A), and body size

(B). The numbers denote the sample size of rotifers the nuclei of which were measured (A), or

the number of measured individuals (B). Least Square means ± 0.95 c.i.

Fig. 4. Results for the fitness measures. The interactions among the hypoxia types (MH – mild hypoxia; SHT – severe hypoxia driven by too high temperature; SHO2 – severe hypoxia driven by inadequately low oxygen concentrations), oxic regime (N – normoxia; H – hypoxia) and clone in the ANCOVA analyses for two fitness measures: population growth rate *r*, with count II number as a dependent variable and count I as a covariate (A), and egg ratio, with egg number as a dependent variable and count II as a covariate (B). N = 4 replicates. Least Square means ± 0.95 c.i.

Fig. 5. Schematic summary of the patterns found as referred to the hypotheses displayed in

Fig. 1A. The arrows (red – warm-preferring clone, green – medium temperature-preferring clone,

blue – cold-preferring clone) represent the positive or (when crossed) negative causative effects

we found while testing across the three experimental stages. Dashed arrows indicate the effect

that we found without the intermediate impact we expected. The lack of specific arrow indicates

that no result was achieved for a given relationship. The dots show the positive or (when crossed)

negative ultimate results in the cases when the causative effect through the intermediate trait was

not known.

Fig. S1. The analysis of raw absorbance data for the reliability of controls. The control samples

perceived as unreliable are crossed out and were not included in the analyses. Absorbance is a

measure proportional to the amount of oxygen dissolved in a sample. MH – mild hypoxia

treatment, SHT – severe hypoxia treatment, driven by elevated temperature, SHO2 – severe

hypoxia treatment driven by alleviated oxygen concentration.