Title

Repression by the *Arabidopsis* TOPLESS corepressor requires association with the core

Mediator complex

Authors

Alexander R. Leydon[1], Wei Wang[2,†], Hardik P. Gala[1], Sabrina Gilmour[1], Samuel Juarez-Solis[1],

Mollye L. Zahler[1], Joseph E. Zemke[1], Ning Zheng[2,3], Jennifer L. Nemhauser[1*]

[1]Department of Biology, University of Washington

[2]Department of Pharmacology & [3]Howard Hughes Medical Institute, University of Washington

[†]Present address: Key Laboratory of Plant Stress Biology, State Key Laboratory of Cotton

Biology, School of Life Science, Jinming Campus, Henan University, Kaifeng, Henan Province,

475004, PR of China.

*Lead Contact.

Abstract

The plant corepressor TOPLESS (TPL) is recruited to a large number of loci that are selectively

induced in response to developmental or environmental cues, yet the mechanisms by which it

inhibits expression in the absence of these stimuli is poorly understood. Previously, we had

used the N-terminus of *Arabidopsis thaliana* TPL to enable repression of a synthetic auxin

response circuit in *Saccharomyces cerevisiae* (yeast). Here, we leveraged the yeast system to

interrogate the relationship between TPL structure and function, specifically scanning for

repression domains. We identified a potent repression domain in Helix 8 located within the CRA

domain, which directly interacted with the Mediator middle domain subunits Med21 and Med10.

Interactions between TPL and Mediator were required to fully repress transcription in both yeast

and plants. In contrast, we found that multimer formation, a conserved feature of many

corepressors, had minimal influence on the repression strength of TPL.

Keywords

Corepressors, Transcriptional Repression, Mediator, Tup1, TOPLESS

Introduction

Control over gene expression is essential for life. This is especially evident during development

when the switching of genes between active and repressed states drives fate determination.

Mutations that interfere with repression lead to or exacerbate numerous cancers (Wong et al.,

2014) and cause developmental defects in diverse organisms (Grbavec et al., 1998; Long et al.,

2006), yet many questions remain about how cells induce, maintain, and relieve transcriptional

repression. Transcriptional repression is controlled in part by a class of proteins known as

corepressors that interact with DNA-binding transcription factors and actively recruit repressive

machinery. Transcriptional corepressors are found in all eukaryotes and include the animal

SMRT (silencing mediator of retinoic acid and thyroid hormone receptor) and NCoR (nuclear

receptor corepressor) complexes (Mottis et al., 2013; Oberoi et al., 2011), the yeast Tup1

(Keleher et al., 1992; Matsumura et al., 2012; Tzamarias and Struhl, 1994), and its homologs

Drosophila Groucho (Gro) and mammalian transducing-like enhancer (TLE) (Agarwal et al.,

2015).

In plants, the role of Gro/TLE-type corepressors is played by TOPLESS (TPL),

TOPLESS-RELATED (TPR1-4), LEUNIG (LUG) and its homolog (LUH), and High Expression of

Osmotically responsive genes 15 (HOS15) (Causier et al., 2012; Lee and Golz, 2012; Liu and

Karmarkar, 2008; Long et al., 2006; Zhu et al., 2008). Plant corepressors share a general

structure, where at the N-terminus a LIS1 homology (LisH) domain contributes to protein

dimerization (Delto et al., 2015; Kim et al., 2004). At the C-terminus, WD40 repeats form beta-

propeller structures that are involved in protein-protein interactions (Collins et al., 2019; Liu et

al., 2019). In TPL family corepressors, the LisH is followed by a C-terminal to LisH (CTLH)

domain that binds transcriptional repressors through an Ethylene-responsive element binding

factor-associated Amphiphilic Repression (EAR) motif found in partner proteins (Causier et al.,

2012; Kagale et al., 2010). The N-terminal domain also contains a CT11-RanBPM (CRA)

domain, which provides a second TPL dimerization interface and stabilizes the LisH domain (Ke

et al., 2015; Martin-Arevalillo et al., 2017). The beta-propellers bind to the non-EAR TPL

recruitment motifs found in a subset of transcriptional regulators (RLFGV- and DLN-type motifs,

(Collins et al., 2019; Liu et al., 2019)), and may control protein interaction with other repressive

machinery. Defects in the TPL family have been linked to aberrant stem cell homeostasis

(Busch et al., 2010), organ development (Gonzalez et al., 2015), and hormone signaling

(Causier et al., 2012; Kagale et al., 2010), especially the plant hormone auxin (Long et al.,

2006).

We have previously demonstrated the recapitulation of the auxin response pathway in

Saccharomyces cerevisiae (yeast) by porting individual components of the *Arabidopsis* auxin

nuclear response (Pierre-Jerome et al., 2014). In this <u>A</u>rabidopsis <u>t</u>haliana <u>A</u>uxin <u>R</u>esponse

<u>C</u>ircuit in <u>S</u>accharomyces <u>c</u>erevisiae (*At*ARC[Sc] - Figure 1A), an auxin responsive transcription

factor (ARF) binds to a promoter driving a fluorescent reporter. In the absence of auxin, the ARF

protein activity is repressed by interaction with a full-length Aux/IAA protein fused to the N-

terminal domain of TPL. Upon the addition of auxin, the TPL-IAA fusion protein is targeted for

degradation through interaction with a member of the Auxin Signaling F-box protein family, and

releases the transcriptional repression of the fluorescent reporter. Reporter activation can be

quantified after auxin addition by microscopy or flow cytometry (Pierre-Jerome et al., 2014). In

the original build and characterization of *At*ARC[Sc], it was noted that the two N-terminal

truncations of TPL (N100 or N300) behave differently (Pierre-Jerome et al., 2014). While both

truncations are able to repress the function of a transcriptional activator fused to an Aux/IAA,

only the TPLN100 fusion shows alleviation of repression after auxin addition. TPLN300 fusions

to Aux/IAAs maintain strong durable repression even under high concentrations of auxin. This

disparity is not due to differential rates of protein degradation, as both proteins appear to be

turned over with equal efficiency after auxin addition (Pierre-Jerome et al., 2014).

The conservation of TPL's repressive function in yeast suggests that the protein partners

that enact repression are conserved across eukaryotes. Consistent with this speculation, the

series of alpha-helices that form the N-terminal portion of TPL (Figure 1B) (Martin-Arevalillo et

al., 2017) is highly reminiscent of naturally-occurring truncated forms of mammalian TLE

(Gasperowicz and Otto, 2005), such as Amino-terminal Enhancer of Split (AES) (Zhang et al.,

2010), the Groucho ortholog LSY-22 (Flowers et al., 2010), and the unrelated mouse repressor

protein MXI1 (Schreiber-Agus et al., 1995). Gro/TLE family members are generally considered

to repress by recruiting histone deacetylases to control chromatin compaction and availability for

transcription (Chen and Courey, 2000; Long et al., 2006). An alternative hypothesis has been

described for Tup1 in yeast, where Tup1 blocks the recruitment of RNA Polymerase II (Pol-II)

(Wong and Struhl, 2011), possibly through contacts with Mediator complex subunits Med21 or

Med3 (Gromöller and Lehming, 2000; Papamichos-Chronakis et al., 2000). However, like many

of these family members, multiple repression mechanisms have been described for TPL at

different genetic loci. For example, TPL has been found to recruit the repressive CDK8 Mediator

complex (Ito et al., 2016), chromatin remodeling enzymes such as Histone Deacetylase 19

(HD19) (Long et al., 2006) and directly bind to histone proteins (Ma et al., 2017).

Here, we leveraged the power of yeast genetics to interrogate the mechanism of TPL

repression. Using AtARC^{Sc}, we discovered that the N-terminal domain of TPL contains two

distinct repression domains that can act independently. We mapped the first, weaker repression

domain to the first 18 amino acids of the LisH domain (Helix 1), and the second, more potent

domain to Helix 8 which falls within the CRA domain. Full repression by Helix 8 required the

Mediator complex, specifically direct interaction with Med21 and Med10. This interaction

required the same Med21 residues that control transcriptional activation of Tup1-regulated

genes in yeast. In addition, we found that multimerization of TPL was not required for repression

in yeast or in plants. Our yeast results were validated with plant assays, and extended to include

evidence that interaction with the middle domain of Mediator was required for TPL repression of

the auxin-regulated development program giving rise to lateral roots. Our findings point to a

conserved functional connection between Tup1/TPL corepressors and the Mediator complex

that together create a repressed state poised for rapid activation.

Results

To understand how TPL represses transcription, we first localized repressive activity within the

protein using the AtARC^{Sc} (Figure 1A). The extent of auxin-induced turnover of TPLN100 and

TPLN300 fusion proteins appears similar, although neither are completely degraded (Pierre-

Jerome et al., 2014). In this way, auxin sensitizes the AtARC^{Sc} to even subtle differences in the

strength of repressive activity by reducing the relative concentration of the TPL fusion proteins.

To pinpoint the region conferring the strong repression of TPL300, we generated a deletion

series of the N-terminus guided by the available structural information (Figure 1B-C, (Ke et al.,

2015; Martin-Arevalillo et al., 2017)).

We started by identifying a shorter truncation, TPLN188, which behaved identically to

TPLN300 (Figure 1D, (Pierre-Jerome et al., 2014)). Subsequently, we deleted each alpha

helical domain starting with Helix 9 (constructs are named in the format Helix x – Helix y or Hx-

Hy). We found that Helix 8 was required for the maximum level of repression activity and for the

maintenance of repression after auxin addition (Figure 1D). All constructs lacking Helix 8

retained the ability to repress transcription, but this repression was lifted in the presence of

auxin (Figure 1D) as had been observed for the original TPLN100 construct (Pierre-Jerome et

al., 2014). In addition to the repressive activity of Helix 8, further deletions revealed that the 18

amino acids of Helix 1 were sufficient to confer repression on their own (H1, Figure 1E). To test

whether Helix 8 activity depended on Helix 1, we analyzed a construct consisting of Helix 3

through Helix 9 (H3-H9, Figure 1D), which was able to repress transcription. Thus, Helix 1

(LisH) and Helix 8 (CRA) could contribute to TPL-mediated repression on their own (Figure 1D).

To identify the minimal domain required for Helix 8-based repression, we generated

additional deletions (Figure 1C,F, Figure 1 - Figure Supplement 1). Helix 8 and the following

linker were not sufficient for repression (Figure 1F), and removal of Helix 9 or of the linker

between Helix 8 and Helix 9 slightly increased sensitivity to auxin compared to TPLN188 (H1-

H8Δ8L, Figure 1 - Figure Supplement 1). A deletion that removed both the LisH and Helix 8

repression domains (H3-H7) was only able to weakly repress reporter expression (Figure 1 -

Figure Supplement 1). These results demonstrate that Helix 8, in combination with the linker

between Helix 8 and Helix 9 (which folds over Helix 1), was required for repression following

addition of auxin. Moreover, the repressive activity of Helix 8 and the linker were only functional

in the context of the larger Helix 3-Helix 8 truncation that carries the CTLH domain and a portion

of the CRA domain.

Figure 1. The N-terminal domain of TPL contains two independent repression domains.

A. Schematic of the ARC[Sc]. The auxin responsive promoter driving the fluorescent protein

Venus carries binding sites for the ARF transcription factor. In the absence of Auxin the IAA-

TPL fusion protein is bound to the ARF and maintains the circuit in a repressed state. Upon

addition of auxin the IAA-TPL protein is targeted for ubiquitination and subsequent protein

degradation, activating transcription of the fluorescent reporter. **B.** TPL domains are LisH (LIS1

homology motif, blue), CTLH (C-terminal LisH motif, orange), CRA (CT11-RanBPM, red -

dimerization and green - foldback), and two WD40, beta-propeller motifs (purple). N-terminal

domains are indicated on the solved structure of the first 202 amino acids ((Martin-Arevalillo et

al., 2017), 5NQS). The termini of the TPL-N100 truncation used in the original ARC^{Sc} studies is

indicated. **C.** Diagram indicating the structure of constructs analyzed in experiments shown in

subsequent panels. Repression Index (Rep.) is a scaled measure of repression strength with 0

set to the level of repression observed with IAA3 and 10 set to the level of repression by

TPLN188-IAA3. Auxin induction level (Aux. ind.) indicates the fold change difference between

reporter expression before auxin addition (time zero) and at the end of an experiment (~500

minutes) **D-F.** Helix 1 and the CRA domain (Helix 3-Helix 8) can act independently to repress

transcription. Each panel represents two independent time course flow cytometry experiments

of the TPL helices indicated, all fused to IAA3. Every point represents the average fluorescence

of 5-10,000 individually measured yeast cells (a.u. - arbitrary units). Auxin (IAA-10µM) was

added at the indicated time (gray bar, + Aux).

To determine which of the many known or predicted TPL-binding partners could mediate

the repression activity of Helix8, we identified known interactors with either TPL or other

Gro/TLE co-repressors, and then introduced the *Arabidopsis* homologs of these genes into the

cytoplasmic split-ubiquitin (cytoSUS) system (Asseck and Grefen, 2018). We chose the

cytoSUS system over yeast two-hybrid, because in cytoSUS the interaction between target

proteins takes place in the cytoplasm, and we had observed that the TPL N-terminus could

repress activation of yeast two hybrid prototrophy reporters (Figure 1D, Figure 2 – Figure

Supplement 1A). Putative direct interactors include histone deacetylases (HDACs - AtHDAC9,

AtHDAC6, (Long et al., 2006)), Histone proteins (Histone H3, Histone H4, (Ma et al., 2017)),

and the Mediator components MED13 (AtMED13, (Ito et al., 2016)) and MED21, which has

been demonstrated to interact with Tup1, the yeast homolog of TPL (Gromöller and Lehming,

2000). We did not observe any interactions between TPL-N188 and the HDACs HDA6 and

HDA9; the histone protein AtHIS4; or the Mediator subunit AtMED13 (Figure 2A, Figure 2 –

Figure Supplement 1B). HDAC interaction with TPL has been previously hypothesized to occur

through indirect interactions with partner proteins (Krogan et al., 2012), however direct

interactions with histones and MED13 have been detected (Ito et al., 2016; Ma et al., 2017). The

absence of interaction between TPL-N188 and these proteins may be due to differences

between methods, or interaction interfaces in the C-terminal WD40 repeats.

Strong interaction was detected between TPL-N188 and AtMED21, a component of the

Mediator middle domain (Figure 2A). MED21 is one of the most highly conserved Mediator

subunits (Bourbon, 2008), and has a particularly highly conserved N-terminus (Figure 2 – Figure

Supplement 2A,C-E). In yeast, Tup1 interacts with the first 31 amino acids of ScMed21, with the

first seven amino acids being absolutely required (Gromöller and Lehming, 2000). We observed

that the equivalent truncation of AtMED21 (AtMED21-N31) was sufficient for interaction with

TPL-N188 (Figure 2A). We next created truncations of the N-terminal domain of AtMED21 to

closely match those that had been made in yeast (Figure 2B, Figure 2 – Figure Supplement 1B)

where deletion of the first five amino acids of ScMed21 (ScΔ5Med21) severely reduce the ability

of the Mediator complex to co-purify with Pol-II and CDK8 kinase complex (Sato et al., 2016).

Interaction between TPLN188 and AtMED21 similarly required the first five amino acids of

AtMED21 (Figure 2B), and, as in yeast, this was not a result of destabilization of the AtMED21

protein (Figure 2B). AtMED21 interaction was specific to the Helix8-based repression domain,

as it interacted with TPLH3-H9 (Figure 2E), and not TPLH1-H5 (Figure 2 – Figure Supplement

1C). Further screening of middle domain Mediator components identified an additional

interaction with AtMED10B, the predominantly expressed MED10 isoform in *Arabidopsis*

(Klepikova et al., 2016) (Figure 2C). Interactions between TPLN188 and both AtMED21 and

AtMED10B were confirmed by immunoprecipitation (Figure 2D).

Figure 2. The Helix 8 repression domain of TPL directly interacts with AtMED21 and

AtMED10B. A-C,E. cytoSUS assays with candidate interacting proteins. Nub-3xHA is the N-

terminal fragment of Ubiquitin expressed with no fusion protein and is used as a negative

control. Each prey protein is from *Arabidopsis*. -WL, -ADE - dropout lacking Trp, Leu and Ade

(growth control), -WLMH, -ADE – dropout lacking Trp, Leu, His, Met, and Ade (selective media).

The plating for each panel was performed at the same day, white lines are provided when plates

were cropped for clarity. **B**. Alignments of the *Arabidopsis* (At) and *Saccharomyces* (Sc) MED21

proteins are shown above cytoSUS assays with the same bait shown in (**A**). Western blots

below the colonies indicated that AtMED21 N-terminal Δ3 and Δ5 are well expressed in assay

conditions. **C.** cytoSUS assays with selected Mediator proteins in the middle module. **D.** The

TPL-ProteinA-TF fusion protein can pull down TPL, AtMED21 and AtMED10B from yeast

extracts using IgG-beads. Detection of the VP16 transcriptional activator demonstrates

enrichment of the fusion protein (αVP16). Each prey protein is detected via the 3xHA tag (αHA),

and efficacy of purification was judged by PGK1 depletion (αPGK1). **E.** A TPL-N truncation

lacking the LisH domain (TPLH2-H9) could still interact with the AtMED21-N31 truncation. This

bait construct interacted with IAA3, but only minimally with the negative control (free Nub-3xHA).

F. Yeast Mediator (bottom, 5N9J) and AtTPL (top, 5NQV) manually juxtaposed to compare

relative domain sizes and feasibility of a TPL-MED21-MED10B interaction. TPL Helix 8-9 is

colored green. MED21 is colored aqua, with the N-terminus colored red, and the EAR peptide in

orange. MED10 is colored teal, with the C-terminus colored purple. The dotted line indicates the

border between TPL and Mediator structures.

A manual juxtaposition of the yeast Mediator structure with the *Arabidopsis* TPL N-

terminal structure shows that Helix 8 and the linker following face away from the tetramer and

are therefore optimally placed to interact with Mediator components (Figure 2F). To pinpoint

which residues of Helix 8 coordinate repression through interaction with MED21, we identified

solution-facing amino acids (Martin-Arevalillo et al., 2017) (Figure 1C). We reasoned that such

residues were most likely not involved in stabilizing the hydrophobic interactions between intra-

TPL helical domains and could therefore be available to interact with partner proteins. Eight

amino acids in Helix 8 were mutated to alanine in the context of the H3-H8-IAA3 fusion protein

(Figure 3A, light green residues) to enable assessment of repression activity in the absence

(Figure 3B) or presence (Figure 3C) of auxin. No single amino acid was essential for repression.

Two mutations (R140A and K148A) slightly increased baseline expression of the reporter

(Figure 3B-C). With the exception of E152A, which behaved similarly to controls, all of the

mutations altered the stability of repression after auxin addition, either by increasing (S138A,

V145A, E146A, K149A) or decreasing (I142A) the final fluorescence level (Figure 3C). Mutating

E146 and K149 also increased the speed with which the reporter responded to auxin (Figure

3C), suggesting that these two neighboring residues could be a critical point of contact with co-

repressive machinery. S138A had a small increase in auxin sensitivity, while I142 reduced auxin

sensitivity (Figure 3E).

We next tested whether the residues in Helix 8 that were required for repression (V145,

E146, K148, K149, Figure 3A-C) were also required for interaction with AtMED21. Single

alanine mutations of these four amino acids in the context of TPLN188 significantly reduced

interaction with AtMED21, while the quadruple mutation (here called Quad^{AAAA}) completely

abrogated AtMED21 binding (Figure 3D). These mutations had little effect on interaction with

AtMED10B (Figure 2 – Figure Supplement 2D). Introduction of Quad^{AAAA} mutations into the

Helix 3 through Helix 8 context (H3-H8-Quad^{AAAA}) in the AtARC^{Sc} largely phenocopied a deletion

of Helix 8 (yellow and pink, Figure 4E, compare to H3-H7, Figure 1 – Figure Supplement 1). In

contrast, TPLN188-Quad^{AAAA} largely retained the repressive activity of wild-type N188 (red and

black, Figure 3E), consistent with the observation that Helix 1 is sufficient for full repression

(Figure 1D). These results indicate that the CRA domain (H3-H8) requires contact with MED21

to repress, and that this is independent of repression via Helix 1.

Figure 3. Identification of critical residues within Helix 8 repression domain. A. Sequence
and structure of Helix 8 (5NQS). Helix 8 is colored green, and amino acids chosen for mutation
are highlighted in light green in both the sequence and the structure. **B.** Repression activity of
indicated single and double alanine mutations. **C.** Time course flow cytometry of selected
mutations of Helix 8 following auxin addition. TPLH3-8-IAA3 fusion proteins (black) were
compared to indicated single mutations to alanine (red). Controls - Helix 1 (H1 – blue), and
TPLN188 (dark green). **D.** A series of alanine mutations (V145A, E146A, K148A, K149A, and
the quadruple mutant Quad^{AAAA} chosen from Figure 2D-F) were introduced into the TPL-N188
bait construct and tested for interaction with wild-type TPL-N188, AtMED21 and controls. Each
single alanine mutation reduces TPL interaction with AtMED21, while the quad mutation
abrogated interaction. **E.** The Helix 8 Quad^{AAAA} mutation was introduced into the TPLN188-
IAA3 and TPLH3-8-IAA3 fusion proteins and compared to wild type N188 in time course flow
cytometry. For all cytometry experiments, the indicated TPL construct is fused to IAA3. Every

point represents the average fluorescence of 5-10,000 individually measured yeast cells (a.u. -

arbitrary units). Auxin (IAA-10µM) was added at the indicated time (gray bar, + Aux). At least

two independent experiments are shown for each construct.

The large Mediator complex stabilizes the pre-initiation complex (PIC) enabling

transcriptional activation (Kornberg, 2005; Nozawa et al., 2017; Roeder, 1996; Schilbach et al.,

2017). The connection we found between TPL, two Mediator components and transcriptional

repression raised the possibility that other parts of the Mediator complex might also contribute to

corepressor function. To test this, we needed a way to measure whether loss of function of

individual Mediator components led to de-repression at an individual locus. In the case of the

yeast corepressor Tup1, a standard approach has been to test the level of transcription at target

genes in the presence of deletion mutations of Tup1-interacting proteins (Gromöller and

Lehming, 2000; Lee et al., 2000; Zhang and Reese, 2004). One challenge to such an approach

for Mediator components is that loss of function mutants can be lethal or exhibit drastic

physiological phenotypes (Biddick and Young, 2005).

To avoid these complications, we turned to the well-established Anchor Away system for

inducible protein depletion (Haruki et al., 2008) and combined it with quantification of

transcriptional activity at the synthetic locus in the AtARC^S (Figure 4B). However, AtARC^{Sc}

integrates components at four genomic locations using prototrophic markers that are not

compatible with those needed for Anchor Away. To overcome this limitation, we re-recreated the

ARC on a single plasmid (we refer to this plasmid as SPARC) using the Versatile Genetic

Assembly System (VEGAS, (Mitchell et al., 2015)). SPARC behaved with similar dynamics to

the original AtARC^{Sc} on both solid and liquid growth conditions (Figure 4 – Figure Supplement

1A-C). As a first test of the Anchor Away system with SPARC, we fused Tup1 and its partner

protein Cyc8 to two copies of the FKBP12-rapamycin-binding (FRB) domain of human mTOR

(Haruki et al., 2008). Rapamycin treatment of strains targeting either of these proteins caused

no release of repression on the SPARC reporter, providing confirmation of orthogonality of the

synthetic system in yeast (Figure 4 – Figure Supplement 1D).

Before testing repressive function, we first performed chromatin immunoprecipitation

using the Anchor Away FRB tag to ask whether Mediator proteins or specific Mediator modules

were detectable at the promoters of TPL-repressed genes. We began by assaying the

integrated AtARC^{Sc} locus and comparing it to a Tup1-repressed locus, SUC2 (Carlson and

Botstein, 1982; Fleming and Pennings, 2007, p. 2; Trumbly, 1992). We generated a yeast strain

with an integrated MED21-FRB fusion protein and an integrated AtARC^{Sc} locus with a 2xHA

epitope-tagged TPLN188-IAA3 repressor protein. We observed enrichment of both MED21-FRB

and the 2xHA-TPLN188-IAA3 fusion protein at the AtARC^{Sc} locus (Figure 4C). We also

observed a modest enrichment (2-fold) of MED21-FRB at the SUC2 locus. Next, we introduced

the fully repressed SPARC plasmid containing TPLN188 (SPARC^{N188}) into a library of Anchor

Away yeast strains that allow specific depletion of Mediator components (see Figure 4A-B)

(Haruki et al., 2008; Petrenko et al., 2017). We tested representatives of the mediator complex

(Tail – Med15, Head – Med18, Middle – Med21 & Med14, Kinase – CDK8, Basal Transcription

Factors – TBP1, TFIIA, and RNA polymerase II – Rbp1) by ChIP-qPCR. We observed

enrichment of all tested core Mediator complex members, as well as Basal Transcription

Factors, at both the SPARC and the SUC2 loci, with very little enrichment of RNA polymerase II

(Figure 4D). Higher enrichment of members of the middle module (i.e. Med21, Med14, Figure

4D) may point to nucleation of the complex by assembly of the middle domain.

We next tested whether the association of Mediator complex components were required

for TPL-mediated repression (Figure 4A-B). Nuclear depletion of Mediator components from the

Tail, Head, and Middle domain triggered clear activation of the SPARC^{N188} reporter (Figure 4E-

I). Depletion of the Mediator kinase module component CDK8 had a more modest effect (Figure

4H). One caveat to this approach is that nuclear depletion of components that are absolutely

required for transcriptional activation, such as the Tail triple Anchor Away (17,3,15 - Figure 4E)

or the RNA Polymerase II (Pol II) Anchor Away (Rpb1 – Figure 4I), cannot be assayed for

impacts on repression using transcription of the reporter as the output.

To further interrogate the impact of Mediator on repression, we next focused on the

other side of the interaction, namely which region of MED21 was required for interaction with

TPL. Deletion of the first 7 amino acids of ScMed21 (Δ7Med21) partially activates genes that

are normally repressed by Tup1 (Gromöller and Lehming, 2000), so we first tested if the same

held true for TPL-mediated repression. We introduced SPARCs with different TPL constructs

into strains where wild-type ScMed21 or N-terminal deletions were targets of Anchor Away.

Importantly, the addition of the FRB tag did not alter ScMED21 function (Figure 4J, Figure 4 –

Figure Supplement 3E-F). We observed that deletion of either five or seven of the N-terminal

residues of ScMed21 increased the expression of the reporter in SPARC^{N188} to a level similar as

what is observed with TPL H1-H5 (Figure 4J, Figure 4 – Figure Supplement 1E-F). No mutation

340 increased the SPARC's sensitivity to auxin. As Δ7ScMed21 had a noticeable impact on growth,

341 as has been reported previously (Gromöller and Lehming, 2000; Hallberg et al., 2006), we

342 removed it from further studies. Δ5ScMed21 had no observable growth defects, although this

343 deletion is known to be sufficient to alter Mediator assembly and disrupt binding of Pol II and the

344 CDK8 kinase module (Hallberg et al., 2006; Sato et al., 2016).

345   The fully repressed SPARC^{N188} in Δ3ScMed21 or Δ5ScMed21 mutants showed elevated

346 reporter transcription when compared to strains carrying wild-type ScMED21 (Figure 4K). The

347 addition of Rapamycin further increased reporter expression, particularly in the Δ3ScMed21

348 strains, suggesting that this deletion could only partially disrupt the TPLN188-Med21 interaction.

349 We used chromatin immunoprecipitation to directly test whether Δ5Med21 showed a change in

350 association with the SPARC^{N188} promoter. While this deletion would be expected to reduce

351 association with TPL, the resulting de-repression of the locus should lead to an increase in

352 association with the activated promoter. Indeed, we observed a 1.6x fold increase in Δ5Med21-

353 FRB promoter binding compared to wild-type (Figure 4L). While this enrichment is modest

354 compared to a repressor-free SPARC^{IAA14} (Figure 4L, gray bar), it is correlated with the

355 magnitude of the increase in transcriptional activation of the reporter in the Δ5Med21 genotype

356 (Figure 4J-K). To confirm that the interaction with TPL was not unique to ScMed21, we replaced

357 the first five amino acids of ScMed21-FRB with the corresponding sequence from AtMED21.

358 The strain carrying this chimeric protein had an identical repression profile as the one with

359 native ScMed21 (Figure 4 – Figure Supplement 2B), and showed no difference in growth or

360 viability (Figure 4 – Figure Supplement 2C).

361   To minimize any possible off-target impact of ScMed21 deletions, we introduced

362 estradiol inducible versions of ScMed21 (iScMed21) into the Anchor Away SPARC^{N188} strains

363 (Figure 4 – Figure Supplement 4, (McIsaac et al., 2013)). The combination of all three synthetic

364 systems--ARC^{Sc}, Anchor Away and estradiol inducibility--made it possible to rapidly deplete the

365 wild type ScMed21-FRB from the nucleus while simultaneously inducing ScMed21 variants and

366 visualizing the impact on a single auxin-regulated locus. Depletion of nuclear ScMed21 by

367 Rapamycin increased levels of the reporter in all genotypes examined (Figure 4G,K) while

368 increasing cell size even in short time-courses, consistent with its essential role in many core

369 pathways (Figure 4 – Figure Supplement 3A, (Gromöller and Lehming, 2000)). When wild-type

370 iScMed21 was induced, there was a rescue of both reporter repression and cell size (Figure 4 –

371 Figure Supplement 4C,D), whereas induction of either Δ3 and Δ5 variants resulted in

significantly less reporter repression (Figure 4 – Figure Supplement 4D). iΔ3Med21 was

induced and accumulated at a comparable level to wild type Med21, while iΔ5Med21 is less

stable (Figure 4 – Figure Supplement 4E-F). In the time courses with both Rapamycin and

estradiol, we did not observe the cell size increases observed in the Rapamycin treatments

alone (populations were evenly distributed around a single mean), suggesting we were

observing the immediate effects of the Med21 deletions (Figure 4 – Figure Supplement 4G-H).

Figure 4. Repression by TPL requires interaction with the N-terminus of MED21 at

promoters. A. Model of the proposed interaction between the TPL N-terminus with Mediator,

where TPL interaction with Mediator 21 and 10 inhibit the recruitment of Pol-II. Proteins in this

complex that were tested by Anchor Away are listed on the right. **B.** Schematic of _At_ARC[Sc]

combined with methods for inducible expression and nuclear depletion of MED21. In Anchor

Away, the yeast ribosomal protein 13A (RPL13A) is fused to the rapamycin binding protein

FKBP. Addition of Rapamycin induces dimerization between FKBP and any target protein fused

to 2xFRB, resulting in removal of the target protein from the nucleus. For these experiments,

AtARC^{Sc} was assembled into a single plasmid (SPARC) rather than being integrated into

separate genomic loci (Figure 4 – Figure Supplement 1). **C.** 2xHA-TPLN188-IAA3 and MED21-

FRB association with the ARC and the ScSUC2 promoters. ChIP was performed with αHA and

αFRB before qPCR was used to quantify enrichment at specified loci. **D.** Association of FRB-

tagged components of Mediator and the transcriptional machinery with the SPARC plasmid and

the ScSUC2 promoters. ChIP was performed with αFRB and qPCR was used to quantify

enrichment at specified loci. **E-I.** Time-course flow cytometry analysis of SPARC^{N188} in Mediator

Anchor Away yeast strains with Rapamycin (orange bar, + Rapa). Two Med21 strains were

compared in the Middle domain (E), 21 (generated in this study) and 21* (generated in

(Petrenko et al., 2017)). Both 21 and 21* demonstrated similar increases in reporter expression.

J. Quantification of Venus fluorescence from SPARC^{N188} in wild type and N-terminal ScMed21

deletions with and without auxin. The x-axis indicates strain and which FRB fusion protein is

being tested. Yeast was grown for 48 hours on SDO media with or without auxin and colony

fluorescence was quantified and plotted with the auxin responsive SPARC^{H1-H5} in wild type as a

reference. Background - red autofluorescence was used as a reference for total cell density. **K.**

Time-course flow cytometry analysis of SPARC^{N188} in wild type and N-terminal ScMed21

deletions with and without Rapamycin. Genotypes are indicated in the colored key inset into the

graph. For (**E-I, K**) a.u. - arbitrary units. Rapamycin was added at the indicated time (orange

bar, + Rapa). Every point represents the average fluorescence of 5-10,000 individually

measured yeast cells. **L.** Association of MED21-FRB or Δ5-MED21-FRB with SPARC plasmids.

ChIP was performed with αFRB and qPCR was used to quantify enrichment at specified loci. **C-**

D,L. A region of the ACT1 gene body was arbitrarily defined as background, and data is

presented as fold enrichment over the control gene. Averages and standard errors of four

replicates are shown.

Several lines of evidence suggest that, in addition to interactions with other partners,

homomultimerization modulates TPL repression potential. First, structures of the N-terminal

domains of TPL (Martin-Arevalillo et al., 2017) and a rice homolog OsTPR2 (Ke et al., 2015)

reveal high conservation of residues that coordinate formation of homotetramers and connect

tetramer formation to Aux/IAA binding. Second, the dominant TPL mutant *tpl-1* altered a single

amino acid in the ninth helix of the TPL-N terminus (N176H) that induces aggregation of TPL

and its homologs (TPR1-4), reducing total activity (Long et al., 2006; Ma et al., 2017). Third,

TPL recruitment motifs found in the rice strigolactone signaling repressor D53 induce higher-

order oligomerization of the TPL N-terminus, which increases histone binding and transcriptional

repression (Ma et al., 2017). Our studies in yeast suggest there may be a more complex

relationship between tetramer formation and repression, as we have measured strong

repressive activity in several constructs that are unlikely (TPLN100 (Pierre-Jerome et al., 2014))

or unable (H1, H3-8; Figure 1C) to form tetramers (compare Figure 1B with Figure 5A). To

quantify the potential for interaction among our constructs, we used the cytoSUS assay (Asseck

and Grefen, 2018). Helix 8 was required for strongest interaction between TPL constructs

(Figure 5 – Figure Supplement 1A), although this assessment was complicated by the fact that

some of the shorter constructs accumulated to significantly lower levels (Figure 5 – Figure

Supplement 1B). The weak interaction we could observe between full-length TPL-N and the

Helix 1 through Helix 3 construct (H1-3), indicated that the TPL LisH domain is sufficient for

dimerization. Therefore, while auxin-insensitive repression may require multimeric TPL, this

higher-order complex was not required for auxin-sensitive repression mediated by Helix 1

(Figure 1E).

To avoid any potential artifacts from analysis of truncated forms of the N-terminus, we

next generated site-specific mutations that disrupted multimerization in the context of TPLN188.

Martin-Arevalillo et al. had previously identified a triple mutation (K102S-T116A-Q117S-E122S)

that abrogated the ability of the CRA domain (Helix 6 and Helix 7) to form inter-TPL interactions

(Martin-Arevalillo et al., 2017). As this mutant form of TPL is only capable of dimerizing through

its LisH domain, we refer to it here as LDimer (Figure 5A). The LDimer mutations in TPLN188

retained the same auxin insensitive repression auxin behavior as wild-type TPLN188 (Figure

5D), supporting the finding from the deletion series.

To make a fully monomeric form of TPL, we introduced mutations into the dimerization

interface of the LisH domain in the context of LDimer. We first mutated one of a pair of

interacting residues (F15) to a series of amino acids (Tyrosine - Y, Alanine - A, Arginine - R, or

Aspartic Acid - D) in the context of either LDimer (Figure 5D), or H1-2 (Figure 5B, 5E).

Conversion of F15 to the polar and charged aspartic acid (D) completely abolished repression

activity, while the positively charged arginine was better tolerated (Figure 5D,E). The conversion

of F15 to tyrosine had no effect on LDimer (Figure 5D), and only a minimal increase in auxin

sensitivity in the context of H1-2 (Figure 5E). We then combined LDimer-F15Y with a mutation

of the coordinating residue L8 to serine with the intention of stabilizing the now solvent-facing

residues. The repressive behavior of this mutant was indistinguishable from that of LDimer

(Figure 5F).

To further push the LDimer towards a monomeric form, we introduced two additional

mutations (S5A, E19S, Figure 5C,G). Size exclusion chromatography confirmed that this

combination of mutations (S5A-L8S-F15Y-E19S-K102S-T116A-Q117S-E122S, hereafter called

Monomer) successfully shifted the majority of the protein into a monomeric state (Figure 5H);

however, this shift had no observable impact on repression strength before or after auxin

addition (Figure 5G). To test whether these mutations had a similar impact on *in vivo* TPL

complexes, we introduced the LDimer and Monomer mutations into the cytoSUS assay. In

contrast to the *in vitro* chromatography results with purified proteins, Monomer expressed in

yeast retained measurable interaction with wild-type TPL, LDimer or Monomer, albeit at a

reduced level than what was observed between wild type TPLN188 constructs (Figure 5I). A

caveat to this apparent difference between assays is that the Monomer mutations led to a

striking increase in protein concentration in yeast (Figure 5 – Figure Supplement 1C), likely

partially compensating for the decrease in affinity.

Figure 5. Multimerization is not required for repression in yeast. A. TPL can form a
homotetramer via the CRA (red) and LisH (blue) domains. Asterisks indicate mutations that
block or diminish these interactions. **B-C**. Locations of critical positions in Helix 1 are
highlighted for two interacting TPL monomers (shown in light and dark blue). Interacting amino
acids share the same color (adapted from 5NQV). **D-G.** Time course flow cytometry analysis of
TPLN-IAA3 fusion proteins carrying selected single point mutations in N188-LDimer-IAA3 (**D**)
and the TPLH1-2 truncation (**E**). The F15Y mutation had little effect on repression activity for
either TPL construct. Double mutations (F15Y, L8S in LDimer) (**F**) or the quadruple Monomer
mutations (S5A, L8S, F15Y, E19S in LDimer) (**G**) showed repression activity that was
indistinguishable from LDimer or wild type N188 fused to IAA3. For all cytometry experiments,
the indicated TPL construct is fused to IAA3. Every point represents the average fluorescence
of 5-10,000 individually measured yeast cells (a.u. - arbitrary units). Auxin (IAA-10µM) was
added at the indicated time (gray bar, + Aux). At least two independent experiments are shown
for each construct. **H**. Size Exclusion Chromatography on TPLN188 wild type (green), LDimer
(purple) and Monomer (orange) tetramerization mutants. **I**. CytoSUS on TPL tetramerization
mutants.

To ascertain which of our findings about TPL required the sensitivity and simplicity of the

synthetic context and which could be observed in the full complexity of intact plant systems, we

performed a set of experiments in *Nicotiana benthamiana* (tobacco) and *Arabidopsis.*

Bimolecular Fluorescence Complementation (BiFC) confirmed the interaction between TPL and

MED21 (Figure 6A), which was further validated by co-immunoprecipitation using tobacco

extracts (Figure 6B). We were also able to pull down MED21 and TPL using MED10B (Figure 6

– Figure Supplement 1A). BiFC also confirmed the importance of the same TPL Helix 8

residues for the TPL-AtMed21 interaction (Figure 6A, TPL^{H8QuadA}). Similarly, the Δ5AtMED21 N-

terminal truncation eliminated interaction with full-length TPL (Figure 6A). We next developed a

quantitative repression assay based on UAS/GAL4-VP16 (Brand and Perrimon, 1993) (Figure

6C). To block potentially confounding interactions with endogenous TPL/TPRs or TIR1/AFBs,

we engineered a variant of IAA14 with mutations in the two EAR domains (EAR^{AAA}) and in the

degron (P306S) (IAA14^{mED} ; Figure 6C). After prototyping the system in yeast (Figure 6 – Figure

Supplement 1B-C), we quantified repression strength of constructs carrying *TPLN-IAA14^{mED}*

variants using the well-characterized synthetic auxin responsive promoter DR5 (Ulmasov et al.,

1997). As expected, DR5 was strongly induced by co-transformation with AtARF19, and this

induction was sharply reduced by the inclusion of *UAS-TPLN188-IAA14^{mED}* and *GAL4-VP16*

(Figure 6 – Figure Supplement 1D). Overall, we observed strong correlation in repression

activity between what was observed in yeast and in tobacco.

To connect the observed differences in repression strength to a developmental context,

we generated transgenic *Arabidopsis* lines where the UAS-TPL-IAA14^{mED} constructs were

activated in the cells where IAA14 normally acts to regulate the initiation of lateral root primordia

(Figure 6D, (Gala et al., 2020; Laplaze et al., 2005)). Expression of functional TPL-IAA14^{mED}

fusion proteins in these xylem pole pericycle cells should strongly suppress production of lateral

roots, phenocopying the solitary root (*slr*) mutant, which carries an auxin-resistant form of IAA14

(Fukaki et al., 2002). Indeed, TPLN188 fusion constructs sharply decreased lateral root density

(Figure 6E), while transformants expressing either IAA14^{mED} (with no TPL fusion) or TPLN188

(with no IAA14 fusion) had no effect on lateral root production (Figure 6E). TPLH3-H9

decreased lateral root density albeit not as effectively as TPLN188, suggesting that Helix 1 is

required for full repression in a native context (Figure 6E). Both LDimer and Monomer

constructs (Figure 6E) were able to repress lateral root development to the same extent as

TPLN188, meaning that multimer formation is not required for TPL-mediated repression in this

context. The fusion containing the Helix 8 quadruple mutant demonstrated a clear loss of

repression, indicating that the TPL-MED21 interaction is critical for repression when expressed

in lateral root forming cells (Figure 6E).

Given this result, we wanted to directly test the role of AtMed21 in auxin-regulated

development in the presence of native isoforms of TPL and Aux/IAAs. This was complicated by

the fact that, as in yeast, AtMED21 is essential in plants. While homozygous loss-of-function

mutations are embryo lethal (Dhawan et al., 2009), plants heterozygous for *Atmed21* mutations

appear wild-type (Figure 6 – Figure Supplement 2A-B). To overcome this obstacle, we took two

approaches that relied on the same xylem pole pericycle driver as described for the TPL

functional assays. First, we expressed N-terminal deletion variants of MED21 that should

weaken or sever interaction with TPL (Figure 6F - Δ3MED21, Δ5 MED21, Δ7 MED21). If

MED21-TPL interaction is critical to maintain normal expression of the lateral root program,

reduced interaction should trigger an increase in lateral root density. This was exactly what we

observed for transgenic lines expressing any of the three deletions. Second, we repressed

transcription of *AtMED21* by introducing a dCAS9-TPLN300 synthetic repressor under the

control of a UAS promoter along with three sgRNAs complementary to the *AtMED21* promoter.

Similar to the predictions above, reduced expression of AtMED21 in xylem pole pericycle cells

should stimulate lateral root development. This is indeed what we observed (Figure 6 – Figure

Supplement 2C-E).

Previous reports connected the Mediator kinase module, and specifically the function of

AtMED13/ GRAND CENTRAL (GCT), to repression by TPL (Ito et al., 2016). We did not find

evidence for this relationship in our yeast assays, and wanted to investigate further. Although

embryo lethality is seen in the majority of homozygous *med13/gct* individuals, a small

percentage do survive. This made it possible to examine the effect of expressing wild type and

monomeric forms of TPLN188 in xylem pole pericycle cells in the absence of MED13/GCT

activity (Figure 6 – Figure Supplement 1E-F). Loss of *med13* function was unable to rescue the

production of lateral roots in these lines, leading to the conclusion that TPLN188 requires

AtMED21 but not AtMED13 for repression, at least in this context.

Figure 6. The TPL CRA repression domain behaves similarly in yeast and plants

A. Bimolecular Fluorescence Complementation assay performed in tobacco. Each image is an epi-fluorescent micrograph taken at identical magnification from tobacco epidermal cells at two days post injection. The YFP image is colored green (left panel). p35S:H2B-RFP was used as a

control and is false colored magenta (right panel). **B.** Co-immunoprecipitation of MED21 and

TPL from tobacco leaves. MED21-YFP-HA was immunoprecipitated using anti-HA, and YFP-

TPL was detected using the YFP fusion. Actin was used to demonstrate that the purification had

removed non-specific proteins. Numbers on the left of blots indicate sizes of protein standards

in kilodaltons. **C.** Design of UAS-TPL-IAA14^{mED} and UAS-MED21 constructs. Mutation of the

conserved Lysine residues in the EAR domain disrupted potential interactions with endogenous

TPL/TPR proteins. The IAA14 degron has been mutated (P306S) to render it auxin insensitive.

UAS – upstream activating sequence, ttRBCS - Rubisco terminator sequence. **D.** Auxin induced

degradation of IAA14 is absolutely required for initiation of lateral root development (cartoon,

left). An enhancer trap line (J0121) expresses GAL4-VP16 and UAS-GFP in in xylem pole

pericycle cells. **E.** N-terminal domains of TPL were sufficient to repress the development of

lateral roots in *Arabidopsis* seedlings. The density of emerged lateral roots was measured in T1

seedlings at 14 days after germination. **F.** N-terminal deletions in *AtMED21* were sufficient to

dominantly increase the development of lateral roots in *Arabidopsis* seedlings. The density of

emerged lateral roots was measured in T1 seedlings at 14 days after germination. **F-G**. Lower

case letters indicate significant difference (ANOVA and Tukey HSD multiple comparison test;

$p<0.001$).

Discussion

A review of the current literature on corepressors gives the conflicting impressions that (a)

corepressor function is broadly conserved, and (b) that every organism (and perhaps even

every corepressor) has a distinct mode for transcriptional repression (Adams et al., 2018; Mottis

et al., 2013; Perissi et al., 2010; Wong and Struhl, 2011). We hoped that the *At*ARC^{Sc} could

facilitate a resolution to this apparent contradiction by targeting repression to a single synthetic

locus. We focused our initial efforts on the analysis of the N-terminal portion of TPL which has

multiple known protein-protein interaction surfaces (Ke et al., 2015; Martin-Arevalillo et al.,

2017). Experiments with the *At*ARC^{Sc} identified two independent repression domains (Figure 1),

and we focused additional study on the stronger of the two localized to the CRA domain. Within

this domain, we were able to identify two interacting partners that are part of the middle domain

of the Mediator complex: MED21 and MED10 (Figure 2). Four amino acids within Helix 8 with R-

groups oriented away from the hydrophobic core of the TPL structure were found to be required

for both Med21-binding and repressive function (Figure 3). Indeed, the entire core Mediator

complex (Head, Middle, Tail) appears to be recruited to TPL-repressed loci and required to

maintain repression (Figure 4). Contrary to our initial hypothesis, the monomeric form of TPL

was sufficient for strong repression in yeast and in plants (Figure 5), leaving open the question

of the role of higher-order TPL complex formation. Finally, we were able to confirm that our

insights from synthetic assays in yeast were relevant to regulation of the auxin-mediated lateral

root development pathway in intact plants (Figure 6).

Corepressors coordinate multiple mechanisms of repression through discrete protein

interactions, leading to robust control over eukaryotic transcription by combining repression

modalities. Corepressor function has variously been linked to (a) altering chromatin

confirmation, often through interaction with histone modifying proteins or histone proteins

themselves, (b) direct interference with transcription factor binding or function, and (c) physical

spreading of long-range oligomeric corepressor complexes across regions of regulatory DNA

(Perissi et al., 2010). Dissection of the importance of each modality in Tup1 repression has

been challenging (Lee et al., 2000; Zhang and Reese, 2004). The tour-de-force of corepressor

mechanism studies in yeast concluded that the primary function of Tup1 was to physically block

activators (Wong and Struhl, 2011). In their work, the authors utilized the Anchor Away

approach to correlate the importance of HDACs, transcriptional machinery and chromatin

remodeling enzymes to the repression state of endogenously repressed Cyc8-Tup1 target

genes. They observed that Tup1 did not block the binding of transcription factors but inhibited

the recruitment of one Mediator component in the tail domain, GAL11/MED15, as well as Pol-II

and the chromatin remodelers Snf2 and Sth1. They additionally observed that HDACs had only

a supportive role in reinforcing Tup1 repression. These results led to their hypothesis that Tup1

blocks the activation domains of transcription factors, and suggested this was through direct

binding to activation domains (Wong and Struhl, 2011).

The synthetic system used here allowed us to build on this model and further refine our

understanding of TPL's repressive activity. In our experiments we see a similar set of

conditions, with TPL recruited to the DNA-bound transcriptional activator (ARF), and several

possible mechanisms of repression. Unlike Tup1, we have subdivided the TPL protein to identify

interactions between TPL and individual protein interactors with no effect on yeast function. In

these experiments we can eliminate the possibility that TPL blocks ARF activation by directly

blocking the transcription factor activation domains because we see a loss of repression only

when TPL-MED21 binding is eliminated through specific point mutations (Figure 4J-L). Our

inducible swap of MED21 isoforms also corroborates these findings (Figure 4 – Figure

Supplement 3), as the SPARC remains genetically identical in these strains, indicating that TPL-

MED21 interaction is regulating Mediator activity not a TPL-ARF interaction. Furthermore, our

results correlate well with findings that repressed targets are reactivated when this portion of

MED21 is deleted in yeast (TPL - Figure 4, Tup1 - (Gromöller and Lehming, 2000)). Therefore,

we suggest that instead of directly binding activation domains that TPL (and likely Tup1) binds

to components of Mediator (MED21, MED10B and possibly others) recruited by the transcription

factor. Indeed, it is easier to rationalize that the repressor binds the same domains of the

Mediator complex recruited by the transcription factor's activation domain (with the same

affinity) as opposed to binding each diverse activation domain (with varying affinity). In this

model, corepressor binding blocks formation of a fully active Mediator complex thereby limiting

Polymerase II recruitment and promoter escape (Petrenko et al., 2017).

The Mediator complex is a multi-subunit complex that connects DNA-bound transcription

factors and the RNA polymerase II complex (Pol-II) to coordinate gene expression (Flanagan et

al., 1991; Kim et al., 1994). The yeast Mediator subunits are organized into four separate

modules; head, middle, tail and kinase, with a strong conservation of module components in

plants (Dolan and Chapple, 2017; Maji et al., 2019; Malik et al., 2017; Samanta and Thakur,

2015). Med21 forms a heterodimer with Med7 to and interacts with Med10, among others, to

create the central region of the Middle region of the Mediator complex. The Med21 N-terminus

is centered on a flexible hinge region (Baumli et al., 2005), which is required for Pol-II

recruitment and CDK8 kinase module recruitment (Sato et al., 2016). The protein interaction

between TPL and MED21 occurs at the N-terminus of MED21, highlighting the importance of

this region as a signaling hub (Sato et al., 2016). Other lines of evidence support this role, as

this region binds the yeast homolog of TPL, Tup1 (Gromöller and Lehming, 2000) through a

completely different protein domain as no homology can be found between TPL Helix 8 and

Tup1 in any region by primary amino acid homology (i.e. BLAST).

As suggested by Ito and colleagues (Ito et al., 2016) and supported by our synthetic

system, auxin-induced removal of TPL is sufficient to induce changes in the composition of the

Mediator complex; however, multiple points of contact likely exist between the Mediator complex

and other parts of the transcriptional machinery in both transcriptionally repressed and active

states. For auxin response, specifically, there are several lines of evidence to support this

model, including documented association between the structural backbone of Mediator, MED14,

and activated and repressed auxin loci in *Arabidopsis* (Ito et al., 2016). In addition, MED12 and

MED13 are required for auxin-responsive gene expression in the root, and MED12 acts

upstream of AUX1 in the root growth response to sugar (Raya-González et al., 2018). MED18 in

the head module represses auxin signaling and positively regulates the viability of the root

meristem (Raya-González et al., 2018). PFT1/MED25 regulates auxin transport and response in

the root (Raya-González et al., 2014). MED7, MED21's partner protein in the hinge domain, is

required for normal root development and loss of MED7 function impacts expression of auxin

signaling components (Kumar et al., 2018). Previous research identified the Mediator CDK8

subunit, specifically MED13 (MAB2), as an interactor with the full-length TPL protein (Ito et al.,

2016). We could not observe interaction between the N-terminal domain of TPL and AtMED13,

AtCYC8, or AtCYCC (Supplemental Figure 1B), suggesting that any direct interactions occur

outside the N-terminal region.

The conserved interaction of both TPL and Tup1 with Mediator has implications for

modeling eukaryotic transcription (e.g., (Estrada et al., 2016)). By stabilizing the Mediator

complex, TPL (and by extension Tup1) may create a 'pre-paused' state that allows rapid

recruitment of Pol-II and activation once TPL is removed. This would be compatible with the

multiple repression mechanisms have been described for TPL at different genetic loci. TPL

recruitment of the repressive CDK8 Mediator complex (Ito et al., 2016), chromatin remodeling

enzymes such as Histone Deacetylase 19 (HD19) (Long et al., 2006) and contact with histone

proteins (Ma et al., 2017) would be removed with TPL upon relief of repression. It will be critical

in the future to understand how these various forms of repression interact, and especially to

map the dynamics of assembly and disassembly of complexes as loci transition from repressed

to active states and back to repressed once again.

RESOURCE AVAILABILITY

Lead Contact

Further information and requests for resources and reagents should be directed to and will be

fulfilled by the Lead Contact, Jennifer, Nemhauser (jn7@uw.edu).

Materials Availability

Plasmids generated in this study will be deposited to [Addgene, Jennifer Nemhauser].

Data and Code Availability

Data and Code Availability Statement

All flow cytometry data is deposited at https://flowrepository.org/. Repository IDS:

FR-FCM-Z2GM, FR-FCM-Z2GR, FR-FCM-Z2GQ, FR-FCM-Z2GX, FR-FCM-Z2GT, FR-

FCM-Z2W2.

All protein interactions will be deposited at IntAct.

All code is available through Github:

https://github.com/achillobator/TPL_Structure_Function/

Method Details

Cloning

Construction of TPL-IAA3 and TPL-IAA14 fusion proteins were performed by golden gate

cloning as described in (Pierre-Jerome et al., 2014). Variant and deletion constructs were

created using PCR-mediated site-directed mutagenesis. Site-directed mutagenesis primers

were designed using NEBasechanger and implemented through Q5® Site-Directed

Mutagenesis (NEB, Cat #E0554S). TPL interactor genes were amplified as cDNAs from wild

type Col-0 RNA using reverse transcriptase (SuperScript™ IV Reverse Transcriptase,

Invitrogen) and gene-specific primers from IDT (Coralville, Iowa), followed by amplification with

Q5 polymerase (NEB). These cDNAs were subsequently cloned into Plasmids for cytoSUS

using a Gibson approach (Gibson et al., 2009), through the Aquarium Biofabrication facility (Ben

Keller et al., 2019). The coding sequence of the genes of interest were confirmed by sequencing

(Genewiz; South Plainfield, NJ). For UAS driven constructs, the TPLN188-IAA14 coding

sequence was amplified with primers containing engineered BsaI sites and introduced into the

pGII backbone with the UAS promoter and RBSC terminator (Siligato et al., 2016) using Golden

Gate cloning (Weber et al., 2011). Subsequent mutations were performed on this backbone

using PCR-mediated site-directed mutagenesis (see above). Construction of C-terminal 2xFRB

fusions for Anchor Away were constructed as described in (Haruki et al., 2008). Inducible

MED21 was constructed as described in (McIsaac et al., 2013). For cell type specific

knockdown mediated by dCas9-TPLN300, Gibson cloning was used modify the pHEE401E

plasmid, replacing the egg-specific promoter and Cas9 from pHEE401E (Wang et al., 2015) with

UAS promoter and dCas9- TPLN300 fusion protein (Khakhar et al., 2018). The resulting plasmid

is used as starting point to clone 3 sgRNAs targeting the *AtMED21* promoter (identified using

CHOP CHOP (Labun et al., 2019). sgRNAs: GACGCAGAGTCTGTTGGGTGTGG,

TTTAAAATGGGCTTTTAAGGTGG, AACACTGAAGTAGAATTGGGTGG ranging from -170 to

+90 region from the TSS) using PCR and Golden Gate cloning strategy described in (Wang et

al., 2015).

Flow Cytometry

Fluorescence measurements were taken using a Becton Dickinson (BD) special order

cytometer with a 514-nm laser exciting fluorescence that is cut off at 525 nm prior to

photomultiplier tube collection (BD, Franklin Lakes, NJ). Events were annotated, subset to

singlet yeast using the FlowTime R package (Wright et al., 2019). A total of 10,000 - 20,000

events above a 400,000 FSC-H threshold (to exclude debris) were collected for each sample

and data exported as FCS 3.0 files for processing using the flowCore R software package and

custom R scripts (Supplemental File 1, (Havens et al., 2012; Pierre-Jerome et al., 2017)). Data

from at least two independent replicates were combined and plotted in R (ggplots2).

Yeast Methods

Standard yeast drop-out and yeast extract–peptone–dextrose plus adenine (YPAD) media were

used, with care taken to use the same batch of synthetic complete (SC) media for related

experiments. A standard lithium acetate protocol (Gietz and Woods, 2002) was used for

transformations of digested plasmids. All cultures were grown at 30°C with shaking at 220 rpm.

Anchor-Away approaches were followed as described in (Haruki et al., 2008), and anchor away

strains were obtained from EURO-SCARF (euroscarf.de). Endogenous genomic fusions of

ScMed21-FRB were designed by fusing MED21 homology to the pFA6a-FRB-KanMX6 plasmid

for chromosomal integration into the parental anchor away strain as in (Petrenko et al., 2017),

selectable through G418 resistance (G418, Geneticin, Thermo Fisher Scientific). Tup1-FRB and

Cyc8-FRB were constructed as described in (Wong and Struhl, 2011). Mediator anchor away

strains were created in (Petrenko et al., 2017) and kindly donated by Dr. Kevin Struhl. SPARC

construction required a redesign of promoters and terminators used in the AtARC^{Sc} to eliminate

any repetitive DNA sequences (See Figure 4 – Figure Supplement 1), using a Golden Gate

cloning approach into level 1 vectors. Subsequent assembly of individual transcriptional units

into a larger plasmid utilized VEGAS assembly which was performed as described in (Mitchell et

al., 2015). To create an acceptor plasmid for the assembled transcriptional units, we

synthesized a custom vector containing VA1 and VA2 homology sites for recombination (Twist

Bioscience, South San Francisco, CA). In between these sites we incorporated a pLac:mRFP

cassette to allow identification of uncut destination plasmid in $E.\ coli$, flanked by EcoRI sites for

linearization. Finally, the CEN6/ARSH4 was transferred from pRG215 (Addgene #64525) into

the acceptor plasmid by Golden Gate reaction using designed BsmBI sites engineered into the

acceptor plasmid and the primers used to amplify the CEN/ARS (See Figure 4 – Figure

Supplement 1). For the cytoplasmic split-ubiquitin protein-protein interaction system, bait and

prey constructs were created using the plasmids pMetOYC and pNX32, respectively (Addgene,

https://www.addgene.org/Christopher_Grefen/). Interaction between bait and prey proteins were

evaluated using a modified version of the split ubiquitin technique (Asseck and Grefen, 2018).

After two days of growth on control and selection plates, images were taken using a flatbed

scanner (Epson America, Long Beach, CA). Inducible ScMed21 strains (iMed21) were grown

overnight, and then diluted back to 100 events per microliter as determined by flow cytometry

and grown at 30C with 250rpm in a deepwell 96-well plate format. Strains were supplemented

with β-estradiol (20μM) for 4 hours followed by Rapamycin addition. Samples were analyzed by

flow cytometry throughout these growth experiments.

Western Blot

Yeast cultures that had been incubated overnight in synthetic complete (SC) media were diluted

to OD600 = 0.6 and incubated until cultures reached OD600 ~1. Cells were harvested by

centrifugation. Cells were lysed by vortexing for 5 min in the presence of 200 μl of 0.5-mm

diameter acid washed glass beads and 200 μl SUMEB buffer (1% SDS, 8 M urea, 10 mM

MOPS pH 6.8, 10 mM EDTA, 0.01% bromophenol blue, 1mM PMSF) per one OD unit of original

culture. Lysates were then incubated at 65° for 10 min and cleared by centrifugation prior to

electrophoresis and blotting. Antibodies: Anti-HA-HRP (REF-12013819001, Clone 3F10,

Roche/Millipore Sigma, St. Louis, MO), Anti-FLAG (F3165, Monoclonal ANTI-FLAG® M2,

Millipore Sigma, St. Louis, MO), Anti-FRB (ALX-215-065-1, Enzo Life Sciences, Farmingdale,

NY, (Haruki et al., 2008)), Anti-VP16 (1-21) (sc-7545, Santa Cruz Biotechnology, Dallas TX),

Anti-GFP (ab290, AbCam, Cambridge, United Kingdom), Anti-MYC (71d10, 2278S, Cell

Signaling, Danvers, MA), Anti-PGK1 (ab113687, AbCam, Cambridge, United Kingdom).

Protein expression and purification

All multimer deficient TPL proteins were expressed in *Escherichia coli* Rosetta 2 strain. Bacteria

cultures were grown at 37ºC until they achieved an $OD^{600}nm$ of 0.6-0.9. Protein expression was

induced with isopropyl-β-D-1-thyogalactopiranoside (IPTG) at a final concentration of 400 μM at

18 °C overnight. Bacteria cultures were centrifuged and the pellets were resuspended in the

buffer A (CAPS 200 mM pH 10.5, NaCl 500 mM, TCEP 1 mM), where cells were lysed by

sonication. His-tagged AtTPL188 (wt and mutants) bacteria pellets were resuspended in buffer

A with EDTA-free antiprotease (Roche). The soluble fractions recovered after sonication were

passed through a Ni-sepharose (GE Healthcare) column previously washed with buffer A and

the bound proteins were eluted with buffer A with 250 mM imidazole. A second purification step

was carried out on Gel filtration Superdex 200 10/300 GL (GE Healthcare) equilibrated with

buffer A.

Co-Immunoprecipitation

Co-IP from yeast was performed using the cytoSUS strains. Cultures were grown to OD600 0.5

(~1E7 cells/ml) using selective media, harvested, and resuspended in 200 μl SUME (1% SDS,

8M Urea, 10mM MOPS, pH 6.8, 10mM EDTA) buffer with protease inhibitors. Cells were lysed

by vortexing 3 X 1 min full speed with 100 μl of 0.5 mm Acid Washed Glass Beads, clarified by

centrifugation (1 min, 1000rpm), and supernatant was mixed with 1ml IP buffer (15mM

Na2HPO4, mw 142; 150mM NaCl, mw 58; 2% Triton X-100, 0.1% SDS, 0.5% DOC, 10mM

EDTA, 0.02% NaN3) with protease inhibitors, and incubated with 100 μl of IgG sepharose at

25°C for 2 hours with rotation. The Beads were washed 1x with IP buffer, and 2x with IP-Wash

buffer (50mM NaCl, mw58; 10mM TRIS, mw 121; 0.02% NaN3) with protease inhibitors. Protein

was eluted with 50 μl of SUME buffer + 0.005% bromophenol blue by incubation at 65°C for 10

minutes and run on handmade 12% acrylamide SDS-PAGE gels, and western blotted

accordingly. Co-IPs from tobacco were performed on leaves two days after injection as

described in (Song et al., 2014), using 35S:MED21-YFP-HA, 35S:TPL-YFP (full length), and

35S:MED10B-MEC-ProtA constructs. For Co-IPs with HA, extracts were incubated with Anti-

HA-Biotin (High Affinity (3F10), Sigma, 12158167001) and Streptavidin conjugated magnetic

beads (Life Technologies, Dynabeads M-280 Streptavidin, 112.05D). For Co-IPs with MED10B-

MEC-ProtA we used IgG Sepharose® 6 Fast Flow (Sigma, GE17-0969-01) beads and

increased washing steps (1xIP buffer, 4xWash buffer, 5 total). The only modification to buffers

was an addition of the detergent NP-40 at 0.1% in the IP and wash buffer. Samples were run on

handmade 10% acrylamide SDS-PAGE gels, and western blotted accordingly.

Bimolecular Fluorescence Complementation (BiFC)

Bimolecular fluorescent complementation experiments were performed on 3-week-old *N.*

benthamiana plants grown at 22°C under long days (16 h light/8 h dark) on soil (Sunshine #4

mix) as per (Martin et al., 2009). pSITE vectors were used to generate BiFC constructs for

MED21, Δ5-MED21, TPL and TPL^{H8QuadA} – proteins (Martin et al., 2009). In all cases the

combinations are N-terminal fusions of either the nEYFP or cEYFP to the cDNA of MED21 or

TPL. RFP fused Histone H2B was used as a nuclear marker (Goodin et al., 2002). Injection of

Agrobacterium strains into tobacco leaves was performed as in (Goodin et al., 2002), but the

OD_{600} of the Agrobacterium culture used was adjusted to 0.5. Two days after transfection, plant

leaves were imaged using an epifluorescence microscope (Leica Biosystems, model: DMI

3000B).

Chromatin Immunoprecipitation.

Yeast were inoculated in liquid YPD and grown at 30 °C with shaking at 225 rpm. After 18 h of

growth, cultures were diluted 1:50 in fresh YPD to a final volume of 200 mL. Four hours post-

dilution, cells were treated with 20 mL of fix solution [11% (vol/vol) formaldehyde, 0.1 M NaCl, 1

mM EDTA, 50 mM Hepes–KOH] for 20 min at room temperature with shaking. Cultures were

further treated with 36 mL of 2.5 M glycine for 5 min to quench the cross-linking. Cells were then

pelleted at 4°C, washed twice with ice-cold TBS, and flash-frozen in liquid nitrogen. Cells were

lysed in breaking buffer [100 mM Tris, pH 8.0, 20% (vol/vol) glycerol, and 1 mM PMSF] on a

bead beater before sonication. Samples were processed in a Bioruptor Plus sonication device at

50% for 30 cycles. Following centrifugation (to pellet cellular debris), the supernatant was used

for the IP reaction. Biotin conjugated Anti-HA (High Affinity (3F10), Sigma, 12158167001)

coupled to Streptavidin coated Dynabeads Streptavidin conjugated magnetic beads (Life

Technologies, Dynabeads M-280 Streptavidin, 112.05D) was used to probe for HA-tagged

TPLN188. Anti-FRB coupled to Protein A Dynabeads (Life Technologies, 100.02D). Following

elution from the beads, samples were incubated overnight at 65 °C to reverse cross-links. DNA

was purified using a Monarch PCR purification kit (NEB). qPCR for three independent replicates

was performed using iQ SYBR Green Supermix (Biorad) in a C100 thermocycler fitted with a

CFX96 Real-Time Detection System (Biorad).

Protein alignments

The MED21 protein sequence was were aligned to homologs using CLC Sequence Viewer 7, a

tree was constructed using a Neighbor-Joining method, and bootstrap analysis performed with

10,000 replicates.

Plant growth

For synthetic repression assays in tobacco Agrobacterium-mediated transient transformation of

Nicotiana benthamiana was performed as per (Yang et al., 2000). 5 ml cultures of

Agrobacterium strains were grown overnight at 30C shaking at 220rpm, pelleted and incubated

in MMA media (10 mM MgCl2, 10 mM MES pH 5.6, 100 µM acetosyringone) for 3 hours at room

temperature with rotation. Strain density was normalized to an OD600 of 1 for each strain in the

final mixture of strains before injection into tobacco leaves. Leaves were removed, and 8

different regions were excised using a hole punch, placed into a 96-well microtiter plate with

100µl of water. Each leaf punch was scanned in a 4x4 grid for yellow and red fluorescence

using a plate scanner (Tecan Spark, Tecan Trading AG, Switzerland). Fluorescence data was

quantified and plotted in R (ggplots). For *Arabidopsis thaliana* experiments using the GAL4-

UAS system (Laplaze et al., 2005), J0121 was introgressed eight times into Col-0 accession

from the C24 accession, and rigorously checked to ensure root growth was comparable to Col-0

before use. UAS-TPL-IAA14^{mED} constructs were introduced to J0121 introgression lines by floral

dip method (Clough and Bent, 1998). T1 seedlings were selected on 0.5X LS (Caisson

Laboratories, Smithfield, UT) + 25µg/ml Hygromycin B (company) + 0.8% phytoagar

(Plantmedia; Dublin, OH). Plates were stratified for 2 days, exposed to light for 6 h, and then

grown in the dark for 3 d following a modification of the method of (Harrison et al., 2006).

Hygromycin resistant seedlings were identified by their long hypocotyl, enlarged green leaves

and long root. Transformants were transferred by hand to fresh 0.5X LS plates + 0.8% Bacto

agar (Thermo Fisher Scientific) and grown vertically for 14 days at 22°C. Plates were scanned

on a flatbed scanner (Epson America, Long Beach, CA) at day 14. *slr* and *med21/MED21*

(WiscDsLox461-464K13) seeds were obtained from the Arabidopsis Biological Resource Center

(Columbus, OH). CRISPR/CAS9 based mutations in AtMED2 were generated as described in

(Wang et al., 2015). We created a novel mutation in *AtMED21* that introduces a single base-pair

insertion of G at nucleotide 214 after the A of the start codon (i214G). This mutation alters the

amino acid sequence starting at residue 25 and creates an early stop codon after 11 random

amino acids (Supplemental Figure 6D).

Data submissions

All flow cytometry data will be deposited at https://flowrepository.org/. All plasmids will be

deposited through Addgene at https://www.addgene.org/Jennifer_Nemhauser/.

Acknowledgements

We would like to thank Prof. Jef Boeke for kindly providing VEGAS adaptor and regulatory

element plasmids; Dr. Jennifer Brophy and Prof. José Dinneny for kindly providing the

pUBQ10:GAL4:VP16 plasmid; Dr. Natalia Petrenko and Dr. Kevin Struhl for kindly providing

Mediator anchor away strains; and Prof. Grant Brown and Prof. Maitreya Dunham for advice on

yeast genetics and approaches. We thank members of the Nemhauser group including Amy

Lanctot, Romi Ramos, Eric Yang, and Dr. Sarah Guiziou for constructive discussions and

comments on this manuscript. We also thank Morgan Hamm for the custom R script used here

to analyze anchor away plates.

Author Contributions

Conceptualization: ARL, JLN; Investigation: ARL, WW, HPG, SG, SJS, MLZ, JEZ; Software:

ARL, HPG; Formal Analysis: ARL, HPG; Visualization: ARL, WW, HPG, SG, SJS, MLZ, JEZ;

Writing-Original Draft: ARL; Writing-Review & Editing: WW, HPG, SG, SJS, MLZ, JEZ, NZ, JLN;

Supervision, Project administration, Funding Acquisition, Resources: JLN, NZ.

Declaration of Interests

The authors declare no competing interests.

894   **Funding**

895   National Institutes of Health (NIH): ARL, HPG, SG, SJS, JEZ, MZ & JLN R01- GM107084.

896   Howard Hughes Medical Institute (HHMI): ARL, SJS, JEZ, MZ & JLN - Faculty Scholar Award.

897   Ning Zheng is a Howard Hughes Medical Institute Investigator.

898   ARL is supported by the Simons Foundation through the Life Science Research Foundation.

References

Adams GE, Chandru A, Cowley SM. 2018. Co-repressor, co-activator and general transcription factor: the many faces of the Sin3 histone deacetylase (HDAC) complex. *Biochem J* **475**:3921–3932. doi:10.1042/BCJ20170314

Agarwal M, Kumar P, Mathew SJ. 2015. The Groucho/Transducin-like enhancer of split protein family in animal development. *IUBMB Life* **67**:472–481. doi:10.1002/iub.1395

Asseck LY, Grefen C. 2018. Detecting Interactions of Membrane Proteins: The Split-Ubiquitin System. *Methods Mol Biol Clifton NJ* **1794**:49–60. doi:10.1007/978-1-4939-7871-7_4

Baumli S, Hoeppner S, Cramer P. 2005. A Conserved Mediator Hinge Revealed in the Structure of the MED7·MED21 (Med7·Srb7) Heterodimer. *J Biol Chem* **280**:18171–18178. doi:10.1074/jbc.M413466200

Ben Keller, Justin Vrana, Abraham Miller, Garrett Newman, Eric Klavins. 2019. Aquarium: The Laboratory Operating System. Zenodo. doi:10.5281/zenodo.2535715

Biddick R, Young ET. 2005. Yeast Mediator and its role in transcriptional regulation. *C R Biol* **328**:773–782. doi:10.1016/j.crvi.2005.03.004

Bourbon H-M. 2008. Comparative genomics supports a deep evolutionary origin for the large, four-module transcriptional mediator complex. *Nucleic Acids Res* **36**:3993–4008. doi:10.1093/nar/gkn349

Brand AH, Perrimon N. 1993. Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. *Dev Camb Engl* **118**:401–415.

Busch W, Miotk A, Ariel FD, Zhao Z, Forner J, Daum G, Suzaki T, Schuster C, Schultheiss SJ, Leibfried A, Haubeiss S, Ha N, Chan RL, Lohmann JU. 2010. Transcriptional control of a plant stem cell niche. *Dev Cell* **18**:849–861. doi:10.1016/j.devcel.2010.03.012

Carlson M, Botstein D. 1982. Two differentially regulated mRNAs with different 5' ends encode secreted with intracellular forms of yeast invertase. *Cell* **28**:145–154. doi:10.1016/0092-8674(82)90384-1

Causier B, Ashworth M, Guo W, Davies B. 2012. The TOPLESS interactome: a framework for gene repression in Arabidopsis. *Plant Physiol* **158**:423–438. doi:10.1104/pp.111.186999

Chen G, Courey AJ. 2000. Groucho/TLE family proteins and transcriptional repression. *Gene* **249**:1–16. doi:10.1016/s0378-1119(00)00161-x

Clough SJ, Bent AF. 1998. Floral dip: a simplified method for Agrobacterium-mediated transformation of Arabidopsis thaliana. *Plant J Cell Mol Biol* **16**:735–743.

Collins J, O'Grady K, Chen S, Gurley W. 2019. The C-terminal WD40 repeats on the TOPLESS co-repressor function as a protein–protein interaction surface. *Plant Mol Biol* **100**:47–58. doi:10.1007/s11103-019-00842-w

Delto CF, Heisler FF, Kuper J, Sander B, Kneussel M, Schindelin H. 2015. The LisH motif of muskelin is crucial for oligomerization and governs intracellular localization. *Struct Lond Engl 1993* **23**:364–373. doi:10.1016/j.str.2014.11.016

Dhawan R, Luo H, Foerster AM, Abuqamar S, Du H-N, Briggs SD, Mittelsten Scheid O, Mengiste T. 2009. HISTONE MONOUBIQUITINATION1 interacts with a subunit of the mediator complex and regulates defense against necrotrophic fungal pathogens in Arabidopsis. *Plant Cell* **21**:1000–1019. doi:10.1105/tpc.108.062364

Dolan WL, Chapple C. 2017. Conservation and Divergence of Mediator Structure and Function: Insights from Plants. *Plant Cell Physiol* **58**:4–21. doi:10.1093/pcp/pcw176

Estrada J, Wong F, DePace A, Gunawardena J. 2016. Information Integration and Energy Expenditure in Gene Regulation. *Cell* **166**:234–244. doi:10.1016/j.cell.2016.06.012

Flanagan PM, Kelleher RJ, Sayre MH, Tschochner H, Kornberg RD. 1991. A mediator required for activation of RNA polymerase II transcription in vitro. *Nature* **350**:436–438. doi:10.1038/350436a0

Fleming AB, Pennings S. 2007. Tup1-Ssn6 and Swi-Snf remodelling activities influence long-
range chromatin organization upstream of the yeast SUC2 gene. *Nucleic Acids Res*
35:5520–5531. doi:10.1093/nar/gkm573
Flowers EB, Poole RJ, Tursun B, Bashllari E, Pe'er I, Hobert O. 2010. The Groucho ortholog
UNC-37 interacts with the short Groucho-like protein LSY-22 to control developmental
decisions in C. elegans. *Dev Camb Engl* **137**:1799–1805. doi:10.1242/dev.046219
Fukaki H, Tameda S, Masuda H, Tasaka M. 2002. Lateral root formation is blocked by a gain-
of-function mutation in the SOLITARY-ROOT/IAA14 gene of Arabidopsis. *Plant J Cell*
Mol Biol **29**:153–168.
Gala HP, Lanctot A, Jean-Baptiste K, Guiziou S, Chu JC, Zemke JE, George W, Queitsch C,
Cuperus JT, Nemhauser JL. 2020. A single cell view of the transcriptome during lateral
root initiation in Arabidopsis thaliana. *bioRxiv* 2020.10.02.324327.
doi:10.1101/2020.10.02.324327
Gasperowicz M, Otto F. 2005. Mammalian Groucho homologs: redundancy or specificity? *J Cell*
Biochem **95**:670–687. doi:10.1002/jcb.20476
Gibson DG, Young L, Chuang R-Y, Venter JC, Hutchison CA, Smith HO. 2009. Enzymatic
assembly of DNA molecules up to several hundred kilobases. *Nat Methods* **6**:343–345.
doi:10.1038/nmeth.1318
Gietz RD, Woods RA. 2002. Transformation of yeast by lithium acetate/single-stranded carrier
DNA/polyethylene glycol method. *Methods Enzymol* **350**:87–96. doi:10.1016/s0076-
6879(02)50957-5
Gonzalez N, Pauwels L, Baekelandt A, De Milde L, Van Leene J, Besbrugge N, Heyndrickx KS,
Cuéllar Pérez A, Durand AN, De Clercq R, Van De Slijke E, Vanden Bossche R,
Eeckhout D, Gevaert K, Vandepoele K, De Jaeger G, Goossens A, Inzé D. 2015. A
Repressor Protein Complex Regulates Leaf Growth in Arabidopsis. *Plant Cell* **27**:2273–
2287. doi:10.1105/tpc.15.00006
Goodin MM, Dietzgen RG, Schichnes D, Ruzin S, Jackson AO. 2002. pGD vectors: versatile
tools for the expression of green and red fluorescent protein fusions in agroinfiltrated
plant leaves. *Plant J Cell Mol Biol* **31**:375–383. doi:10.1046/j.1365-313x.2002.01360.x
Grbavec D, Lo R, Liu Y, Stifani S. 1998. Transducin-like Enhancer of split 2, a mammalian
homologue of Drosophila Groucho, acts as a transcriptional repressor, interacts with
Hairy/Enhancer of split proteins, and is expressed during neuronal development. *Eur J*
Biochem **258**:339–349.
Gromöller A, Lehming N. 2000. Srb7p is a physical and physiological target of Tup1p. *EMBO J*
19:6845–6852. doi:10.1093/emboj/19.24.6845
Hallberg M, Hu G-Z, Tronnersjö S, Shaikhibrahim Z, Balciunas D, Björklund S, Ronne H. 2006.
Functional and physical interactions within the middle domain of the yeast mediator. *Mol*
Genet Genomics MGG **276**:197–210. doi:10.1007/s00438-006-0135-7
Harrison SJ, Mott EK, Parsley K, Aspinall S, Gray JC, Cottage A. 2006. A rapid and robust
method of identifying transformed Arabidopsis thaliana seedlings following floral dip
transformation. *Plant Methods* **2**:19. doi:10.1186/1746-4811-2-19
Haruki H, Nishikawa J, Laemmli UK. 2008. The anchor-away technique: rapid, conditional
establishment of yeast mutant phenotypes. *Mol Cell* **31**:925–932.
doi:10.1016/j.molcel.2008.07.020
Havens KA, Guseman JM, Jang SS, Pierre-Jerome E, Bolten N, Klavins E, Nemhauser JL.
2012. A synthetic approach reveals extensive tunability of auxin signaling. *Plant Physiol*
160:135–142. doi:10.1104/pp.112.202184
Ito J, Fukaki H, Onoda M, Li L, Li C, Tasaka M, Furutani M. 2016. Auxin-dependent
compositional change in Mediator in ARF7- and ARF19-mediated transcription. *Proc*
Natl Acad Sci U S A **113**:6562–6567. doi:10.1073/pnas.1600739113

Kagale S, Links MG, Rozwadowski K. 2010. Genome-wide analysis of ethylene-responsive
element binding factor-associated amphiphilic repression motif-containing transcriptional
regulators in Arabidopsis. *Plant Physiol* **152**:1109–1134. doi:10.1104/pp.109.151704
Ke J, Ma H, Gu X, Thelen A, Brunzelle JS, Li J, Xu HE, Melcher K. 2015. Structural basis for
recognition of diverse transcriptional repressors by the TOPLESS family of corepressors.
Sci Adv **1**:e1500107. doi:10.1126/sciadv.1500107
Keleher CA, Redd MJ, Schultz J, Carlson M, Johnson AD. 1992. Ssn6-Tup1 is a general
repressor of transcription in yeast. *Cell* **68**:709–719. doi:10.1016/0092-8674(92)90146-4
Khakhar A, Leydon AR, Lemmex AC, Klavins E, Nemhauser JL. 2018. Synthetic hormone-
responsive transcription factors can monitor and re-program plant development. *eLife* **7**.
doi:10.7554/eLife.34702
Kim MH, Cooper DR, Oleksy A, Devedjiev Y, Derewenda U, Reiner O, Otlewski J, Derewenda
ZS. 2004. The Structure of the N-Terminal Domain of the Product of the Lissencephaly
Gene Lis1 and Its Functional Implications. *Structure* **12**:987–998.
doi:10.1016/j.str.2004.03.024
Kim YJ, Björklund S, Li Y, Sayre MH, Kornberg RD. 1994. A multiprotein mediator of
transcriptional activation and its interaction with the C-terminal repeat domain of RNA
polymerase II. *Cell* **77**:599–608. doi:10.1016/0092-8674(94)90221-6
Klepikova AV, Kasianov AS, Gerasimov ES, Logacheva MD, Penin AA. 2016. A high resolution
map of the Arabidopsis thaliana developmental transcriptome based on RNA-seq
profiling. *Plant J* **88**:1058–1070. doi:10.1111/tpj.13312
Kornberg RD. 2005. Mediator and the mechanism of transcriptional activation. *Trends Biochem
Sci* **30**:235–239. doi:10.1016/j.tibs.2005.03.011
Krogan NT, Hogan K, Long JA. 2012. APETALA2 negatively regulates multiple floral organ
identity genes in Arabidopsis by recruiting the co-repressor TOPLESS and the histone
deacetylase HDA19. *Dev Camb Engl* **139**:4180–4190. doi:10.1242/dev.085407
Kumar KRR, Blomberg J, Björklund S. 2018. The MED7 subunit paralogs of Mediator function
redundantly in development of etiolated seedlings in Arabidopsis. *Plant J* **96**:578–594.
doi:10.1111/tpj.14052
Labun K, Montague TG, Krause M, Torres Cleuren YN, Tjeldnes H, Valen E. 2019.
CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. *Nucleic
Acids Res* **47**:W171–W174. doi:10.1093/nar/gkz365
Laplaze L, Parizot B, Baker A, Ricaud L, Martinière A, Auguy F, Franche C, Nussaume L,
Bogusz D, Haseloff J. 2005. GAL4-GFP enhancer trap lines for genetic manipulation of
lateral root development in Arabidopsis thaliana. *J Exp Bot* **56**:2433–2442.
doi:10.1093/jxb/eri236
Lee JE, Golz JF. 2012. Diverse roles of Groucho/Tup1 co-repressors in plant growth and
development. *Plant Signal Behav* **7**:86–92. doi:10.4161/psb.7.1.18377
Lee M, Chatterjee S, Struhl K. 2000. Genetic analysis of the role of Pol II holoenzyme
components in repression by the Cyc8-Tup1 corepressor in yeast. *Genetics* **155**:1535–
1542.
Liu X, Galli M, Camehl I, Gallavotti A. 2019. RAMOSA1 ENHANCER LOCUS2-Mediated
Transcriptional Repression Regulates Vegetative and Reproductive Architecture. *Plant
Physiol* **179**:348–363. doi:10.1104/pp.18.00913
Liu Z, Karmarkar V. 2008. Groucho/Tup1 family co-repressors in plant development. *Trends
Plant Sci* **13**:137–144. doi:10.1016/j.tplants.2007.12.005
Long JA, Ohno C, Smith ZR, Meyerowitz EM. 2006. TOPLESS regulates apical embryonic fate
in Arabidopsis. *Science* **312**:1520–1523. doi:10.1126/science.1123841
Ma H, Duan J, Ke J, He Y, Gu X, Xu T-H, Yu H, Wang Y, Brunzelle JS, Jiang Y, Rothbart SB,
Xu HE, Li J, Melcher K. 2017. A D53 repression motif induces oligomerization of

TOPLESS corepressors and promotes assembly of a corepressor-nucleosome complex. *Sci Adv* **3**:e1601217. doi:10.1126/sciadv.1601217

Maji S, Dahiya P, Waseem M, Dwivedi N, Bhat DS, Dar TH, Thakur JK. 2019. Interaction map of Arabidopsis Mediator complex expounding its topology. *Nucleic Acids Res* **47**:3904–3920. doi:10.1093/nar/gkz122

Malik N, Agarwal P, Tyagi A. 2017. Emerging functions of multi-protein complex Mediator with special emphasis on plants. *Crit Rev Biochem Mol Biol* **52**:475–502. doi:10.1080/10409238.2017.1325830

Martin K, Kopperud K, Chakrabarty R, Banerjee R, Brooks R, Goodin MM. 2009. Transient expression in Nicotiana benthamiana fluorescent marker lines provides enhanced definition of protein localization, movement and interactions in planta. *Plant J Cell Mol Biol* **59**:150–162. doi:10.1111/j.1365-313X.2009.03850.x

Martin-Arevalillo R, Nanao MH, Larrieu A, Vinos-Poyo T, Mast D, Galvan-Ampudia C, Brunoud G, Vernoux T, Dumas R, Parcy F. 2017. Structure of the Arabidopsis TOPLESS corepressor provides insight into the evolution of transcriptional repression. *Proc Natl Acad Sci U S A*. doi:10.1073/pnas.1703054114

Matsumura H, Kusaka N, Nakamura T, Tanaka N, Sagegami K, Uegaki K, Inoue T, Mukai Y. 2012. Crystal structure of the N-terminal domain of the yeast general corepressor Tup1p and its functional implications. *J Biol Chem* **287**:26528–26538. doi:10.1074/jbc.M112.369652

McIsaac RS, Oakes BL, Wang X, Dummit KA, Botstein D, Noyes MB. 2013. Synthetic gene expression perturbation systems with rapid, tunable, single-gene specificity in yeast. *Nucleic Acids Res* **41**:e57. doi:10.1093/nar/gks1313

Mitchell LA, Chuang J, Agmon N, Khunsriraksakul C, Phillips NA, Cai Y, Truong DM, Veerakumar A, Wang Y, Mayorga M, Blomquist P, Sadda P, Trueheart J, Boeke JD. 2015. Versatile genetic assembly system (VEGAS) to assemble pathways for expression in S. cerevisiae. *Nucleic Acids Res* **43**:6620–6630. doi:10.1093/nar/gkv466

Mottis A, Mouchiroud L, Auwerx J. 2013. Emerging roles of the corepressors NCoR1 and SMRT in homeostasis. *Genes Dev* **27**:819–835. doi:10.1101/gad.214023.113

Nozawa K, Schneider TR, Cramer P. 2017. Core Mediator structure at 3.4 Å extends model of transcription initiation complex. *Nature* **545**:248–251. doi:10.1038/nature22328

Oberoi J, Fairall L, Watson PJ, Yang J-C, Czimmerer Z, Kampmann T, Goult BT, Greenwood JA, Gooch JT, Kallenberger BC, Nagy L, Neuhaus D, Schwabe JWR. 2011. Structural basis for the assembly of the SMRT/NCoR core transcriptional repression machinery. *Nat Struct Mol Biol* **18**:177–184. doi:10.1038/nsmb.1983

Papamichos-Chronakis M, Conlan RS, Gounalaki N, Copf T, Tzamarias D. 2000. Hrs1/Med3 is a Cyc8-Tup1 corepressor target in the RNA polymerase II holoenzyme. *J Biol Chem* **275**:8397–8403. doi:10.1074/jbc.275.12.8397

Perissi V, Jepsen K, Glass CK, Rosenfeld MG. 2010. Deconstructing repression: evolving models of co-repressor action. *Nat Rev Genet* **11**:109–123. doi:10.1038/nrg2736

Petrenko N, Jin Y, Wong KH, Struhl K. 2017. Evidence that Mediator is essential for Pol II transcription, but is not a required component of the preinitiation complex in vivo. *eLife* **6**:e28447. doi:10.7554/eLife.28447

Pierre-Jerome E, Jang SS, Havens KA, Nemhauser JL, Klavins E. 2014. Recapitulation of the forward nuclear auxin response pathway in yeast. *Proc Natl Acad Sci U S A* **111**:9407–9412. doi:10.1073/pnas.1324147111

Pierre-Jerome E, Wright RC, Nemhauser JL. 2017. Characterizing Auxin Response Circuits in Saccharomyces cerevisiae by Flow Cytometry. *Methods Mol Biol Clifton NJ* **1497**:271–281. doi:10.1007/978-1-4939-6469-7_22

Raya-González J, Oropeza-Aburto A, López-Bucio JS, Guevara-García ÁA, de Veylder L, López-Bucio J, Herrera-Estrella L. 2018. MEDIATOR18 influences Arabidopsis root

architecture, represses auxin signaling and is a critical factor for cell viability in root meristems. *Plant J Cell Mol Biol* **96**:895–909. doi:10.1111/tpj.14114

Raya-González J, Ortiz-Castro R, Ruíz-Herrera LF, Kazan K, López-Bucio J. 2014. PHYTOCHROME AND FLOWERING TIME1/MEDIATOR25 Regulates Lateral Root Formation via Auxin Signaling in Arabidopsis. *Plant Physiol* **165**:880–894. doi:10.1104/pp.114.239806

Roeder RG. 1996. The role of general initiation factors in transcription by RNA polymerase II. *Trends Biochem Sci* **21**:327–335.

Samanta S, Thakur JK. 2015. Importance of Mediator complex in the regulation and integration of diverse signaling pathways in plants. *Front Plant Sci* **6**:757. doi:10.3389/fpls.2015.00757

Sato S, Tomomori-Sato C, Tsai K-L, Yu X, Sardiu M, Saraf A, Washburn MP, Florens L, Asturias FJ, Conaway RC, Conaway JW. 2016. Role for the MED21-MED7 Hinge in Assembly of the Mediator-RNA Polymerase II Holoenzyme. *J Biol Chem* **291**:26886–26898. doi:10.1074/jbc.M116.756098

Schilbach S, Hantsche M, Tegunov D, Dienemann C, Wigge C, Urlaub H, Cramer P. 2017. Structures of transcription pre-initiation complex with TFIIH and Mediator. *Nature* **551**:204–209. doi:10.1038/nature24282

Schreiber-Agus N, Chin L, Chen K, Torres R, Rao G, Guida P, Skoultchi AI, DePinho RA. 1995. An amino-terminal domain of Mxi1 mediates anti-Myc oncogenic activity and interacts with a homolog of the yeast transcriptional repressor SIN3. *Cell* **80**:777–786.

Siligato R, Wang X, Yadav SR, Lehesranta S, Ma G, Ursache R, Sevilem I, Zhang J, Gorte M, Prasad K, Wrzaczek M, Heidstra R, Murphy A, Scheres B, Mähönen AP. 2016. MultiSite Gateway-Compatible Cell Type-Specific Gene-Inducible System for Plants. *Plant Physiol* **170**:627–641. doi:10.1104/pp.15.01246

Song YH, Estrada DA, Johnson RS, Kim SK, Lee SY, MacCoss MJ, Imaizumi T. 2014. Distinct roles of FKF1, GIGANTEA, and ZEITLUPE proteins in the regulation of CONSTANS stability in Arabidopsis photoperiodic flowering. *Proc Natl Acad Sci* **111**:17672–17677. doi:10.1073/pnas.1415375111

Trumbly RJ. 1992. Glucose repression in the yeast Saccharomyces cerevisiae. *Mol Microbiol* **6**:15–21. doi:10.1111/j.1365-2958.1992.tb00832.x

Tzamarias D, Struhl K. 1994. Functional dissection of the yeast Cyc8-Tup1 transcriptional co-repressor complex. *Nature* **369**:758–761. doi:10.1038/369758a0

Ulmasov T, Hagen G, Guilfoyle TJ. 1997. ARF1, a transcription factor that binds to auxin response elements. *Science* **276**:1865–1868. doi:10.1126/science.276.5320.1865

Wang Z-P, Xing H-L, Dong L, Zhang H-Y, Han C-Y, Wang X-C, Chen Q-J. 2015. Egg cell-specific promoter-controlled CRISPR/Cas9 efficiently generates homozygous mutants for multiple target genes in Arabidopsis in a single generation. *Genome Biol* **16**:144. doi:10.1186/s13059-015-0715-0

Weber E, Engler C, Gruetzner R, Werner S, Marillonnet S. 2011. A Modular Cloning System for Standardized Assembly of Multigene Constructs. *PLOS ONE* **6**:e16765. doi:10.1371/journal.pone.0016765

Wong KH, Struhl K. 2011. The Cyc8-Tup1 complex inhibits transcription primarily by masking the activation domain of the recruiting protein. *Genes Dev* **25**:2525–2539. doi:10.1101/gad.179275.111

Wong MM, Guo C, Zhang J. 2014. Nuclear receptor corepressor complexes in cancer: mechanism, function and regulation. *Am J Clin Exp Urol* **2**:169–187.

Wright RC, Bolten N, Pierre-Jerome E. 2019. flowTime: Annotation and analysis of biological dynamical systems using flow cytometry. Bioconductor version: Release (3.10). doi:10.18129/B9.bioc.flowTime

Yang Y, Li R, Qi M. 2000. In vivo analysis of plant promoters and transcription factors by
agroinfiltration of tobacco leaves. *Plant J* **22**:543–551. doi:10.1046/j.1365-
313x.2000.00760.x
Zhang Y, Gao S, Wang Z. 2010. Structural and functional analysis of amino-terminal enhancer
of split in androgen-receptor-driven transcription. *Biochem J* **427**:499–511.
doi:10.1042/BJ20091631
Zhang Z, Reese JC. 2004. Redundant Mechanisms Are Used by Ssn6-Tup1 in Repressing
Chromosomal Gene Transcription in Saccharomyces cerevisiae. *J Biol Chem*
279:39240–39250. doi:10.1074/jbc.M407159200
Zhu J, Jeong JC, Zhu Y, Sokolchik I, Miyazaki S, Zhu J-K, Hasegawa PM, Bohnert HJ, Shi H,
Yun D-J, Bressan RA. 2008. Involvement of Arabidopsis HOS15 in histone deacetylation
and cold tolerance. *Proc Natl Acad Sci* **105**:4945–4950. doi:10.1073/pnas.0801029105

1162
1163
1164
1165

1166
1167 **Figure 1 – Figure Supplement 1. Helix 1 and the CRA domain (Helix 3-Helix 8) can act**

1168 **independently to repress transcription**. Each color represents two independent time course

1169 flow cytometry experiments of the TPL helices indicated, all fused to IAA3 or IAA3 alone (black

1170 line). Every point represents the average fluorescence of 5-10,000 individually measured yeast

1171 cells (a.u. - arbitrary units). Auxin (IAA-10μM) was added at the indicated time (gray bar, + Aux).

1172
1173

1174

Figure 2 – Figure Supplement 1. The TPL-N terminal domain (TPLN188) interacts with the

N-terminus of AtMED21. **A.** Identifying TPL-N terminal domain interactor proteins through

Yeast Two Hybrid screening identifies TPL as a problematic Bait protein, as it may silence the

activation despite successful binding of a prey protein (see second row from bottom where

N188 and MED21 show very weak reporter activity on 3AT). 3AT - 3-Amino-1,2,4-triazole.

Plates were measured after 3 days to allow TPL-MED21 interactions to grow. **B.** Identifying

TPL-N terminal domain interactor proteins through cytoplasmic split ubiquitin protein interaction

assay. We tested the N-terminal and C-terminal portions of MED13 separately and divided the

coding sequence at amino acid 967 (MED13N = aa1-967, MED13C = aa968-1908). Each bait

tested is the *Arabidopsis* homolog cloned from cDNA from the Col-0 accession, with the

exception of *AtMED13*, which was synthesized *de novo* via Twist

(https://www.twistbioscience.com/). Plates were scanned at 3 days after plating to allow weaker

interactions to develop if they were present. **C.** TPL interacts with MED21 through an interaction

within Helices 8-9. Plates were scanned at 2 days after plating. **D.** The Helix 8 Quadruple

mutation (V145A, E146A, K148A, K149A) does not affect AtMED10B binding to TPL. Plates

were scanned at 3 days after plating. **B-D.** The relative position of the N-terminal portion of

ubiquitin (Nub) is indicated for each bait protein.

1192

Figure 2 – Figure Supplement 2. Homology and structure of the MED21 subunit of the

mediator complex. A. Protein levels of AtMED21 cytoSUS constructs in yeast. Two different

exposure times are shown to demonstrate the lower abundance of the truncation with only the

first 31 amino acids of the AtMED21 (N31). Asterisks indicate the size predicted for the

indicated protein. **B.** Protein alignment of selected MED21 homologs from various species. Dr -

Drosophila melanogaster, Tr - *Takifugu rubripes*, Ms - *Mus musculus*, Gg - *Gallus gallus*, Hs -

Homo sapiens, Sp - *Strongylocentrotus purpuratus*, Sc - *Saccharomyces cerevisiae*, At -

Arabidopsis thaliana, Os - *Oryza sativa*, Sm - *Selaginella moellendorffii*, Pp - *Physcomitrella*

patens. Alignment was performed in CLC sequence viewer 7, using a neighbor joining method.

C-D. Structure of the MED21 (Cyan) & MED7 (Blue) hetero dimer, adapted from 1YKH, (Baumli

et al., 2005). The amino acids in the N-terminus that were solved are highlighted in red up to the

7th amino acid of the yeast MED21. **C.** The cartoon visualization, **D.** Surface visualization. **E.**

Core mediator (5N9J, (Nozawa et al., 2017)) with the location of MED21 and MED7 indicated

with the same colors from (C-D). In this structure the location of the MED21 N-terminus is again

indicated in red, demonstrating its close proximity to the Knob region (dotted circle).

1208

A Repressor/Substrate (VA1-pTDH3-TPLN-IAA14-ttACS2-VA3) F-Box Receptor (VA3-pRPS2-AFB2-ttCIT1-VA4) Selectable Prototrophy (VA4-LEU2-VA5) Transcription Factor (VA5-pADH1-ARF19-ttADH1-VA6) Reporter (VA6-pP3_2X-Venus-ttCYC1-VA2)

B Anchor Away Parental Strain: MATα tor1-1 fpr1::NAT RPL13A-2×FKBP12::TRP1

SPARC^IAA14 — IAA14, AFB2, ARF19, p3-2x-UbiVenus

SPARC^N188 — TPLN188-IAA14, AFB2, ARF19, p3-2x-UbiVenus,

SPARC^H1-5 — TPLH1-H5-IAA14, AtAFB2, ARF19, p3-2x-UbiVenus

1209

Figure 4 – Figure Supplement 1. Construction and characterization of the Single Locus

Auxin Response Circuit (SPARC). A. Design schematic of the approach utilized to create the

SPARC through a VEGAS assembly approach. Each individual transcriptional unit (TU) was

checked to replace promoters or terminators that utilized identical sequences and replaced with

an alternative sequence indicated by a purple underline. These TUs were assembled into level

1 plasmids by Golden Gate reaction. Subsequently, they were amplified by PCR using primers

specific for the Vegas Adaptor (VA) sequences specific for their TU cassette. In example, for the

first Repressor/Substrate TU the TU was amplified using primers for VA1 and VA3 and purified

by a PCR cleanup column (NEB). The acceptor plasmid was cut with EcoRI and both TU and

acceptor plasmid was transformed into yeast and recombinant plasmids were selected on

synthetic drop out (SDO) plates lacking Leucine. **B.** Primary SPARC transformants were struck

out onto fresh SDO -Leu and imaged for Venus expression, demonstrating varying levels of

reporter expression that correlate with TPL repressor domains. Plasmid DNA was purified from

these strains and sequenced to confirm the proper recombination of TU cassettes. **C.** Time

course flow cytometry of SPARC strains following auxin addition. For all cytometry experiments,

the indicated TPL construct is fused to IAA14, because this IAA works better in haploid yeast

strains that IAA3. Every point represents the average fluorescence of 5-10,000 individually

measured yeast cells (a.u. - arbitrary units). Auxin (IAA-10μM) was added at the indicated time

(gray bar, + Aux). Four independent experiments are shown for each construct. **D.** The yeast

TPL homolog Tup1 and its partner protein Cyc8 do not repress the SPARC. Quantified

fluorescence from the single locus auxin response circuit (SPARC) introduced into Tup1 and

Cyc8 anchor away lines demonstrates no increased fluorescence from the reporter upon

depletion of Tup1 or Cyc8 from the nucleus. Anchor away depletion of Tup1 or Cyc8 results in

slower yeast growth. To normalize for this disparity in growth, Venus fluorescence was

normalized to red autofluorescence, where each pixel was normalized to the corresponding red

autofluorescence collected for that position and plotted as a boxplot. Two individual biological

replicates (two separate experiments) were evaluated, and the data was pooled. **E-F.** Med21 N

terminal deletions are viable in *Saccharomyces* and demonstrate altered SPARC transcriptional

states. **E.** A representative grayscale image of fellow fluorescence of spot plates of yeast strains

carrying SPARC plasmids in Med21 N-terminal deletions. Each is plated at an OD600 of 0.1 on

SDO with or without auxin (10μM IAA). **F.** Venus fluorescence from (**E**) was normalized to red

background (autofluorescence), where each pixel was normalized to the corresponding red

autofluorescence collected for that position and plotted as a boxplot. Two individual biological

replicates (two separate experiments) were evaluated, and the data was pooled and is

presented as boxplots.

1245

1246

Figure 4 – Figure Supplement 2. N-terminal ScMed21 deletions impair auxin responsive

transcriptional activation A. Time-course flow cytometry analysis of SPARC[H1-H5] in wild type

and n-terminal ScMed21 deletions with and without auxin. Genotypes are indicated in the

colored key inset into the graph. Auxin (IAA-10μM) was added at the indicated time (gray bar, +

Aux). **B**. Conversion of the first five amino acids of ScMed21 to the corresponding sequence

from AtMED21 results in an identical repression profile. Time course flow cytometry of SPARC

strains following auxin addition. For all cytometry experiments, the indicated TPL construct is

fused to IAA14, because this IAA performs better in haploid yeast strains that IAA3. Every point

represents the average fluorescence of 5-10,000 individually measured yeast cells (a.u. -

arbitrary units). Auxin (IAA-10µM) was added at the indicated time (gray bar, + Aux). Two

independent experiments are shown for each construct. **C.** Cell growth of the strains in (**B**)

indicate the swap of the N-terminal region had no effect on yeast growth or viability. Data

presented is events per microliter over the time-course of the cytometry experiments.

A.

MED21   MED21-FRB   Δ3-MED21-FRB   Δ5-MED21-FRB   Δ7-MED21-FRB

SSC.A

-Rapa
+Rapa

FSC.H

B.

+ Rapamycin

RPL13A — FKBP
FRB
MED21

RPL13A — MED21

Nuclear
export

Auxin Promoter — Venus

MED21-FLAG — Inducible Promoter

+β-estradiol

C.

-Rapa
-β-Estradiol   +Rapa,
                 +β-Estradiol

iMED21-FLAG   iΔ3MED21-FLAG   iΔ5MED21-FLAG

FSC.H

D.

SPARC^TPLN188

+Mock   +Rapa, + β-estradiol

Fluorescence (a.u. x10,000)

a   a   b   a   b   c

iMED21-FLAG: Wt   Δ3   Δ5   Wt   Δ3   Δ5
+ MED21-FRB

E.

Med21-FRB   Wt   Δ3   Δ5   Δ7

αFRB

αPGK1

F.

Med21-FLAG   Wt   Δ3   Δ5   Wt   Δ3   Δ5
β-estradiol   -   -   -   +   +   +

αFLAG

αPGK1

G.

MED21
MED21-FRB
Δ3-MED21-FRB
Δ5-MED21-FRB

SDO

+Rapa

Count Distribution (Density)

Fluorescence (a.u.)

H.

MED21-FRB +
iMED21-FLAG
iΔ3-MED21-FLAG
iΔ5-MED21-FLAG

SDO

+Rapa
+β-estra

Fluorescence (a.u.)

1260

Figure 4 – Figure Supplement 3. Inducible MED21 rescues rapamycin induced yeast

growth defects. A. Depletion of nuclear ScMed21 by Rapamycin increased cell size even in

short time-courses, consistent with its essential role in many core pathways. Scatterplots of side

scatter area by forward scatter height (SSC.A x FSC.H) indicate large scale increases in cell

size in populations of yeast with (blue) or without (red) Rapamycin treatment. **B.** Schematic of

AtARCSc combined with methods for inducible expression and nuclear depletion of MED21. In

Anchor Away, the yeast ribosomal protein 13A (RPL13A) is fused to the rapamycin binding

protein FKBP. Addition of Rapamycin induces dimerization between FKBP and any target

protein fused to FRB, resulting in removal of the target protein from the nucleus. For these

experiments, AtARCSc was assembled into a single plasmid (SPARC) rather than being

integrated into separate genomic loci (Figure 4 – Figure Supplement 1). Estradiol-inducible

ScMed21 (iMed21) made it possible to replace wild-type MED21 with targeted deletions or

mutations. **C.** Inducible Med21 (iMed21) wild type and variants cell size were examined before

(red) and after (blue) treatment with Rapamycin and b-estradiol to simultaneously deplete the

wild-type Med21-FRB fusion, and induce the transcription of the Med21 variant. Scatterplots of

side scatter area by forward scatter height (SSC.A x FSC.H) demonstrate a less disrupted cell

size compared to Anchor Away strains in (**A**). **D.** Rapid replacement of Med21-FRB with

inducible Med21-FLAG demonstrated the requirement for the ScMed21 N-terminus in TPL

repression. iMed21 isoforms were induced by addition of β-estradiol (20μM) for 4 hours followed

by Rapamycin addition. Fluorescence was quantified by cytometry after 300 minutes. Lower

case letters indicate significant difference (ANOVA and Tukey HSD multiple comparison test;

$p < 0.001$). **E-F.** Protein abundance of ScMed21 variants was tested by SDS-PAGE & western

blot **E.** Stably integrated Med21-FRB variants. **F.** Induced iMed21-FLAG variants. **G-H.**

Histograms of Venus fluorescence in inducible Med21 (iMed21) strains demonstrate that

populations were evenly distributed around a single mean, suggesting we were observing the

immediate effects of the Med21 deletions. The histograms were built using ggplots Density

function to create a visualization of count distribution. These samples were tested at 300

minutes (as in Figure 4D) and plotted to visualize cells at the equivalent stage of growth, Med21

depletion, and induction.

1290

1291

1292

1293

Figure 5 – Figure Supplement 1. TPL multimerization requires Helix 8. A. Cytoplasmic split

ubiquitin interaction (cytoSUS) assay on serial deletions of TPL. Interaction of bait and prey

proteins reconstitute split ubiquitin, release a synthetic transcription factor that allows growth on

media lacking Histidine and Adenine. The expression level of the bait protein can be repressed

through increased Methionine in the media. **B**. Protein levels of Nub-TPL fusions were tested by

PAGE and western blotting for the c-terminal 3xHA epitope tag included in all constructs.

Deletions longer than H1-4 are detectable at higher levels (left panel), whereas shorter isoforms

required longer exposure times to detect (right panel). Total protein loading levels were tested

by the housekeeping gene PGK1 (bottom panel). **C.** Protein expression analysis by western

blotting of tetramerization mutants expressed in yeast for cytoSUS interaction assay in Figure

5I. Prey constructs are C-terminally fused to 2xHA. Total protein loading levels were tested by

blotting against the housekeeping gene PGK1 (bottom panel).

1306
1307

Figure 6 – Figure Supplement 1. The TPL-MED21 interaction is required for repression in

plants. A. MED21 and TPL Co-Immunoprecipitated with AtMED10B from tobacco extracts.

Each construct was expressed under the viral 35S promoter, and tissues were harvested after 2

days of injection. MED10B was purified by incubation with IgG sepharose beads (see methods),

and the presence of interacting proteins was determined by western blotting. Actin was used as

a control to determine the efficacy of washing. **B-C.** Engineering and prototyping a variant of

TPLN-IAA14^{mED} which carries mutations in the EAR domain (EAR^{AAA}) and in the degron

(P306S) in yeast. **B.** Cartoon schematic of the mutations tested during prototyping of the

TPLN188-IAA14mED construct. In each case the identical glycine-serine linker (GS) was used

as the flexible linker between the 2xHA-TPLN188 protein and the portion of IAA14 retained in

the construct. **C.** Time course flow cytometry of TPLN-IAA14^{mED} strains following auxin addition.

Strains containing the TPLN-IAA14^{mED} was tested in both haploid and diploid strains and

demonstrated similar repression profiles. Every point represents the average fluorescence of 5-

10,000 individually measured yeast cells (a.u. - arbitrary units). Auxin (IAA-10μM) was added at

the indicated time (gray bar, + Aux). Two independent experiments are shown for each

construct. **D.** Transient expression of indicated TPL constructs in tobacco. DR5:Venus - the

synthetic DR5 auxin promoter (Ulmasov et al., 1997) driving Venus, ARF19 - p35S:AtARF19-

1xFLAG, GAL4:VP16 - pUBQ10:GAL4-VP16, TPLN-X-UAS-TPL-IAA14^{mED} with various TPL

truncations or mutations. **E-F.** The TPL-N terminus functions as a repressor independently of

the MED13 component of the Mediator CDK8 Kinase module in *Arabidopsis*. The dominant

lateral root repression constructs (UAS-TPLN188-IAA14, UAS-Monomer-IAA14) and the

dominant *slr* mutant were crossed to the *med13* mutant (*gct-5*, a.k.a. *mab2-2*, *SAIL_1169H11*).

We performed time-course lateral root density assays and genotyped the progeny in the F1 and

F2 generations. We observed a ~1% transmission rate of *gct-5* homozygotes. **E.** Time-course of

the mean of lateral root density over days 10-14 post germination plus and minus standard

error. **F.** Boxplots of lateral root density for day 14 (**E.**). Lower case letters indicate significant

difference (ANOVA and Tukey HSD multiple comparison test; $p < 0.001$).

1336

1337

1338

Figure 6 – Figure Supplement 2. The essential gene MED21 is required for normal lateral root development in plants. A. Identification and characterization of a novel CAS9-based insertional mutation in *MED21*. The *MED21* genomic locus (AT4G04780) is shown as a cartoon, with a zoom in on the beginning of the coding sequence highlighted with the amino acid sequence. The location of the *med21/MED21* mutant (WiscDsLox461-464K13, see triangle), and the sgRNA we employed (see green annotation and NGG PAM site) highlighted. The

insertion of a G at nucleotide position +214 after the transcriptional start site abrogates the

sgRNA site (red annotation above with i214G). A representative sequencing trace demonstrates

the position where the heterozygote carries i215G, and the predicted effect to the coding

sequence is shown at the top right – a red arrow indicates the first codon affected by the i214G

mutation. The Inset pictures at the top left demonstrate the embryo lethality phenotype in

med21^{i214G}/MED21 heterozygote siliques. White asterisks indicate the embryos that have begun

to degenerate. These aborted seeds are visibly brown indicating that fertilization took place

allowing the seed coat to form before development failed. **B.** *med21*/*MED21* heterozygotes are

haplo-sufficient for lateral root development. Lateral root density (number of lateral roots /

primary root length) was calculated at 10 days post germination. Lower case letters indicate

significant difference (ANOVA and Tukey HSD multiple comparison test; $p < 0.005$). **C.** MED21-

CAS9 repressors targeted to MED21 display increased lateral root densities. Lateral root

density (number of lateral roots / primary root length) was calculated at 10 days post

germination. Numbers below boxplots are p-values for pairwise comparisons with control using

a Wilcoxon rank sum test. **D.** Ratio of lateral root lengths to total root lengths (lateral root

lengths + primary root length) in dCAS9 repressor lines targeting *MED21* calculated at 10 days

post germination. Statistical tests (ANOVA and Wilcox test) are reported above the graph. **G.**

Representative root traces of dCAS9 repressor lines targeting *MED21* calculated at 10 days

post germination.

1364