**Title**

Structure-function analysis of *Arabidopsis* TOPLESS reveals fundamental conservation of

repression mechanisms across eukaryotes

**Authors**

Alexander R. Leydon[1], Wei Wang[2,†], Hardik P. Gala[1], Sabrina Gilmour[1], Samuel Juarez-Solis[1],

Mollye L. Zahler[1], Joseph E. Zemke[1], Ning Zheng[2,3], Jennifer L. Nemhauser[1]

[1]Department of Biology, University of Washington

[2]Department of Pharmacology & [3]Howard Hughes Medical Institute, University of Washington

[†]Present address: Key Laboratory of Plant Stress Biology, State Key Laboratory of Cotton

Biology, School of Life Science, Jinming Campus, Henan University, Kaifeng, Henan Province,

475004, PR of China.

## Summary

The plant corepressor TOPLESS (TPL) is recruited to a large number of loci that are selectively induced in response to developmental or environmental cues, yet the mechanisms by which it inhibits expression in the absence of these stimuli is poorly understood. Previously, we had used the N-terminus of *Arabidopsis thaliana* TPL to enable repression of a synthetic auxin response circuit in *Saccharomyces cerevisiae* (yeast). Here, we leveraged the yeast system to interrogate the relationship between TPL structure and function, specifically scanning for repression domains. We identified a potent repression domain in Helix 8 located within the CRA domain, which directly interacted with the Mediator middle domain subunits Med21 and Med10. Interactions between TPL and Mediator were required to fully repress transcription in both yeast and plants. In contrast, we found that multimer formation, a conserved feature of many corepressors, had minimal influence on the repression strength of TPL.

## Keywords

Corepressors, Transcriptional Repression, Mediator, Tup1, TOPLESS

**Introduction**

Control over gene expression is essential for life. This is especially evident during development when the switching of genes between active and repressed states drives fate determination. Mutations that interfere with repression lead to or exacerbate numerous cancers (Wong et al., 2014) and cause developmental defects in diverse organisms (Grbavec et al., 1998; Long et al., 2006), yet many questions remain about how cells induce, maintain, and relieve transcriptional repression. Transcriptional repression is controlled in part by a class of proteins known as corepressors that interact with DNA-binding transcription factors and actively recruit repressive machinery. Transcriptional corepressors are found in all eukaryotes and include the animal SMRT (silencing mediator of retinoic acid and thyroid hormone receptor) and NCoR (nuclear receptor corepressor) complexes (Mottis et al., 2013; Oberoi et al., 2011), the yeast Tup1 (Matsumura et al., 2012), and its homologs Drosophila Groucho (Gro) and mammalian transducing-like enhancer (TLE) (Agarwal et al., 2015).

In plants, TOPLESS (TPL), TOPLESS-RELATED (TPR1-4), LEUNIG (LUG) and its homolog (LUH), High Expression of Osmotically responsive genes 15 (HOS15) all act as Gro/TLE-type corepressors (Causier et al., 2012; Lee and Golz, 2012; Liu and Karmarkar, 2008; Long et al., 2006; Zhu et al., 2008). Defects in the TPL family have been linked to aberrant stem cell homeostasis (Busch et al., 2010), organ development (Gonzalez et al., 2015), and hormone signaling (Causier et al., 2012; Kagale et al., 2010), especially the plant hormone auxin (Long et al., 2006). Plant corepressors share a general structure, where at the N-terminus a LIS1 homology (LisH) domain contributes to protein dimerization (Delto et al., 2015; Kim et al., 2004). At the C-terminus, WD40 repeats form beta-propeller structures that are involved in protein-protein interactions (Collins et al., 2019; Liu et al., 2019). In TPL family corepressors, the LisH is followed by a C-terminal to LisH (CTLH) domain that binds transcriptional repressors through an Ethylene-responsive element binding factor-associated Amphiphilic Repression (EAR) motif found in partner proteins (Causier et al., 2012; Kagale et al., 2010). The N-terminal domain also contains a CT11-RanBPM (CRA) domain, which provides a second TPL dimerization interface and stabilizes the LisH domain (Ke et al., 2015; Martin-Arevalillo et al., 2017). While there is speculation that the beta-propellers may control protein interaction with other repressive machinery, it is well-established that they bind to the non-EAR TPL recruitment motifs found in transcriptional regulators (RLFGV- and DLN-type motifs, (Liu et al., 2019)).

We have previously demonstrated the recapitulation of the auxin response pathway in

*Saccharomyces cerevisiae* (yeast) by porting individual components of the *Arabidopsis* auxin

nuclear response (Pierre-Jerome et al., 2014). In this <u>A</u>rabidopsis <u>t</u>haliana <u>A</u>uxin <u>R</u>esponse

<u>C</u>ircuit in <u>S</u>accharomyces <u>c</u>erevisiae (*At*ARC$^{Sc}$), an auxin responsive transcription factor (ARF)

binds to a promoter driving a fluorescent reporter. In the absence of auxin, the ARF protein

activity is repressed by interaction with a full-length Aux/IAA protein fused to the N-terminal

domain of TPL. Upon the addition of auxin, the TPL-IAA fusion protein is targeted for

degradation through interaction with a member of the Auxin Signaling F-box protein family (TIR1

or AFB2), and releases the transcriptional repression of the fluorescent reporter. Reporter

activation can be quantified after auxin addition by microscopy or flow cytometry (Pierre-Jerome

et al., 2014). In the original build and characterization of *At*ARC$^{Sc}$, it was noted that the two N-

terminal truncations of TPL (N100 or N300) behave differently (Pierre-Jerome et al., 2014).

While both truncations are able to repress the function of a transcriptional activator fused to an

Aux/IAA, only the TPLN100 fusion shows alleviation of repression after auxin addition. TPLN300

fusions to Aux/IAAs maintain strong durable repression even under high concentrations of

auxin. This disparity is not due to differential rates of protein degradation, as both proteins

appear to be turned over with equal efficiency after auxin addition (Pierre-Jerome et al., 2014).

Structures of the N-terminal domains of TPL rice homolog OsTPR2 (Ke et al., 2015) and

the *Arabidopsis* TPL (Martin-Arevalillo et al., 2017) have recently been solved. These structures

reveal high conservation of protein folds in the N-terminus, as well as the residues that

coordinate formation of homotetramers (Figure 1A). Several lines of evidence suggest that the

multimeric TPL modulates repression potential. First, the dominant TPL mutant *tpl-1* altered a

single amino acid in the ninth helix of the TPL-N terminus (N176H) that induces aggregation of

TPL and its homologs (TPR1-4), reducing total activity (Long et al., 2006; Ma et al., 2017).

Second, TPL recruitment motifs found in the rice strigolactone signaling repressor D53 induce

higher-order oligomerization of the TPL N-terminus, which increases histone binding and

transcriptional repression (Ma et al., 2017). Third, structural studies of *Arabidopsis* TPL

demonstrated interdependency of the TPL tetramer formation and Aux/IAA binding (Martin-

Arevalillo et al., 2017). One contrary piece of evidence is the strong repressive activity of the

TPL N100 construct which lacks the majority of the CRA domain ((Martin-Arevalillo et al., 2017),

Figure 1A) and is unlikely able to form tetramers.

The conservation of TPL's repressive function in yeast suggests that the protein partners

that enact the repression are conserved across eukaryotes. Consistent with this speculation, the

series of alpha-helices that form the N-terminal portion of TPL is highly reminiscent of naturally-

occurring truncated forms of mammalian TLE (Gasperowicz and Otto, 2005), such as Amino-

terminal Enhancer of Split (AES) (Zhang et al., 2010), the Groucho ortholog LSY-22 (Flowers et

al., 2010), and the unrelated mouse repressor protein MXI1 (Schreiber-Agus et al., 1995).

Gro/TLE family members are generally considered to repress by recruiting histone deacetylases

to control chromatin compaction and availability for transcription (Chen and Courey, 2000; Long

et al., 2006). An alternative hypothesis has been described for Tup1 in yeast, where Tup1

blocks the recruitment of RNA Polymerase II (Pol-II) (Wong and Struhl, 2011), possibly through

contacts with Mediator complex subunits Med21 or Med3 (Gromöller and Lehming, 2000;

Papamichos-Chronakis et al., 2000). However, like many of these family members, multiple

repression mechanisms have been described for TPL at different genetic loci. For example, TPL

has been found to recruit the repressive CDK8 Mediator complex (Ito et al., 2016), chromatin

remodeling enzymes such as Histone Deacetylase 19 (HD19) (Long et al., 2006) and directly

bind to histone proteins (Ma et al., 2017).

Here, we leveraged the power of yeast genetics to interrogate the mechanism of TPL

repression. Using *At*ARC$^{Sc}$, we discovered that the N-terminal domain of TPL contains two

distinct repression domains that can act independently. We mapped the first, weaker repression

domain to the first 18 amino acids of the LisH domain (Helix 1), and the second, more potent

domain to Helix 8 which falls within the CRA domain. Full repression by Helix 8 required direct

interaction with the Mediator complex, through interactions with Med21 and Med10. This

interaction required the same Med21 residues that control transcriptional activation of Tup1-

regulated genes in yeast. In addition, we found that multimerization of TPL was not required for

repression in yeast or in plants. Our yeast results were validated with plant assays, and

extended to include evidence that interaction with the middle domain of Mediator was required

for TPL repression of the auxin genes governing lateral root development. Our findings point to

a conserved functional connection between Tup1/TPL corepressors and the Mediator complex

that together create a repressed state with the capacity for rapid activation.

**Results**

To understand how TPL represses transcription, we first sought to localize repressive activity

within the protein. In the *At*ARC[Sc], the extent of auxin-induced turnover of TPLN100 and

TPLN300 fusions appear similar, although neither are completely degraded (Pierre-Jerome et

al., 2014). One interpretation is that auxin addition increases the sensitivity of the assay to

detect subtle differences in the strength of repressive activity of each fusion protein by reducing

its relative concentration. In this light, we would argue that TPLN300 is a stronger repressor

than TPLN100. To further exploit this synthetic repression assay, we began by generating a

deletion series of the N-terminus guided by the available structural information (Figure 1A-B,

(Ke et al., 2015; Martin-Arevalillo et al., 2017)). We started with a TPLN188-IAA3 fusion protein

construct, which behaves identically to TPLN300 (Figure 1B, (Pierre-Jerome et al., 2014)), and

subsequently deleted each alpha helical domain starting with Helix 9 (constructs are named in

the format Helix x – Helix y or Hx-Hy). We found that Helix 8 was required for the maximum

level of repression activity and for the maintenance of repression after auxin addition (Figure

1C). All constructs lacking Helix 8 retained the ability to repress transcription, but this repression

was lifted in the presence of auxin (Figure 1C) as had been observed for the original TPLN100

construct (Pierre-Jerome et al., 2014). Further deletions revealed that including only the 18

amino acids of Helix 1 was sufficient to confer repression (H1, Figure 1D). To test whether Helix

8 activity depended on Helix 1, we tested an additional construct consisting solely of Helix 3

through Helix 9 (H3-H9, Figure 1D). This construct was also able to repress ARF activity, thus

demonstrating that both Helix 1 (LisH) and Helix 8 (CRA) can act independently of one another

(Figure 1D). To identify the minimal domain needed for Helix 8-based repression, we generated

new deletions (Figure 1B,E-F). Helix 8 and the following linker were not sufficient for repression

(Figure 1E), and removal of Helix 9 or of the linker between Helix 8 and Helix 9 slightly

increased sensitivity to auxin compared to TPLN188 (H1-H8Δ8L, Figure 1F). A deletion that

removed both the LisH and Helix 8 repression domains (H3-H7) was only able to weakly

repress reporter expression (Figure 1F). Together, these results demonstrate that Helix 1 and

Helix 8 could act as repression domains, and that the linker between Helix 8 and Helix 9 (which

folds over Helix 1) was required for repression following addition of auxin. Helix 1 alone in the

LisH domain was sufficient to act on its own as a modular repression domain. The repressive

activity of Helix 8 was only functional in the context of the larger Helix 3-Helix 8 truncation that

carries the CTLH domain and a portion of the CRA domain.

**Figure 1. The N-terminal domain of TPL contains two independent repression domains.**

**A.** TPL domains are LisH (LIS1 homology motif, blue), CTLH (C-terminal LisH motif, orange),

CRA (CT11-RanBPM, red - dimerization and green - foldback), and two WD40, beta-propeller

motifs (purple). N-terminal domains are indicated on the solved structure of the first 202 amino

acids ((Martin-Arevalillo et al., 2017), 5NQS). The termini of the TPL-N100 truncation used in

the original ARC$^{Sc}$ studies is indicated. **B.** Diagram indicating the structure of constructs

analyzed in experiments shown in subsequent panels. Repression Index (Rep.) is a scaled

measure of repression strength with 0 set to the level of repression observed with IAA3 and 10

set to the level of repression by TPLN188-IAA3. Auxin induction level (Aux.) indicates the fold

change difference between reporter expression before auxin addition (time zero) and at the end

of an experiment (~500 minutes) **C-F.** Helix 1 and the CRA domain (Helix 3-Helix 8) can act

independently to repress transcription. Each panel represents two independent time course flow

cytometry experiments of the TPL helices indicated, all fused to IAA3. Every point represents

the average fluorescence of 5-10,000 individually measured yeast cells (a.u. - arbitrary units).

Auxin (IAA-10µM) was added at the indicated time (gray bar, + Aux).

To determine which of the many known or predicted TPL-binding partners could mediate

the repression activity of Helix1 and Helix8, we identified known interactors with either TPL or

other Gro/TLE co-repressors, and then introduced the *Arabidopsis* homologs of these genes

into the cytoSUS system (Asseck and Grefen, 2018). We chose the cytoSUS system as it would

remove the interaction between target proteins to the cytoplasm, as we observed that anchoring

the TPL N-terminus had the capacity to repress transcriptional activation of both our synthetic

auxin reporter and the activation of yeast two hybrid prototrophy reporters (Figure 1,

Supplemental Figure 1A). Putative direct interactors include histone deacetylases (HDACs -

AtHDAC9, AtHDAC6, (Long et al., 2006)), Histone proteins (Histone H3, Histone H4, (Ma et al.,

2017)), and the Mediator components MED13 (AtMED13, (Ito et al., 2016)) and MED21, which

has been demonstrated to interact with Tup1, the yeast homolog of TPL (Gromöller and

Lehming, 2000). We did not observe any interactions between TPL-N188 and the HDACs HDA6

and HDA9; the histone protein AtHIS4; or the Mediator subunit AtMED13 (Figure 2A,

Supplemental Figure 1B). HDAC interaction with TPL has been previously hypothesized to

occur through indirect interactions with partner proteins (Krogan et al., 2012), however direct

interactions with histones and MED13 have been detected (Ito et al., 2016; Ma et al., 2017). The

absence of interaction between TPL-N188 and these proteins may be due to differences

between methods, or interaction interfaces in the C-terminal WD40 repeats.

Strong interaction was detected between TPL-N188 and AtMED21, a component of the

Mediator middle domain (Figure 2A). MED21 is one of the most highly conserved Mediator

subunits (Bourbon, 2008), and has a particularly highly conserved N-terminus (Supplemental

Figure 2A,C-E). In yeast, Tup1 interacts with the first 31 amino acids of ScMed21, with the first

seven amino acids being absolutely required (Gromöller and Lehming, 2000). We observed that

the equivalent truncation of AtMED21 (AtMED21-N31) was sufficient for interaction with TPL-

N188 (Figure 2A). We next created truncations of the N-terminal domain of AtMED21 to closely

match those that had been made in yeast (Figure 2B, Supplemental Figure 2B) where deletion

of the first five amino acids of ScMed21 (ScΔ5Med21) severely reduce the ability of the

Mediator complex to co-purify with Pol-II and CDK8 kinase complex (Sato et al., 2016).

Interaction between TPLN188 and AtMED21 similarly required the first five amino acids of

AtMED21 (Figure 2B), and this truncation did not significantly impact protein levels (Figure 2B).

We used cytoSUS screening to ask whether TPL-N188 interacted with other middle

domain Mediator components, and identified an interaction with AtMED10B (Figure 2C). There

are two MED10 isoforms in *Arabidopsis*; AtMED10A and AtMED10B with 76.5% amino acid

identity between isoforms, and *AtMED10B* is nearly always expressed at a higher level than

*AtMED10A* (Klepikova et al., 2016). Consistent with the cytoSUS results, TPLN188 was able to

pull down both AtMED21 and AtMED10B from yeast extracts (Figure 2D). AtMED21 interaction

was specific to the Helix8-based repression domain, as it interacted with TPLH3-H9 (Figure 2E),

and not TPLH1-H5 (Supplemental Figure 1C). A manual juxtaposition of the yeast Mediator

structure with the *Arabidopsis* TPL N-terminal structure shows that Helix 8 and 9 of TPL face

away from the tetramer, and are therefore optimally placed to interact with Mediator

components (Figure 2F).

**Figure 2. The Helix 8 repression domain of TPL directly interacts with AtMED21 and AtMED10B. A-C,E.** cytoSUS assays with candidate interacting proteins. Nub-3xHA is the N-terminal fragment of Ubiquitin expressed with no fusion protein and is used as a negative control. Each prey protein is from *Arabidopsis*. -WL, -ADE - dropout lacking Trp, Leu and Ade

(growth control), -WLMH, -ADE – dropout lacking Trp, Leu, His, Met, and Ade (selective media).

**B**. Alignments of the *Arabidopsis* (At) and *Saccharomyces* (Sc) MED21 proteins are shown

above cytoSUS assays with the same bait shown in (**A**). Western blots below the colonies

indicated that AtMED21 N-terminal Δ3 and Δ5 are well expressed in assay conditions. **C.**

cytoSUS assays with selected Mediator proteins in the middle module.  **D.** The TPL-ProteinA-TF

fusion protein can pull down TPL, AtMED21 and AtMED10B from yeast extracts using IgG-

beads. Detection of the VP16 transcriptional activator demonstrates enrichment of the fusion

protein (αVP16). Each prey protein is detected via the 3xHA tag (αHA), and efficacy of

purification was judged by PGK1 depletion (αPGK1). **E.** A TPL-N truncation lacking the LisH

domain (TPLH2-H9) could still interact with the AtMED21-N31 truncation. This bait construct

interacted with IAA3, but only minimally with the negative control (free Nub-3xHA). **F.**  Yeast

Mediator (bottom, 5N9J) and AtTPL (top, 5NQV) manually docked to compare relative domain

sizes and feasibility of a TPL-MED21-MED10B interaction. TPL Helix 8-9 is colored green.

MED21 is colored aqua, with the N-terminus colored red, and the EAR peptide in orange.

MED10 is colored teal, with the C-terminus colored purple.

To pinpoint which residues of Helix3-9 could coordinate repression through interaction

with MED21, we identified likely solution-facing amino acids from Helix 8, as this helix had the

greatest impact on repression strength (Figure 1C). We hypothesized that these amino acids

were not involved in stabilizing the hydrophobic interactions between intra-TPL helical domains

and might be available to interact with partner proteins. In the context of H3-H8-IAA3, eight

amino acids in Helix 8 were mutated to alanine (Figure 3A, light green residues). Repression

activity was assessed in the absence (Figure 3B) or presence (Figure 3C) of auxin. No single

amino acid was essential for repression (Figure 3B). Two mutations (R140A and K148A) slightly

increased baseline expression of the reporter (Figure 2B-C). All mutations, except E152A that

behaved similarly to controls, altered the stability of repression after auxin addition, either by

increasing (S138A,V145A, E146A, K149A) or decreasing (I142A) the final fluorescence level

(Figure 2C). Mutating E146 and K149 also increased the speed with which the reporter

responded to auxin (Figure 2C), suggesting that these two neighboring residues could be a

critical point of contact with co-repressive machinery. S138A had a small increase in auxin

sensitivity, while I142 reduced auxin sensitivity (Figure 2E). TPL/TPR corepressors are recruited

to transcription factors through an Ethylene-responsive element binding factor-associated

Amphiphilic Repression (EAR) motif (Causier et al., 2012; Kagale et al., 2010), which binds to

the TPL in a pocket adjacent to Helix 8 (Ke et al., 2015; Martin-Arevalillo et al., 2017). While

these two residues (S138A, I142) do not contact the conserved leucine residues of the EAR

motif (LxLxL), in the AtTPL structure, the C-terminal portion of the IAA27 EAR domain makes

contact with these residues (Martin-Arevalillo et al., 2017). As we are using a TPL-IAA fusion

protein, repression does not depend on EAR-TPL interaction, therefore making it difficult to fully

assess any role this interaction normally plays in recruiting repression machinery.

We tested whether the residues in Helix 8 that were required for repression (V145,

E146, K148, K149, Figure 3A-C) were also required for interaction with AtMED21. Single

alanine mutations of these four amino acids in the context of TPLN188 significantly reduced

interaction with AtMED21, while the quadruple mutation (here called Quad$^{AAAA}$) completely

abrogated AtMED21 binding (Figure 3D). These mutations had little effect on interaction with

AtMED10B (Supplemental Figure 1D). Additionally, we observed that most of these single

mutations decreased binding between TPL and IAA3, consistent with their position along the

EAR binding pocket (Figure 3D). When tested in the *At*ARC$^{Sc}$, TPLN188-Quad$^{AAAA}$ resembled

the repressive activity of wild-type N188 (red and black, Figure 3E), consistent with the

observation that Helix 1 is present in this construct and is sufficient for repression (Figure 1D).

Introduction of Quad$^{AAAA}$ mutations into the Helix 3 through Helix 8 context (H3-H8-Quad$^{AAAA}$)

largely phenocopied a Helix 3 through Helix 7 truncation (H3-H7, Figure 1F) with a drastically

reduced repression strength and rapid alleviation of repression by auxin addition (yellow and

pink, Figure 4E). Residual repression in H3-H7 likely comes from binding to MED10B, which

binds outside amino acids 145-149 (Supplemental Figure 1D). These results indicate that the

CRA domain (H3-H8) requires contact with MED21 to repress, and that this is independent of

the repression via Helix 1.

**Figure 3. Identification of critical residues within Helix 8 repression domain. A.** Sequence and structure of Helix 8 (5NQS). Helix 8 is colored green, and amino acids chosen for mutation are highlighted in light green in both the sequence and the structure. **B.** Repression activity of indicated single and double alanine mutations. **C.** Time course flow cytometry of selected mutations of Helix 8 following auxin addition. TPLH3-8-IAA3 fusion proteins (black) were compared to indicated single mutations to alanine (red). Controls - Helix 1 (H1 – blue), and TPLN188 (dark green). **D.** A series of alanine mutations (V145A, E146A, K148A, K149A, and the quadruple mutant Quad$^{AAAA}$ chosen from Figure 2D-F) were introduced into the TPL-N188 bait construct and tested for interaction with wild-type TPL-N188, IAA3 and AtMED21. Each single alanine mutation reduces TPL interaction with AtMED21, while the quad mutation abrogated interaction. These mutations also reduced the binding strength of TPLN with IAA3. **E.** The Helix 8 Quad$^{AAAA}$ mutation was introduced into the TPLN188-IAA3 and TPLH3-8-IAA3

fusion proteins and compared to wild type N188 in time course flow cytometry. For all cytometry

experiments, the indicated TPL construct is fused to IAA3. Every point represents the average

fluorescence of 5-10,000 individually measured yeast cells (a.u. - arbitrary units). Auxin (IAA-

10µM) was added at the indicated time (gray bar, + Aux). At least two independent experiments

are shown for each construct.

The TPL N-terminus (N188) interacted with at least two components of the Mediator

complex, AtMED21 and AtMED10B, and interaction between residues in Helix 8 (V145, E146,

K148, K149) were necessary for repression (Figure 3). These results suggest that the Mediator

complex is required for the TPL N-terminal domain to control repression (Figure 4A). To

determine whether a protein-protein interaction is required for corepressor function, it is critical

to demonstrate that reciprocal loss of function mutations activate repressed genes. In the case

of Tup1, a standard approach has been to test deletion mutations of Tup1-interacting proteins to

determine whether their interaction represents a true repression modality on the target gene

(Gromöller and Lehming, 2000; Lee et al., 2000; Zhang and Reese, 2004). We sought to test

whether loss of function in the Mediator complex would result in activation of the TPL-repressed

reporter in the $AtARC^{Sc}$. However, Mediator loss of function mutants can be lethal or exhibit

drastic physiological phenotypes (Biddick and Young, 2005). Instead, we combined the Anchor

Away system for rapid, chemically conditional protein depletion (Haruki et al., 2008) with the

auxin response circuit (Figure 4B).

$AtARC^{Sc}$ integrates components at four genomic locations using prototrophic markers

that are not compatible with those needed for Anchor Away. To overcome this limitation, we re-

recreated the entire ARC on a single plasmid (SLARC) using the Versatile Genetic Assembly

System (VEGAS, (Mitchell et al., 2015)). SLARC behaved with similar dynamics to the original

$AtARC^{Sc}$ on both solid and liquid growth conditions (Supplemental Figure 3A-C). As a first test of

the Anchor Away system with SLARC, we fused Tup1 and its partner protein Cyc8 to two copies

of the FKBP12-rapamycin-binding (FRB) domain of human mTOR (Haruki et al., 2008).

Rapamycin treatment of strains targeting either of these proteins caused no release of

repression, providing confirmation that the ARC acts orthogonally to the yeast corepressor

(Supplemental Figure 3D).

To test whether Mediator proteins or specific modules were required for TPL repression

we introduced the fully repressed SLARC plasmid containing TPLN188 ($SLARC^{N188}$) into a

library of Anchor Away yeast strains that allow specific depletion of Mediator components (see

Figure 4A-B) developed in (Haruki et al., 2008; Petrenko et al., 2017). Upon nuclear depletion of

the target genes we saw a striking activation of the auxin reporter when Mediator components

from the Tail, Head, and Middle domains were depleted (Figure 4C-E). We observed a less

pronounced activation upon depletion of CDK8 (Figure 4F), suggesting that this module plays

an ancillary role compared to the core Mediator complex. It is critical to note that Anchor Away

of components that are absolutely required for transcriptional activation will show no activation,

such as the Tail triple Anchor Away (17,3,15 - Figure 4C) or the RNA Polymerase II (Pol II)

Anchor Away (Rpb1 – Figure 4G). These observations support the interpretation that the entire

Mediator core (Tail, Head, Middle) is required for robust repression by the TPL N terminus,

which is recruited to the middle domain by MED10 and MED21.

Because we have pinpointed TPL N-terminal repression and protein interaction to the

specific region of AtMED21 we chose to focus in on this region in the yeast Med21 protein.

Deletion of the first 7 amino acids of ScMed21 ($\Delta$7Med21) partially releases genes that are

normally repressed by Tup1 into a transcriptionally active state (Gromöller and Lehming, 2000).

We hypothesized that TPL repression would also be partially alleviated by such a deletion.

ScMed21 is an essential gene, and yeast carrying deletions like $\Delta$7Med21 grow more slowly

than the wild type (Gromöller and Lehming, 2000; Hallberg et al., 2006), which itself alters

expression levels of reporters. In fact, even a $\Delta$5Med21 yeast mutant has been demonstrated to

alter Mediator assembly as the first 5 amino acids of ScMed21 are required for binding of Pol II

and the CDK8 kinase module to the Mediator core (Hallberg et al., 2006; Sato et al., 2016).

We introduced SLARCs with different TPL constructs into strains where ScMed21 wild-

type or N-terminal deletions were stably expressed and targets of Anchor Away. Strains

expressing ScMed21 with or without FRB fusions behaved similarly (Figure 4H, Supplemental

Figure 3E-F). We compared the transcriptional output of the fully repressed SLARC$^{N188}$ in

MED21 N-terminal mutants lacking the first three, five or seven amino acids. We observed that

all three deletions significantly increased the expression of the reporter in SLARC$^{N188}$, while no

mutation increased the SLARC's sensitivity to auxin (Figure 4H, Supplemental Figure 3E-F). As

$\Delta$7ScMed21 did not increase reporter expression when compared to $\Delta$5ScMed21 yet did have a

noticeable impact on growth, we did not use it in further analyses. Deletions of ScMed21 N-

terminal residues had no effect on repression in SLARC$^{H1-H5}$ but they do impair auxin responsive

transcriptional activation (Figure 4I), consistent with the role of this region in promoting Pol-II

recruitment (Sato et al., 2016). The fully repressed SLARC[N188] in Δ3ScMed21 or Δ5ScMed21

mutants showed elevated transcription of the reporter and when depleted demonstrate an

increase in reporter activity (Figure 4J), indicating that there is still residual TPLN188-Mediator

interaction taking place. Conversion of the first five amino acids of ScMed21 to the

corresponding sequence from AtMED21 resulted in an identical repression profile

(Supplemental Figure 4E), and had no effect on yeast growth or viability (Supplemental Figure

4F), further highlighting the conservation of this repression mechanism between the two

organisms.

The stably expressed N-terminal deletions of ScMed21 likely alter the expression of

multiple yeast genes, and this state change could confound our interpretations of the

importance of ScMed21 on TPL repression. To minimize the off-target impact of ScMed21

deletions, we introduced estradiol inducible versions of ScMed21 (iScMed21) into the Anchor

Away SLARC[N188] strains (Figure 4B, (McIsaac et al., 2013)). The combination of all three

synthetic systems made it possible to rapidly deplete the wild type ScMed21-FRB from the

nucleus while simultaneously inducing ScMed21 variants and visualizing the impact on a single

auxin-regulated locus. Depletion of nuclear ScMed21 by Rapamycin increased levels of the

reporter in all genotypes examined (Figure 4J) while also increasing cell size even in short time-

courses, consistent with its essential role in many core pathways (Supplemental Figure 4A,

(Gromöller and Lehming, 2000)). When wild-type iScMed21 was induced, there was a rescue of

both phenotypes (Figure 4K, black), whereas induction of either Δ3 and Δ5 variants

recapitulated the reporter activation seen in the stably expressed mutant versions (Figure 4K,

green and blue, Supplemental Figure 4B). iΔ3Med21 was induced and accumulated at a

comparable level to wild type Med21, while Δ5 appears to be less stable (Figure 4L-M). In these

short time courses, we did not observe the cell size increases observed in the Rapamycin

treatments (populations were evenly distributed around a single mean, suggesting we were

observing the immediate effects of the Med21 deletions (Supplemental Figure 4C-D)).

**Figure 4. Repression by TPL requires interaction with the N-terminus of MED21.**

**A.** Model of the proposed interaction between the TPL N-terminus with Mediator, where TPL interaction with Mediator 21 and 10 inhibit the recruitment of Pol-II. Proteins in this complex that were tested by Anchor Away are listed on the right. **B.** Schematic of _At_ARC<sup>Sc</sup> combined with

**Figure 4. Repression by TPL requires interaction with the N-terminus of MED21.**

**A.** Model of the proposed interaction between the TPL N-terminus with Mediator, where TPL interaction with Mediator 21 and 10 inhibit the recruitment of Pol-II. Proteins in this complex that were tested by Anchor Away are listed on the right. **B.** Schematic of _At_ARCSc combined with

methods for inducible expression and nuclear depletion of MED21. In Anchor Away, the yeast

ribosomal protein 13A (RPL13A) is fused to the rapamycin binding protein FKBP. Addition of

Rapamycin induces dimerization between FKBP and any target protein fused to FRB, resulting

in removal of the target protein from the nucleus. For these experiments, $AtARC^{Sc}$ was

assembled into a single plasmid (SLARC) rather than being integrated into separate genomic

loci (Figure 4 – Figure Supplement 1). Estradiol-inducible ScMed21 (iMed21) made it possible

to replace wild-type MED21 with targeted deletions or mutations. **C-G.** Time-course flow

cytometry analysis of $SLARC^{N188}$ in Mediator Anchor Away yeast strains with Rapamycin

(orange bar, + Rapa). Two Med21 strains were compared in the Middle domain (E), 21

(generated in this study) and 21* (generated in (Petrenko et al., 2017)). Both 21 and 21*

demonstrated similar increases in reporter expression.  **H.** Quantification of Venus fluorescence

from $SLARC^{N188}$ in wild type and N-terminal ScMed21 deletions with and without auxin. Yeast

was grown for 48 hours on SDO media with or without auxin and colony fluorescence was

quantified and plotted with the auxin responsive $SLARC^{H1-H5}$ in wild type as a reference.

Background - red autofluorescence was used as a reference for total cell density. **I.** Time-course

flow cytometry analysis of $SLARC^{H1-H5}$ in wild type and n-terminal ScMed21 deletions with and

without auxin. Genotypes are indicated in the colored key inset into the graph. Auxin (IAA-

10µM) was added at the indicated time (gray bar, + Aux). **J.** Time-course flow cytometry

analysis of $SLARC^{N188}$ in wild type and N-terminal ScMed21 deletions with and without

Rapamycin. Genotypes are indicated in the colored key inset into the graph. **K.** Rapid

replacement of Med21-FRB with inducible Med21-FLAG demonstrated the requirement for the

ScMed21 N-terminus in TPL repression. iMed21 isoforms were induced by addition of β-

estradiol (20µM) for 4 hours followed by Rapamycin addition. Fluorescence was quantified by

cytometry after 300 minutes (indicated by the dashed box in **D**). Lower case letters indicate

significant difference (ANOVA and Tukey HSD multiple comparison test; p<0.001). **L-M.** Protein

abundance of ScMed21 variants was tested by SDS-PAGE & western blot. For (**C-G, J**) a.u. -

arbitrary units. Rapamycin was added at the indicated time (orange bar, + Rapa). Every point

represents the average fluorescence of 5-10,000 individually measured yeast cells.

Our data indicated that truncated TPL repression domains were sufficient for interaction

with Mediator complex and transcriptional repression, however these truncations (i.e. TPLH3-

H8), are unable to form tetramers. This observation re-opened the question as to whether TPL

multimerization affects repression, as both the N-terminus of AtTPL and OsTPR2 adopt a

tetrameric form when crystallized (Figure 5A (Ke et al., 2015; Martin-Arevalillo et al., 2017)). We

used the cytoplasmic split ubiquitin (cytoSUS) protein-protein interaction assay (Asseck and

Grefen, 2018) to test this hypothesis. We observed that Helix 8 was required for strongest

interaction between TPL constructs (Supplemental Figure 5A), although this assessment was

complicated by the fact that some of the shorter constructs accumulated to significantly lower

levels (Supplemental Figure 5B). The weak interaction we could observe between full-length

TPL-N and the Helix 1 through Helix 3 construct (H1-3), indicated that the TPL LisH domain is

sufficient for dimerization. Therefore, while auxin-insensitive repression may require multimeric

TPL, this higher-order complex was not required for auxin-sensitive repression mediated by

Helix 1 (Figure 1C-D). In their study of the AtTPL structure, Martin-Arevalillo et al. identified a

triple mutation (K102S-T116A-Q117S-E122S) that abrogated the ability of the CRA domain

(Helix 6 and Helix 7) to form inter-TPL interactions (Martin-Arevalillo et al., 2017). As this mutant

form of TPL is only capable of dimerizing through its LisH domain, we refer to it here as LDimer

(Figure 5A). The LDimer mutations in TPLN188 retained the same auxin insensitive repression

auxin behavior as wild-type TPLN188 (Figure 5D), supporting the finding from the deletion

series.

To make a fully monomeric form of TPL, we introduced mutations into the dimerization

interface of the LisH domain in the context of LDimer. We first mutated one of a pair of

interacting residues (F15) to a series of amino acids (Tyrosine - Y, Alanine - A, Arginine - R, or

Aspartic Acid - D) in the context of either LDimer (Figure 5D), or H1-2 (Figure 5B, 5E). We

observed that conversion of F15 to the polar and charged aspartic acid (D) completely

abolished repression activity, while the polar and positively charged arginine was better

tolerated (Figure 5D,E). The conversion of F15 to tyrosine had no effect on LDimer (Figure 5D),

and only a minimal increase in auxin sensitivity in the context of H1-2 (Figure 5E). We then

combined LDimer-F15Y with a mutation of the coordinating residue L8 to serine with the

intention of stabilizing the now solvent-facing residues. The repressive behavior of this mutant

was indistinguishable from that of LDimer (Figure 5F). To further push the LDimer towards a

monomeric form, we introduced two additional mutations (S5A, E19S, Figure 5C,G). Size

exclusion chromatography confirmed that this combination of mutations (S5A-L8S-F15Y-E19S-

K102S-T116A-Q117S-E122S, hereafter called Monomer) successfully shifted the majority of the

protein into a monomeric state (Figure 5H); however, this shift had no observable impact on

repression strength before or after auxin addition (Figure 5G). To test whether these mutations

had a similar impact on *in vivo* TPL complexes, we introduced the LDimer and Monomer

mutations into the cytoSUS assay. In contrast to the *in vitro* chromatography results with purified

proteins, Monomer expressed in yeast retained measurable interaction with wild-type TPL,

LDimer or Monomer, albeit at a reduced level than what was observed between two wild type

TPLN188 constructs (Figure 5I). A caveat to this apparent difference between assays is that the

Monomer mutations led to a striking increase in protein concentration in yeast (Figure 5 – Figure

Supplement 1C), likely partially compensating for the decrease in affinity.

**Figure 5. Multimerization is not required for repression in yeast. A.** TPL can form a

homotetramer via the CRA (red) and LisH (blue) domains. Asterisks indicate mutations that

block or diminish these interactions.  **B-C**. Locations of critical positions in Helix 1 are

highlighted for two interacting TPL monomers (shown in light and dark blue). Interacting amino

acids share the same color (adapted from 5NQV).

**D-G.** Time course flow cytometry analysis of TPLN-IAA3 fusion proteins carrying selected single

point mutations in N188-LDimer-IAA3 (**D**) and the TPLH1-2 truncation (**E**). The F15Y mutation

had little effect on repression activity for either TPL construct. Double mutations (F15Y, L8S in

LDimer) (**F**) or the quadruple Monomer mutations (S5A, L8S, F15Y, E19S in LDimer) (**G**)

showed repression activity that was indistinguishable from LDimer or wild type N188 fused to

IAA3. For all cytometry experiments, the indicated TPL construct is fused to IAA3. Every point

represents the average fluorescence of 5-10,000 individually measured yeast cells (a.u. -

arbitrary units). Auxin (IAA-10µM) was added at the indicated time (gray bar, + Aux). At least

two independent experiments are shown for each construct. **H**. Size Exclusion Chromatography

on TPLN188 wild type (green), LDimer (purple) and Monomer (orange) tetramerization mutants.

**I**. CytoSUS on TPL tetramerization mutants.

480            While the $At$ARC$^{Sc}$ relies on the strong conservation of the core regulatory machinery in

yeast and plants, there are important differences between the synthetic and native systems. For

example, while the TPL-N188 construct (and other variants that contain Helix 8 and the linker

between Helix 8 and Helix 9) repress transcription in yeast even after auxin addition, this is not

the case for the many auxin-regulated genes in plants. To uncover these differences, we first

tested for TPL and MED21 interaction using Bimolecular Fluorescence Complementation (BiFC)

in *Nicotiana benthamiana* (tobacco). Full-length TPL and MED21 fusions with YFP were

localized to the nucleus and did interact with one another by BiFC (Figure 6A). The same TPL

Helix 8 residues identified as necessary for a strong interaction with AtMED21 in yeast (Figure 3

- V145A, E146A, K148A, K149A –TPL$^{H8QuadA}$) were also important for interaction in plants

(Figure 6A). Similarly, the Δ5AtMED21 N-terminal truncation completely eliminated interaction

with full-length TPL (Figure 6A). Co-immunoprecipitation assays confirmed the AtMED21 and

TPL interaction (Figure 6B). We were also able to pull down MED21 and TPL using MED10B

from tobacco extracts, supporting our previous interaction assays from yeast (Supplemental

Figure 6A).

495            We next developed a UAS/GAL4-VP16-based (Brand and Perrimon, 1993) conditional,

quantitative repression system for plants that relied on induction of a gene encoding a TPLN-

IAA14 fusion protein (Figure 6A). To sensitize the output of this assay, we used a variant of

IAA14 with mutations in the two EAR domains (EAR$^{AAA}$) and in the degron (P306S) to block

interference from the endogenous TPL/TPRs or auxin-mediated degradation by the TIR1/AFBs,

respectively (IAA14$^{mED}$ ; Figure 6C). After prototyping the system in yeast (Supplemental Figure

6B), we transiently transformed constructs carrying *TPLN-IAA14$^{mED}$* variants into tobacco to test

repression of the well-characterized auxin response promoter DR5 (Ulmasov et al., 1997). DR5

was strongly induced by co-transformation with the transcriptional activator AtARF19. This

induction was sharply reduced if the reporter and activator were co-transformed with *UAS-*

*TPLN188-IAA14$^{mED}$* and *GAL4-VP16* (Figure 6D). In general, we observed strong correlation in

repression activity between what was observed in yeast and in tobacco. Truncations containing

Helix 1 (H1) or Helix 8 (H3-H9), as well as the full N-terminus (N188), repressed the DR5

reporter (Figure 6D). LDimer and Monomer variants retained a similarly strong repressive

activity in tobacco as what had been observed in yeast (Figure 6D). These results are

consistent with previous evidence that loss of dimerization through the CRA domain has no

effect on transcriptional repression of the DR5 promoter in protoplasts (Martin-Arevalillo et al.,

2017). We also observed congruity between yeast and plant data in that the Helix 8 quadruple

alanine mutation showed similar repression strength as wild type N188, possibly due to

presence of Helix 1 (N188$^{H8QuadA}$ - Figure 6D).

We were concerned that the lack of effect of the Helix 8 mutation and multimerization

mutants (LDimer and Monomer) on repression strength could reflect artifacts from heterologous

overexpression. To test the effect of these mutants in a native context, we generated transgenic

*Arabidopsis* lines where the UAS-TPL-IAA14$^{mED}$ constructs were activated selectively in xylem

pole pericycle cells, where IAA14 normally acts to regulate the initiation of lateral root primordia

(Figure 6E, (Laplaze et al., 2005)). Transgenic plants expressing functional TPL variants in

these cells should make very few if any lateral roots, phenocopying the original solitary root (*slr*)

mutant, which carries an auxin-resistant form of IAA14 (Fukaki et al., 2002). As expected,

transformants expressing either IAA14$^{mED}$ (with no TPL fusion) or TPLN188 (with no IAA14

fusion) had no effect on lateral root production (Figure 6F). In contrast, TPLN188 fusion

constructs sharply decreased lateral root density (Figure 6F). The TPLH3-H9 truncation

decreased lateral root density but was not as efficient as TPLN188 stressing the role of the LisH

in repression (Figure 6F). Both LDimer and Monomer constructs (Figure 6F) were able to

repress lateral root development, meaning that multimer formation is not required for TPL

repression of native auxin-induced genes in this context. The fusion containing the Helix 8

quadruple mutant demonstrated a clear loss of repression, indicating that the TPL-MED21

interaction is critical for repression when expressed at endogenous levels (Figure 6F).

AtMED21 loss-of-function mutations are embryo lethal when homozygous (Dhawan et

al., 2009; Supplemental Figure 6D), and plants heterozygous for *Atmed21* mutations appear

wild-type, including exhibiting normal lateral root density (Supplemental Figure 6E). These

phenotypes make testing the specific function of AtMED21interactions with TPL challenging. As

an alternative means to probe AtMED21 function in regulating auxin responses, we devised a

method to specifically repress transcription of *AtMED21* in xylem pole pericycle cells, similar to

the approach used for testing the function of TPL variants. To do this, we introduced a dCAS9-

TPLN300 synthetic repressor under the control of a UAS promoter into J0121[Col-0] along with

three sgRNAs complementary to the *AtMED21* promoter. If AtMED21 is required for TPL-based

repression, we would expect loss of AtMED21 in xylem pole pericycle cells, which would result

in an increase in lateral root number and length. This is indeed what we observed (Figure 6G,

Supplemental Figure 6F-G).

**Figure 6. The TPL CRA repression domain behaves similarly in yeast and plants**

**A.** Bimolecular Fluorescence Complementation assay performed in tobacco. Each image is an

epi-fluorescent micrograph taken at identical magnification from tobacco epidermal cells at two

days post injection. The YFP image is colored green (left panel). p35S:H2B-RFP was used as a

control and is false colored magenta (right panel). **B.** Co-immunoprecipitation of MED21 and

TPL from tobacco leaves. MED21-YFP-HA was immunoprecipitated using anti-HA, and YFP-

TPL was detected using the YFP fusion. Actin was used to demonstrate that the purification had

removed non-specific proteins. Numbers on the left of blots indicate sizes of protein standards

in kilodaltons. **C.** Design of UAS-TPL-IAA14$^{mED}$ constructs. Mutation of the conserved Lysine

residues in the EAR domain disrupted potential interactions with endogenous TPL/TPR

proteins. The IAA14 degron has been mutated (P306S) to render it auxin insensitive. UAS –

upstream activating sequence, ttRBCS - Rubisco terminator sequence. **D.** Transient expression

of indicated TPL constructs in tobacco. DR5:Venus - the synthetic DR5 auxin promoter

(Ulmasov et al., 1997) driving Venus, ARF19 - p35S:AtARF19-1xFLAG, GAL4:VP16 -

pUBQ10:GAL4-VP16, TPLN-X-UAS-TPL-IAA14$^{mED}$ with various TPL truncations or mutations.

**E.** Auxin induced degradation of IAA14 is absolutely required for initiation of lateral root

development (cartoon, left). An enhancer trap line (J0121) expresses GAL4-VP16 and UAS-

GFP in in xylem pole pericycle cells. **F.** N-terminal domains of TPL were sufficient to repress the

development of lateral roots in *Arabidopsis* seedlings. The density of emerged lateral roots was

measured in T1 seedlings at 14 days after germination. **G.** MED21-CAS9 repressors targeted to

MED21 display increased lateral root densities. Lateral root density (number of lateral roots /

primary root length) was calculated at 10 days post germination. Numbers below boxplots are p-

values for pairwise comparisons with control using a Wilcoxon rank sum test. **F-G**. Lower case

letters indicate significant difference (ANOVA and Tukey HSD multiple comparison test;

p<0.001).

**Discussion**

A review of the current literature on corepressors gives the conflicting impressions that (a)

corepressor function is broadly conserved, and (b) that every organism (and perhaps even each

corepressor) has a distinct mode for transcriptional repression (Adams et al., 2018; Mottis et al.,

2013; Perissi et al., 2010; Wong and Struhl, 2011). We hoped that the *At*ARC$^{Sc}$ could facilitate a

resolution to this apparent contradiction by targeting repression to a single synthetic locus. We

focused our initial efforts on the analysis of the N-terminal portion of TPL which has multiple

known protein-protein interaction surfaces (Ke et al., 2015; Martin-Arevalillo et al., 2017).

Experiments with the *At*ARC[Sc] identified two repression domains [Helix 1 and the CRA domain

(Figure 1)] within the N-terminus of TPL, both of which were subsequently confirmed to repress

transcription in plants as well. This led us to hypothesize that they are therefore likely contact

points for other proteins (Figure 2). In the case of the more potent CRA domain, we were able to

identify two of these partners MED21 and MED10 (Figure 2). Amino acids within Helix 8 (in the

CRA domain) that were critical for repression activity have their R-groups oriented away from

the hydrophobic core of TPL structure and are required for Med21-binding and repression

(Figure 3). The Mediator subunit Med21 also interacts with the yeast corepressor Tup1, which

suggests a fundamental conservation in at least one corepressor mechanism across species.

Indeed, the whole core Mediator complex (Head, Middle, Tail) appears to be required for

repression through TPL (Figure 4). Contrary to our initial hypothesis, the monomeric form of

TPL was sufficient for strong repression in yeast and in plants, leaving open the question of the

role of higher-order TPL complex formation (Figure 5). Finally, we were able to show that the

TPL-MED21 interaction has functional significance for auxin response in plants (Figure 6).

The Mediator complex is a multi-subunit complex that connects DNA-bound transcription

factors and the RNA polymerase II complex (Pol-II) to coordinate gene expression (Flanagan et

al., 1991; Kim et al., 1994). The yeast Mediator subunits are organized into four separate

modules; head, middle, tail and kinase, with a strong conservation of module components in

plants (Dolan and Chapple, 2017; Maji et al., 2019; Malik et al., 2017; Samanta and Thakur,

2015). Med21 forms a heterodimer with Med7 to and interacts with Med10, among others, to

create the central region of the Middle region of the Mediator complex. The Med21 N-terminus

is centered on a flexible hinge region (Baumli et al., 2005), which is required for Pol-II

recruitment and CDK8 kinase module recruitment (Sato et al., 2016). The protein interaction

between TPL and MED21 occurs at the N-terminus of MED21, highlighting the importance of

this region as a signaling hub (Sato et al., 2016). Other lines of evidence support this role, as

this region binds the yeast homolog of TPL, Tup1 (Gromöller and Lehming, 2000) through a

completely different protein domain as no homology can be found between TPL Helix 8 and

Tup1 in any region by primary amino acid homology (i.e. BLAST).

Corepressors coordinate multiple mechanisms of repression through discrete protein

interactions, leading to robust control over eukaryotic transcription by combining repression

modalities. Corepressor function has variously been linked to (a) altering chromatin

confirmation, often through interaction with histone modifying proteins or histone proteins

themselves, (b) direct interference with transcription factor binding or function and (c) physical

spreading of long-range oligomeric corepressor complexes across regions of regulatory DNA

(Perissi et al., 2010). Dissection of the importance of each modality in Tup1 repression has

been challenging (Lee et al., 2000; Zhang and Reese, 2004). The tour-de-force of corepressor

mechanism studies in yeast concluded that the primary function of Tup1 was to physically block

activators (Wong and Struhl, 2011). In their work, the authors utilized the Anchor Away

approach to correlate the importance of HDACs, transcriptional machinery and chromatin

remodeling enzymes to the repression state of endogenously repressed Cyc8-Tup1 target

genes. They observed that Tup1 did not block the binding of transcription factors but inhibited

the recruitment of one Mediator component in the tail domain, GAL11/MED15, as well as Pol-II

and the chromatin remodelers Snf2 and Sth1. They additionally observed that HDACs had only

a supportive role in reinforcing Tup1 repression. These results led to their hypothesis that Tup1

blocks the activation domains of transcription factors, and suggested this was through direct

binding to activation domains (Wong and Struhl, 2011).

The synthetic system used here allowed us to build on this model and further refine our

understanding of TPL's repressive activity. In our experiments we see a similar set of

conditions, with TPL recruited to the DNA-bound transcriptional activator (ARF), and several

possible mechanisms of repression. Unlike Tup1, we have subdivided the TPL protein to identify

interactions between TPL and individual protein interactors with no effect on yeast function. In

these experiments we can eliminate the possibility that TPL blocks ARF activation by directly

blocking the transcription factor activation domains because we see a loss of repression only

when TPL-MED21 binding is eliminated through specific point mutations (Figure 4H). Our

inducible swap of MED21 isoforms also corroborates these findings (Figure 4), as the SLARC

remains genetically identical in these strains, indicating that TPL-MED21 interaction is

regulating activity not a TPL-ARF interaction. Furthermore, our results correlate well with

findings that repressed targets are reactivated when this portion of MED21 is deleted in yeast

(TPL - Figure 4, Tup1 - (Gromöller and Lehming, 2000)). Therefore, we suggest that instead of

directly binding activation domains that TPL (and likely Tup1) binds to components of Mediator

(MED21, MED10B and possibly others) recruited by the transcription factor. Indeed, it is easier

to rationalize that the repressor binds the same domains of the Mediator complex recruited by

the transcription factor's activation domain (with the same affinity) as opposed to binding each

diverse activation domain (with varying affinity). In this model, corepressor binding blocks

formation of a fully active Mediator complex thereby limiting Polymerase II recruitment and

promoter escape (Petrenko et al., 2017).

Multiple points of contact likely exist between the Mediator complex and other parts of

the transcriptional machinery in both transcriptionally repressed and active states. For auxin

response, specifically, there are several lines of evidence to support this model, including

documented association between the structural backbone of Mediator, MED14, and activated

and repressed auxin loci in *Arabidopsis* (Ito et al., 2016). In addition, MED12 and MED13 are

required for auxin-responsive gene expression in the root, and MED12 acts upstream of AUX1

in the root growth response to sugar (Raya-González et al., 2018). MED18 in the head module

represses auxin signaling and positively regulates the viability of the root meristem (Raya-

González et al., 2018). PFT1/MED25 regulates auxin transport and response in the root (Raya-

González et al., 2014). MED7, MED21's partner protein in the hinge domain, is required for

normal root development and loss of MED7 function impacts expression of auxin signaling

components (Kumar et al., 2018). Previous research identified the Mediator CDK8 subunit,

specifically MED13 (MAB2), as an interactor with the full-length TPL protein (Ito et al., 2016).

We could not observe interaction between the N-terminal domain of TPL and AtMED13,

AtCYC8, or AtCYCC (Supplemental Figure 1B), suggesting that any direct interactions occur

outside the N188 fragment. As suggested by Ito and colleagues (Ito et al., 2016) and supported

by our synthetic system, auxin-induced removal of TPL is sufficient to induce changes in the

composition of the Mediator complex, facilitating both rapid activation and rapid return to a

repressed basal state.

The conserved interaction of both TPL and Tup1 with Mediator has implications for

modeling the dynamics of eukaryotic transcription (e.g., (Estrada et al., 2016)). By stabilizing the

Mediator complex, TPL (and by extension Tup1) may create a 'pre-paused' state that allows

rapid recruitment of Pol-II and activation once TPL is removed. Alternatively, TPL could allow for

recruitment of a poised or paused Pol-II, allowing for rapid activation (and re-activation) of a

repressed locus. Support for this idea comes from the genome-wide correlation between

Groucho corepressor binding and polymerase pausing in *Drosophila* (Kaul et al., 2014).

However, in non-metazoan lineages (i.e. plants and yeast) the existence of bona fide pausing is

disputed, mainly due to the absence of the pausing regulator Negative Elongation Factor

complex (NELF, (Gaertner and Zeitlinger, 2014)). One possibility is that plants and yeast have

an ancestral or alternative form of pausing that is NELF-independent. Support for this idea

comes from studies of *Arabidopsis* where Pol-II was found to be markedly enriched at the five-

prime ends of genes, the same pattern seen with metazoan pausing (Zhu et al., 2018). In yeast,

Tup1 has been implicated in a form of transcriptional memory that involves recruiting a poised

form of preinitiation Pol-II to allow rapid reactivation of genes involved in sugar utilization (Sood

et al., 2017).

**Methods**

**Cloning**

Construction of TPL-IAA3 and TPL-IAA14 fusion proteins were performed as described in

(Pierre-Jerome et al., 2014). Variant and deletion constructs were created using PCR-mediated

site-directed mutagenesis. Site-directed mutagenesis primers were designed using

NEBasechanger and implemented through Q5® Site-Directed Mutagenesis (NEB, Cat

#E0554S). TPL interactor genes were amplified as cDNAs from wild type Col-0 RNA using

reverse transcriptase (SuperScript™ IV Reverse Transcriptase, Invitrogen) and gene-specific

primers from IDT (Coralville, Iowa), followed by amplification with Q5 polymerase (NEB). These

cDNAs were subsequently cloned into Plasmids for cytoSUS using a Gibson approach (Gibson

et al., 2009), through the Aquarium Biofabrication facility (Ben Keller et al., 2019). The coding

sequence of the genes of interest were confirmed by sequencing (Genewiz; South Plainfield,

NJ). For UAS driven constructs, the TPLN188-IAA14 coding sequence was amplified with

primers containing engineered BsaI sites and introduced into the pGII backbone with the UAS

promoter and RBSC terminator (Siligato et al., 2016) using Golden Gate cloning (Weber et al.,

2011). Subsequent mutations were performed on this backbone using PCR-mediated site-

directed mutagenesis (see above). Construction of C-terminal 2xFRB fusions for Anchor Away

were constructed as described in (Haruki et al., 2008). Inducible MED21 was constructed as

described in (McIsaac et al., 2013). For cell type specific knockdown mediated by dCas9-

TPLN300, Gibson cloning was used modify the pHEE401E plasmid, replacing the egg-specific

promoter and Cas9 from pHEE401E (Wang et al., 2015) with UAS promoter and dCas9-

TPLN300 fusion protein (Khakhar et al., 2018). The resulting plasmid is used as starting point to

clone 3 sgRNAs targeting the *AtMED21* promoter (identified using CHOP CHOP (Labun et al.,

2019). sgRNAs: GACGCAGAGTCTGTTGGGTGTGG, TTTAAAATGGGCTTTTAAGGTGG,

AACACTGAAGTAGAATTGGGTGG ranging from -170 to +90 region from the TSS) using PCR

and Golden Gate cloning strategy described in (Wang et al., 2015).

**Flow Cytometry**

Fluorescence measurements were taken using a Becton Dickinson (BD) special order

cytometer with a 514-nm laser exciting fluorescence that is cut off at 525 nm prior to

photomultiplier tube collection (BD, Franklin Lakes, NJ). Events were annotated, subset to

singlet yeast using the FlowTime R package (Wright et al., 2019). A total of 10,000 - 20,000

events above a 400,000 FSC-H threshold (to exclude debris) were collected for each sample

and data exported as FCS 3.0 files for processing using the flowCore R software package and

custom R scripts (Supplemental File 1, (Havens et al., 2012; Pierre-Jerome et al., 2017)). Data

from at least two independent replicates were combined and plotted in R (ggplots2).

**Yeast Methods**

Standard yeast drop-out and yeast extract–peptone–dextrose plus adenine (YPAD) media were

used, with care taken to use the same batch of synthetic complete (SC) media for related

experiments. A standard lithium acetate protocol (Gietz and Woods, 2002) was used for

transformations of digested plasmids. All cultures were grown at 30°C with shaking at 220 rpm.

Anchor-Away approaches were followed as described in (Haruki et al., 2008), and anchor away

strains were obtained from EURO-SCARF (euroscarf.de). Endogenous genomic fusions of

ScMed21-FRB were designed by fusing MED21 homology to the pFA6a-FRB-KanMX6 plasmid

for chromosomal integration into the parental anchor away strain as in (Petrenko et al., 2017),

selectable through G418 resistance (G418, Geneticin, Thermo Fisher Scientific). Tup1-FRB and

Cyc8-FRB were constructed as described in (Wong and Struhl, 2011). Mediator anchor away

strains were created in (Petrenko et al., 2017) and kindly donated by Dr. Kevin Struhl. SLARC

construction required a redesign of promoters and terminators used in the *At*ARC[Sc] to eliminate

any repetitive DNA sequences (See Figure 4 – Figure Supplement 1), using a Golden Gate

cloning approach into level 1 vectors. Subsequent assembly of individual transcriptional units

into a larger plasmid utilized VEGAS assembly which was performed as described in (Mitchell et

al., 2015). To create an acceptor plasmid for the assembled transcriptional units, we

synthesized a custom vector containing VA1 and VA2 homology sites for recombination (Twist

Bioscience, South San Francisco, CA). In between these sites we incorporated a pLac:mRFP

cassette to allow identification of uncut destination plasmid in *E. coli*, flanked by EcoRI sites for

linearization. Finally, the CEN6/ARSH4 was transferred from pRG215 (Addgene #64525) into

the acceptor plasmid by Golden Gate reaction using designed BsmBI sites engineered into the

acceptor plasmid and the primers used to amplify the CEN/ARS (See Figure 4 – Figure

Supplement 1). For the cytoplasmic split-ubiquitin protein-protein interaction system, bait and

prey constructs were created using the plasmids pMetOYC and pNX32, respectively (Addgene,

https://www.addgene.org/Christopher_Grefen/). Interaction between bait and prey proteins were

evaluated using a modified version of the split ubiquitin technique (Asseck and Grefen, 2018).

After two days of growth on control and selection plates, images were taken using a flatbed

scanner (Epson America, Long Beach, CA). Inducible ScMed21 strains (iMed21) were grown

overnight, and then diluted back to 100 events per microliter as determined by flow cytometry

and grown at 30C with 250rpm in a deepwell 96-well plate format. Strains were supplemented

with β-estradiol (20μM) for 4 hours followed by Rapamycin addition. Samples were analyzed by

flow cytometry throughout these growth experiments.

**Western Blot**

Yeast cultures that had been incubated overnight in synthetic complete (SC) media were diluted

to OD600 = 0.6 and incubated until cultures reached OD600 ~1. Cells were harvested by

centrifugation. Cells were lysed by vortexing for 5 min in the presence of 200 μl of 0.5-mm

diameter acid washed glass beads and 200 μl SUMEB buffer (1% SDS, 8 M urea, 10 mM

MOPS pH 6.8, 10 mM EDTA, 0.01% bromophenol blue, 1mM PMSF) per one OD unit of original

culture. Lysates were then incubated at 65° for 10 min and cleared by centrifugation prior to

electrophoresis and blotting. Antibodies: Anti-HA-HRP (REF-12013819001, Clone 3F10,

Roche/Millipore Sigma, St. Louis, MO), Anti-FLAG (F3165, Monoclonal ANTI-FLAG® M2,

Millipore Sigma, St. Louis, MO), Anti-FRB (ALX-215-065-1, Enzo Life Sciences, Farmingdale,

NY, (Haruki et al., 2008)), Anti-VP16 (1-21) (sc-7545, Santa Cruz Biotechnology, Dallas TX),

Anti-GFP (ab290, AbCam, Cambridge, United Kingdom), Anti-MYC (71d10, 2278S, Cell

Signaling, Danvers, MA), Anti-PGK1 (ab113687, AbCam, Cambridge, United Kingdom).

**Protein expression and purification**

All multimer deficient TPL proteins were expressed in *Escherichia coli* Rosetta 2 strain. Bacteria

cultures were grown at 37ºC until they achieved an OD$^{600}$nm of 0.6-0.9. Protein expression was

induced with isopropyl-β-D-1-thyogalactopiranoside (IPTG) at a final concentration of 400 µM at

18 °C overnight. Bacteria cultures were centrifuged and the pellets were resuspended in the

buffer A (CAPS 200 mM pH 10.5, NaCl 500 mM, TCEP 1 mM), where cells were lysed by

sonication. His-tagged AtTPL188 (wt and mutants) bacteria pellets were resuspended in buffer

A with EDTA-free antiprotease (Roche). The soluble fractions recovered after sonication were

passed through a Ni-sepharose (GE Healthcare) column previously washed with buffer A and

the bound proteins were eluted with buffer A with 250 mM imidazole. A second purification step

was carried out on Gel filtration Superdex 200 10/300 GL (GE Healthcare) equilibrated with

buffer A.

**Co-Immunoprecipitation**

Co-IP from yeast was performed using the cytoSUS strains. Cultures were grown to OD600 0.5

(~1E7 cells/ml) using selective media, harvested, and resuspended in 200 µl SUME (1% SDS,

8M Urea, 10mM MOPS, pH 6.8, 10mM EDTA) buffer with protease inhibitors. Cells were lysed

by vortexing 3 X 1 min full speed with 100 µl of 0.5 mm Acid Washed Glass Beads, clarified by

centrifugation (1 min, 1000rpm), and supernatant was mixed with 1ml IP buffer (15mM

Na2HPO4, mw 142; 150mM NaCl, mw 58; 2% Triton X-100, 0.1% SDS, 0.5% DOC, 10mM

EDTA, 0.02% NaN3) with protease inhibitors, and incubated with 100 $\mu$l of IgG sepharose at

25°C for 2 hours with rotation. The Beads were washed 1x with IP buffer, and 2x with IP-Wash

buffer (50mM NaCl, mw58; 10mM TRIS, mw 121; 0.02% NaN3) with protease inhibitors. Protein

was eluted with 50 µl of SUME buffer + 0.005% bromophenol blue by incubation at 65°C for 10

minutes and run on handmade 12% acrylamide SDS-PAGE gels, and western blotted

accordingly. Co-IPs from tobacco were performed on leaves two days after injection as

described in (Song et al., 2014). For Co-IPs with HA, extracts were incubated with Anti-HA-

Biotin (High Affinity (3F10), Sigma, 12158167001) and Streptavidin conjugated magnetic beads

(). For Co-IPs with MED10B-MEC-ProtA we used IgG Sepharose® 6 Fast Flow (Sigma, GE17-

0969-01) beads and increased washing steps (1xIP buffer, 4xWash buffer, 5 total). The only

modification to buffers was an addition of the detergent NP-40 at 0.1% in the IP and wash

buffer. Samples were run on handmade 10% acrylamide SDS-PAGE gels, and western blotted

accordingly.

**Bimolecular Fluorescence Complementation (BiFC)**

Bimolecular fluorescent complementation experiments were performed on 3-week-old *N.*

*benthamiana* plants grown at 22°C under long days (16 h light/8 h dark) on soil (Sunshine #4

mix) as per (Martin et al., 2009). pSITE vectors were used to generate BiFC constructs for

MED21, Δ5-MED21, TPL and TPL[H8QuadA] – proteins (Martin et al., 2009). In all cases the

combinations are N-terminal fusions of either the nEYFP or cEYFP to the cDNA of MED21 or

TPL. RFP fused Histone H2B was used as a nuclear marker (Goodin et al., 2002). Injection of

Agrobacterium strains into tobacco leaves was performed as in (Goodin et al., 2002), but the

$OD_{600}$ of the Agrobacterium culture used was adjusted to 0.5. Two days after transfection, plant

leaves were imaged using an epifluorescence microscope (Leica Biosystems, model: DMI

3000B).

**Protein alignments**

The MED21 protein sequence was were aligned to homologs using CLC Sequence Viewer 7, a

tree was constructed using a Neighbor-Joining method, and bootstrap analysis performed with

10,000 replicates.

**Plant growth**

For synthetic repression assays in tobacco Agrobacterium-mediated transient transformation of

*Nicotiana benthamiana* was performed as per (Yang et al., 2000). 5 ml cultures of

Agrobacterium strains were grown overnight at 30C shaking at 220rpm, pelleted and incubated

in MMA media (10 mM MgCl2, 10 mM MES pH 5.6, 100 µM acetosyringone) for 3 hours at room

temperature with rotation. Strain density was normalized to an OD600 of 1 for each strain in the

final mixture of strains before injection into tobacco leaves. Leaves were removed, and 8

different regions were excised using a hole punch, placed into a 96-well microtiter plate with

100µl of water. Each leaf punch was scanned in a 4x4 grid for yellow and red fluorescence

using a plate scanner (Tecan Spark, Tecan Trading AG, Switzerland). Fluorescence data was

quantified and plotted in R (ggplots). For *Arabidopsis thaliana* experiments using the GAL4-UAS

system (Laplaze et al., 2005), J0121 was introgressed eight times into Col-0 accession from the

C24 accession, and rigorously checked to ensure root growth was comparable to Col-0 before

use. UAS-TPL-IAA14$^{mED}$ constructs were introduced to J0121 introgression lines by floral dip

method (Clough and Bent, 1998). T1 seedlings were selected on 0.5X LS (Caisson

Laboratories, Smithfield, UT) + 25μg/ml Hygromycin B (company) + 0.8% phytoagar

(Plantmedia; Dublin, OH). Plates were stratified for 2 days, exposed to light for 6 h, and then

grown in the dark for 3 d following a modification of the method of (Harrison et al., 2006).

Hygromycin resistant seedlings were identified by their long hypocotyl, enlarged green leaves

and long root. Transformants were transferred by hand to fresh 0.5X LS plates + 0.8% Bacto

agar (Thermo Fisher Scientific) and grown vertically for 14 days at 22°C. Plates were scanned

on a flatbed scanner (Epson America, Long Beach, CA) at day 14. *slr* and *med21/MED21*

(WiscDsLox461-464K13) seeds were obtained from the Arabidopsis Biological Resource Center

(Columbus, OH). CRISPR/CAS9 based mutations in AtMED2 were generated as described in

(Wang et al., 2015). We created a novel mutation in *AtMED21* that introduces a single base-pair

insertion of G at nucleotide 214 after the A of the start codon (i214G). This mutation alters the

amino acid sequence starting at residue 25 and creates an early stop codon after 11 random

amino acids (Supplemental Figure 6D).

**Data submissions**

All flow cytometry data will be deposited at https://flowrepository.org/. All plasmids will be

deposited through Addgene at https://www.addgene.org/Jennifer_Nemhauser/.

**Acknowledgements**

We would like to thank Prof. Jef Boeke for kindly providing VEGAS adaptor and regulatory

element plasmids; Dr. Jennifer Brophy and Prof. José Dinneny for kindly providing the

pUBQ10:GAL4:VP16 plasmid; Dr. Natalia Petrenko and Dr. Kevin Struhl for kindly providing

Mediator anchor away strains; and Prof. Grant Brown and Prof. Maitreya Dunham for advice on

yeast genetics and approaches. We thank members of the Nemhauser group including Amy

Lanctot, Romi Ramos, Eric Yang, and Dr. Sarah Guiziou for constructive discussions and

comments on this manuscript. We also thank Morgan Hamm for the custom R script used here

to analyze anchor away plates.

**Author Contributions**

Conceptualization: ARL, JLN; Investigation: ARL, WW, HPG, SG, SJS, MLZ, JEZ; Software:

ARL, HPG; Formal Analysis: ARL, HPG; Visualization: ARL, WW, HPG, SG, SJS, MLZ, JEZ;

Writing-Original Draft: ARL; Writing-Review & Editing: WW, HPG, SG, SJS, MLZ, JEZ, NZ, JLN;

Supervision, Project administration, Funding Acquisition, Resources: JLN, NZ.

**Declaration of Interests**

The authors declare no competing interests.

**Funding**

National Institutes of Health (NIH): ARL, HPG, SG, SJS, JEZ, MZ & JLN R01- GM107084.

Howard Hughes Medical Institute (HHMI): ARL, SJS, JEZ, MZ & JLN - Faculty Scholar Award.

Ning Zheng is a Howard Hughes Medical Institute Investigator.

ARL is supported by the Simons Foundation through the Life Science Research Foundation.

**References**

Adams, G.E., Chandru, A., and Cowley, S.M. (2018). Co-repressor, co-activator and general transcription factor: the many faces of the Sin3 histone deacetylase (HDAC) complex. Biochem. J. *475*, 3921–3932.

Agarwal, M., Kumar, P., and Mathew, S.J. (2015). The Groucho/Transducin-like enhancer of split protein family in animal development. IUBMB Life *67*, 472–481.

Asseck, L.Y., and Grefen, C. (2018). Detecting Interactions of Membrane Proteins: The Split-Ubiquitin System. Methods Mol. Biol. Clifton NJ *1794*, 49–60.

Baumli, S., Hoeppner, S., and Cramer, P. (2005). A Conserved Mediator Hinge Revealed in the Structure of the MED7·MED21 (Med7·Srb7) Heterodimer. J. Biol. Chem. *280*, 18171–18178.

Ben Keller, Justin Vrana, Abraham Miller, Garrett Newman, and Eric Klavins (2019). Aquarium: The Laboratory Operating System (Zenodo).

Biddick, R., and Young, E.T. (2005). Yeast Mediator and its role in transcriptional regulation. C. R. Biol. *328*, 773–782.

Bourbon, H.-M. (2008). Comparative genomics supports a deep evolutionary origin for the large, four-module transcriptional mediator complex. Nucleic Acids Res. *36*, 3993–4008.

Brand, A.H., and Perrimon, N. (1993). Targeted gene expression as a means of altering cell fates and generating dominant phenotypes. Dev. Camb. Engl. *118*, 401–415.

Busch, W., Miotk, A., Ariel, F.D., Zhao, Z., Forner, J., Daum, G., Suzaki, T., Schuster, C., Schultheiss, S.J., Leibfried, A., et al. (2010). Transcriptional control of a plant stem cell niche. Dev. Cell *18*, 849–861.

Causier, B., Ashworth, M., Guo, W., and Davies, B. (2012). The TOPLESS interactome: a framework for gene repression in Arabidopsis. Plant Physiol. *158*, 423–438.

Chen, G., and Courey, A.J. (2000). Groucho/TLE family proteins and transcriptional repression. Gene *249*, 1–16.

Clough, S.J., and Bent, A.F. (1998). Floral dip: a simplified method for Agrobacterium-mediated
transformation of Arabidopsis thaliana. Plant J. Cell Mol. Biol. *16*, 735–743.

Collins, J., O'Grady, K., Chen, S., and Gurley, W. (2019). The C-terminal WD40 repeats on the
TOPLESS co-repressor function as a protein–protein interaction surface. Plant Mol. Biol. *100*,
47–58.

Delto, C.F., Heisler, F.F., Kuper, J., Sander, B., Kneussel, M., and Schindelin, H. (2015). The
LisH motif of muskelin is crucial for oligomerization and governs intracellular localization. Struct.
Lond. Engl. 1993 *23*, 364–373.

Dhawan, R., Luo, H., Foerster, A.M., Abuqamar, S., Du, H.-N., Briggs, S.D., Mittelsten Scheid,
O., and Mengiste, T. (2009). HISTONE MONOUBIQUITINATION1 interacts with a subunit of the
mediator complex and regulates defense against necrotrophic fungal pathogens in Arabidopsis.
Plant Cell *21*, 1000–1019.

Dolan, W.L., and Chapple, C. (2017). Conservation and Divergence of Mediator Structure and
Function: Insights from Plants. Plant Cell Physiol. *58*, 4–21.

Estrada, J., Wong, F., DePace, A., and Gunawardena, J. (2016). Information Integration and
Energy Expenditure in Gene Regulation. Cell *166*, 234–244.

Flanagan, P.M., Kelleher, R.J., Sayre, M.H., Tschochner, H., and Kornberg, R.D. (1991). A
mediator required for activation of RNA polymerase II transcription in vitro. Nature *350*, 436–
438.

Flowers, E.B., Poole, R.J., Tursun, B., Bashllari, E., Pe'er, I., and Hobert, O. (2010). The
Groucho ortholog UNC-37 interacts with the short Groucho-like protein LSY-22 to control
developmental decisions in C. elegans. Dev. Camb. Engl. *137*, 1799–1805.

Fukaki, H., Tameda, S., Masuda, H., and Tasaka, M. (2002). Lateral root formation is blocked
by a gain-of-function mutation in the SOLITARY-ROOT/IAA14 gene of Arabidopsis. Plant J. Cell
Mol. Biol. *29*, 153–168.

Gaertner, B., and Zeitlinger, J. (2014). RNA polymerase II pausing during development.
Development *141*, 1179–1183.

Gasperowicz, M., and Otto, F. (2005). Mammalian Groucho homologs: redundancy or
specificity? J. Cell. Biochem. *95*, 670–687.

Gibson, D.G., Young, L., Chuang, R.-Y., Venter, J.C., Hutchison, C.A., and Smith, H.O. (2009).
Enzymatic assembly of DNA molecules up to several hundred kilobases. Nat. Methods *6*, 343–
345.

Gietz, R.D., and Woods, R.A. (2002). Transformation of yeast by lithium acetate/single-stranded
carrier DNA/polyethylene glycol method. Methods Enzymol. *350*, 87–96.

Gonzalez, N., Pauwels, L., Baekelandt, A., De Milde, L., Van Leene, J., Besbrugge, N.,
Heyndrickx, K.S., Cuéllar Pérez, A., Durand, A.N., De Clercq, R., et al. (2015). A Repressor
Protein Complex Regulates Leaf Growth in Arabidopsis. Plant Cell *27*, 2273–2287.

Goodin, M.M., Dietzgen, R.G., Schichnes, D., Ruzin, S., and Jackson, A.O. (2002). pGD
vectors: versatile tools for the expression of green and red fluorescent protein fusions in
agroinfiltrated plant leaves. Plant J. Cell Mol. Biol. *31*, 375–383.

Grbavec, D., Lo, R., Liu, Y., and Stifani, S. (1998). Transducin-like Enhancer of split 2, a
mammalian homologue of Drosophila Groucho, acts as a transcriptional repressor, interacts
with Hairy/Enhancer of split proteins, and is expressed during neuronal development. Eur. J.
Biochem. *258*, 339–349.

Gromöller, A., and Lehming, N. (2000). Srb7p is a physical and physiological target of Tup1p.
EMBO J. *19*, 6845–6852.

Hallberg, M., Hu, G.-Z., Tronnersjö, S., Shaikhibrahim, Z., Balciunas, D., Björklund, S., and
Ronne, H. (2006). Functional and physical interactions within the middle domain of the yeast
mediator. Mol. Genet. Genomics MGG *276*, 197–210.

Haruki, H., Nishikawa, J., and Laemmli, U.K. (2008). The anchor-away technique: rapid,
conditional establishment of yeast mutant phenotypes. Mol. Cell *31*, 925–932.

Havens, K.A., Guseman, J.M., Jang, S.S., Pierre-Jerome, E., Bolten, N., Klavins, E., and
Nemhauser, J.L. (2012). A synthetic approach reveals extensive tunability of auxin signaling.
Plant Physiol. *160*, 135–142.

Ito, J., Fukaki, H., Onoda, M., Li, L., Li, C., Tasaka, M., and Furutani, M. (2016). Auxin-

dependent compositional change in Mediator in ARF7- and ARF19-mediated transcription. Proc.

Natl. Acad. Sci. U. S. A. *113*, 6562–6567.

Kagale, S., Links, M.G., and Rozwadowski, K. (2010). Genome-wide analysis of ethylene-

responsive element binding factor-associated amphiphilic repression motif-containing

transcriptional regulators in Arabidopsis. Plant Physiol. *152*, 1109–1134.

Kaul, A., Schuster, E., and Jennings, B.H. (2014). The Groucho co-repressor is primarily

recruited to local target sites in active chromatin to attenuate transcription. PLoS Genet. *10*,

e1004595.

Ke, J., Ma, H., Gu, X., Thelen, A., Brunzelle, J.S., Li, J., Xu, H.E., and Melcher, K. (2015).

Structural basis for recognition of diverse transcriptional repressors by the TOPLESS family of

corepressors. Sci. Adv. *1*, e1500107.

Khakhar, A., Leydon, A.R., Lemmex, A.C., Klavins, E., and Nemhauser, J.L. (2018). Synthetic

hormone-responsive transcription factors can monitor and re-program plant development. ELife

*7*.

Kim, M.H., Cooper, D.R., Oleksy, A., Devedjiev, Y., Derewenda, U., Reiner, O., Otlewski, J.,

and Derewenda, Z.S. (2004). The Structure of the N-Terminal Domain of the Product of the

Lissencephaly Gene Lis1 and Its Functional Implications. Structure *12*, 987–998.

Kim, Y.J., Björklund, S., Li, Y., Sayre, M.H., and Kornberg, R.D. (1994). A multiprotein mediator

of transcriptional activation and its interaction with the C-terminal repeat domain of RNA

polymerase II. Cell *77*, 599–608.

Klepikova, A.V., Kasianov, A.S., Gerasimov, E.S., Logacheva, M.D., and Penin, A.A. (2016). A

high resolution map of the Arabidopsis thaliana developmental transcriptome based on RNA-

seq profiling. Plant J. *88*, 1058–1070.

Krogan, N.T., Hogan, K., and Long, J.A. (2012). APETALA2 negatively regulates multiple floral

organ identity genes in Arabidopsis by recruiting the co-repressor TOPLESS and the histone

deacetylase HDA19. Dev. Camb. Engl. *139*, 4180–4190.

Kumar, K.R.R., Blomberg, J., and Björklund, S. (2018). The MED7 subunit paralogs of Mediator
function redundantly in development of etiolated seedlings in Arabidopsis. Plant J. *96*, 578–594.

Labun, K., Montague, T.G., Krause, M., Torres Cleuren, Y.N., Tjeldnes, H., and Valen, E.
(2019). CHOPCHOP v3: expanding the CRISPR web toolbox beyond genome editing. Nucleic
Acids Res. *47*, W171–W174.

Laplaze, L., Parizot, B., Baker, A., Ricaud, L., Martinière, A., Auguy, F., Franche, C., Nussaume,
L., Bogusz, D., and Haseloff, J. (2005). GAL4-GFP enhancer trap lines for genetic manipulation
of lateral root development in Arabidopsis thaliana. J. Exp. Bot. *56*, 2433–2442.

Lee, J.E., and Golz, J.F. (2012). Diverse roles of Groucho/Tup1 co-repressors in plant growth
and development. Plant Signal. Behav. *7*, 86–92.

Lee, M., Chatterjee, S., and Struhl, K. (2000). Genetic analysis of the role of Pol II holoenzyme
components in repression by the Cyc8-Tup1 corepressor in yeast. Genetics *155*, 1535–1542.

Liu, Z., and Karmarkar, V. (2008). Groucho/Tup1 family co-repressors in plant development.
Trends Plant Sci. *13*, 137–144.

Liu, X., Galli, M., Camehl, I., and Gallavotti, A. (2019). RAMOSA1 ENHANCER LOCUS2-
Mediated Transcriptional Repression Regulates Vegetative and Reproductive Architecture.
Plant Physiol. *179*, 348–363.

Long, J.A., Ohno, C., Smith, Z.R., and Meyerowitz, E.M. (2006). TOPLESS regulates apical
embryonic fate in Arabidopsis. Science *312*, 1520–1523.

Ma, H., Duan, J., Ke, J., He, Y., Gu, X., Xu, T.-H., Yu, H., Wang, Y., Brunzelle, J.S., Jiang, Y., et
al. (2017). A D53 repression motif induces oligomerization of TOPLESS corepressors and
promotes assembly of a corepressor-nucleosome complex. Sci. Adv. *3*, e1601217.

Maji, S., Dahiya, P., Waseem, M., Dwivedi, N., Bhat, D.S., Dar, T.H., and Thakur, J.K. (2019).
Interaction map of Arabidopsis Mediator complex expounding its topology. Nucleic Acids Res.
*47*, 3904–3920.

Malik, N., Agarwal, P., and Tyagi, A. (2017). Emerging functions of multi-protein complex

Mediator with special emphasis on plants. Crit. Rev. Biochem. Mol. Biol. *52*, 475–502.

Martin, K., Kopperud, K., Chakrabarty, R., Banerjee, R., Brooks, R., and Goodin, M.M. (2009).

Transient expression in Nicotiana benthamiana fluorescent marker lines provides enhanced

definition of protein localization, movement and interactions in planta. Plant J. Cell Mol. Biol. *59*,

150–162.

Martin-Arevalillo, R., Nanao, M.H., Larrieu, A., Vinos-Poyo, T., Mast, D., Galvan-Ampudia, C.,

Brunoud, G., Vernoux, T., Dumas, R., and Parcy, F. (2017). Structure of the Arabidopsis

TOPLESS corepressor provides insight into the evolution of transcriptional repression. Proc.

Natl. Acad. Sci. U. S. A.

Matsumura, H., Kusaka, N., Nakamura, T., Tanaka, N., Sagegami, K., Uegaki, K., Inoue, T., and

Mukai, Y. (2012). Crystal structure of the N-terminal domain of the yeast general corepressor

Tup1p and its functional implications. J. Biol. Chem. *287*, 26528–26538.

McIsaac, R.S., Oakes, B.L., Wang, X., Dummit, K.A., Botstein, D., and Noyes, M.B. (2013).

Synthetic gene expression perturbation systems with rapid, tunable, single-gene specificity in

yeast. Nucleic Acids Res. *41*, e57.

Mitchell, L.A., Chuang, J., Agmon, N., Khunsriraksakul, C., Phillips, N.A., Cai, Y., Truong, D.M.,

Veerakumar, A., Wang, Y., Mayorga, M., et al. (2015). Versatile genetic assembly system

(VEGAS) to assemble pathways for expression in S. cerevisiae. Nucleic Acids Res. *43*, 6620–

6630.

Mottis, A., Mouchiroud, L., and Auwerx, J. (2013). Emerging roles of the corepressors NCoR1

and SMRT in homeostasis. Genes Dev. *27*, 819–835.

Oberoi, J., Fairall, L., Watson, P.J., Yang, J.-C., Czimmerer, Z., Kampmann, T., Goult, B.T.,

Greenwood, J.A., Gooch, J.T., Kallenberger, B.C., et al. (2011). Structural basis for the

assembly of the SMRT/NCoR core transcriptional repression machinery. Nat. Struct. Mol. Biol.

*18*, 177–184.

Papamichos-Chronakis, M., Conlan, R.S., Gounalaki, N., Copf, T., and Tzamarias, D. (2000).
Hrs1/Med3 is a Cyc8-Tup1 corepressor target in the RNA polymerase II holoenzyme. J. Biol.
Chem. *275*, 8397–8403.

Perissi, V., Jepsen, K., Glass, C.K., and Rosenfeld, M.G. (2010). Deconstructing repression:
evolving models of co-repressor action. Nat. Rev. Genet. *11*, 109–123.

Petrenko, N., Jin, Y., Wong, K.H., and Struhl, K. (2017). Evidence that Mediator is essential for
Pol II transcription, but is not a required component of the preinitiation complex in vivo. ELife *6*,
e28447.

Pierre-Jerome, E., Jang, S.S., Havens, K.A., Nemhauser, J.L., and Klavins, E. (2014).
Recapitulation of the forward nuclear auxin response pathway in yeast. Proc. Natl. Acad. Sci. U.
S. A. *111*, 9407–9412.

Pierre-Jerome, E., Wright, R.C., and Nemhauser, J.L. (2017). Characterizing Auxin Response
Circuits in Saccharomyces cerevisiae by Flow Cytometry. Methods Mol. Biol. Clifton NJ *1497*,
271–281.

Raya-González, J., Ortiz-Castro, R., Ruíz-Herrera, L.F., Kazan, K., and López-Bucio, J. (2014).
PHYTOCHROME AND FLOWERING TIME1/MEDIATOR25 Regulates Lateral Root Formation
via Auxin Signaling in Arabidopsis. Plant Physiol. *165*, 880–894.

Raya-González, J., Oropeza-Aburto, A., López-Bucio, J.S., Guevara-García, Á.A., de Veylder,
L., López-Bucio, J., and Herrera-Estrella, L. (2018). MEDIATOR18 influences Arabidopsis root
architecture, represses auxin signaling and is a critical factor for cell viability in root meristems.
Plant J. Cell Mol. Biol. *96*, 895–909.

Samanta, S., and Thakur, J.K. (2015). Importance of Mediator complex in the regulation and
integration of diverse signaling pathways in plants. Front. Plant Sci. *6*, 757.

Sato, S., Tomomori-Sato, C., Tsai, K.-L., Yu, X., Sardiu, M., Saraf, A., Washburn, M.P., Florens,
L., Asturias, F.J., Conaway, R.C., et al. (2016). Role for the MED21-MED7 Hinge in Assembly of
the Mediator-RNA Polymerase II Holoenzyme. J. Biol. Chem. *291*, 26886–26898.

Schreiber-Agus, N., Chin, L., Chen, K., Torres, R., Rao, G., Guida, P., Skoultchi, A.I., and
DePinho, R.A. (1995). An amino-terminal domain of Mxi1 mediates anti-Myc oncogenic activity
and interacts with a homolog of the yeast transcriptional repressor SIN3. Cell *80*, 777–786.

Siligato, R., Wang, X., Yadav, S.R., Lehesranta, S., Ma, G., Ursache, R., Sevilem, I., Zhang, J.,
Gorte, M., Prasad, K., et al. (2016). MultiSite Gateway-Compatible Cell Type-Specific Gene-
Inducible System for Plants. Plant Physiol. *170*, 627–641.

Song, Y.H., Estrada, D.A., Johnson, R.S., Kim, S.K., Lee, S.Y., MacCoss, M.J., and Imaizumi,
T. (2014). Distinct roles of FKF1, GIGANTEA, and ZEITLUPE proteins in the regulation of
CONSTANS stability in Arabidopsis photoperiodic flowering. Proc. Natl. Acad. Sci. *111*, 17672–
17677.

Sood, V., Cajigas, I., D'Urso, A., Light, W.H., and Brickner, J.H. (2017). Epigenetic
Transcriptional Memory of GAL Genes Depends on Growth in Glucose and the Tup1
Transcription Factor in Saccharomyces cerevisiae. Genetics *206*, 1895–1907.

Ulmasov, T., Hagen, G., and Guilfoyle, T.J. (1997). ARF1, a transcription factor that binds to
auxin response elements. Science *276*, 1865–1868.

Wang, Z.-P., Xing, H.-L., Dong, L., Zhang, H.-Y., Han, C.-Y., Wang, X.-C., and Chen, Q.-J.
(2015). Egg cell-specific promoter-controlled CRISPR/Cas9 efficiently generates homozygous
mutants for multiple target genes in Arabidopsis in a single generation. Genome Biol. *16*, 144.

Weber, E., Engler, C., Gruetzner, R., Werner, S., and Marillonnet, S. (2011). A Modular Cloning
System for Standardized Assembly of Multigene Constructs. PLOS ONE *6*, e16765.

Wong, K.H., and Struhl, K. (2011). The Cyc8-Tup1 complex inhibits transcription primarily by
masking the activation domain of the recruiting protein. Genes Dev. *25*, 2525–2539.

Wong, M.M., Guo, C., and Zhang, J. (2014). Nuclear receptor corepressor complexes in cancer:
mechanism, function and regulation. Am. J. Clin. Exp. Urol. *2*, 169–187.

Wright, R.C., Bolten, N., and Pierre-Jerome, E. (2019). flowTime: Annotation and analysis of
biological dynamical systems using flow cytometry (Bioconductor version: Release (3.10)).

Yang, Y., Li, R., and Qi, M. (2000). In vivo analysis of plant promoters and transcription factors
by agroinfiltration of tobacco leaves. Plant J. *22*, 543–551.

Zhang, Z., and Reese, J.C. (2004). Redundant Mechanisms Are Used by Ssn6-Tup1 in
Repressing Chromosomal Gene Transcription in Saccharomyces cerevisiae. J. Biol. Chem.
*279*, 39240–39250.

Zhang, Y., Gao, S., and Wang, Z. (2010). Structural and functional analysis of amino-terminal
enhancer of split in androgen-receptor-driven transcription. Biochem. J. *427*, 499–511.

Zhu, J., Jeong, J.C., Zhu, Y., Sokolchik, I., Miyazaki, S., Zhu, J.-K., Hasegawa, P.M., Bohnert,
H.J., Shi, H., Yun, D.-J., et al. (2008). Involvement of Arabidopsis HOS15 in histone
deacetylation and cold tolerance. Proc. Natl. Acad. Sci. *105*, 4945–4950.

Zhu, J., Liu, M., Liu, X., and Dong, Z. (2018). RNA polymerase II activity revealed by GRO-seq
and pNET-seq in Arabidopsis. Nat. Plants *4*, 1112–1123.

1100

1101

1102
1103

## Supplemental Figures

1105

**Supplemental Figure 1. The TPL-N terminal domain (TPLN188) interacts with the N-terminus of AtMED21**. **A.** Identifying TPL-N terminal domain interactor proteins through Yeast Two Hybrid screening identifies TPL as a problematic Bait protein, as it may silence the activation despite successful binding of a prey protein (see second row from bottom where N188 and MED21 show very weak reporter activity on 3AT). 3AT - 3-Amino-1,2,4-triazole. Plates were measured after 3 days to allow TPL-MED21 interactions to grow. **B.** Identifying TPL-N terminal domain interactor proteins through cytoplasmic split ubiquitin protein interaction assay. We tested the N-terminal and C-terminal portions of MED13 separately and divided the coding sequence at amino acid 967 (MED13N = aa1-967, MED13C = aa968-1908). Each bait tested is the *Arabidopsis* homolog cloned from cDNA from the Col-0 accession, with the

exception of *AtMED13*, which was synthesized *de novo* via Twist

(https://www.twistbioscience.com/). Plates were scanned at 3 days after plating to allow weaker

interactions to develop if they were present. **C.** TPL interacts with MED21 through an interaction

within Helices 8-9. Plates were scanned at 2 days after plating. **D.** The Helix 8 Quadruple

mutation (V145A, E146A, K148A, K149A) does not affect AtMED10B binding to TPL. Plates

were scanned at 3 days after plating. **B-D.** The relative position of the N-terminal portion of

ubiquitin (Nub) is indicated for each bait protein.

1123

**Supplemental Figure 2. Homology and structure of the MED21 subunit of the mediator**

**complex. A.** Protein levels of AtMED21 cytoSUS constructs in yeast. Two different exposure

times are shown to demonstrate the lower abundance of the truncation with only the first 31

amino acids of the AtMED21 (N31). Asterisks indicate the size predicted for the indicated

protein. **B.** Protein alignment of selected MED21 homologs from various species. Dr -

*Drosophila melanogaster*, Tr - *Takifugu rubripes*, Ms - *Mus musculus*, Gg - *Gallus gallus*, Hs -

*Homo sapiens*, Sp - *Strongylocentrotus purpuratus*, Sc - *Saccharomyces cerevisiae*, At -

*Arabidopsis thaliana*, Os - *Oryza sativa*, Sm - *Selaginella moellendorffii*, Pp - *Physcomitrella*

*patens*. Alignment was performed in CLC sequence viewer 7, using a neighbor joining method.

**C-D.** Structure of the MED21 (Cyan) & MED7 (Blue) hetero dimer, adapted from 1YKH, (Baumli

et al., 2005). The amino acids in the N-terminus that were solved are highlighted in red up to the

7th amino acid of the yeast MED21. **C.** The cartoon visualization, **D.** Surface visualization. **E.**

Core mediator (5N9J, (Nozawa et al., 2017)) with the location of MED21 and MED7 indicated

with the same colors from (C-D). In this structure the location of the MED21 N-terminus is again

indicated in red, demonstrating its close proximity to the Knob region (dotted circle).

1139

1140

**Supplemental Figure 3. Construction and characterization of the Single Locus Auxin**

**Response Circuit (SLARC). A.** Design schematic of the approach utilized to create the SLARC

through a VEGAS assembly approach. Each individual transcriptional unit (TU) was checked to

replace promoters or terminators that utilized identical sequences and replaced with an

alternative sequence indicated by a purple underline. These TUs were assembled into level 1

plasmids by Golden Gate reaction. Subsequently, they were amplified by PCR using primers

specific for the Vegas Adaptor (VA) sequences specific for their TU cassette. In example, for the

first Repressor/Substrate TU the TU was amplified using primers for VA1 and VA3 and purified

by a PCR cleanup column (NEB). The acceptor plasmid was cut with EcoRI and both TU and

acceptor plasmid was transformed into yeast and recombinant plasmids were selected on

synthetic drop out (SDO) plates lacking Leucine. **B.** Primary SLARC transformants were struck

out onto fresh SDO -Leu and imaged for Venus expression, demonstrating varying levels of

reporter expression that correlate with TPL repressor domains. Plasmid DNA was purified from

these strains and sequenced to confirm the proper recombination of TU cassettes. **C.** Time

course flow cytometry of SLARC strains following auxin addition. For all cytometry experiments,

the indicated TPL construct is fused to IAA14, because this IAA works better in haploid yeast

strains that IAA3. Every point represents the average fluorescence of 5-10,000 individually

measured yeast cells (a.u. - arbitrary units). Auxin (IAA-10µM) was added at the indicated time

(gray bar, + Aux). Four independent experiments are shown for each construct. **D.** The yeast

TPL homolog Tup1 and its partner protein Cyc8 do not repress the SLARC. Quantified

fluorescence from the single locus auxin response circuit (SLARC) introduced into Tup1 and

Cyc8 anchor away lines demonstrates no increased fluorescence from the reporter upon

depletion of Tup1 or Cyc8 from the nucleus. Anchor away depletion of Tup1 or Cyc8 results in

slower yeast growth. To normalize for this disparity in growth, Venus fluorescence was

normalized to red autofluorescence, where each pixel was normalized to the corresponding red

autofluorescence collected for that position and plotted as a boxplot. Two individual biological

replicates (two separate experiments) were evaluated, and the data was pooled. **E-F.** Med21 N

terminal deletions are viable in *Saccharomyces* and demonstrate altered SLARC transcriptional

states. **E.** A representative grayscale image of fellow fluorescence of spot plates of yeast strains

carrying SLARC plasmids in Med21 N-terminal deletions. Each is plated at an OD600 of 0.1 on

SDO with or without auxin (10µM IAA). **F.** Venus fluorescence from (**E**) was normalized to red

background (autofluorescence), where each pixel was normalized to the corresponding red

autofluorescence collected for that position and plotted as a boxplot. Two individual biological

replicates (two separate experiments) were evaluated, and the data was pooled and is

presented as boxplots.

1176

1177

**Supplemental Figure 4. Inducible MED21 rescues rapamycin induced yeast growth**

**defects. A.** Depletion of nuclear ScMed21 by Rapamycin increased cell size even in short time-

courses, consistent with its essential role in many core pathways. Scatterplots of side scatter

area by forward scatter height (SSC.A x FSC.H) indicate large scale increases in cell size in

populations of yeast with (blue) or without (red) Rapamycin treatment. **B.** Inducible Med21

(iMed21) wild type and variants cell size were examined before (red) and after (blue) treatment

with Rapamycin and b-estradiol to simultaneously deplete the wild-type Med21-FRB fusion, and

induce the transcription of the Med21 variant. Scatterplots of side scatter area by forward

scatter height (SSC.A x FSC.H) demonstrate a less disrupted cell size compared to Anchor

Away strains in (**A**). **C-D.** Histograms of Venus fluorescence in inducible Med21 (iMed21)

strains demonstrate that populations were evenly distributed around a single mean, suggesting

we were observing the immediate effects of the Med21 deletions. The histograms were built

using ggplots Density function to create a visualization of count distribution. These samples

were tested at 300 minutes (as in Figure 4D), and plotted to visualize cells at the equivalent

stage of growth, Med21 depletion, and induction. **C.** Effect of anchor away of Med21-FRB

variants alone and **D.** Depletion of ScMed21-FRB after induction (b-estradiol added 4 hours

before Rapamycin treatment) of iMed21. **E**. Conversion of the first five amino acids of ScMed21

to the corresponding sequence from AtMED21 results in an identical repression profile. Time

course flow cytometry of SLARC strains following auxin addition. For all cytometry experiments,

the indicated TPL construct is fused to IAA14, because this IAA performs better in haploid yeast

strains that IAA3. Every point represents the average fluorescence of 5-10,000 individually

measured yeast cells (a.u. - arbitrary units). Auxin (IAA-10μM) was added at the indicated time

(gray bar, + Aux). Two independent experiments are shown for each construct. **F.** Cell growth of

the strains in (**E**) indicate the swap of the N-terminal region had no effect on yeast growth or

viability. Data presented is events per microliter over the time-course of the cytometry

experiments. **C.** Protein expression analysis by western blotting of strains used in A & B. In this

ARC, TPLN188-IAA3 is N-terminally fused to 2xHA. Total protein loading levels were tested by

blotting against the housekeeping gene PGK1 (bottom panel).

1206

1207

1208

**Supplemental Figure 5. TPL multimerization requires Helix 8. A.** Cytoplasmic split ubiquitin
interaction (cytoSUS) assay on serial deletions of TPL. Interaction of bait and prey proteins
reconstitute split ubiquitin, release a synthetic transcription factor that allows growth on media
lacking Histidine and Adenine. The expression level of the bait protein can be repressed through
increased Methionine in the media. **B**. Protein levels of Nub-TPL fusions were tested by PAGE
and western blotting for the c-terminal 3xHA epitope tag included in all constructs. Deletions
longer than H1-4 are detectable at higher levels (left panel), whereas shorter isoforms required
longer exposure times to detect (right panel). Total protein loading levels were tested by the
housekeeping gene PGK1 (bottom panel). **C.** Protein expression analysis by western blotting of
tetramerization mutants expressed in yeast for cytoSUS interaction assay in Figure 5I. Prey
constructs are C-terminally fused to 2xHA. Total protein loading levels were tested by blotting
against the housekeeping gene PGK1 (bottom panel).

1223

**Supplemental Figure 6. The TPL-MED21 interaction is required for repression in plants.**

**A.** MED21 and TPL Co-Immunoprecipitated with AtMED10B from tobacco extracts. Each

construct was expressed under the viral 35S promoter, and tissues were harvested after 2 days

of injection. MED10B was purified by incubation with IgG sepharose beads (see methods), and

the presence of interacting proteins was determined by western blotting. Actin was used as a

control to determine the efficacy of washing. **B-C.** Engineering and prototyping a variant of

TPLN-IAA14$^{mED}$ which carries mutations in the EAR domain (EAR$^{AAA}$) and in the degron

(P306S) in yeast. **B.** Cartoon schematic of the mutations tested during prototyping of the

TPLN188-IAA14mED construct. In each case the identical glycine-serine linker (GS) was used

as the flexible linker between the 2xHA-TPLN188 protein and the portion of IAA14 retained in

the construct. **C.** Time course flow cytometry of TPLN-IAA14$^{mED}$ strains following auxin addition.

Strains containing the TPLN-IAA14$^{mED}$ was tested in both haploid and diploid strains and

demonstrated similar repression profiles. Every point represents the average fluorescence of 5-

10,000 individually measured yeast cells (a.u. - arbitrary units). Auxin (IAA-10µM) was added at

the indicated time (gray bar, + Aux). Two independent experiments are shown for each

construct. **D.** Identification and characterization of a novel CAS9-based insertional mutation in

*MED21*. The *MED21* genomic locus (AT4G04780) is shown as a cartoon, with a zoom in on the

beginning of the coding sequence highlighted with the amino acid sequence. The location of the

*med21/MED21* mutant (WiscDsLox461-464K13, see triangle), and the sgRNA we employed

(see green annotation and NGG PAM site) highlighted. The insertion of a G at nucleotide

position +214 after the transcriptional start site abrogates the sgRNA site (red annotation above

with i214G). A representative sequencing trace demonstrates the position where the

heterozygote carries i215G, and the predicted effect to the coding sequence is shown at the top

right – a red arrow indicates the first codon affected by the i214G mutation. The Inset pictures at

the top left demonstrate the embryo lethality phenotype in *med21$^{i214G}$/MED21* heterozygote

siliques. White asterisks indicate the embryos that have begun to degenerate. These aborted

seeds are visibly brown indicating that fertilization took place allowing the seed coat to form

before development failed. **E.** *med21/MED21* heterozygotes are haplo-sufficient for lateral root

development. Lateral root density (number of lateral roots / primary root length) was calculated

at 10 days post germination. Lower case letters indicate significant difference (ANOVA and

Tukey HSD multiple comparison test; $p<0.001$). **F.** Ratio of lateral root lengths to total root

lengths (lateral root lengths + primary root length) in dCAS9 repressor lines targeting *MED21*

calculated at 10 days post germination. Statistical tests (ANOVA and Wilcox test) are reported

above the graph. **G.** Representative root traces of dCAS9 repressor lines targeting *MED21*

calculated at 10 days post germination.

1259