**Genome-wide data inferring the evolution and population**

**demography of the novel pneumonia coronavirus (SARS-CoV-2)**

Bin Fang[1*], Linlin Liu[1*], Xiao Yu[1*], Xiang Li[1*], Guojun Ye[1], Juan Xu[3], Ling Zhang[3] ,

Faxian Zhan[1], Guiming Liu[4], Tao Pan[2#] , Yilin Shu[2#] , Yongzhong Jiang[1#]

1. Influenza Reference Laboratory , Institute of Health Inspection and Testing , Hubei

Provincial Center for Disease Control and Prevention, Wuhan 430079, China

2. Anhui Province Key Laboratory for Conservation and Exploitation of Biological

Resource, College of Life Sciences, Anhui Normal University, Wuhu 241000, Anhui,

China

3. Hubei Province Key Laboratory of Occupational Hazard Identification and Control,

School of Public Health, Wuhan University of Science and Technology, Wuhan,

Hubei, 430065, China

4. Beijing Agro-Biotechnology Research Center, Beijing Academy of Agriculture and

Forestry Sciences, 100097, Beijing, China

*These authors have equal contribution to this study.

#Correspondence: Tao Pan, pantao20070778@163.com; Yilin Shu, sjshuyilin@163.com;

Yongzhong Jiang, hbcdcxd@163.com.

**Other Author's Email:**

Bin Fang: nicolfang@163.com

Guojun Ye: gjy0822@sina.comv

Lin-lin Liu: mice0809@163.com

Xiang Li: li_xiang75@yeah.net

Xiao Yu: fish-yuxiao@hotmail.com

Juan Xu: xujuan9270@163.com

Ling Zhang: zhangling@wust.edu.cn

Faxian Zhan: zhanfx@163.com

Guiming Liu: mingguiliu@aliyun.com

**Abstract**

Since December 2019, coronavirus disease 2019 (COVID-19) emerged in

Wuhan, Central China and rapidly spread throughout China. Up to March 3, 2020,

SARS-CoV-2 has infected more than 89,000 people in China and other 66 countries

across six continents. In this study, we used 10 new sequenced genomes of

SARS-CoV-2 and combined 136 genomes from GISAID database to investigate the

genetic variation and population demography through different analysis approaches

(e.g. Network, EBSP, Mismatch, and neutrality tests) in the previous three months.

The results showed that eighty haplotypes had 183 substitution sites, including 27

parsimony-informative and 156 singletons. Sliding window analyses of genetic

diversity suggested a certain range in mutations and variation in genomic abundance

of SARS-CoV-2, which could be explaining the existing widespread and high

adaptation of the deadly virus. Phylogenetic analysis showed that pangolin may not be

an intermediate host. The network indicated that, in the original haplotype (H14), one

patient sample lived near the Huanan seafood market (approximately 2 km), which

indicating high possibility of the patient having a history of unconscious contact with

this market. However, based on this clue, we cannot accurately conclude that

whether this market was the origin center of SARS-CoV-2. Additionally, 16 genomes,

collected from this market, assigned to 10 haplotypes, indicated a circulating infection

within the market in a short term, which would have led to the outbreak of

SARS-CoV-2 in Wuhan and other areas. The EBSP results showed that the first

estimated expansion date began from 7 December 2019, which indicated that the

transmission of SARS-CoV-2 could have begun from person to person in mid to late

November.

**Key words** SARS-CoV-2; metagenomic next-generation sequencing; virus

evolution;   population demography; phylogenetic relationship

**Introduction**

As the largest non-segmented genomes among all the RNA viruses (about 30 kb

in length), Coronaviruses (CoVs) own the plasticity due to the mutation and

recombination, which increased the potential risks of spread across species [1, 2]. The

COVID-19 (original named 2019-nCOV) is the seventh member of enveloped RNA

coronavirus (subgenus Sarbecovirus, Orthocoronavirinae subfamily) [3]. In early

December, 2019, an unexplained pneumonia associated with the severe acute

respiratory syndrome coronavirus 2 (SARS-CoV-2) emerged in Wuhan, China [4, 5].

The emergence and rapid spread of the SARS-CoV-2 between infected and healthy

people became so devastating as a large population within Wuhan was getting

infected [6, 7]. There's evidence that SARS-CoV-2 may have originated from bats, but

there is no clear information about the intermediate host that transferred it to humans

[8-10]. On 30 January 2020, the World Health Organization (WHO) declared the

outbreak of COVID-19 to be a Public Health Emergency of International Concern. Up

to March 3, 2020, SARS-CoV-2 has infected more than 89,000 people in China and

other 66 countries across the six continents (source: World Health Organization

report).

Despite the worldwide rapid spread, the genomic variation dynamics,

evolutionary rate, and virus transmission dynamics of SARS-CoV-2 are not yet well

understood. In several recent studies, phylogenetic relationships, variations,

evolutionary rates, and propagation dynamics were analyzed using limited genomic

data from the SARS-CoV-2 [5, 9-17]. As the epidemic progresses, many research

institutes around the world have obtained and submitted SARS-CoV-2 genome

sequences, increasing the number of SARS-CoV-2 genome sequences in the Data

Centers (i.e, GISAID). Given the extremely rapid spread of SARS-CoV-2, an updated

analysis with significantly larger sample sizes by incorporating cases throughout the

world is urgently needed. This would allow for more accurate identification of

SARS-CoV-2 variant dynamics, evolutionary rates, virus transmission dynamics, and

epidemic history in order to effectively implement public health measures, promote

drugs and vaccines development and to efficiently prevent similar epidemics in the

future.

Recent research findings have suggested bat and other non-bat intermediate

mammals (such as pangolin) as potential intermediate host for the SARS-CoV-2 that

have so far been widely transmitted to humans [5, 9, 18]. According to the medical

information of the first patients to be infected in Wuhan, 27 out of 41 patients were

found to be traders of wild mammals in the Huanan seafood market, which directly

suggests that the Huanan seafood market as the possible source of origin of the

SARS-CoV-2, and later got transmitted it to other areas by the infected people [4, 19].

However, due to some infected persons having no relation or access to the Huanan

food market, some researchers are skeptical of the market as the only actual origin or

the source of SARS-CoV-2 transmission to humans [4, 19, 20]. Therefore, up to date,

in the absence of key potential intermediary host, the origin and transmission pattern

of SARS-CoV-2 still need to be thoroughly probed for a more reliable and accurate

information on the origin and transmission mechanisms of the virus.

In this study, 56 full genome sequences of outgroups and 136 genomes of

SARS-CoV-2 from GISAID EpiFluTM database (access date 22 February 2020) were

collected. Ten new sequenced genomes of SARS-CoV-2 were obtained by

macrogenomic sequencing. The time origin, genetic diversity, transmission dynamics

and evolutionary history of SARS-CoV-2 were analyzed based on the genomic data to

provide understanding on the origin, transmission pathway, and evolutionary

characteristics of SARS-CoV-2 outbreak.

**Materials and methods**

*Patients and samples*

In this study, ten patients with unexplained viral pneumonia from five hospitals

in Hubei Province were included. Four samples were collected in December, four in

January as well as two samples in February. The December samples were examined

by LightCycle 480II fluorescent PCR instrument (Roche, Basel, Switzerland) for

investigating 26 respiratory pathogens and detection of the virus using SARS-Cov and

MERS-Cov primer probes. All the ten samples were detected by the SARS-CoV-2

primer probes. Two samples were used in the virus isolation. The 10 samples were

also subjected to metagenomic next-generation sequencing. Specific sample

information is presented in Table 1.

*Virus isolation*

Virus were isolated by HUH7 cells from one lavage fluid sample and one

pharynx swab sample. The cells were monitored daily for cytopathic effects by light

microscopy. HUH7 cells were harvested after 6 days of culturing at 37℃. The

supernatant was collected to extract the total RNA nucleic acid through EZ1 virus

mini kit v2.0 (955134; Qiagen, Heiden, Germany), detection was performed by Light

Cycle 480II fluorescent PCR instrument (Roche, Basel, Switzerland) followed by

subsequent sequencing.

*Library preparation and sequencing*

Total RNA extracted from 10 samples were subjected to metagenomic next

generation sequencing testing by EZ1 virus mini kit v.2.0 (955134; Qiage, Heiden,

Germany). TruSeq Stranded Total RNA Library Prep Kit (Illumina, San Diego, CA,

USA) was used to remove rRNA, reverse transcription and synthesize the

double-stranded DNA. The fragmenting, modifying the ends, connecting the joints

and enriching of DNA were done by NexteraXT library prepkit (Illumina, San Diego,

CA, USA), then the DNA library was obtained. Iseq$^{TM}$ 100 i1 Cartridge, Miseq v2

reagent kit, and High output Reagent Cartridge (illumina, San Diego, CA, USA) were

used for deep sequencing in Iseq100, Miseq, and Miniseq platforms (illumina, San

Diego, CA, USA), respectively. About 11.5 GB data were obtained for each sample.

*Virus genome analysis*

Using CLC Genomics Workbench 12 and Geneious 12.0.1 software (QIAGEN

Bioinformatics, Redwood City, CA, USA) and using reference sequence

BetaCoV/Wuhan-Hu-2019 (EPI_ISL_402125), the raw sequencing data of illumina

were analyzed. The open reading frames of the verified genome sequences were

predicted using Geneious and annotated using the Conserved Domain Database [21].

*Phylogenetic reconstructions*

In this study, for confirmation of the evolutionary history of SARS-CoV-2

(Table S1), 136 complete genomes were retrieved from GISAID (Table S1, access on

22 February, 2020), and combined with our sequenced 10 genomes of this virus. In

total, 56 outgroups were collected following the previous study [5]. Mafft v.7.450 was

used to align the sequence of SARS-CoV-2 with reference sequences [22].

Phylogenetic analyses of the complete genome and major coding regions were done

with RAxML v.8.2.9 [23] in 1000 bootstrap replicates, employing the general time

reversible nucleotide substitution model. The tree was visualized with FigTree v.1.4.3

[24].

*Population structure*

On this analysis technique, to precisely decode the evolutionary history of

SARS-CoV-2, the genome EPI_ISL_402131 (bat-RaTG13-CoV) from GISAID was

also included as the outgroup following the procedure outlined in the previous study

[25], because it is the closest sister betacoronavirus to SARS-CoV-2.3. The 136

complete genome sequences of SARS-CoV-2 and an outgroup (bat-RaTG13-CoV)

were aligned using MAFFT then the alignment was manually checked using Geneious.

For probing the haplotypes relationships among localities, the Network v.4.6.1

(Fluxus Technology, Suffolk, UK) was used to construct the minimum-spanning

network based on the full median-joining algorithm [26]. During the alignment, 10

genomes containing ambiguous sites or "N" bases or more degenerated bases were

excluded in this study (Table S1). In addition, 4 genome sequences (EPI_ISL_409067,

EPI_ISL_412981, EPI_ISL_407071 and EPI_ISL_408489) with one degenerated

bases were included, which were divided into two sequences without degenerated

bases. For ensuring the accuracy of this analysis, the 5'UTR and 3'UTR contain

missing and ambiguous sites of both regions were excluded in the following

alignment analyses. DnaSP v.5.10 [27] was used to convert the relevant format.

Population genetic indices in was estimated in DnaSP, including nucleotide diversity

($\pi$) and haplotype diversity (*Hd*). $F_{ST}$ between haplotypes was calculated in Mega 6.0

based on haplotype frequency differences with 10,000 permutations [28].

Additionally, a sliding window of 500 bp and steps of 50 bp were used to estimate

nucleotide diversity ($\pi$) for the entire alignment this data. Nucleotide diversity for the

entire alignment was plotted against midpoint positions of each window. To indicate

the relative position of the mutations in the genome, we selected the EPI_ISL_402124

sequence as the reference genome.

*Population demography*

The alignment was then imported into DnaSP for haplotype analyses. Population

size changes were estimated based on a constant population size hypothesis using

DnaSP, in combination with neutrality tests (Tajima's *D* and *Fu*'s Fs). The mismatch

distribution was estimated based on the genomes of SARS-CoV-2 in Arlequin to test

the hypothesis of recent population growth. The Harpending's raggedness index (Rag)

and the sum of squares deviation (SSD) were used to determine the smoothness of

observed mismatch distribution and the degree of fit between observed and simulated

data [29]. Because of the sensitivity to demographic changes of neutral tests, Tajima's

*D* [30] and *Fu*'s Fs [31] were estimated using 10,000 coalescent simulations to assess

the significance in Arlequin. In addition, the Extended bayesian skyline plot (EBSP)

analysis was conducted to examine past population dynamics of SARS-CoV-2 based

on 146 genomes by BEAST v.1.8 [32]. The divergence times were estimated in

BEAST using a Bayesian Markov chain Monte Carlo (MCMC) method with a strict

clock. In this study, the substitution rate was set as $0.92 \times 10^{-3}$ (95% CI,

$0.33 \times 10^{-3}$-$1.46 \times 10^{-3}$) substitution/site/year based on the most recent estimation for

SARS-CoV-2 [25]. The other parameters were set as follows: extended Bayesian

skyline process, 10 million MCMC generations, sampling every 1,000th iteration, the

initial 25% burn-in. Tracer was used to check the convergence of the MCMC analyses

(ESS values >200). Convergence of the two independent MCMC runs was assessed in

Tracer, as was convergence of model parameter values (effective sample size [ESS])

to ensure ESS values >200.

### Results and discussion

*Genomic variations of SARS-CoV-2*

According to the database, the genome size of SARS-CoV-2 varied from 29,409

bp to 29,911 bp. In the network dataset, the aligned matrix was 29,130 bp in length

including 183 variable sites, of which 27 were parsimony-informative and 156 were

singletons, which were classified as 80 haplotypes (Table S2). Nucleotide diversity ($\pi$)

was $0.16 \times 10^{-3} \pm 0.02 \times 10^{-3}$ (Standard Deviation, SD) and haplotype diversity (*Hd*) was

$0.954 \pm 0.013$ (SD) and variance of *Hd* was $0.16 \times 10^{-3}$ (Table S3). In this study, our

new sequenced 10 genomes of SARS-CoV-2 represented 10 haplotypes (H3, H8, H10,

H14, H16, H28, H29, H30, H76 and H78; Table S1). Among these 10 genomes, four

samples were collected from those patients who had been exposed to the Huanan

seafood market in Wuhan. Additionally, the HBCDC-HB-03/2019 belonged to the

core H3 haplotype and most of them represented new haplotype. Sliding window

analysis of SARS-CoV-2 revealed significant regional variation across the alignment

(Fig. 1). The plot readily showed a relative high degree of nucleotide variation

amongst the aligned SARS-CoV-2 genomes for any given window of 500 bp and

steps of 50 bp, with the $\pi$ ranging from 0.00005 to 0.0014. Additionally, based on the

curve, there were several relative high variation area, compared with other fragments

(Fig. 1). The $F_{ST}$ analyses among 80 haplotypes ranged from 0.00003 (e.g, H3 and H9;

H14 and H15; H3 and H20) to 0.00134 (H19 and H80), which indicated genetic

differentiation among these haplotypes (Fig. 2, Table S4). Overall, the H19 showed

relatively large genetic distance with other haplotypes, while the H3 showed opposite

pattern, and was also confirmed by the network figure (Fig. 2, Table S4). Overall,

SARS-CoV-2 maintain the relatively rich mutations that has occurred across the world,

which may be the main reason of the numerous subgenomic RNAs generated during

the viral replication [3, 9]. This phenomenon also provides the possibility of

widespread and adaptation for this virus. Fortunately, under the strict quarantine

policy in China since 23 January 2020, the circulation and spreading of some

haplotypes may have relatively reduced in circulation frequency relatively, compared

with the early stage of COVID-19.

*Phylogenetic relationship of SARS-CoV-2*

The virus's popularity and its intermediate host has been a topic of great concern.

Some researchers have suggested that the SARS-CoV-2 originated from the bats [5,9].

However, some other researchers have proposed that the non-bat intermediate

mammals (e.g. pangolins) may be the transmission path of this genus [18]. However,

other studies have also failed to confirm the pangolins as the intermediate host to this

viral genus [33]. In this study, based on the phylogenetic tree, bat-RaTG13-CoV was

the sister group to the SARS-CoV-2 and the two pangolin samples collected from

Guangdong which showed a relatively large genetic distance from the SARS-CoV-2

(Fig. 3 and S1). Based on this finding, , the study suggested that pangolin may not be

the intermediate host, as had been reiterated by other previous study [33]. As the

Huanan seafood market was closed on 1 January 2020, this created many challenges

in identifying the first intermediate hosts (whether people or animal) of SARS-CoV-2.

Therefore, more sampling and analysis of the sample from this area may suggest

clearer results for more concrete conclusions in future studies.

*Evolutionary relationships of SARS-CoV-2 haplotypes*

The evolutionary network of 80 haplotypes of SARS-CoV-2, with bat-RaTG13-CoV

as the outgroup, is shown in Figure 4. The network analysis showed typical star

network, with several core haplotype node (H3, H14, H15) and edge haplotype nodes.

In the network, fifty-five satellite haplotypes and H14 connected to the H3 haplotype;

eighteen satellite haplotypes and H15+H3 connected to the H14 haplotype; and five

satellite haplotypes and H14 connected to H15. Most of these haplogroups were

separated by one to six mutations, except two haplotypes (H19 and H80, Fig. 4). The

evolutionary network showed that bat-RaTG13-CoV to be connected through a

hypothesized haplotype (mv6) to the H15 and H31 haplotypes by single mutations

(Fig. 4). However, the mutations between mv6 and bat-RaTG13-CoV was more than

800, which indicated that SARS-CoV-2 still has a relative distant kinship with the

outgroup (bat-RaTG13-CoV).

As the most abundant haplotype, H3 included 28 samples, while 55 satellite

haplotypes are directly derived from the H3 haplotype (Fig. 4 and Table S1).

Moreover, many haplotypes from other countries should also be derived from the H3

haplotype. The current samplings showed that the H3 haplotype has been found to be

in 28 samples, but 12 of 28 samples were collected from Wuhan in Hubei Province.

Fifteen of the 55 satellite haplotypes were also collected from Hubei Province (Fig. 4).

One possible explanation was that a common haplotype from the Huanan seafood

market (Fig. 4) was rapidly circulated at an early stage of human-to-human

transmissions. It is worth noting that in the H14, there are two samples

(EPI_ISL_406801 and EPI_ISL_ 412979) from Wuhan. Although these two hosts

didn't directly link with the Huanan market, one of the host (EPI_ISL_ 412979)

having lived in a residential area about 2 kilometers from the Huanan seafood market.

This also provides the possibility of them having indirect possible contact with the

Huanan seafood market.

Additionally, the networks of 24[th] December, 2019 - 30[th] December, 2019 and

24<sup>th</sup> December, 2019 - 6<sup>th</sup> January, 2020 also suggested that the most frequent

haplotype (H3) occurred. All the haplotypes were collected from Wuhan, which

indicated that Wuhan may have acted as the important origin center (Fig. 4). On the

other hand, in this study, 16 sequences of Huanan seafood market (4 new sequenced

and 12 GISAID data) were collected. Fifteen of out of the sixteen samples were

collected before 1<sup>st</sup> Januray, 2020 and one samples was collected on 8<sup>th</sup> Januray, 2020.

These samples were represented ten haplotypes (H2, H3, H4, H5, H6, H7, H8, H10,

H11, H16), which indicated the high proportion of haplotype diversity in Huanan

seafood market. Among those haplotypes, the H3 haplotype, the most abundant, was

present in 6 samples from Huanan seafood market, while the other haplotypes were

directly derived from the H3 haplotype. All the samples from the Huanan seafood

market had the H3 haplotype and other 9 derived haplotypes (Fig. 4), indicating that

there were circulated infections within the market in a short term. Noteworthy, in the

network, a total of 65 virus samples from 15 other countries were assigned to 43

haplotypes. Among them, 27 haplotypes were satellite haplotypes of H3 haplotype, 11

haplotypes were satellite haplotypes of H14 haplotype, and 3 haplotypes were satellite

haplotype of H15 haplotype. Indeed, most of haplotypes originated from the H3,

which may hint the phenomenon was related to the input of virus carriers from

Wuhan.

Based on the outgroup (bat-RaTG13-CoV) having a possible direct connection

with outgroup H15 and H31, it indicates that both the H15 and H31 were the

suggested ancestral haplotypes (Fig. 4), as had also been suggested by other previous

study [25]. The H15 was only recovered from five Shenzhen (Guangdong) samples,

while H31 included 3 United States samples and 1 Fujian sample from China (Fig. 4).

Based on the epidemiological statistics, all the patients from the H15 were ever

traveled to Wuhan [16]. As to the H31, three genomes from the same patient in Unite

State [34] and 1 genome from Fujian of China (Table S1) were included. For the

patient in Unite State, He may have infected during the period of visiting his family in

China [25]. The travel history of Fujian sample was unclear, but we cannot deny that

it has a history of unconscious Wuhan contacts. Based on the theory in the previous

study [25], this current study also supported two main evolutionary paths, that the

available haplotypes could be from H15 through H14 to H3, or from H31 through

H14 to H3 (Fig. 4). Both these two path demonstrate that H14 was the key connection

from an ancestral haplotype to H3. However, unlike the previous research, in H14,

two individuals (EPI_ISL_406801 and EPI_ISL_ 412979) were collected from Wuhan.

Additionally, there was another new haplotype (H28, EPI_ISL_412980), which

originated from the H14, and was also collected from Wuhan. Although the three

hosts of these two haplotypes didn't directly link with the Huanan seafood market,

one of the hosts (EPI_ISL_ 412979) lived in a residential area about 2 kilometers

from the Huanan seafood market. This also provides the possibility for that host to

have unconsciously indirect contact with the Huanan seafood market. Overall, it

cannot further be proved the SARS-CoV-2 in the Huanan seafood market had been

transmitted from other places or that Huanan seafood market did not host the original

source of SARS-CoV-2. Due to the closing up of the Huanan seafood market on 1$^{st}$ of

January, 2020, it made it difficult to determine whether there were intermediate hosts

in the market. To accurately determine whether this market was the central origin,

further collection of early SARS-CoV-2 samples within/around the Huanan seafood

market is needed to provide genomic information and epidemiological survey data

needed in integration analysis. However, based on the facts provided in the current

study undertaken in the early days of the outbreak of the pneumonia infection in

Wuhan, the Huanan seafood market promoted the spread of SARS-CoV-2, and since

then, infected travelers have spread to throughout China and other countries.

*Population size expansion of SARS-CoV-2*

As to population demography of SARS-CoV-2, the EBSP results revealed that it

experienced an expansion of effective population size for the last 66 days prior to $11^{th}$

of February, 2020 (Fig. 5). With the latest one being sampled on $11^{th}$ February 2020,

the first expansion date is estimated to had begun from $7^{th}$ December 2019. The

mismatch in the distribution of the total populations was also clearly shown through

unimodal (Fig. 6) and the neutral tests (Tajima's D and Fu's Fs) revealed statistically

significant population expansion of SARS-CoV-2 (Table S3), which also supported

the EBSP results. In addition, with the change of time, multiple evidence (i.e.

haplotype number, Tajima's D, Fu's FS) indicated a relatively highest population

growth at around $21^{st}$ - $27^{th}$ of January (Table S3). The demographic expansion of

SARS-CoV-2 during this period was consistent with the patients first diagnosed on $8^{th}$

December, 2019 [4,19] and the previously suggested most recent common ancestor

(TMRCA) dates for SARS-CoV-2 at $6^{th}$ December, 2019) [17]. Additionally,

considering that the virus has longest reported incubation period of up to 24 days [6],

the virus may had first infected humans in mid to late November, which is basically

consistent with the results of previous studies [25]. Therefore, in this study, EBSP

revealed that SARS-CoV-2 experienced an effective population size expansion since

$7^{th}$ December 2019, which was also supported by a star-like network, EBSP, the

neutral tests (Fu's and Tajima's D test) and mismatch analysis.

**Contributors**

Yilin Shu, Tao Pan, and Yongzhong Jiang conceived the research, analyzed the data,

interpreted the results, and wrote the draft manuscript; Bin Fang, Linlin Liu, Xiao Yu

and Xiang Li performed macrogenome sequencing experiments. Xiao Yu and Xiang

Li performed virus isolation experiments. Guojun Ye, Bin Fang, Juan Xu, Ling,

Zhang, Yilin Shu and Faxian Zhan collected data. All authors reviewed and approved

the final version of the manuscript.

**Declaration of interests**

We declare no competing interests.

**Acknowledgements**

We thank scientists and researchers for depositing the whole genomic sequences of

Novel Pneumonia Coronavirus (SARS-CoV-2) in the Global Initiative on Sharing All

Influenza Data (GISAID) EpiFlu™; and the GISAID database for allowing us access

to sequence for non-commercial scientific research. We also are grateful to Dr. Oscar

Omondi Donde (Egerton University, Kenya) for polishing the manuscript language.

Thanks to Yifa Zhu, Yangyang Tao and Xierong Li (Tianmen Center for Disease

Control and Prevention) for collecting samples and thanks to Maoyi Chen, Jie Hu,

Chunlin Mao (Jingzhou Center for Disease Control and Prevention) for sampling.

**Reference**

[1].   Cui J, Li F, Shi ZL. Origin and evolution of pathogenic coronaviruses. Nat

Rev Microbiol. 2019;17(3):181-192.

[2].   Gorbalenya AE. Severe acute respiratory syndrome-related coronavirus–The

species and its viruses, a statement of the Coronavirus Study Group. BioRxiv.

2020;DOI: https://doi.org/10.1101/2020.02.07.937862.

[3].   Zhu N, Zhang DY, Wang WL, et al. A novel coronavirus from patients with

pneumonia in China, 2019. New Engl J Med. 2020;382(8):727-733.

[4].   Huang C, Wang Y, Li X, et al. Clinical features of patients infected with 2019

novel coronavirus in Wuhan, China. The Lancet. 2020;395(10223):497-506

[5].   Lu RJ, Zhao X, Li J, et al. Genomic characterisation and epidemiology of

2019 novel coronavirus: implications for virus origins and receptor binding.

Lancet. 2020; DOI: 10.1016/S0140-6736(20)30251-8.

[6].   Guan WJ, Ni ZY, Hu Y, et al. Clinical characteristics of 2019 novel

coronavirus    infection    in    China.    N    Engl    J    Med. 2020;    DOI:

10.1056/NEJMoa2002032.

[7].   Yang Y, Lu QB, Liu MJ, et al. Epidemiological and clinical features of the

2019    novel    coronavirus    outbreak    in    China.    MedRxiv.

2020;DOI: https://doi.org/10.1101/2020.02.10.20021675.

[8].   Wassenaar TM, Zou Y. 2019_nCoV: Rapid classification of betacoronaviruses

and    identification    of    traditional    Chinese    medicine    as    potential    origin    of

zoonotic coronaviruses. Lett Appl Microbiol. 2020; DOI: 10.1111/lam.13285.

394 [9]. Zhou P, Yang XL, Wang XG, et al. A pneumonia outbreak associated with a

395   new coronavirus of probable bat origin. Nature. 2020; DOI:

396   10.1038/s41586-020-2012-7.

397 [10]. Wu F, Zhao S, Yu B, et al. A new coronavirus associated with human

398   respiratory disease in China. Nature. 2020; DOI: 10.1038/s41586-020-2008-3.

399 [11]. Benvenuto D, Giovannetti M, Ciccozzi A, et al. The 2019-new coronavirus

400   epidemic: evidence for virus evolution. J Med Virol. 2020;

401   DOI: 10.1101/2020.01.24.915157

402 [12]. Ceraolo C, Giorgi FM. Genomic variance of the 2019-nCoV coronavirus. J

403   Med Virol. 2020; DOI:10.1002/jmv.25700.

404 [13]. Chen LJ, Liu WY, Zhang Q, et al. RNA based mNGS approach identifies a

405   novel human coronavirus from two individual pneumonia cases in 2019

406   Wuhan outbreak. Emerg Microbes Infec. 2020; 9(1):313-319.

407 [14]. Paraskevis D, Kostaki EG, Magiorkinis G, et al. Full-genome evolutionary

408   analysis of the novel corona virus (2019-nCoV) rejects the hypothesis of

409   emergence as a result of a recent recombination event. Infect Genet Evol. 2020;

410   79:104212.

411 [15]. Chan JF, Kok KH, Zhu Z, et al. Genomic characterization of the 2019 novel

412   human-pathogenic coronavirus isolated from a patient with atypical

413   pneumonia after visiting Wuhan. Emerg Microbes Infec. 2020;9(1):221-236.

414 [16]. Chan JFW, Yuan SF, Kok KH, et al. A familial cluster of pneumonia associated

415   with the 2019 novel coronavirus indicating person-to-person transmission: a

study of a family cluster. Lancet. 2020; 395(10223):514-523.

[17]. Li XW, Wang W, Zhao XF, et al. Transmission dynamics and evolutionary

history of 2019-nCoV. J Med Virol. 2020; DOI: 10.1002/jmv.25701.

[18]. Lam TTY, Shum MHH, Zhu HC, et al. Identification of 2019-nCoV related

coronaviruses in Malayan pangolins in southern China. bioRxiv. 2020;

DOI: https://doi.org/10.1101/2020.02.13.945485.

[19]. Li Q, Guan XH, Wu P, et al. Early transmission dynamics in Wuhan, China, of

novel coronavirus–infected pneumonia. New Engl J Med. 2020;

DOI: 10.1056/NEJMoa2001316.

[20]. Cohen J. Wuhan seafood market may not be source of novel virus spreading

globally. Science. 2020;10.1126/science.abb0611.

[21]. Marchler-Bauer A, Bo Y, Han LY, et al. CDD/SPARCLE: functional

classification of proteins via subfamily domain architectures. Nucleic Acids

Res. 2017;45(D1):D200-D203.

[22]. Katoh K, Standley DM. MAFFT multiple sequence alignment software

version 7: improvements in performance and usability. Mol Biol Evol.

2013;30(4):772-780.

[23]. Stamatakis A. RAxML version 8: a tool for phylogenetic analysis and

post-analysis of large phylogenies. Bioinformatics. 2014;30:1312–1313.

[24]. Rambaut A. FigTree v1.4.3; 2016. Available at:

http://tree.bio.ed.ac.uk/software/figtree/ 2016.

[25]. Yu WB, Tang GD, Zhang L, et al. Decoding the evolution and transmissions of

the novel pneumonia coronavirus (SARS-CoV-2) using whole genomic data.

ChinaXiv. 2020;202002.00033v2.

[26].  Bandelt H, Forster P, Röhl A. Median-joining networks for inferring

intraspecific phylogenies. Mol Biol Evol. 1999;16(1):37-48.

[27].  Librado P, Rozas J. DnaSP v5: a software for comprehensive analysis of DNA

polymorphism data. Bioinformatics. 2009;25(11):1451–1452.

[28].  Tamura K, Stecher G, Peterson D, et al. MEGA6: molecular evolutionary

genetics analysis version 6.0. Mol Biol Evol. 2013;30(12):2725–2729.

[29].  Harpending HC. Signature of ancient population growth in a low-resolution

mitochondrial DNA mismatch distribution. Hum Biol. 1994 ;66(4):591-600.

[30].  Tajima F. The effect of change in population size on DNA polymorphism.

Genetics. 1989;123(3):597–601.

[31].  Fu YX. Statistical tests of neutrality of mutations against population growth,

hitchhiking and background selection. Genetics. 1997;147(2):915–925.

[32].  Drummond AJ, Suchard MA, Xie D, et al. Bayesian phylogenetics with

BEAUti and the BEAST 1.7. Mol Biol Evol. 2012;29(8):1969–1973.

[33].  Liu P, Jiang JZ, Wan XF, et al. Are pangolins the intermediate host of the 2019

novel coronavirus (2019-nCoV) ? bioRxiv. 2020;DOI:

https://doi.org/10.1101/2020.02.18.954628.

[34].  Holshue ML, DeBolt C, Lindquist S, et al. First Case of 2019 Novel

Coronavirus in the United States. N Engl J Med 2020;

10.1056/NEJMoa2001191.

**Table 1. Sample information for 10 SARS-Cov-2 infected patients.**

| Sample name | Sample type | Collection date | Collection site | Exposure to huanan seafood marke |
|---|---|---|---|---|
| BetaCoV/Wuhan/HBCDC-HB-02/2019 | Bronchoalveolar lavage fluid | 30th Dec,2019 | Wuhan, Hubei Province | Yes |
| BetaCoV/Wuhan/HBCDC-HB-01/2019 | Bronchoalveolar lavage fluid | 30th Dec,2019 | Wuhan, Hubei Province | Yes |
| BetaCoV/Wuhan/HBCDC-HB-03/2019 | Bronchial scraping | 30th Dec,2019 | Wuhan, Hubei Province | Yes |
| BetaCoV/Wuhan/HBCDC-HB-04/2019 | Sputum | 30th Dec,2019 | Wuhan, Hubei Province | No |
| BetaCoV/Wuhan/HBCDC-HB-04/2020 | Throat swab | 18th Jan,2020 | Wuhan, Hubei Province | No |
| BetaCoV/Wuhan/HBCDC-HB-03/2020 | Throat swab | 18th Jan,2020 | Wuhan, Hubei Province | No |
| BetaCoV/Wuhan/HBCDC-HB-02/2020 | Throat swab | 17th Jan,2020 | Wuhan, Hubei Province | No |
| BetaCoV/Jingzhou/HBCDC-HB-01/2020 | Throat swab and cultured virus | 8th Jan,2020 | Jingzhou,Hubei Province | Yes |
| BetaCoV/Wuhan/HBCDC-HB-06/2020 | Faeces | 7th Feb,2020 | Wuhan, Hubei Province | No |
| BetaCoV/Tianmen/HBCDC-HB-07/2020 | Throat swab | 8th Feb,2020 | Tianmen, Hubei Province | No |

**Figure Legend**

**Fig. 1. Sliding window analyses showing the nucleotide diversity based on**

**alignment of genomes of SARS-CoV-2.** The red line shows the value of nucleotide

diversity ($\pi$) in a sliding window analysis of window size 500 bp with step size 50, the

value is inserted at its mid-point. EPI_ISL_402124 sequence was selected as the

reference genome to indicate the mutation position.

**Fig. 2. Pairwise $F_{ST}$ among 80 haplotypes o**f SARS-CoV-2. Consistent with the

Table S4.

**Fig. 3. Simplify phylogenetic tree of SARS-CoV-2 form Fig. S1.** The nodal

numbers are ML bootstrap values.

**Fig. 4. Median-joining network with node sizes proportional to the frequencies of**

**haplotypes in SARS-CoV-2 (A, B and C).** (A)All the individuals collected during

24[th] December to 30[th] December in 2019; (B) All the individuals collected during 24[th]

December in 2019 to 6[th] January in 2020; (C) All the individuals collected before 11[th]

February in 2019. The numbers of mutations separating the haplotypes are shown on

the branches, except for the lower than seven-step mutations. The little red diamond

nodes indicate undetected haplotypes. The sampling areas are indicated by different

colors .

**Fig. 5. Population fluctuation infered by Extended bayesian skyline plot (EBSP)**

**of SARS-CoV-2.** The x-axis indicates time in days BP, and the y-axis indicates the

effective population size divided by generation time in units of $N_{e\tau}$ (the product of

effective population size and generation time in days). The blue areas represent 95 %

highest posterior density. Time is expressed in days.

**Fig. 6: Mismatch distributions analyses for SARS-CoV-2 in different period (A,**

**B, C, D, E, F, G).** (A) All the individuals collected during 24[th] December to 30[th]

December, 2019; (B) 24[th] December in 2019 to 6[th] January in 2020; (C) 24[th]

December in 2019 to 13[th] January in 2020; (D) 24[th] December in 2019 to 20[th]

January in 2020; (E) 24[th] December in 2019 to 27[th] January in 2020; (F) 24[th]

December in 2019 to 3[th] February in 2020; (G) 24[th] December in 2019 to 11[th]

February in 2020. The x coordinate represents the number of differences in each pair

of sequence comparisons; the y coordinate represents the frequencies of pairwise

differences. The blue histogram are the observed frequencies of pairwise divergences

among sequences and the red line refers to the expectation under the model of

population expansion.

**Fig. S1. Phylogenetic tree of** SARS-CoV-2**.** The nodal numbers are ML bootstrap

values.