1 **Queen pheromone modulates the expression of epigenetic modifier genes in the brain of**
2 **honeybee workers**
3

4 Carlos Antônio Mendes Cardoso-Junior[1,2*], Isobel Ronai[2], Klaus Hartfelder[1], Benjamin P.
5 Oldroyd[2]
6
7 [1] Departamento de Biologia Celular e Bioagentes Patogênicos, Faculdade de Medicina de
8 Ribeirão Preto, Universidade de São Paulo, Ribeirão Preto, Brazil.
[2] Behaviour, Ecology and Evolution (BEE) laboratory, Macleay Building A12, University of
Sydney, Sydney NSW 2006, Australia.
9
10
[*] **Corresponding author**: carloscardoso.bio@usp.br

**Contact Information**

Carlos A. M. Cardoso-Júnior – carloscardoso.bio@usp.br; Phone: +55 16 3315 3063

Isobel Ronai – isobel.ronai@sydney.edu.au

Klaus Hartfelder – klaus@fmrp.usp.br

Benjamin Oldroyd – benjamin.oldroyd@sydney.edu.au

**Running title:** Epigenetics in honeybee pheromone communication

**Abstract**

Pheromones are used by many insects to mediate social interactions. In the highly eusocial honeybee (*Apis mellifera*) queen mandibular pheromone (QMP) is involved in the regulation of reproduction and behaviour of workers. The molecular mechanisms by which QMP acts are largely unknown. Here we investigate how genes responsible for epigenetic modifications to DNA, RNA and histones respond to the presence of QMP. We show that several of these genes are upregulated in the honeybee brain when workers are exposed to QMP. This provides a plausible mechanism by which pheromone signalling may influence gene expression in the brain of honeybee workers. We propose that pheromonal communication systems, such as those used by social insects, evolved to respond to environmental signals by making use of existing epigenomic machineries.

Keywords: DNMT, HAT, HDAC, SIRT, honey bee

## 1. Introduction

The transition to social living in the eusocial insects required that the reproductive interests of individual workers be subsumed by the collective interests of the colony [1–3]. In particular, workers are functionally sterile, whereas queens are highly fecund [4]. For such systems to evolve it was also necessary that tasks can be distributed among workers in ways that enhance colony-level productivity [5–8]. Both, the regulation of worker fertility and the efficient allocation of tasks among workers required the evolution of effective inter- and intra-caste communication systems that can rapidly respond to the changing needs of the colony. Communication between nestmates most often occurs via pheromones [9,10], chemical signals that are produced by one individual, and cause changes in the behaviour or physiology of another [11,12].

In the western honeybee (*Apis mellifera*) a key pheromone is the queen mandibular pheromone (QMP), which is a blend of fatty acids secreted by the head glands of the queen. It affects several important traits of workers, including their reproduction [13], retinue response to queens [14,15], learning capacity [16], nestmate recognition [17] and age at onset of foraging [18]. Phenotypic variation in these traits is associated with differential gene expression in the brains of workers [19,20]. Nonetheless, little is known about the intermediate steps between QMP production and release by the queen, the regulation of gene expression in workers, and changes in their behaviour [21–24].

Epigenetic mechanisms are a likely mediator between a worker's social environment and global gene expression responses [25]. Several classes of epigenetic mechanism described in honeybees are potentially associated with environmental cues [26–29]. For example, DNA methylation, a reversible chemical modification of cytosines in CpG contexts, is associated with behavioural maturation in the brains of honeybee nurses and foragers [30,31]. DNA methylation is catalysed and maintained by the DNA methyltransferase (DNMT) family of enzymes [32,33]. Interestingly, in honeybees, the expression of genes associated with the maintenance of DNA methylation levels after DNA replication (*Dnmt1a* and *Dnmt1b*) are modulated by different social stimuli to *Dnmt3*, an enzyme that establishes DNA methylation patterns *de novo* [25,34–36]. In addition, the expression of *Dnmt2* (also called *Trdmt1*), a gene whose enzyme product methylates RNA substrates [37], is affected by different social contexts [25,30]. These studies suggest that epigenetic machineries associated with nucleotide modification are affected by several environmental cues.

Another epigenetic mechanism, histone post-translational modifications (HPTMs), change chromatin structure by altering the physicochemical affinity between DNA and histones and thereby affect gene expression [38]. HPTMs are catalysed by histone modifier proteins [33], which can be divided into three functional classes: writers, erasers and readers. "Writer" enzymes add chemical radicals to histone tails by covalent modification. For example, lysine acetyltransferases (KATs) promote acetylation of lysine residues [33], which reduces the affinity between DNA and nucleosomes. Histone acetylation induces chromatin relaxation and is often associated with increased gene expression [39]. In contrast, "Eraser" enzymes remove such chemical radicals from histone tails. Classical eraser enzymes are the histone deacetylases (HDACs) and Sirtuins, which remove acetyl groups from lysine residues, resulting in chromatin compaction and, consequently, inhibition of gene expression [33,40]. Finally, "Reader" enzymes recognise epigenetic modifications and induce chromatin remodelling through the recruitment of protein complexes [33]. A honey bee proteome study [27] has shown that histone tails are extensively modified by epigenetic marks, indicating that writer, eraser and reader enzymes are present in the honey bee. Furthermore, differential accumulation of HPTMs has been associated with caste differentiation and behaviour in bees and ants [27,41–44].

107     Given that QMP affects behaviours in honey bee workers [14,15,18,45] we
108 hypothesised that the expression of genes associated with epigenetic modification to
109 nucleotide and histones would respond to QMP exposure in the brain of honey bee workers.
110 These epigenetic mechanisms can, thus, serve as proxies to understand the regulation of
111 global changes in gene expression in a complex social environment.
112
113 **2. Material and Methods**
114
**(a) Biological material**
To obtain age-matched adult workers we collected brood frames from four queenright *A. m.*
*ligustica* source colonies and kept them in an incubator overnight at 34.5 °C. From each
source colony, workers were randomly allocated to two cages (n = 150 bees per cage, eight
cages in total). One cage from each colony (QMP$^+$) was furnished with a 0.5 queen
equivalent per day QMP strip (Phero Tech Inc. Canada), which is an effective queen mimic
in cage experiments with young workers [21,46]. The other cage from each colony (QMP$^-$)
contained no QMP strip. Pollen, honey and water were provided *ad libitum*. Food was
replenished when necessary, and the number of dead workers was recorded each day, which
was nearly the same in the QMP$^+$ and QMP$^-$ cages (data not shown). Cages were kept in an
incubator at 34.5 °C for four days. Workers were collected on dry ice at Day 0 (directly from
the brood comb), Day 1 and Day 4. Day 1 was chosen to identify genes with a quick response
to the QMP treatment, and Day 4 was chosen to identify the genes that are still influenced by
the QMP exposure after prolonged exposure. We then dissected the brains of the workers on
dry ice [47].
130
**(b) Identification of the honeybee DNA methyltransferases and histone modifiers**
We identified the nucleotide and histone modifier genes in the honey bee genome (Amel_4.5)
[48], searching manually for the names of each epigenetic gene in GenBank (NCBI
Resources Coordinators 2018) (Table S1) based on a large list of histone-modifier genes
present in eukaryotes [50,51]. We filtered this list by selecting those associated with
acetylation and deacetylation processes. From this list we identified the proteins that are
predicted to reside in the nucleus using ProtComp v9.0 (Softberry, Inc.). The genes and their
respective proteins were characterised following a previously described workflow [13].
139
**(c) Gene expression quantification and bioinformatics analysis**
Each sample consisted of a single brain. We extracted total RNA from the brain through
maceration in TRIzol (Invitrogen) and a Direct-zol$^{TM}$ RNA Miniprep kit (Zymo Research).
The RNA was treated with Turbo DNase (Thermo Fisher Scientific) and quantified with a
Qubit 2.0 Fluorometer (Invitrogen). cDNA was synthesised from 600 ng of RNA using a
SuperScript$^{TM}$ III Reverse Transcriptase Kit (Invitrogen) with oligo(dT) primer and
suspended in ultrapure water (5 ng cDNA/µL).
The expression of four nucleotide modifier genes and 11 histone modifier genes
(Table S1) was quantified by reverse transcription quantitative real-time PCR (RT-qPCR)
[46,52]. Assays were set up with 2.5 µL SsoAdvanced™ Universal SYBR$^®$ Green Supermix
(Bio Rad), 1.25 pmol of each primer, 1 µL diluted cDNA (5 ng) in a total volume of 5 µL
using a CFX384 Real-Time System (Bio-Rad). For each experimental sample (four source
colonies, three ages and two treatments) three technical replicates were conducted. Cycle
conditions were as follows: 95 °C for 10 min followed by 40 cycles of 95 °C for 10 s, 60 °C
for 10 s and 72 °C for 15 s. At the end of the RT-qPCR protocol a melting curve analysis was
run to confirm a single amplification peak. Primer efficiencies (Table S2) were calculated
based on an amplification curve of 10 points obtained through serial dilution of mixed cDNA

samples. The expression of the genes of interest was normalised against the expression of two reference genes (*Rpl32* and *Ef1α*), whose expression was found stable according to *BestKeeper* [53]. Relative expression levels were calculated [52], using a formula that normalises gene expression to the reference genes taking into account the efficiency of each primer set. The genes, primer sequences and efficiencies are listed in Table S2.

**(d) Statistical analysis**

To compare the expression of the QMP$^+$ and QMP$^-$ treatments at Day 1 and Day 4 we used a generalised linear mixed model (GLMM) with 'colony' as random effect and 'treatment' and 'age' as fixed effects. To model the gene expression data, we used link = identity, family = Gaussian. Where necessary, a transformation $\log_{10}$ function was applied (see Table S3 for details). We used Day-0 data as a baseline for gene expression. GLMM analyses were performed in R [54] loading the packages lme4, car and emmeans. An adjusted *p*-value (Tukey correction for each gene) lower than 0.05 was considered significant for all statistical tests.

## 3. Results

Using the protein sequences of the 15 genes studied, we first acquired *in-silico* evidence (e.g. subcellular location, predicted domains and homology with other species) that each gene was a *bona fide* epigenetic modifier of DNA, RNA or histones (Table S1). In 1-day old workers the expression of twelve genes associated with epigenetic processes (*Dnmt1b*, *Dnmt2*, *Dnmt3*, *Kat2a*, *Kat3b*, *Kat6b*, *Kat8*, *Hdac1*, *Hdac3*, *Sirt 1, Sirt7* and *Rcs1*) was affected by exposure to QMP (GLMM, $p < 0.05$, Figures 1 and 2, Table S3). However, only four genes (*Dnmt1b*, *Dnmt2*, *Kat3b* and *Sirt7*) continued to be differentially expressed at the age of four days (GLMM, $p < 0.05$, Figures 1 and 2, Table S3). Age was statistically significant for 13 of the 15 genes (GLMM, $p < 0.05$, Figure 1 and Table S3), the exceptions being *Kat 7* and *Dnmt3*. A significant interaction between treatment and age was found for three genes: *Hdac1*, *Sirt1* and *Kat6b* ($p < 0.05$, Table S3).

## 4. Discussion

Our study shows that QMP affects the expression of 12 of 15 genes that are associated with epigenetic processes in the brain of honeybee workers. As predicted, our data indicates that epigenetic mechanisms are likely mediators between queen pheromone signalling and the regulation of worker gene expression. Given that QMP alters worker behaviour [14,15,18,45], we suggest that pheromonal communication evolved by making use of existing epigenetic mechanisms that orchestrate transcriptomic changes necessary to propagate pheromonal information.

Some expression responses are particularly worthy of note. For instance, the expression of *Dnmt3*, the *de novo* methylator of DNA, is regulated by queen pheromones in brains of honeybee workers (this study) and whole-body RNA extracts of honeybee workers [35] and two ant species (*Lasius flavus* and *Lasius niger*) [55]. Expression of *Kat 8* is upregulated in the brains of QMP-treated honeybee workers. This gene is differentially spliced in *L. flavus* ants treated with queen pheromone [55]. Together, these results suggest an evolutionary conservation in the epigenetic pathways responsive to queen pheromones in social insects.

We detected that several histone modifiers genes associated with acetylation/deacetylation processes are differentially expressed in the brains of adult workers. This finding suggests that queen signals influence the modification of histones to promote chromatin reorganisation and thereby altering gene expression in worker brains. In line with this hypothesis, histone acetylation contributes to the regulation of foraging behaviour in ants

[43]. Interestingly, it was recently shown that honeybee queens regulate worker fertility
through polycomb repressive complex 2 (PRC2) activity and differential histone methylation
marks [56]. We propose that queens, via QMP, influence modifications to histones to
regulate behavioural plasticity in the brains of honeybee workers, just as they do in ovaries.

Pheromonal modulation of gene expression in honeybee workers changes over time
[19,20]. Gene expression in QMP$^-$ workers is relatively stable from Day 0 to Day 1 when
compared to QMP$^+$ workers, suggesting QMP *actively* promotes expression of several
epigenetic modifier genes already within 24 hours. Only four of these continued to be
differentially expressed after 4 days of QMP exposure, indicating that the expression of the
majority epigenetic modifiers is dynamically switched on and off [19].

Our study provides evidence that many genes associated with epigenetic modification
are differentially expressed in the brains of honeybee workers in response to the presence of
queen pheromone. These changes wrought by the genes studied here likely drive changes in
gene expression in the brains of adult workers, providing a plausible mechanism by which a
queen can influence both the rate of behavioural maturation and reproductive behaviour of
her workers. This property of QMP would explain why it acts both as a short-term 'releaser'
pheromone that merely indicates queen presence, as well as a long-term 'primer' pheromone
that regulates behavioural maturation and reproductive behaviour.

225  
226  **Data accessibility**
227  The data that support this study are available in the supplementary material.

228  
229  **Authors' contributions**
230  CAM designed the study, carried out the cage experiments, dissections, performed molecular
231  lab work, analysed data and wrote the manuscript draft. IR carried out the protein
232  characterisation and wrote the manuscript. BPO and KH supervised the work and revised the
233  manuscript. All authors gave final approval for publication.

234  
**Competing interests**
The authors declare no conflict of interests.

**Funding**
This research was funded by the Australian Research Council DP180101696 and Fundação
de Amparo à Pesquisa do Estado de São Paulo (FAPESP - #2016/15881-0 and 2017/09269-3
to CAM). The funders had no role in study design, data collection and analysis, decision to
publish, or preparation of the manuscript.

243  
244  
245  
**Figure 1.** Relative expression of four nucleotide modifier (DNA methyltransferase) genes in
the brains of 0-, 1- and 4-days old honeybee workers, exposed to queen mandibular
pheromone (QMP$^+$) or not (QMP$^-$). Each box shows the interquartile range ($25^{th}$-$75^{th}$
percentiles) and the median (line), while whiskers represent the farthest points of $2.5^{th}$-$97.5^{th}$
percentiles. Relative expression was calculated for each gene at all three ages. Day 0 was
used as the baseline for gene expression. Statistical information: GLMM test with Tukey
correction for multiple pairwise comparisons, * $p < 0.05$, ** $p < 0.01$, *** $p < 0.001$, N=32;
N=8 from each of the four colonies).

254  
255  **Figure 2.** Relative expression of 11 histone modifier genes in the brains of 0-, 1- and 4-days
256  old honey bee workers, exposed to queen mandibular pheromone (QMP$^+$) or not (QMP$^-$). (a)

Relative expression of histone acetyltransferases genes (writer enzymes). (b) Relative expression of histone deacetylases and Sirtuin genes (eraser enzymes). (c) Relative expression of the *Rsc1* gene (reader enzyme). Each box shows the interquartile range (25th-75th percentiles) and the median (line), while whiskers represent the farthest points of 2.5th-97.5th percentiles. Relative expression was calculated for each gene at all three ages. Day 0 was used as the baseline for gene expression. Statistical information is as in figure 1.

**References**

1.  Ratnieks FLW, Foster KR, Wenseleers T. 2006 Conflict resolution in insect societies. *Annu. Rev. Entomol.* **51**, 581–608. (doi:10.1146/annurev.ento.51.110104.151003)

2.  Ratnieks FLW. 1988 Reproductive harmony via mutual policing by workers in eusocial Hymenoptera. *Am. Nat.* **132**, 217–236. (doi:10.1086/284846)

3.  Maynard-Smith J, Szathmáry E. 1995 *The major transitions in evolution.* Oxford: Oxford University Press.

4.  Ronai I, Vergoz V, Oldroyd BP. 2016 The mechanistic, genetic, and evolutionary basis of worker sterility in the social Hymenoptera. *Adv. Study Behav.* **48**, 251–317. (doi:10.1016/bs.asb.2016.03.002)

5.  Gordon DM. 1996 The organization of work in social insect colonies. *Nature* **380**, 121–124. (doi:10.1038/380121a0)

6.  Page RE. 1997 The evolution of insect societies. *Endeavour* **21**, 114–120. (doi:10.1016/S0160-9327(97)80220-7)

7.  Bonabeau E, Theraulaz G, Deneubourg JL, Aron S, Camazine S. 1997 Self-organization in social insects. *Trends Ecol. Evol.* **12**, 188–193. (doi:10.1016/S0169-5347(97)01048-3)

8.  Visscher PK. 2003 How self-organization evolves. *Nature* **421**, 799–800. (doi:10.1038/421799a)

9.  Free JB. 1987 *Pheromones of social bees*. London: Chapman and Hall.

10. Meer RKV, Preston CA. 2008 Social insect pheromones. In *Encyclopedia of Entomology*, pp. 223–262. Springer, Dordrecht. (doi:https://doi.org/10.1007/978-1-4020-6359-6)

11. Holman L. 2010 Queen pheromones. *Commun. Integr. Biol.* **3**, 558–560. (doi:10.4161/cib.3.6.12976)

12. Billen J, Morgan ED. 1998 *Pheromone communication in social insects: sources and secretions*. Westview Press.

13. Ronai I, Oldroyd BP, Vergoz V. 2016 Queen pheromone regulates programmed cell death in the honey bee worker ovary. *Insect Mol. Biol.* **25**, 646–652. (doi:10.1111/imb.12250)

14. Slessor KN, Kaminski LA, King GGS, Borden JH, Winston ML. 1988 Semiochemical basis of the retinue response to queen honey bees. *Nature* **332**, 354–356. (doi:10.1038/332354a0)

15. Keeling CI, Slessor KN, Higo HA, Winston ML. 2003 New components of the honey bee (*Apis mellifera* L.) queen retinue pheromone. *Proc. Natl. Acad. Sci. U. S. A.* **100**, 4486–4491. (doi:10.1073/pnas.0836984100)

16. Vergoz V, Schreurs HA, Mercer AR. 2007 Queen pheromone blocks aversive learning in young worker bees. *Science.* **317**, 384–386. (doi:10.1126/science.1142448)

17. Fan Y, Richard FJ, Rouf N, Grozinger CM. 2010 Effects of queen mandibular pheromone on nestmate recognition in worker honeybees, *Apis mellifera. Anim. Behav.* **79**, 649–656. (doi:10.1016/j.anbehav.2009.12.013)

18.  Pankiw T, Winston ML, Huang Z, Robinson GE. 1998 Queen mandibular gland
pheromone influences worker honey bee (*Apis mellifera* L.) foraging ontogeny and
juvenile hormone titers. *J. Insect Physiol.* **44**, 685–692. (doi:10.1016/S0022-
1910(98)00040-7)
19.  Grozinger CM, Sharabash NM, Whitfield CW, Robinson GE. 2003 Pheromone-
mediated gene expression in the honey bee brain. *Proc. Natl. Acad. Sci. U. S. A.* **100**
**Suppl**, 14519–14525. (doi:10.1073/pnas.2335884100)
20.  Fussnecker BL, McKenzie AM, Grozinger CM. 2011 cGMP modulates responses to
queen mandibular pheromone in worker honey bees. *J. Comp. Physiol. A* **197**, 939–
948. (doi:10.1007/s00359-011-0654-5)
21.  Ronai I, Barton DA, Oldroyd BP, Vergoz V. 2015 Regulation of oogenesis in honey
bee workers via programed cell death. *J. Insect Physiol.* **81**, 36–41.
(doi:10.1016/j.jinsphys.2015.06.014)
22.  Ronai I, Oldroyd BP, Barton DA, Cabanes G, Lim J, Vergoz V. 2016 Anarchy is a
molecular signature of worker sterility in the honey bee. *Mol. Biol. Evol.* **33**, 134–142.
(doi:10.1093/molbev/msv202)
23.  Huang ZY, Plettner E, Robinson GE. 1998 Effects of social environment and worker
mandibular glands on endocrine-mediated behavioral development in honey bees. *J.*
*Comp. Physiol. - A Sensory, Neural, Behav. Physiol.* **183**, 143–152.
(doi:10.1007/s003590050242)
24.  Brockmann A, Brückner D, Crewe RM. 1998 The EAG response spectra of workers
and drones to queen honeybee mandibular gland components: The evolution of a
social signal. *Naturwissenschaften* **85**, 283–285. (doi:10.1007/s001140050500)
25.  Cardoso-Junior CAM, Eyer M, Dainat B, Hartfelder K, Dietemann V. 2018 Social
context influences the expression of DNA methyltransferase genes in the honeybee.
*Sci. Rep.* **8**, 11076. (doi:10.1038/s41598-018-29377-8)
26.  Wang Y, Jorda M, Jones PL, Maleszka R, Ling X, Robertson HM, Mizzen CA,
Peinado MA, Robinson GE. 2006 Functional CpG methylation system in a social
insect. *Science* **314**, 645–647. (doi:10.1126/science.1135213)
27.  Dickman MJ, Kucharski R, Maleszka R, Hurd PJ. 2013 Extensive histone post-
translational modification in honey bees. *Insect Biochem. Mol. Biol.* **43**, 125–137.
(doi:10.1016/j.ibmb.2012.11.003)
28.  Yan H, Simola DF, Bonasio R, Liebig J, Berger SL, Reinberg D. 2014 Eusocial insects
as emerging models for behavioural epigenetics. *Nat. Rev. Genet.* **15**, 677–688.
(doi:10.1038/nrg3787)
29.  Yan H, Bonasio R, Simola DF, Liebig J, Berger SL, Reinberg D. 2015 DNA
Methylation in social insects: how epigenetics can control behavior and longevity.
*Annu. Rev. Entomol.* **60**, 435–452. (doi:10.1146/annurev-ento-010814-020803)
30.  Herb BR, Wolschin F, Hansen KD, Aryee MJ, Langmead B, Irizarry R, Amdam G V,
Feinberg AP. 2012 Reversible switching between epigenetic states in honeybee
behavioral subcastes. *Nat. Neurosci.* **15**, 1371–1373. (doi:10.1038/nn.3218)
31.  Lockett GA, Kucharski R, Maleszka R. 2012 DNA methylation changes elicited by
social stimuli in the brains of worker honey bees. *Genes, Brain Behav.* **11**, 235–242.
(doi:10.1111/j.1601-183X.2011.00751.x)
32.  Bestor TH. 2000 The DNA methyltransferases of mammals. *Hum Mol Genet* **9**, 2395–
2402.
33.  Allis CD, Jenuwein T. 2016 The molecular hallmarks of epigenetic control. *Nat. Rev.*
*Genet.* **17**, 487–500. (doi:10.1038/nrg.2016.59)
34.  Cardoso-Júnior CAM, Guidugli-Lazzarini KR, Hartfelder K. 2018 DNA methylation
affects the lifespan of honey bee (*Apis mellifera* L.) workers – Evidence for a

regulatory module that involves vitellogenin expression but is independent of juvenile
hormone function. *Insect Biochem. Mol. Biol.* **92**, 21–29.
(doi:10.1016/j.ibmb.2017.11.005)

35.  Holman L, Trontti K, Helantera H. 2016 Queen pheromones modulate DNA
methyltransferase activity in bee and ant workers. *Biol Lett* **12**, 20151038.
(doi:10.1098/rsbl.2015.1038)

36.  Kucharski R, Maleszka J, Foret S, Maleszka R. 2008 Nutritional control of
reproductive status in honeybees via DNA methylation. *Science* **319**, 1827–1830.
(doi:10.1126/science.1153069)

37.  Jeltsch A, Ehrenhofer-Murray A, Jurkowski TP, Lyko F, Reuter G, Ankri S, Nellen W,
Schaefer M, Helm M. 2016 Mechanism and biological role of *Dnmt2* in nucleic acid
methylation. *RNA Biol* **14**, 1108–1123.

38.  Allis DC, Caparros M-L, Jenuwein T, Reinberg D. 2007 Overview and concepts. In
*Epigenetics*, pp. 47–116. Cold Spring Harbor Laboratory Press.

39.  Albaugh BN, Arnold KM, Denu JM. 2011 KAT(ching) metabolism by the tail: Insight
into the links between lysine acetyltransferases and metabolism. *ChemBioChem* **12**,
290 – 298. (doi:10.1002/cbic.201000438)

40.  Pasyukova EG, Vaiserman AM. 2017 HDAC inhibitors: A new promising drug class
in anti-aging research. *Mech. Ageing Dev.* **166**, 6–15. (doi:10.1016/j.mad.2017.08.008)

41.  Simola DF, Ye C, Mutti NS, Dolezal K, Bonasio R, Liebig J, Reinberg D, Berger SL.
2013 A chromatin link to caste identity in the carpenter ant *Camponotus floridanus*.
*Genome Res.* **23**, 486–496. (doi:10.1101/gr.148361.112)

42.  Cardoso-Júnior CAM, Fujimura PT, Santos-Júnior CD, Borges NA, Ueira-Vieira C,
Hartfelder K, Goulart LR, Bonetti AM. 2017 Epigenetic modifications and their
relation to caste and sex determination and adult division of labor in the stingless bee
*Melipona scutellaris*. *Genet. Mol. Biol.* **40**, 61–68. (doi:10.1590/1678-4685-GMB-
2016-0242)

43.  Simola DF *et al.* 2016 Epigenetic (re)programming of caste-specific behavior in the
ant *Camponotus floridanus*. *Science* **351**, aac6633. (doi:10.1126/science.aac6633)

44.  Wojciechowski M, Lowe R, Maleszka J, Conn D, Maleszka R, Hurd PJ. 2018
Phenotypically distinct female castes in honey bees are defined by alternative
chromatin states during larval development. *Genome Res.* **28**, 1532–1542.
(doi:10.1101/gr.236497.118)

45.  Beggs KT, Glendining KA, Marechal NM, Vergoz V, Nakamura I, Slessor KN,
Mercer AR. 2007 Queen pheromone modulates brain dopamine function in worker
honey bees. *Proc. Natl. Acad. Sci. U. S. A.* **104**, 2460–2464.
(doi:10.1073/pnas.0608224104)

46.  Cardoso-Júnior CAM, Oldroyd B, Ronai I. 2019 Vitellogenin expression in the ovaries
of adult honeybee workers provides insights into the evolution of reproductive and
social traits. *BioRxiv Prepr.* (doi:10.1101/547760)

47.  Vergoz V, Lim J, Duncan M, Cabanes G, Oldroyd BP. 2012 Effects of natural mating
and CO2 narcosis on biogenic amine receptor gene expression in the ovaries and brain
of queen honey bees, *Apis mellifera*. *Insect Mol. Biol.* **21**, 558–567.
(doi:10.1111/j.1365-2583.2012.01159.x)

48.  Elsik CG *et al.* 2014 Finding the missing honey bee genes: Lessons learned from a
genome upgrade. *BMC Genomics* **15**. (doi:10.1186/1471-2164-15-86)

49.  Coordinators R. 2018 Database resources of the National Center for Biotechnology
Information. *Nucleic Acids Res.* **46**, D8–D13. (doi:10.1093/nar/gkn741)

50.  Jęśko H, Wencel P, Strosznajder RP, Strosznajder JB. 2017 Sirtuins and their roles in
brain aging and neurodegenerative disorders. *Neurochem. Res.* **42**, 876–890.

(doi:10.1007/s11064-016-2110-y)

51.   Yun M, Wu J, Workman JL, Li B. 2011 Readers of histone modifications. *Cell Res* **21**,
564–578. (doi:10.1038/cr.2011.42)

52.   Brito RM, McHale M, Oldroyd BP. 2010 Expression of genes related to reproduction
and pollen foraging in honey bees (*Apis mellifera*) narcotized with carbon dioxide.
*Insect Mol. Biol.* **19**, 451–461. (doi:10.1111/j.1365-2583.2010.01003.x)

53.   Pfaffl MW, Tichopad A, Prgomet C, Neuvians TP. 2004 Determination of stable
housekeeping genes, differentially regulated target genes and sample integrity:
BestKeeper - Excel-based tool using pair-wise correlations. *Biotechnol. Lett.* **26**, 509–
515. (doi:10.1023/B:BILE.0000019559.84305.47)

54.   Team RC. 2018 R: A language and environment for statistical computing. R
foundation for statistical computing, Vienna, Austria. URL: http://www.R-project.org/.

55.   Holman L, Helanterä H, Trontti K, Mikheyev AS. 2019 Comparative transcriptomics
of social insect queen pheromones. *Nat. Commun.* **10**, 1593. (doi:10.1038/s41467-019-
09567-2)

56.   Duncan EJ, Leask MP, Dearden PK. 2020 Genome architecture facilitates phenotypic
plasticity in the honeybee (*Apis mellifera*). *Mol. Biol. Evol.* **[Epub]**.
(doi:10.1093/molbev/msaa057)