TITLE

Lysine63-linked ubiquitin chains earmark GPCRs for BBSome-

mediated removal from cilia

AUTHORS: Swapnil Rohidas Shinde, Andrew R. Nager[#] and Maxence V. Nachury[*]

AFFILIATION:

Department of Ophthalmology, University of California San Francisco, CA 94143, USA

* Correspondence: maxence.nachury@ucsf.edu

Current address: Cancer Immunology Discovery, Pfizer Inc., San Diego, CA, 92121, USA

ABSTRACT (160 words)

G-protein coupled receptors (GPCRs) undergo regulated trafficking into and out of cilia to

control cilium-based signaling pathways. β-arrestin 2, a molecular sensor of activated GPCRs,

and the BBSome are required for the signal-dependent exit of ciliary GPCRs but it is not known

how β-arrestin 2 relays the activation state of GPCRs to the ciliary exit machinery. Here we find

that, upon activation, the ciliary GPCRs SSTR3 and GPR161 become tagged with K63-linked

ubiquitin (K63Ub) chains in a β-arrestin 2-dependent manner prior to BBSome-mediated exit.

Removal of ubiquitin acceptor residues from SSTR3 and GPR161 demonstrates that

ubiquitination of ciliary GPCRs is required for their regulated exit from cilia. Furthermore,

targeting a K63Ub-specific deubiquitinase to cilia blocks the exit of GPR161, SSTR3 and SMO

from cilia. Finally, ubiquitinated proteins accumulate in cilia of mammalian photoreceptors and

Chlamydomonas cells. We conclude that K63Ub chains mark GPCRs and other unwanted ciliary

proteins for recognition by the ciliary exit machinery.

INTRODUCTION

The regulated trafficking of signaling receptors in and out of cilia is a near-universal feature of cilia-based signaling pathways that is germane to the transduction of signals (Nachury and Mick, 2019; Anvarian et al., 2019; Mykytyn and Askwith, 2017). For example, in Hedgehog signaling, pathway activation leads to the disappearance of the Hedgehog receptor Patched 1 and of the G protein-coupled receptor (GPCR) GPR161 from cilia and to the concurrent accumulation of the GPCR Smoothened (SMO) in cilia (Rohatgi et al., 2007; Corbit et al., 2005; Mukhopadhyay et al., 2013). In all three cases, regulated exit from cilia plays a major role in the on-demand redistribution of signaling molecules (Nachury and Mick, 2019). Signal-dependent exit has been generalized to every ciliary GPCRs studied to date. The Somatostatin Receptor 3 (SSTR3), Dopamine Receptor 1 (D1R), Melanocortin concentrating hormone receptor 1 (MCHR1), the neuropeptide receptor 2 (NPY2R) all disappear from cilia upon exposure to agonist (Domire et al., 2011; Loktev and Jackson, 2013; Nager et al., 2017; Green et al., 2015). A major question is how GPCRs are selected for removal from cilia only under specific conditions (typically upon activation).

Two major trafficking modules participate directly in signal-dependent exit from cilia. First, the BBSome is an evolutionarily conserved complex of eight Bardet-Biedl Syndrome (BBS) proteins that directly recognizes intracellular determinants in GPCRs and ferries them out of cilia (Nachury, 2018; Wingfield et al., 2018). The BBSome associates with the intraflagellar transport (IFT) machinery and has been proposed to act as adaptor between the motor-driven intraflagellar transport trains and the membrane proteins that are to be removed from cilia. Second, β-arrestin 2 is a well-established molecular sensor of the activation state of GPCRs that is required for the signal-dependent exit of GPR161 and SSTR3 (Pal et al., 2016; Green et al., 2015; Nager et al., 2017). No association of β-arrestin 2 with ciliary trafficking complexes has been reported to date and it remains unclear how β-arrestin 2 relays information regarding the state of activation of ciliary GPCRs to the ciliary exit machinery.

An emerging player in ciliary exit is ubiquitination (Shearer and Saunders, 2016). Ubiquitin (Ub), a 76-amino acid polypeptide, becomes conjugated to acceptor lysine residues on substrate

proteins by ubiquitin ligases and routes substrates for degradation or other regulatory fates (Yau

and Rape, 2016; Swatek and Komander, 2016). A role for ubiquitin in promoting ciliary exit is

suggested by multiple lines of evidence. First, interfering with Patched1 ubiquitination blocks its

signal-dependent exit from cilia (Kim et al., 2015; Yue et al., 2014). Second, fusing ubiquitin to

the cytoplasmic tails of PKD-2, of the olfactory receptor ODR-10 or of the TRP channel OSM-9

results in the disappearance of these proteins from cilia (Hu et al., 2007; Xu et al., 2015). The

accumulation of these ubiquitin-protein fusions inside cilia when BBSome function is

compromised suggests that the BBSome might sort ubiquitinated signaling receptors out of cilia,

in line with a reported association of BBSome with ubiquitinated proteins in trypanosomes

(Langousis et al., 2016). Interestingly, TRIM32 is a ubiquitin ligase that is mutated in BBS

patients in one consanguineous family and interference with TRIM32 function in zebrafish

produces typical BBS-related phenotypes (Chiang et al., 2006). Third, the ubiquitin ligase Cbl is

recruited to cilia upon activation of the ciliary tyrosine kinase receptor PDGFRαα and Cbl is

required for termination of PDGFRαα signaling (Schmid et al., 2018). Last, a dramatic rise in

ubiquitination of the ciliary proteome is observed when *Chlamydomonas* cilia disassemble (Huang

et al., 2009). In particular, α-tubulin becomes ubiquitinated on K304 upon cilia disassembly and

expression of α-tubulin[K304R] slows down cilia disassembly (Wang et al., 2019).

Together, these data led us to investigate the interplay between ubiquitination, β-arrestin 2 and

the ciliary exit machinery. Given the well-established function of β-arrestin 2 in directing

activation-dependent ubiquitination of GPCRs on the surface of endosomes (Henry et al., 2012;

Bhandari et al., 2007; Shenoy et al., 2008), we posited that β-arrestin 2 may direct the activation-

dependent ubiquitination of ciliary GPCRs and that ubiquitinated GPCRs are selected by the

BBSome for removal from cilia.

RESULTS

Signal-dependent ubiquitination of ciliary GPCRs is required for their regulated exit from cilia

A central prediction of the above model is that the activation of ciliary GPCRs will lead to their ubiquitination inside cilia prior to BBSome-mediated retrieval. We blocked BBSome-dependent exit by deleting its cognate GPTase ARL6/BBS3 and monitored ciliary ubiquitin levels by immunostaining. When SSTR3 was expressed in $Arl6^{-/-}$ IMCD3 cells, addition of its agonist somatostatin (sst) led to a drastic increase in the ciliary levels of ubiquitin as reported by the well-characterized FK2 (**Figure 1A-B)** and FK1 (**Figure S1A-B**) monoclonal antibodies (Fujimuro et al., 1994; Emmerich and Cohen, 2015; Haglund et al., 2003). As the two antibodies have identical selectivity for ubiquitin species, all further experiments were conducted with the FK2 antibody. Because the sst-dependent increase in ciliary ubiquitin signal was no longer observed in the absence of SSTR3 expression (**Figure 1A-B and S1C-D**), these results indicate that SSTR3 itself becomes ubiquitinated inside cilia upon activation. We noted that ciliary ubiquitin was already detectable in the absence of sst in $Arl6^{-/-}$ cells. This ciliary ubiquitin signal did not depend upon expression of SSTR3 (compare 3rd and 5th violin plots in **Figure 1B**) and suggests that a GPCR endogenous to IMCD3 cells may become activated and ubiquitinated inside $Arl6^{-/-}$ cilia under our culture conditions. No ubiquitin signal was detected in cilia when BBSome function was intact (**Figure 1A-B and S1A-D**), indicating that the BBSome efficiently removes ubiquitinated GPCRs from cilia.

One signaling pathway that is natively expressed in IMCD3 cells is the Hedghog pathway. Prior experiments have detected normal trafficking dynamics of SMO and GPR161 in IMCD3 cells (Ye et al., 2018; Mukhopadhyay et al., 2013) and **Figure 4B**). GPR161 and SMO both accumulate in cilia of $Arl6^{-/-}$ cells (Liew et al., 2014; Zhang et al., 2011) and it is conceivable that the ubiquitin signal detected in cilia corresponds to ubiquitin conjugated to SMO or GPR161. Activation of SMO by small molecule agonists normally triggers exit of GPR161 but this exit is frustrated when BBSome function goes awry. We detected a significant elevation of the ciliary

ubiquitin levels in *Arl6^-/-* cells when treated with the SMO agonist SAG (**Figure 1C-D**),

suggesting that GPR161 may get ubiquitinated prior to its BBSome-dependent exit. Collectively,

these data suggest that GPCRs are ubiquitinated in a regulated fashion inside cilia and

subsequently retrieved into the cell by the BBSome.

To test the functional importance of GPCR ubiquitination in BBSome-mediated exit from cilia,

we removed all ubiquitination sites from SSTR3 and GPR161 by mutating all cytoplasm-

exposed lysine residues to arginine (cK0 variants). In both cases, the cK0 variant underwent

markedly slower signal-dependent exit from cilia than the WT allele (**Figure 1E-F**).

Measurement of exit rates revealed that exit of SSTR3cK0 was ~2-fold slower than SSTR3 and

that exit of GPR161cK0 was ~2.5-fold slower than GPR161. These results strongly support a

role for GPCR ubiquitination in promoting signal-dependent exit from cilia

K63-linked ubiquitin chains inside cilia are required for GPCR exit

We next sought to characterize the type of ubiquitin conjugates that are attached to ciliary

proteins. A single ubiquitin can be conjugated to substrates (monoubiquitination) or lysine

residues on ubiquitin can be conjugated to another ubiquitin molecule for building ubiquitin

chains (polyubiquitination). Lys48 and Lys 63 are the most prevalent ubiquitin chain linkages

with K48 serving as a targeting signal for proteasomal degradation of soluble proteins and K63

serving as a lysosomal sorting signal for membrane proteins or a signal for regulated interactions

(Yau and Rape, 2016; Swatek and Komander, 2016). The FK1 and FK2 antibodies used above

recognizes ubiquitin either as single adduct or in chain but does not recognize free ubiquitin

(Fujimuro et al., 1994; Emmerich and Cohen, 2015). The availability of antibodies specific for

K48 and K63 ubiquitin linkages (Newton et al., 2008) enabled us to determine the types of

ubiquitin conjugates that are attached to ciliary GPCRs. While we found no detectable signal for

K48-linked ubiquitin chains (UbK48) in cilia, the for K63-linked ubiquitin (UbK63) chains

mirrored the ciliary response observed with the FK2 antibody in IMCD3-[SSTR3] cells

subjected to agonist treatment (**Figure 2 and S1E-F**). We conclude that K63Ub chains are

added onto SSTR3 inside cilia upon activation.

To determine the functional importance of UbK63 linkages in tagging GPCRs for exit from cilia,

we devised a strategy that cleaves UbK63 linkages inside cilia (**Figure 3A**). Structural and

biochemical studies have demonstrated that the deubiquitinase AMSH possesses a near-absolute

specificity for UbK63 linkages and does not cleave other Ub chain linkages or Ub-substrates

linkages (Sato et al., 2008; McCullough et al., 2004). We targeted AMSH to cilia using the well-

validated ciliary targeting signal from the ciliopathy protein NPHP3 (Nakata et al., 2012; Wright

et al., 2011, 3) and analyzed the exit of three distinct GPCRs: SMO, GPR161 and SSTR3. To

alleviate concerns related to the interaction of AMSH with the ESCRT-0 protein STAM

(McCullough et al., 2006), we deleted the STAM-interacting domain of AMSH and only

expressed the catalytic domain of AMSH. The signal-dependent exit of SSTR3 was assessed by

treating IMCD3-[SSTR3] cells with sst for 2h. In cells expressing cilia-AMSH (a fusion of

NPHP3[1-200] to GFP and the catalytic domain of AMSH), sst failed to bring a significant

reduction in the ciliary levels of SSTR3 (**Figure 3B-C**). Care was taken to only analyze cells that

expressed modest levels of cilia-AMSH as judged by GFP fluorescence. In contrast, SSTR3 exit

proceeded normally in cells that expressed NPHP3[1-200]-GFP alone or a catalytically dead

variant of cilia-AMSH that we named cilia-ASMH[†] (**Figure 3B-C**). These controls demonstrate

that cilia-AMSH exerts its effects on SSTR3 exit by cleaving UbK63 chains inside cilia.

Similarly, the exit of endogenous GPR161 from RPE cell cilia upon activation of the Hedgehog

pathway was abolished in cells expressing moderate levels of cilia-AMSH (**Figure 3D-E**).

Neither NPHP3[1-200]-GFP nor cilia-AMSH[†] blocked the signal-dependent exit of GPR161

from cilia (**Figure 3D-E**).

Unlike GPR161 and other ciliary GPCRs such as the Somatostatin receptor 3 (SSTR3) that are

removed from cilia by the BBSome only when they become activated, SMO undergoes

constitutive BBSome-dependent exit from cilia in the absence of Hedgehog pathway stimulation.

Signal-dependent accumulation of SMO in cilia is achieved, at least in part, by suppression of its

exit (Milenkovic et al., 2015; Nachury and Mick, 2019; Ye et al., 2018). In cells expressing cilia-

AMSH, endogenous SMO spontaneously accumulated inside cilia in the absence of Hedgehog

pathway activation (**Figure 3F-G**). Again, neither NPHP3[1-200]-GFP nor cilia-AMSH[†]

influenced the ciliary accumulation of SMO (**Figure 3F-G**).

Together, these results indicate that K63-linked ubiquitin chains inside cilia, presumably built

onto each ciliary GPCRs under a specific signaling stimulus, are required for GPCR removal

from cilia.

β-arrestin 2 mediates signal-dependent ubiquitination of ciliary GPCRs

As β-arrestin 2 is required for signal-dependent exit of GPR161 and SSTR3 from cilia (Green et

al., 2015; Pal et al., 2016; Ye et al., 2018), we sought to determine the functional relationship

between β-arrestin 2 and the BBSome. As previously described (Green et al., 2015), β-arrestin 2

is initially undetectable inside cilia and becomes recruited to cilia within minutes of SSTR3

agonist addition (**Figure 4A-C**). The ciliary β-arrestin 2 signal reaches its maximum in 10 min

and the $t_{1/2}$ for ciliary accumulation is less than 5 min (**Figure 4C**). Meanwhile the $t_{1/2}$ for

BBSome recruitment to the tip of cilia was 10 min and the first signs of detectable SSTR3 exit

were seen between 10 and 15 min (**Figure 4C**). These kinetics suggested the following sequence

of events: β-arrestin 2 first becomes recruited to activated ciliary SSTR3, then the BBSome

ferries activated SSTR3 out of cilia. When ARL6 was depleted, the basal levels of ciliary β-

arrestin 2 were elevated but the kinetics of β-arrestin 2 accumulation in cilia upon SSTR3

activation were indistinguishable from control-depleted cells (**Figure 4A-B**). These data suggest

that a fraction of SSTR3 that accumulates in cilia of ARL6-depleted cells is tonically active and

recruit β-arrestin 2. From these data, we conclude that β-arrestin 2 functions either upstream of,

or in parallel with the BBSome in signal-dependent retrieval of GPCRs. Similarly, the kinetics of

GPR161 exit and β-arrestin 2 ciliary recruitment upon activation of the Hh pathway indicated

that β-arrestin 2 reaches its maximal ciliary level before the first signs of GPR161 exit become

evident (**Figure 4D**).

We previously proposed that β-arrestin 2 may directly stabilize the interaction of the BBSome

with its GPCR cargoes (Ye et al., 2018). Yet, we have thus far failed to detect a biochemical

interaction between β-arrestin 2 and the BBSome. Furthermore, a direct binding to the GPCR-

BBSome complexes predicts that β-arrestin 2 co-localizes with IFT trains inside cilia. Against this

hypothesis, β-arrestin 2 was evenly distributed inside cilia (**Figure 4A**), reminiscent of the

distribution pattern of SSTR3 and distinct from the discontinuous pattern of the BBSome inside

cilia.

These data lead us to pursue the hypothesis that β-arrestin 2 licenses activated GPCRs for

recognition by the BBSome by promoting activation-dependent ubiquitination of GPCRs. The

rationale for this hypothesis is two-fold. First, β-arrestin has been shown to function in the signal-

dependent targeting of GPCRs from the plasma membrane to the lysosome by directing the

recruitment of ubiquitin ligases to activated GPCRs (Henry et al., 2012; Bhandari et al., 2007;

Shenoy et al., 2008). Ubiquitinated GPCRs are sorted to the degradative route via ESCRT

complexes and associated ubiquitin recognition modules. Second, ubiquitination promotes ciliary

exit of the signaling receptor and BBSome cargoes Patched 1 (Kim et al., 2015; Yue et al., 2014),

GPR161, SSTR3 and SMO (**Figures 1E-F and 3B-G**).

We thus considered the model that β-arrestin 2 recruits ubiquitin ligases to activated GPCR and

ubiquitinated GPCRs are then selected by the BBSome for removal from cilia. A central

prediction of this model is that the activation of ciliary GPCRs will lead to their β-arrestin 2-

dependent ubiquitination inside cilia prior to BBSome-mediated retrieval. To test this prediction,

we leveraged IMCD3 cell lines deleted for *Arl6* alone or for both *Arl6* and *β-arrestin 2* (Nager et

al., 2017). When IMCD3 *Arl6⁻ᐟ⁻* cells stably expressing SSTR3 were treated with sst, SSTR3 exit

from cilia failed and ubiquitin staining in cilia concomitantly rose (**Figures 1A-B and 4E-F**),

indicating that SSTR3 undergoes signal-dependent ubiquitination. In cells deleted for both *Arl6*

and *β-arrestin 2*, SSTR3 exit still failed but the rise in ciliary ubiquitin signal upon sst addition was

no longer observed (**Figures 4E-F**). Similarly, the increase in ciliary ubiquitin signal seen in *Arl6⁻*

ᐟ⁻ cells treated with the Hedgehog pathway agonist SAG was no longer observed in the absence of

β-arrestin 2 (**Figure 4G-H**). These data indicate that, in the absence of BBSome function, ciliary

GPCRs become ubiquitinated in response to activation in a *ß-arrestin 2*-dependent manner.

Taken together with the requirements for β-arrestin 2 and for ubiquitination in SSTR3 and

GPR161 exit, these data suggest the following order of action: GPCR activation, β-arrestin 2

recruitment, assembly of K63Ub chains on the GPCR, and selection of ubiquitinated GPCRs for

BBSome-mediated retrieval (**Figure 5E**).

Constitutively retrieved BBSome cargoes are ubiquitinated prior to exit

Besides removing GPCRs from cilia in a regulated fashion, the BBSome also clears cilia of

unwanted proteins that accidentally enter cilia. Phospholipase D, an AMP kinase family member

and a truncated hemoglobin are the major proteins that accumulate in cilia of *Chlamydomonas*

Bbs4 mutants (Lechtreck et al., 2009) and it is likely that a range of less abundant proteins may

accumulate in *Bbs4* cilia. When stained for ubiquitin, *Bbs4 Chlamydomonas* showed a 2-fold

enrichment in ciliary signal compared to WT (**Fig. 5A-B**). Isolation of cilia revealed the

accumulation of ubiquitin conjugates above 80 kDa in *Bbs4* mutant *Chlamydomonas* cilia (**Fig.**

5C). These results are consistent with the hypothesis that ciliary proteins subject to constitutive

retrieval are ubiquitinated prior to their removal from cilia by the BBSome.

In the mammalian retina, proteomics studies have found over 100 proteins accumulating in the

outer segment (the equivalent of the cilium) of *Bbs* photoreceptors compared to WT

photoreceptors (Datta et al., 2015). When we stained sections of mouse retina for ubiquitin, we

found a drastic accumulation of ubiquitin signal in the outer segment of *Bbs4^-/-^* photoreceptors

compared to WT photoreceptors (**Fig. 5D**). These results suggest that variety of proteins that

accidentally flow into the photoreceptors outer segments –possibly caught on the tremendous flux

of rhodopsin trafficking to the outer segment (Baker et al., 2008)– are recognized as foreign to the

cilium, ubiquitinated on site and selected for removal by the BBSome (**Figure 5E**).

DISCUSSION

Recognition of K63-linked ubiquitin chains by the ciliary exit machinery

Our results indicate that K63-linked ubiquitin chains tag activated GPCRs and other unwanted proteins for BBSome-mediated removal from cilia. Prior findings in trypanosomes suggest that the BBSome may directly recognize ubiquitin (Langousis et al., 2016), although direct biochemical evidence and information regarding ubiquitin chain specificity is still missing. It is also known that the BBSome directly recognizes cytoplasmic determinants in the GPCRs it ferries out of cilia (Jin et al., 2010; Klink et al., 2017; Ye et al., 2018; Seo et al., 2011; Domire et al., 2011). We propose that the summation of weak molecular interactions between the BBSome and UbK63 chains on one hand and the BBSome and GPCRs cytoplasmic loops on the other hand increases the BBSome-cargo interaction to enable sorting of ubiquitinated GPCRs out of cilia. In support of this bivalent recognition of ubiquitinated cargoes by the BBSome, the affinity of the BBSome for cytoplasmic determinant of GPCRs have been measured to be in the μM range (Klink et al., 2017) and affinities of ubiquitin-binding proteins for ubiquitin typically reside in the sub mM range (Husnjak and Dikic, 2012). The nature of the biochemical entity that directly recognizes ubiquitin, and in particular UbK63 chains remains to be determined: while the BBSome was pulled down on Ub-agarose resin from trypanosome lysate (Langousis et al., 2016), no known Ub-binding domain is found encoded within the BBSome polypeptides. Furthermore, the IFT-A subunit IFT-139 associates with ubiquitinated α-tubulin in ciliary extracts from *Chlamydomonas*, suggesting a possible recognition of ubiquitin by IFT139 (Wang et al., 2019). Again, the absence of known Ub-binding domain in the sequence of IFT139 presents a challenge to this hypothesis.

The suppression of GPR161, SSTR3 and SMO exit by cilia-AMSH (**Figure 3**) indicate that UbK63 chains are recognized inside cilia to direct trafficking out of cilia. Similarly, (Wang et al., 2019) found that α-tubulin is modified by UbK63 chains and not UbK48 chain upon ciliary disassembly and the ubiquitin ligases that have been associated with ciliary clearance of Patched 1 and of PDGFRαα specifically assemble UbK63 chains (Kim et al., 2015; Yue et al., 2014; Schmid et al., 2018). The recognition of UbK63 chains within the cilium contrasts with the role

of UbK63 chains attached to signaling receptors at the plasma membrane. In the latter case,

UbK63 chains are not required for sorting events at the plasma membrane but UbK63 chains

become increasingly necessary as cargoes progress along the degradative route. At the terminal

step of cargo sorting into the lumen of multivesicular endosomes, a strict requirement for K63-

linked ubiquitin chain is observed (Shields and Piper, 2011; Stringer and Piper, 2011; Huang et

al., 2013).

The contrast between sites of action of molecular regulators extends to β arrestin-directed

ubiquitination. While β arrestin functions in cilia to direct the addition of ubiquitin onto

activated GPCRs (**Figure 4**), β arrestin-mediated ubiquitination of plasma membrane GPCRs

selectively affects sorting at the level of endosomes rather that at the plasma membrane

(Bhandari et al., 2007; Henry et al., 2012; Shenoy et al., 2008). In this context, the recognition of

K63-linked ubiquitin chain inside cilia may befit the endosomal origin of primary cilia (Sorokin,

1962; Westlake et al., 2011; Mitchell, 2017).

Finally, an interesting consideration is the relationship between ciliary exit and endocytosis. It has

been proposed that endocytosis of signaling receptors is intimately coupled to their exit from cilia

(Pedersen et al., 2016). However, single molecule imaging of GPR161 exiting cilia suggests that

activated GPR161 diffuses into the plasma membrane after its exit from the ciliary compartment

(Ye et al., 2018). Prior findings of translational ubiquitin fusion with ciliary proteins accumulating

in cilia of nematodes mutants for BBSome or the ESCRT machinery (Hu et al., 2007; Xu et al.,

2015) suggest the provocative possibility that ciliary exit may be directly coupled to degradative

sorting and more work will be require to determine the route taken by ubiquitinated GPCRs out

of cilia.

Ubiquitination and control of the ciliary proteome

A role for ubiquitination in selecting unwanted proteins for removal from cilia may unify the

functions of the BBSome in signal-dependent retrieval of GPCRs and the constitutive clearance

of proteins that accidentally enter cilia. In both of these cases, unwanted ciliary proteins need to

be recognized as 'non-self' by a ciliary surveillance machinery. For GPCRs, β arrestin orchestrate

the recognition of activated GPCRs as unwanted by directing their ubiquitination. A fascinating

question for future investigations is how the more than 100 proteins that accumulate in outer

segments of *Bbs* photoreceptors are recognized as 'non-self' and tagged with ubiquitin.

REFERENCES

Anvarian, Z., K. Mykytyn, S. Mukhopadhyay, L.B. Pedersen, and S.T. Christensen. 2019. Cellular signalling by primary cilia in development, organ function and disease. *Nat. Rev. Nephrol.* doi:10.1038/s41581-019-0116-9.

Baker, S.A., M. Haeri, P. Yoo, S.M. Gospe, N.P. Skiba, B.E. Knox, and V.Y. Arshavsky. 2008. The outer segment serves as a default destination for the trafficking of membrane proteins in photoreceptors. *J. Cell Biol.* 183:485–498. doi:10.1083/jcb.200806009.

Bhandari, D., J. Trejo, J.L. Benovic, and A. Marchese. 2007. Arrestin-2 interacts with the ubiquitin-protein isopeptide ligase atrophin-interacting protein 4 and mediates endosomal sorting of the chemokine receptor CXCR4. *J. Biol. Chem.* 282:36971–36979. doi:10.1074/jbc.M705085200.

Chiang, A.P., J.S. Beck, H. Yen, M.K. Tayeh, T.E. Scheetz, R.E. Swiderski, D.Y. Nishimura, T.A. Braun, K.-Y.A. Kim, J. Huang, K. Elbedour, R. Carmi, D.C. Slusarski, T.L. Casavant, E.M. Stone, and V.C. Sheffield. 2006. Homozygosity mapping with SNP arrays identifies TRIM32, an E3 ubiquitin ligase, as a Bardet-Biedl syndrome gene (BBS11). *Proc. Natl. Acad. Sci. U. S. A.* 103:6287–6292. doi:10.1073/pnas.0600158103.

Corbit, K.C., P. Aanstad, V. Singla, A.R. Norman, D.Y.R. Stainier, and J.F. Reiter. 2005. Vertebrate Smoothened functions at the primary cilium. *Nature.* 437:1018–1021. doi:10.1038/nature04117.

Craige, B., J.M. Brown, and G.B. Witman. 2013. Isolation of Chlamydomonas Flagella. *Curr. Protoc. Cell Biol.* 59:3.41.1-3.41.9. doi:10.1002/0471143030.cb0341s59.

Datta, P., C. Allamargot, J.S. Hudson, E.K. Andersen, S. Bhattarai, A.V. Drack, V.C. Sheffield, and S. Seo. 2015. Accumulation of non-outer segment proteins in the outer segment underlies photoreceptor degeneration in Bardet-Biedl syndrome. *Proc Natl Acad Sci U A.* 112:E4400–9. doi:10.1073/pnas.1510111112.

Domire, J.S., J.A. Green, K.G. Lee, A.D. Johnson, C.C. Askwith, and K. Mykytyn. 2011. Dopamine receptor 1 localizes to neuronal cilia in a dynamic process that requires the Bardet-Biedl syndrome proteins. *Cell. Mol. Life Sci. CMLS.* 68:2951–2960. doi:10.1007/s00018-010-0603-4.

Emmerich, C.H., and P. Cohen. 2015. Optimising methods for the preservation, capture and identification of ubiquitin chains and ubiquitylated proteins by immunoblotting. *Biochem. Biophys. Res. Commun.* 466:1–14. doi:10.1016/j.bbrc.2015.08.109.

Fujimuro, M., H. Sawada, and H. Yokosawa. 1994. Production and characterization of
monoclonal antibodies specific to multi-ubiquitin chains of polyubiquitinated proteins.
FEBS Lett. 349:173–180. doi:10.1016/0014-5793(94)00647-4.

Gorman, D.S., and R.P. Levine. 1965. Cytochrome f and plastocyanin: their sequence in the
photosynthetic electron transport chain of Chlamydomonas reinhardi. *Proc. Natl. Acad. Sci.*
U. S. A. 54:1665–1669. doi:10.1073/pnas.54.6.1665.

Green, J.A., C.L. Schmid, E. Bley, P.C. Monsma, A. Brown, L.M. Bohn, and K. Mykytyn. 2015.
Recruitment of β-Arrestin into Neuronal Cilia Modulates Somatostatin Receptor
Subtype 3 Ciliary Localization. *Mol. Cell. Biol.* 36:223–235. doi:10.1128/MCB.00765-15.

Haglund, K., S. Sigismund, S. Polo, I. Szymkiewicz, P.P. Di Fiore, and I. Dikic. 2003. Multiple
monoubiquitination of RTKs is sufficient for their endocytosis and degradation. *Nat. Cell*
Biol. 5:461–466. doi:10.1038/ncb983.

Henry, A.G., J.N. Hislop, J. Grove, K. Thorn, M. Marsh, and M. von Zastrow. 2012.
Regulation of endocytic clathrin dynamics by cargo ubiquitination. *Dev. Cell*. 23:519–532.
doi:10.1016/j.devcel.2012.08.003.

Howarth, M., and A.Y. Ting. 2008. Imaging proteins in live mammalian cells with biotin ligase
and monovalent streptavidin. *Nat. Protoc.* 3:534–545. doi:10.1038/nprot.2008.20.

Hu, J., S.G. Wittekind, and M.M. Barr. 2007. STAM and Hrs down-regulate ciliary TRP
receptors. *Mol. Biol. Cell*. 18:3277–3289. doi:10.1091/mbc.E07-03-0239.

Huang, F., X. Zeng, W. Kim, M. Balasubramani, A. Fortian, S.P. Gygi, N.A. Yates, and A.
Sorkin. 2013. Lysine 63-linked polyubiquitination is required for EGF receptor
degradation. *Proc. Natl. Acad. Sci.* 110:15722–15727. doi:10.1073/pnas.1308014110.

Huang, K., D.R. Diener, and J.L. Rosenbaum. 2009. The ubiquitin conjugation system is
involved in the disassembly of cilia and flagella. *J. Cell Biol.* 186:601–613.
doi:10.1083/jcb.200903066.

Husnjak, K., and I. Dikic. 2012. Ubiquitin-binding proteins: decoders of ubiquitin-mediated
cellular functions. *Annu. Rev. Biochem.* 81:291–322. doi:10.1146/annurev-biochem-
051810-094654.

Jin, H., S.R. White, T. Shida, S. Schulz, M. Aguiar, S.P. Gygi, J.F. Bazan, and M.V. Nachury.
2010. The conserved Bardet-Biedl syndrome proteins assemble a coat that traffics
membrane proteins to cilia. *Cell*. 141:1208–1219. doi:10.1016/j.cell.2010.05.015.

Kim, J., E.Y. C Hsia, A. Brigui, A. Plessis, P.A. Beachy, and X. Zheng. 2015. The role of ciliary
trafficking in Hedgehog receptor signaling. *Sci. Signal.* 8:ra55–ra55.
doi:10.1126/scisignal.aaa5622.

Klink, B.U., E. Zent, P. Juneja, A. Kuhlee, S. Raunser, and A. Wittinghofer. 2017. A
recombinant BBSome core complex and how it interacts with ciliary cargo. *eLife.* 6.
doi:10.7554/eLife.27434.

Langousis, G., M.M. Shimogawa, E.A. Saada, A.A. Vashisht, R. Spreafico, A.R. Nager, W.D.
Barshop, M.V. Nachury, J.A. Wohlschlegel, and K.L. Hill. 2016. Loss of the BBSome
perturbs endocytic trafficking and disrupts virulence of Trypanosoma brucei. *Proc Natl*
Acad Sci U A. 113:632–637. doi:10.1073/pnas.1518079113.

Lechtreck, K.-F., E.C. Johnson, T. Sakai, D. Cochran, B.A. Ballif, J. Rush, G.J. Pazour, M.
Ikebe, and G.B. Witman. 2009. The Chlamydomonas reinhardtii BBSome is an IFT
cargo required for export of specific signaling proteins from flagella. *J. Cell Biol.*
187:1117–1132. doi:10.1083/jcb.200909183.

Liew, G.M., F. Ye, A.R. Nager, J.P. Murphy, J.S. Lee, M. Aguiar, D. Breslow, S.P. Gygi, and
M.V. Nachury. 2014. The intraflagellar transport protein IFT27 promotes BBSome exit
from cilia through the GTPase ARL6/BBS3. *Dev. Cell.* 31:265–278.
doi:10.1016/j.devcel.2014.09.004.

Loktev, A.V., and P.K. Jackson. 2013. Neuropeptide Y family receptors traffic via the Bardet-
Biedl syndrome pathway to signal in neuronal primary cilia. *Cell Rep.* 5:1316–1329.
doi:10.1016/j.celrep.2013.11.011.

McCullough, J., M.J. Clague, and S. Urbé. 2004. AMSH is an endosome-associated ubiquitin
isopeptidase. *J. Cell Biol.* 166:487–492. doi:10.1083/jcb.200401141.

McCullough, J., P.E. Row, Ó. Lorenzo, M. Doherty, R. Beynon, M.J. Clague, and S. Urbé.
2006. Activation of the Endosome-Associated Ubiquitin Isopeptidase AMSH by STAM,
a Component of the Multivesicular Body-Sorting Machinery. *Curr. Biol.* 16:160–165.
doi:10.1016/j.cub.2005.11.073.

Michel, M.A., P.R. Elliott, K.N. Swatek, M. Simicek, J.N. Pruneda, J.L. Wagstaff, S.M.V.
Freund, and D. Komander. 2015. Assembly and specific recognition of k29- and k33-
linked polyubiquitin. *Mol. Cell.* 58:95–109. doi:10.1016/j.molcel.2015.01.042.

Milenkovic, L., L.E. Weiss, J. Yoon, T.L. Roth, Y.S. Su, S.J. Sahl, M.P. Scott, and W.E.
Moerner. 2015. Single-molecule imaging of Hedgehog pathway protein Smoothened in
primary cilia reveals binding events regulated by Patched1. *Proc Natl Acad Sci U A.*
112:8320–8325. doi:10.1073/pnas.1510094112.

Mitchell, D.R. 2017. Evolution of Cilia. *Cold Spring Harb. Perspect. Biol.* 9:a028290.
doi:10.1101/cshperspect.a028290.

Mukhopadhyay, S., X. Wen, N. Ratti, A. Loktev, L. Rangell, S.J. Scales, and P.K. Jackson.
2013. The ciliary G-protein-coupled receptor Gpr161 negatively regulates the Sonic
hedgehog pathway via cAMP signaling. *Cell.* 152:210–223.
doi:10.1016/j.cell.2012.12.026.

Mykytyn, K., and C. Askwith. 2017. G-Protein-Coupled Receptor Signaling in Cilia. *Cold Spring*
Harb. Perspect. Biol. a028183. doi:10.1101/cshperspect.a028183.

Mykytyn, K., R.F. Mullins, R.F. Mullins, M. Andrews, M. Andrews, A.P. Chiang, A.P. Chiang,
R.E. Swiderski, R.E. Swiderski, B. Yang, B. Yang, T. Braun, T. Braun, T. Casavant, T.
Casavant, E.M. Stone, E.M. Stone, and V.C. Sheffield. 2004. Bardet-Biedl syndrome
type 4 (BBS4)-null mice implicate Bbs4 in flagella formation but not global cilia assembly.
Proc. Natl. Acad. Sci. U. S. A. 101:8664–8669. doi:10.1073/pnas.0402354101.

Nachury, M.V. 2018. The molecular machines that traffic signaling receptors into and out of
cilia. *Curr. Opin. Cell Biol.* 51:124–131. doi:10.1016/j.ceb.2018.03.004.

Nachury, M.V., and D.U. Mick. 2019. Establishing and regulating the composition of cilia for
signal transduction. *Nat. Rev. Mol. Cell Biol.* 20:389–405. doi:10.1038/s41580-019-0116-4.

Nager, A.R., J.S. Goldstein, V. Herranz-Pérez, D. Portran, F. Ye, J.M. García-Verdugo, and
M.V. Nachury. 2017. An Actin Network Dispatches Ciliary GPCRs into Extracellular
Vesicles to Modulate Signaling. *Cell.* 168:252–263.e14. doi:10.1016/j.cell.2016.11.036.

Nakata, K., D. Shiba, D. Kobayashi, and T. Yokoyama. 2012. Targeting of Nphp3 to the
primary cilia is controlled by an N-terminal myristoylation site and coiled-coil domains.
Cytoskelet. Hoboken NJ. 69:221–234. doi:10.1002/cm.21014.

Newton, K., M.L. Matsumoto, I.E. Wertz, D.S. Kirkpatrick, J.R. Lill, J. Tan, D. Dugger, N.
Gordon, S.S. Sidhu, F.A. Fellouse, L. Komuves, D.M. French, R.E. Ferrando, C. Lam,
D. Compaan, C. Yu, I. Bosanac, S.G. Hymowitz, R.F. Kelley, and V.M. Dixit. 2008.
Ubiquitin chain editing revealed by polyubiquitin linkage-specific antibodies. *Cell.*
134:668–678. doi:10.1016/j.cell.2008.07.039.

Pal, K., S. Hwang, B. Somatilaka, H. Badgandi, P.K. Jackson, K. DeFea, and S.
Mukhopadhyay. 2016. Smoothened determines β-arrestin-mediated removal of the G
protein-coupled receptor Gpr161 from the primary cilium. *J. Cell Biol.* 212:861–875.
doi:10.1083/jcb.201506132.

Pedersen, L.B., J.B. Mogensen, and S.T. Christensen. 2016. Endocytic Control of Cellular
425   Signaling at the Primary Cilium. *Trends Biochem. Sci.* 41:784–797.
426   doi:10.1016/j.tibs.2016.06.002.

427   Rohatgi, R., L. Milenkovic, and M.P. Scott. 2007. Patched1 regulates hedgehog signaling at the
428   primary cilium. *Sci. N. Y. NY.* 317:372–376. doi:10.1126/science.1139740.

429   Sato, Y., A. Yoshikawa, A. Yamagata, H. Mimura, M. Yamashita, K. Ookata, O. Nureki, K.
430   Iwai, M. Komada, and S. Fukai. 2008. Structural basis for specific cleavage of Lys 63-
431   linked polyubiquitin chains. *Nature.* 455:358–362. doi:10.1038/nature07254.

432   Schmid, F.M., K.B. Schou, M.J. Vilhelm, M.S. Holm, L. Breslin, P. Farinelli, L.A. Larsen, J.S.
433   Andersen, L.B. Pedersen, and S.T. Christensen. 2018. IFT20 modulates ciliary PDGFRα
434   signaling by regulating the stability of Cbl E3 ubiquitin ligases. *J. Cell Biol.* 217:151–161.
435   doi:10.1083/jcb.201611050.

436   Seo, S., Q. Zhang, K. Bugge, D. Breslow, C.C. Searby, M.V. Nachury, and V.C. Sheffield.
437   2011. A novel protein LZTFL1 regulates ciliary trafficking of the BBSome and
438   Smoothened. *PLoS Genet.* 7:e1002358. doi:10.1371/journal.pgen.1002358.

439   Shaner, N.C., G.G. Lambert, A. Chammas, Y. Ni, P.J. Cranfill, M.A. Baird, B.R. Sell, J.R.
440   Allen, R.N. Day, M. Israelsson, M.W. Davidson, and J. Wang. 2013. A bright
441   monomeric green fluorescent protein derived from Branchiostoma lanceolatum. *Nat.
442   Methods.* 10:407–409. doi:10.1038/nmeth.2413.

443   Shankar, H., H. Shankar, A. Michal, A. Michal, R.C. Kern, R.C. Kern, D.S. Kang, D.S. Kang,
444   V.V. Gurevich, V.V. Gurevich, and J.L. Benovic. 2010. Non-visual arrestins are
445   constitutively associated with the centrosome and regulate centrosome function. *J. Biol.
446   Chem.* 285:8316–8329. doi:10.1074/jbc.M109.062521.

447   Shearer, R.F., and D.N. Saunders. 2016. Regulation of primary cilia formation by the ubiquitin-
448   proteasome system. *Biochem. Soc. Trans.* 44:1265–1271. doi:10.1042/BST20160174.

449   Shenoy, S.K., K. Xiao, V. Venkataramanan, P.M. Snyder, N.J. Freedman, and A.M. Weissman.
450   2008. Nedd4 mediates agonist-dependent ubiquitination, lysosomal targeting, and
451   degradation of the beta2-adrenergic receptor. *J. Biol. Chem.* 283:22166–22176.
452   doi:10.1074/jbc.M709668200.

453   Shields, S.B., and R.C. Piper. 2011. How ubiquitin functions with ESCRTs. *Traffic Cph. Den.*
454   12:1306–1317. doi:10.1111/j.1600-0854.2011.01242.x.

455   Sorokin, S. 1962. Centrioles and the formation of rudimentary cilia by fibroblasts and smooth
456   muscle cells. *J. Cell Biol.* 15:363–377. doi:10.1083/jcb.15.2.363.

Stringer, D.K., and R.C. Piper. 2011. A single ubiquitin is sufficient for cargo protein entry into
MVBs in the absence of ESCRT ubiquitination. *J. Cell Biol.* 192:229–242.
doi:10.1083/jcb.201008121.

Swatek, K.N., and D. Komander. 2016. Ubiquitin modifications. *Cell Res.* 26:399–422.
doi:10.1038/cr.2016.39.

Wang, Q., Z. Peng, H. Long, X. Deng, and K. Huang. 2019. Polyubiquitylation of α-tubulin at
K304 is required for flagellar disassembly in Chlamydomonas. *J. Cell Sci.* 132.
doi:10.1242/jcs.229047.

Westlake, C.J., L.M. Baye, M.V. Nachury, K.J. Wright, K.E. Ervin, L. Phu, C. Chalouni, J.S.
Beck, D.S. Kirkpatrick, D.C. Slusarski, V.C. Sheffield, R.H. Scheller, and P.K. Jackson.
2011. Primary cilia membrane assembly is initiated by Rab11 and transport protein
particle II (TRAPPII) complex-dependent trafficking of Rabin8 to the centrosome. *Proc*
Natl Acad Sci U A. 108:2759–2764. doi:10.1073/pnas.1018823108.

Wingfield, J.L., K.-F. Lechtreck, and E. Lorentzen. 2018. Trafficking of ciliary membrane
proteins by the intraflagellar transport/BBSome machinery. *Essays Biochem.* 62:753–763.
doi:10.1042/EBC20180030.

Wright, K.J., L.M. Baye, A. Olivier-Mason, S. Mukhopadhyay, L. Sang, M. Kwong, W. Wang,
P.R. Pretorius, V.C. Sheffield, P. Sengupta, D.C. Slusarski, and P.K. Jackson. 2011. An
ARL3-UNC119-RP2 GTPase cycle targets myristoylated NPHP3 to the primary cilium.
Genes Dev. 25:2347–2360. doi:10.1101/gad.173443.111.

Xu, Q., Y. Zhang, Q. Wei, Y. Huang, Y. Li, K. Ling, and J. Hu. 2015. BBS4 and BBS5 show
functional redundancy in the BBSome to regulate the degradative sorting of ciliary
sensory receptors. *Sci. Rep.* 5:11855. doi:10.1038/srep11855.

Yau, R., and M. Rape. 2016. The increasing complexity of the ubiquitin code. *Nat. Cell Biol.*
18:579–586. doi:10.1038/ncb3358.

Ye, F., A.R. Nager, and M.V. Nachury. 2018. BBSome trains remove activated GPCRs from
cilia by enabling passage through the transition zone. *J. Cell Biol.* 217:1847–1868.
doi:10.1083/jcb.201709041.

Yue, S., L.-Y. Tang, Y. Tang, Y. Tang, Q.-H. Shen, J. Ding, Y. Chen, Z. Zhang, T.-T. Yu, Y.E.
Zhang, and S.Y. Cheng. 2014. Requirement of Smurf-mediated endocytosis of Patched1
in sonic hedgehog signal reception. *eLife.* 3:128. doi:10.7554/eLife.02555.

Zhang, Q., D. Nishimura, S. Seo, T. Vogel, D.A. Morgan, C. Searby, K. Bugge, E.M. Stone, K.
Rahmouni, and V.C. Sheffield. 2011. Bardet-Biedl syndrome 3 (Bbs3) knockout mouse

model reveals common BBS-associated phenotypes and Bbs3 unique phenotypes. *Proc*
Natl Acad Sci U A. 108:20678–20683. doi:10.1073/pnas.1113220108.

ACKNOWLEDGMENTS

We thank Drs. Kirk Mykytyn, Val Sheffield, Alice Ting, and Mark Scott for the gifts of cDNAs; Kathryn Anderson, Vishva Dixit (Genentech), and Felice Dunn for the gifts of antibodies; Val Sheffield for the gift of the *Bbs4*$^{-/-}$ mice, George B. Witman and the Chlamydomonas Resource Center for the gifs of *Chlamydomonas* strains; Yien-Ming Kuo for help with cryosections of mouse retina and core support; Dhivya Kumar and Tyler Picariello for the assistance with *Chlamydomonas* cultures; Nicholas Morante and Jeremy Reiter for providing *Bbs4*$^{-/-}$ mouse eyes; Fan Ye for generating cell lines used in the study; Irene Ojeda Naharros for help with single cilia tracking and the Nachury lab for helpful discussions. This work was funded by NIGMS (R01-GM089933, M.V.N.). This work was made possible, in part, by NEI EY002162 - Core Grant for Vision Research and by the Research to Prevent Blindness Unrestricted Grant (M.V.N.). Authors contributions: SRS conducted all experimental work except for the experiments in Figure 4A-D, which were conducted by ARN. MVN supervised research. SRS and MVN wrote the manuscript.

Figure 1

Figure 1. Signal-dependent ubiquitination of ciliary GPCRs drives their BBSome-dependent exit.

A. SSTR3 fused to the fluorescent protein mNeonGreen (NG) at its intracellular C-terminus and a biotinylation Acceptor Peptide (AP) tag at its extracellular N-terminus was expressed under the control of an attenuated EF1α promoter (pEF1α$^{\Delta}$) in wild type (WT) and *Arl6$^{-/-}$* IMCD3 cells. Cells were treated with or without somatostatin-14 (sst) for 2 h, then fixed and stained for acetylated tubulin (acTub, magenta) and ubiquitin (Ub, yellow in main panels and gray in insets). ^{AP}SSTR3^{NG} was imaged through the intrinsic fluorescence of NG. Channels were shifted in this and all subsequent micrographs to facilitate visualization of ciliary signals. Scale bar: 5µm (main panel), 2µm (inset). In WT cells, the ciliary SSTR3 signal decrease over the experimental time course. In *Arl6$^{-/-}$* cells, SSTR3 fails to exit cilia and an increase in the ciliary Ub level is detected. As a control, *Arl6$^{-/-}$* cells that did not express SSTR3-NG were tested; no increase in ciliary Ub levels were observed upon addition of sst. **B.** The fluorescence intensity of the Ub channel in the cilium was measured in each condition and the data are represented as violin plots. The thick bar indicates the median and the dotted lines the first and third quartiles. An 11-fold increase in ciliary Ub signal is observed upon addition of sst to SSTR3-expressing cells. Asterisks indicate ANOVA significance value. ********, p= <0.0001. **C.** IMCD3 cells of the indicated genotypes were treated with the Smoothened agonist SAG or the vehicle DMSO for 2h. Cells were then fixed and stained for acTub and Ub. Scale bar 5µm (main panel), 2µm (inset). Activation of Hh signaling promotes a detectable increase in ciliary Ub levels only in *Arl6$^{-/-}$* cells. **D.** Violin plots of the fluorescence intensity of the Ub channel in the cilium in each condition are shown. Asterisks indicate ANOVA significance value. ********, p= <0.0001. **E.** IMCD3-[pEF1α$^{\Delta}$-^{AP}SSTR3] stably expressed the ER-localized biotin ligase BirA to enables the biotinlylation of ^{AP}SSTR3. SSTR3 was either the wild-type allele (WT) or a variant where all five cytoplasm-facing lysine residues (listed in Methods) were mutated to arginine (cK0). Ciliary ^{AP}SSTR3 was pulse-labeled by addition of Alexa647-conjugated mSA (mSA647) to the medium for 5-10 min before addition of sst. Far red fluorescence was tracked in individual cilia at 10 min capture intervals. For each individual cilia, fluorescence intensities were normalized to the value at t = 0. A comparison of

the ciliary levels of SSTR3 at t=0 is shown in **Fig. S2A**. Data were plotted and fitted to a line.

Error bars: 95% CI. $\mathcal{N}$ =18-22 cilia. **F.** GPR161 fused to three tandem repeats of NG at its C-

terminus was expressed under the control of the ultra-weak δ-crystallin promoter (pCrys) in

IMCD3 cells. GPR161 was either the wild-type allele (WT) or a variant where all eighteen

cytoplasm-facing lysine residues (listed in Methods) where mutated to arginine (cK0). IMCD3-

[pCrys-GPR161^{3NG}] cells were treated with SAG for 2h. During the course of the experiment,

NG fluorescence was tracked in individual cilia at 10 min capture intervals. Fluorescence data

were acquired and analyzed as in panel **(E)**. A comparison of the ciliary levels of GPR161 at t=0

is shown in **Fig. S2B**. Error bars: 95% CI. $\mathcal{N}$ =19-10 cilia.

A IMCD3-[pEF1α^{Δ}-^{AP}SSTR3^{NG}] +sst

acTub UbK48 NG   acTub UbK63 NG

WT

Arl6$^{-/-}$

B   UbK48   UbK63

Ciliary ubiquitin levels (au)

n=51

n=37

n=41   n=44

****
****

WT   *Arl6$^{-/-}$*   WT   *Arl6$^{-/-}$*

[pEF1α^{Δ}-SSTR3^{NG}] + sst

Figure 2

Figure 2. Signal-dependent accumulation of K63-linked ubiquitin chains inside

cilia of *Bbs* mutant cells.

A. IMCD3 cells of the indicated genotypes expressing SSTR3^{NG} were treated with somatostatin-

14 for 2 h. Cells were fixed and stained for AcTub (magenta) and with antibodies specific for the

lysine 63 (UbK63) or lysine 48 (UbK48) Ub chain linkages (yellow in main panel and gray in

inset). SSTR3^{NG} was imaged through the intrinsic fluorescence of NG. Scale bar 5µm (main

panel), 2µm (inset). **B.** The fluorescence intensity of the UbK48 and UbK63 channels in the

cilium are represented as violin plots. A 14-fold increase in ciliary Ub abundance is detected with

the K63Ub linkage-specific antibody. Asterisks indicate ANOVA significance value. ********, p=

<0.0001. No ciliary signal is detected with the K48Ub linkage-specific antibody.

Figure 3

Figure 3. Ciliary K63-linked ubiquitin chains are required for the removal of BBSome cargoes from cilia.

A. Diagram of the working model and the experimental strategy. **B.** IMCD3-[pEF1α^A-^{AP}SSTR3; pEF1α-BirA•ER] were transfected with plasmids expressing the ciliary targeting signal (CTS) of NPHP3 fused to GFP only (left panels), to GFP and the catalytic domain of the K63-specific deubiquitinase AMSH (middle panels), or to GFP and the catalytically inactive E280A variant of AMSH catalytic domain (AMSH†, right panels). Ciliary ^{AP}SSTR3 was pulse-labeled with mSA647 for 5-10 min and cells were then treated with or without sst for 2h, before fixation and staining for acetylated tubulin (acTub, magenta) and DNA (blue). The NPHP3^{CTS} fusions were visualized through the intrinsic fluorescence of GFP (cyan) and ^{AP}SSTR3 was visualized via mSA647 (yellow in the main frames and gray in the insets). Scale bar 5µm (main panel), 2µm (inset). **C.** The fluorescence intensities of ciliary $^{mSA647-AP}$SSTR3 are represented as violin plots. Asterisks indicate Kruskal-Wallis test significance values. *******, p= <0.001; ******, p= <0.01. Addition of sst triggers the exit of SSTR3 from cilia in cells transfected with the controls NPHP3^{CTS} and NPHP3^{CTS}-AMSH† but NPHP3^{CTS}-AMSH blocks ciliary exit of SSTR3. **D.** RPE1-hTERT (RPE) cells transfected with the indicated constructs were treated with SAG or vehicle (DMSO) for 2h, then fixed and stained for acetylated tubulin (magenta) and GPR161 (yellow in the main frames and gray in the insets). The NPHP3^{CTS} fusions were visualized through the intrinsic fluorescence of GFP (cyan). Scale bar 5µm (main panel), 2µm (inset). **E.** The fluorescence intensities of ciliary GPR161 are represented as violin plots. Asterisks indicate ANOVA significance value. ********, p= <0.0001. NPHP3^{CTS}-AMSH specifically blocks the SAG-dependent exit of GPR161 from cilia. **F.** IMCD3 cells transfected with the indicated constructs were fixed and stained for acetylated tubulin (magenta) and SMO (yellow in the main frames and gray in the insets). The NPHP3^{CTS} fusions were visualized through the intrinsic fluorescence of GFP (cyan). **G.** The fluorescence intensities of ciliary SMO are represented as violin plots. Asterisks indicate ANOVA significance value. ********, p= <0.0001. NPHP3^{CTS}-AMSH specifically blocks the constitutive exit of SMO from cilia and promotes accumulation of SMO in cilia in the absence of pathway activation.

A

SSTR3 β-Arrestin2

siControl   siArl6

Time with sst (min)

base
tip

base
tip

B

siARL6
$t_{1/2}$=5.0±0.8 min

siControl
$t_{1/2}$=4.6±0.6 min

Ciliary β-Arrestin2 (RFU)

Time (min)

C

Ciliary fluorescence (% of t_0)

SSTR3

Time (min)

20   40

Fluorescence

$F(t_f)$

$F(t_0)$

β-Arrestin2 (cilium)

BBS5 (tip)

D

Ciliary fluorescence (% of t_0)

GPR161

Time (min)

20   40

Ciliary fluorescence

$F(t_f)$

$F(t_0)$

SMO

β-Arrestin2

E   IMCD3-[pEF1α^Δ-SSTR3^{NG}] + sst

WT   Arl6$^{-/-}$ Arl6$^{-/-}$/
                              β-arrestin2$^{-/-}$

acTub Ub NG

F

Ciliary ubiquitin levels (au)

****

n=50

n=47

n=49

WT   Arl6$^{-/-}$ Arl6$^{-/-}$/
                         β-arrestin2$^{-/-}$

IMCD3-[pEF1α^Δ-SSTR3^{NG}] + sst

G   IMCD3 + SAG

WT   Arl6$^{-/-}$ Arl6$^{-/-}$/
                              β-arrestin2$^{-/-}$

acTub Ub

H

Ciliary ubiquitin levels (au)

****   **

n=48

n=48

n=32

WT   Arl6$^{-/-}$ Arl6$^{-/-}$/
                         β-arrestin2$^{-/-}$

IMCD3 + SAG

Figure 4

Figure 4. β-arrestin 2 directs signal-dependent ubiquitination of ciliary GPCRs.

A. β-Arrestin2 is rapidly recruited to cilia upon activation of SSTR3. siRNA-treated IMCD3-[^{AP}SSTR3, β-Arrestin2^{GFP}] cells were pulse-labeled with mSA647 before addition of sst. As shown previously (Shankar et al., 2010), β-Arrestin2 is at the basal body in unstimulated cells. Scale bar: 2 µm. **B.** Kinetics of β-Arrestin2^{GFP} recruitment to cilia upon sst stimulation in control- and ARL6-depleted IMCD3-[^{AP}SSTR3, β-Arrestin2^{GFP}] cells. Data were fit to a simple exponential. (n=10 cilia). **C-D.** Kinetics of early trafficking events for retrieval of SSTR3 **(C)** and GPR161 **(D)**. The top panels show removal of stably expressed ^{AP}SSTR3^{NG} or ^{AP}GPR161^{NG3} following addition of sst **(C)** or SAG **(D)**. The bottom panels show the kinetics of β-Arrestin2^{GFP} entry **(C-D)**, ^{AP}Smoothened^{NG} entry **(D)**, and ^{NG3}BBS5 tip accumulation **(C)** measured in IMCD3 cells stably expressing the indicated proteins. Fluorescence values are normalized to the initial (t_0) and final (t_f) values. Error bars: SEM. (n=6-11 cilia). **E.** IMCD3-[pEF1α^Δ-^{AP}SSTR3^{NG}] WT, knockout for *Arl6* only and double knockout for *Arl6* and *β-Arrestin2* were treated with sst for 2h before fixation and staining for acetylated tubulin (acTub, magenta) and ubiquitin (Ub, yellow in the main frames and gray in the insets). SSTR3^{NG} was visualized through the intrinsic fluorescence of NG. SSTR3 exit is blocked in *Arl6$^{-/-}$* cells regardless of the *β-Arrestin2* genotype but the ciliary Ub signal is only detected in *Arl6$^{-/-}$* cells when β-Arrestin2 function is intact. Scale bar: 5µm (main panel), 1µm (inset). **F.** Violin plots representing the ciliary levels of Ub under the indicated conditions. Asterisks indicate ANOVA significance value. ********, p= <0.0001. **G.** IMCD3 of the indicated genotypes were treated with SAG for 2h before fixation and staining for acetylated tubulin (acTub, magenta) and ubiquitin (Ub, yellow in the main frames and gray in the insets). Scale bar: 5µm (main panel), 1µm (inset). **H.** Violin plots representing the ciliary Ub levels. Asterisks indicate ANOVA significance value. ********, p= <0.0001; ******, p= <0.01. The appearance of a Ub signal in cilia in *Arl6$^{-/-}$* cells depends on β-Arrestin2.

Figure 5

Figure 5. Accumulation of ubiquitinated proteins in cilia of *Chlamydomonas* cells and mammalian photoreceptors.

A. *Chlamydomonas* cells of the indicated genotypes were fixed and stained for acetylated tubulin (acTub, magenta) and ubiquitin (Ub, yellow). Scale bar: 5 µm. A weak ubiquitin signal detected in WT cells is increased in *bbs4* mutant cells. **B.** Violin plots of the ciliary Ub levels in WT and *bbs4 Chlamydomonas* cells. Asterisks indicate Mann-Whitney test significance values. ********, p= <0.0001. The ciliary levels of Ub are increased more than 2-fold in *bbs4* cells as compared to WT. **C.** Cilia purified from WT and *bbs4* cells via deciliation with dibucaine and differential centrifugation were solubilized, and the protein contents resolved by SDS-PAGE. Immunoblots for Ub (using the P4D1 antibody) and tubulin are shown. A rise in ubiquitin conjugates is observed in *bbs4 Chlamydomonas* cells compared to WT. **D.** Mouse retina were fixed, embedded and sectioned before staining for ubiquitin (Ub, yellow), cone arrestin (a marker of the photoreceptor outer segment layer) and DNA (cyan). OS: outer segment; IS: inner segment; ONL: outer nuclear layer; OPL: outer plexiform layer; INL: inner nuclear layer. Scale bar: 10µm. **E.** Model for the role of ubiquitin chains in selecting cargoes for signal-dependent and constitutive exit via BBSome-mediated transport.

Supplementary Figure 1

Figure S1. Signal-dependent ubiquitination of ciliary GPCRs

A. IMCD3 cells of the indicated genotypes expressing ^{AP}SSTR3^{NG} were treated with

somatostatin (sst) for 2 h. Cells were fixed and stained for ubiquitin with the FK1 monoclonal

antibody (Ub (FK1), yellow in main panels and gray in insets) and acetylated tubulin (acTub,

magenta). ^{AP}SSTR3^{NG} was imaged through the intrinsic fluorescence of NG. Scale bar: 5µm

(main panel), 2µm (inset). **B.** Violin plots of the fluorescence intensity of the Ub channel in the

cilium in each condition are shown. A 3-fold increase in ciliary Ub signal is observed upon

addition of sst to SSTR3-expressing cells. Asterisks indicate ANOVA significance value. ********, p=

<0.0001. **C.** IMCD3 cells of the indicated genotypes were treated with or without somatostatin

for 2 h. Cells were fixed and stained for acetylated tubulin (acTub, magenta) and ubiquitin (Ub,

yellow in main panels and gray in insets). Scale bar 5µm (main panel), 2µm (inset). No significant

increase in ciliary Ub levels were observed upon addition of sst to *Arl6$^{-/-}$* cells. **D.** The

fluorescence intensity of the Ub channel in the cilium was measured in each condition and the

data are represented as violin plots. **E.** IMCD3 cells of the indicated genotypes expressing

^{AP}SSTR3^{NG} were treated with somatostatin-14 for 2 h. Cells were fixed and stained for AcTub

(magenta) and UbK63 (yellow in main panel and gray in inset). ^{AP}SSTR3^{NG} was imaged through

the intrinsic fluorescence of NG. Scale bar 5µm (main panel), 2µm (inset). **F.** The fluorescence

intensity of the UbK63 channel in the cilium is represented as violin plots. Asterisks indicate

ANOVA significance value. ********, p= <0.0001.

Supplementary Figure 2

Figure S2. Fluorescence intensities of WT and cK0 GPCR variants at t=0.

A. Ciliary [AP]SSTR3 was pulse-labeled by addition of Alexa647-conjugated mSA (mSA647) to the

medium for 5-10 min before addition of sst. Violin plots of the fluorescence intensities for WT

and cK0 SSTR3 imaged in the far red channel at t=0 are shown. **B.** Violin plots of the

fluorescence intensities of NG fluorescence corresponding to WT and cK0 GPR161[NG] at t=0 are

shown.

MATERIALS AND METHODS

Cell culture

The mouse IMCD3 cell lines used in the study were generated from a parental IMCD3-FlpIn

cell line (gift from P.K. Jackson, Stanford University, Stanford, CA). IMCD3-FlpIn cells were

cultured in DMEM/F12 (11330-057; Gibco) supplemented with 10% FBS (100-106; Gemini

Bio-products), 100 U/ml penicillin-streptomycin (400-109; Gemini Bio-products), and 2 mM L-

glutamine (400-106; Gemini Bio-products). The RPE1-hTERT cell line (ATCC CRL-4000) was

cultured in DMEM/F12 supplemented with 10% FBS, 100 U/ml penicillin-streptomycin, 2 mM

L-glutamine and 0.26% sodium bicarbonate (25080; Gibco).

Ciliation was induced by serum starvation in media containing 0.2% FBS for 16 to 24 h.

Plasmid construction and Generation of stable cell lines

Stable isogenic IMCD3 cell lines were generated using the Flp-In system (ThermoFisher

Scientific). Low-expression promoters and additional expression cassettes were introduced into

pEF5/FRT plasmid as described (Nager et al., 2017; Ye et al., 2018). To reduce expression

levels, the EF1α promoter was attenuated by mutating the TATA-box (pEF1α$^\Delta$) or replaced with

a minimal chicken lens δ-crystallin promoter (pCrys).

Coding sequences were amplified from plasmids encoding mouse GPR161 (BC028163;

Mammalian Gene Collection [MGC]; GE Healthcare), mouse SSTR3 (gift from Kirk Mykytyn,

Ohio State University, Columbus, OH), and human BBS5 (gifts from V. Sheffield, University of

Iowa, Iowa City, IA), BirA-ER (gift from Alice Ting). β-arrestin 2 (a gift from Mark Scott) was

stably expressed from a pCMV-based plasmid (pEGFP-N). SSTR3 and SMO expression were

driven by pEF1α$^\Delta$, GPR161 expression by pCrys, and BBS5 expression by pEF1α. Coding

sequences were fused to GFP, mNeonGreen (NG, (Shaner et al., 2013), or an acceptor peptide

for the biotin ligase BirA (AP) (Howarth and Ting, 2008). Multiple rounds of site-directed

mutagenesis were performed to generate SSTR3-cK0, a variant with all the cytoplasm-facing

lysine residues (K233, K330, K356, K407, and K421) mutated to arginine. GPR161-cK0, a

variant with 18 cytoplasm-exposed lysine residues (K83, K84, K93, K167, K247, K250, K269,

K296, K358, K362, K455, K469, K473, K481, K486, K497, K504, and K548) mutated to

arginine, was gene synthesized (GenScript). Cilia-AMSH was generated by fusing the catalytic

domain of mouse AMSH (gift from David Komander; Addgene plasmid #66712; (Michel et al.,

2015) with NPHP3(1-200) and GFP to create NPHP3[1-200]-GFP-AMSH[243-424]. A

catalytically dead version of AMSH was generated by mutating the water-activating Glu280

residue to Ala (Sato et al., 2008; Huang et al., 2013).

CRISPR-edited *Arl6*$^{-/-}$ and *Arl6*$^{-/-}$*/Arrb2*$^{-/-}$ cell lines were described previously (Liew et al., 2014;

Nager et al., 2017). The genotypes are *Arl6*, NM_001347244.1:c.- 10_25del; c.3_6del; and *Arrb2*,

NM_001271358.1:c.1 12del;c112_113del.

Transfection

For generation of all stable cell lines except for the ones expressing β-arrestin 2^{GFP}, FRT-based

plasmids were reverse transfected using XtremeGene9 (Roche) into IMCD3 Flp-In cells along

with a plasmid encoding the Flp recombinase (pOG44) and stable transformants were selected by

blasticidin resistance (4 ug/ml). The transfection mixture was then added to the cells. pEGFP-N•

β-arrestin 2-GFP was transfected into IMCD3 cells using Lipofectamine 2000, and clones were

selected using neomycin resistance.

Antibodies and drug treatments

The following monoclonal antibodies were used for immunofluorescence: anti-acetylated tubulin

(mouse; clone 6-11B-1; Sigma-Aldrich; 1:500), anti-ubiquitin clone FK2 was used in all

experiments except where indicated (mouse; D058-3; Medical and Biological Laboratories;

1:500), anti-ubiquitin FK1 in **Fig. S1A-B** (mouse; D071-3; Medical and Biological Laboratories;

1:500), anti-ubiquitin Lys48 linkage (rabbit; clone Apu2; 05-1307; Millipore-Genentech; 1:500),

anti-ubiquitin Lys63 linkage (human; clone Apu3; a gift from Genentech; 1:500). The following

monoclonal antibodies were used for immunoblotting: anti-ubiquitin (mouse; clone P4D1; 3936;

Cell Signaling; 1:1000), anti-α tubulin (mouse; clone DM1A; MS-581-P1ABX; Thermo

Scientific;1:1000). The following polyclonal antibodies were used for immunofluorescence: anti-

GPR161 (rabbit; 13398-1-AP; Proteintech; 1:100), anti-Smoothened (rabbit; a gift from Kathryn

Anderson, Memorial Sloan Kettering Cancer Center, New York, NY; 1:500), anti-cone-arrestin

(rabbit; AB15282; Millipore; 1:500). Biotinylated SSTR3 and GPR161 were detected using

Alexa Fluor 647–labeled monovalent streptavidin (mSA647) (Ye et al., 2018). The following

reagents were used at the indicated concentrations: 200 nM SAG and 10 μM somatostatin 14.

Somatostatin 14 stocks were made in DMEM/F12 media, HEPES, no phenol red (11039-021,

GIBCO). SAG was dissolved in DMSO (276855, Sigma-Aldrich).

Imaging and microscopy

Cells were imaged either on a DeltaVision system (Applied Precision) equipped with a PlanApo

60x/1.40 objective lens (Olympus), CoolSNAP HQ2 camera (Photometrics), and solid-state

illumination module (InsightSSI), or on a confocal LSM 700 (Zeiss) microscope equipped with

40x Plan-Apochromat 1.3 DIC oil objective. Z stacks were acquired at 0.5 µm interval (LSM

700, Figs. 1, 2, 5, S1 and S2) or 0.2 µm interval (DeltaVision, Figs. 3 and 4).

For fixed imaging, 60,000 cells were seeded on acid-washed coverglass (12 mm #1.5; 12-545-81;

Thermo Fisher Scientific). Cells were grown for 24 h and then starved for 16-24 h in 0.2% FBS

media before experimental treatment. After treatment, cells were fixed with 4%

paraformaldehyde in PBS for 15 min at 37°C and extracted in -20°C methanol for 5 min. Cells

were then permeabilized in IF buffer (PBS supplemented with 0.1% Triton X-100, 5% normal

donkey serum (017-000-121; Jackson ImmunoResearch Laboratories), and 3% bovine serum

albumin (BP1605-100; Thermo Fisher Scientific)), incubated at room temperature for 1h with

primary antibodies diluted in IF buffer, washed three times in IF buffer, incubated with

secondary antibodies (Jackson ImmunoResearch Laboratories) diluted in IF buffer for 30 min,

and washed three times with IF buffer. DNA was stained with Hoechst 33258 (H1398; Molecular

Probes), cells were washed twice more with PBS, and coverglass mounted on slides using

fluoromount-G (17984-25; Electron Microscopy Sciences).

For live-cell imaging, 300,000 cells were seeded on acid-washed 25mm cover glass (Electron

Microscopy Sciences). After 24 h of growth, cells were serum starved for 16 h and transferred to

the DeltaVision stage for imaging at 37°C within an environmental chamber that encloses the

microscope and the stage. Cells were imaged in DMEM/F12 media, HEPES, no phenol red

(11039-021, GIBCO). To measure SSTR3 exit, cells expressing [AP]SSTR3 were first washed three

times with PBS and then pulse-labeled with 2µg/ml mSA647 for 5 min at 37°C. Cells were then

imaged after addition of sst (for SSTR3) or SAG (for GPR161) for 2 h with an acquisition every

10 min. Each acquisition consisted of a 3-plane Z-stack with each plane separated by 0.5 µm.

Transmittance was set to 5% and exposure time to 500 ms.

WT or *Bbs4*[-/-] mice (Mykytyn et al., 2004) eyes were enucleated at P21. Whole eyes were fixed in

4% paraformaldehyde/PBS overnight at 4°C and then washed three times with PBS. The eyes

were infiltrated in 30% sucrose/PBS at 4 °C overnight. Eyes were placed in OCT compound

embedding medium (4583; Tissue-Tek), frozen on dry ice, and stored at −80 °C. 14 µm sections

were cut on a cryostat (CM 1850; Leica). Sections were processed for immunofluorescence as

follows. Sections were blocked in blocking solution (50 mM Tris pH7.4, 100 mM NaCl, 0.1%

Triton-X 100, 3% normal donkey serum, and 0.1% BSA) for 1 h at room temperature, then

incubated with primary antibodies overnight at 4°C, followed by three washes in wash solution

(50 mM Tris pH7.4, 100 mM NaCl, and 0.1% TX100). After rinsing in wash solution, sections

were incubated with secondary antibodies for 1 h at room temperature. DNA was counterstained

using Hoechst 33258 dye, sections washed twice more with PBS, and mounted on slides using

Fluoromount-G. Sections were imaged on Confocal LSM 700 (Zeiss) microscope.

Image analysis Bulk GPCR exit assays

Files were imported from the Deltavision or LSM700 workstations into ImageJ/ Fiji (National

Institutes of Health) for analysis. For the quantification of ciliary ubiquitin signal in fixed cells,

maximum intensity projections were used. The ciliary ubiquitin intensity was measured using the

following equation:

$F_{Ub\text{-}cilia} = F_{Ub\text{-}cilia\_measured} - F_{background}$

$F_{Ub\text{-}cilia\_measured}$ is the total ciliary ubiquitin fluorescence detected, $F_{background}$ is the background

ubiquitin fluorescence measured in the adjacent area. For all measurements, the fluorescence

integrated density was used. No gamma adjustment was applied during figure preparation; all the

representative micrographs for respective panels are displayed within the same dynamic range.

For some representative micrographs, the most in-focus plane was used rather than the

maximum intensity projection (Fig.2, and Fig. 4, panel E and G).

To measure SSTR3 exit, for each time point, the integrated Alexa Fluor 647 fluorescence density

in the maximum intensity projection was measured using ImageJ. The cilia-adjacent fluorescence

was subtracted as the background, and a mathematical photobleaching correction was applied:

$F_{cilia} = (F_{mSA647\_measured}/F_{mSA647\_1}) + ((1 - e^{-\lambda}) * (n - 1))$, where λ is the photobleaching decay

constant, n is the number of images taken, $F_{mSA647\_measured}$ is the integrated mSA647 fluorescence

measured for image n, F_{mSA657_1} is the measurement for the first time point, and F_{cilia} is the

reported fluorescence. In this equation, F_{cilia} is reported in relative fluorescence units (RFUs).

$^{AP}GPR161^{3NG}$ exit was followed similarly with the difference that NG fluorescence intensity was

measured.

***Chlamydomonas* culture, isolation of flagella and immunofluorescence**

Chlamydomonas WT- g1 (nit1, agg1, mt+) [gift from George B. Witman, University of

Massachusetts Medical School, Worcester, Massachusetts]and *Bbs4*- CC-4377 ptx5-1/bbs4-1::

NIT1 agg1 mt+ (Chlamydomonas Resource Center) strains were cultured asynchronously under

light in TAP media (Gorman and Levine, 1965) for 72 h. Flagellar fractions were prepared from

two liters of cultures of WT or *Bbs4* cells as described (Craige et al., 2013). Briefly, cells were

harvested by centrifugation for 5 min at 2,000 rpm (1,100 x *g*) at room temperature. Cells were

resuspended in 10 mM HEPES, pH 7.4 and centrifuged for 5 min at 2,000 rpm (1,100 x *g*). Next,

the cell pellet was gently resuspended in ice-cold HMDS (10 mM HEPES, pH 7.4, 5 mM

$MgSO_4$, 1 mM DTT, and 4% (w/v) sucrose). Deflagellattion was achieved by addition of 5 mM

dibucaine to each tube of cells and by pipetting up and down ~10 times using a 10-ml plastic

serological pipette. The suspension was spun for 5 min at 1,800 x *g* at 4°C. The supernatant

containing flagella was underlaid with 9 ml ice-cold HMDS-25% sucrose (10 mM HEPES, pH

7.4, 5 mM $MgSO_4$,1 mM DTT, and 25% (w/v) sucrose), followed by a centrifugation step for 10

min at 2,800 rpm (2,400 x *g*), 4°C. The supernatant above the sucrose interface was collected

using a 25-ml serological pipette and flagella were pelleted by centrifugation for 20 min at 30,000

x *g* (16,000 rpm), 4°C. The flagellar pellet was resuspended in 100 µl HMDS buffer. Protein

concentrations were measured by Bradford and 25 ug were resolved by SDS-PAGE for

immunoblotting.

Chlamydomonas immunofluorescence was performed as follows. Cells were fixed with 4 %

paraformaldehyde in MT buffer* (30 mM HEPES, pH7, 5 mM EGTA, 5 mM MgSO4, and

25mM KCl) for 20 minutes in suspension. The cells were centrifuged at 1000 rpm for 5 minutes,

resuspended in 100 µl of fixative and transferred onto slides coated with 1 mg/ml poly-L-lysine.

After 5 minutes, the unadhered cells were washed off by rinsing with PBS. Cells were

permeabilized in 0.5% Triton-X 100 for 20 minutes followed by blocking for 1 h in blocking

buffer (3% fish skin gelatin, 1% BSA, 0.1% Tween20 in PBS) at room temperature. Cells were

incubated at 4°C overnight with primary antibodies diluted in blocking buffer, washed five times

in PBS and incubated with secondary antibodies for 2 h at room temperature. After five washes

in PBS, coverglasses were mounting on slides using Fluoromount-G. Cells were imaged on the

LSM700 confocal microscope.