**TITLE**

**Lysine63-linked ubiquitin chains earmark GPCRs for BBSome-**

**mediated removal from cilia**

**AUTHORS:** Swapnil Rohidas Shinde, Andrew R. Nager[#] and Maxence V. Nachury[*]

**AFFILIATION:**

Department of Ophthalmology, University of California San Francisco, CA 94143, USA

[*] Correspondence: maxence.nachury@ucsf.edu

[#] Current address: Cancer Immunology Discovery, Pfizer Inc., San Diego, CA, 92121, USA

## ABSTRACT (158 words)

Regulated trafficking of G-protein coupled receptors (GPCRs) controls cilium-based signaling pathways. β-arrestin, a molecular sensor of activated GPCRs, and the BBSome, a complex of Bardet-Biedl Syndrome (BBS) proteins, are required for the signal-dependent exit of ciliary GPCRs but the functional interplay between β-arrestin and the BBSome remains elusive. Here we find that, upon activation, ciliary GPCRs become tagged with K63-linked ubiquitin (K63Ub) chains in a β-arrestin-dependent manner prior to BBSome-mediated exit. Removal of ubiquitin acceptor residues from the somatostatin receptor 3 (SSTR3) and from the orphan GPCR GPR161 demonstrates that ubiquitination of ciliary GPCRs is required for their regulated exit from cilia. Furthermore, targeting a K63Ub-specific deubiquitinase to cilia blocks the exit of GPR161, SSTR3 and Smoothened (SMO) from cilia. Finally, ubiquitinated proteins accumulate in cilia of mammalian photoreceptors and *Chlamydomonas* cells when BBSome function is compromised. We conclude that K63Ub chains mark GPCRs and other unwanted ciliary proteins for recognition by the ciliary exit machinery.

## INTRODUCTION

The regulated trafficking of signaling receptors in and out of cilia is a central regulatory mechanism of many cilia-based signaling pathways (Nachury and Mick, 2019; Anvarian et al., 2019; Mykytyn and Askwith, 2017; Gigante and Caspary, 2020). For example, upon activation of the Hedgehog signaling pathway, the Hedgehog receptor Patched 1 and the G protein-coupled receptor (GPCR) GPR161 disappear from cilia while GPCR Smoothened (SMO) accumulates in cilia (Rohatgi et al., 2007; Corbit et al., 2005; Mukhopadhyay et al., 2013). In all three cases, regulated exit from cilia plays a major role in the on-demand redistribution of signaling molecules (Nachury and Mick, 2019). Signal-dependent exit is likely to be a general characteristic of ciliary GPCRs as the Somatostatin Receptor 3 (SSTR3), the Dopamine Receptor 1 (D1R), the Melanocortin concentrating hormone receptor 1 (MCHR1) and the neuropeptide receptor 2 (NPY2R) all disappear from cilia upon exposure to agonist (Domire et al., 2011; Loktev and Jackson, 2013; Nager et al., 2017; Green et al., 2015). A major question is how GPCRs are selected for removal from cilia in an activity-dependent manner.

The major trafficking entities mediating signal-dependent exit from cilia are the BBSome and β-arrestin 2. The BBSome is an evolutionarily conserved complex of eight Bardet-Biedl Syndrome (BBS) proteins that directly recognizes intracellular determinants in GPCRs and ferries them out of cilia (Nachury, 2018; Wingfield et al., 2018). The BBSome associates with the intraflagellar transport (IFT) machinery and has been proposed to act as adaptor between the motor-driven intraflagellar transport trains and the membrane proteins that are to be removed from cilia. Because the BBSome is not known to have the ability to discriminate active from inactive GPCRs, there must exist another layer of regulation that commits activated GPCRs for exit. β-arrestin 2 is a well-established molecular sensor of the activation state of GPCRs that is required for the signal-dependent exit of GPR161 and SSTR3 (Pal et al., 2016; Green et al., 2015; Nager et al., 2017). No association of β-arrestin 2 with ciliary trafficking complexes has been reported to date and it remains unclear how β-arrestin 2 relays information regarding the state of activation of ciliary GPCRs to the ciliary exit machinery.

An emerging player in ciliary exit is ubiquitination (Shearer and Saunders, 2016). Ubiquitin (Ub),

a 76-amino acid polypeptide, becomes conjugated to acceptor lysine residues on substrate

proteins by ubiquitin ligases and tags substrates for degradation or other regulatory fates (Yau

and Rape, 2016; Swatek and Komander, 2016). A role for ubiquitin in promoting ciliary exit is

suggested by multiple lines of evidence. First, interfering with Patched1 ubiquitination blocks its

signal-dependent exit from cilia (Kim et al., 2015; Yue et al., 2014). Second, fusing ubiquitin to

PKD-2, the olfactory receptor ODR-10 or the TRP channel OSM-9 at their cytoplasmic end

results in the disappearance of these proteins from cilia (Hu et al., 2007; Xu et al., 2015). The

accumulation of these ubiquitin fusions inside cilia when BBSome function is compromised

suggests that the BBSome might sort ubiquitinated signaling receptors out of cilia, in line with a

reported association of BBSome with ubiquitinated proteins in trypanosomes (Langousis et al.,

2016). Interestingly, TRIM32 is a ubiquitin ligase mutated in BBS patients (Chiang et al., 2006).

Third, the ubiquitin ligase Cbl is recruited to cilia upon activation of the ciliary tyrosine kinase

receptor PDGFRαα and Cbl is required for termination of PDGFRαα signaling (Schmid et al.,

2018). Last, a dramatic rise in ubiquitination of the ciliary proteome is observed when

*Chlamydomonas* cilia disassemble (Huang et al., 2009). In particular, α-tubulin becomes

ubiquitinated on K304 upon cilia disassembly and expression of α-tubulin[K304R] slows down

cilia disassembly (Wang et al., 2019).

After conjugation onto substrates, ubiquitin itself often serves as a substrate for ubiquitination.

The resulting ubiquitin chains are characterized by the acceptor lysine residue on ubiquitin with

lysine 48-linked Ub (UbK48) chains targeting soluble proteins for proteasomal degradation while

lysine 63-linked Ub (UbK63) chains assembled onto membrane proteins constitute the major

driver for sorting from the limiting membrane of late endosomes into intralumenal vesicles and

ultimately lysosomal degradation (Piper et al., 2014; Yau and Rape, 2016; Swatek and

Komander, 2016). Intriguingly, in shortening *Chlamydomonas* cilia K63Ub linkages predominate

at least 5-fold over K48 linkages and only K63Ub linkages are detected on α-tubulin. A role for

K63Ub chains in ciliary trafficking remains to be determined.

80 Together, these data led us to investigate the interplay between ubiquitination, β-arrestin 2 and

81 the ciliary exit machinery. Using a combination of cellular imaging, biochemical studies and

82 cilia-specific manipulations, we find that β-arrestin 2 directs the activation-dependent addition of

83 K63-linked ubiquitin chains onto ciliary GPCRs that are then selected by the BBSome for

84 removal from cilia.

## RESULTS

## Signal-dependent ubiquitination of ciliary GPCRs is required for their regulated exit from cilia

As a first step in investigating the interplay between ubiquitination and the ciliary exit machinery, we sought to determine whether activation of ciliary GPCRs leads to their ubiquitination inside cilia prior to retrieval into the cell. We blocked BBSome-dependent exit by deleting its cognate GTPase ARL6/BBS3 in mouse inner medulla collecting duct (IMCD3) cells. IMCD3 cells represent a validated cell culture system to study ciliary trafficking and the IMCD3 $Arl6^{-/-}$ line has been previously characterized (Liew et al., 2014; Ye et al., 2018). As in previous studies (Ye et al., 2018; Nager et al., 2017), all GPCRs were expressed at near-endogenous levels by driving expression with extremely weak promoters. The fluorescent protein mNeonGreen (NG) fused to the cytoplasm-facing C-terminus of GPCRs allowed direct visualization. A biotinylation Acceptor Peptide (AP) fused to the extracellular N-terminus biotinylated by an ER-localized biotin ligase (BirA-ER) enabled pulse labeling of surface-exposed molecules and denaturing purifications. When SSTR3 was expressed in $Arl6^{-/-}$ IMCD3 cells, addition of its agonist somatostatin (sst) led to a drastic increase in the ciliary levels of ubiquitin observed upon immunostaining with the well-characterized FK2 (**Figure 1A-B)** and FK1 (**Figure S1A-B**) monoclonal antibodies (Fujimuro et al., 1994; Emmerich and Cohen, 2015; Haglund et al., 2003). All further experiments were conducted with the FK2 antibody. $Arl6^{-/-}$ cells accumulated Ub signals inside cilia in the absence of SSTR3 expression (**Figure 1A-B**), demonstrating that some endogenous proteins become ubiquitinated inside the cilia of $Arl6^{-/-}$ IMCD3 cells. No ubiquitin signal was detected inside cilia when BBSome function was intact (**Figure 1A-B and S1A-D**), indicating that the BBSome efficiently removes ubiquitinated proteins from cilia. Because the sst-dependent increase in ciliary ubiquitin signal depends upon SSTR3 expression (**Figure 1A-B**), these results suggest that either SSTR3 itself or a downstream effector of SSTR3 become ubiquitinated inside cilia upon activation.

To determine whether SSTR3 becomes ubiquitinated in response to sst, we biochemically

isolated $^{AP}$SSTR3$^{NG}$ under denaturing conditions via the biotinylated AP tag and probed for

ubiquitin via an HA tag on transfected HA-Ub. As typical for a glycosylated protein, SSTR3

migrated as a broad band centered around 100 kDa (**Figure 1C**). The amount of SSTR3

recovered did not change appreciably between WT and *Arl6$^{-/-}$* cells or upon stimulation with sst

(**Figure 1C**). Remarkably, while only very faint Ub signals were detected associated with SSTR3

in WT cells, a Ub smear extending from 100 kDa upward was detected in the SSTR3 pull-downs

from *Arl6$^{-/-}$* cells. Furthermore, stimulation with sst resulted in a modest but reproducible

increase in SSTR3 ubiquitination in *Arl6$^{-/-}$* cells (**Figure 1D**). We conclude that the

BBSome/ARL6 system is required for the degradation of ubiquitinated SSTR3 and that SSTR3

becomes ubiquitinated in response to stimulation with sst.

We next sought to test whether signaling receptors endogenous to IMCD3 cells are ubiquitinated

inside cilia in an activity-dependent manner. One signaling pathway that is natively expressed in

IMCD3 cells is the Hedgehog pathway. Prior experiments have detected normal trafficking

dynamics of SMO and GPR161 in IMCD3 cells (Ye et al., 2018; Mukhopadhyay et al., 2013)

and **Figure 4D**). As GPR161 and SMO both accumulate in cilia of *Arl6$^{-/-}$* cells without Hh

pathway stimulation (Liew et al., 2014; Zhang et al., 2011), the ubiquitin signal detected in *Arl6$^{-/-}$*

cilia may correspond to ubiquitin conjugated to SMO or GPR161. We detected a significant

elevation of the ciliary ubiquitin levels in *Arl6$^{-/-}$* cells when treated with the SMO agonist SAG

(**Figure 1E-F**). Because SMO ubiquitination was previously shown to decrease upon its

activation (Xia et al., 2012; Jiang et al., 2019), it is likely that the elevated ciliary ubiquitin signal

detected in SAG-treated cells is associated with GPR161. As activation of SMO triggers exit of

GPR161 but this exit is frustrated when BBSome function goes awry, these data suggest that

GPR161 becomes ubiquitinated prior to its BBSome-dependent exit. Collectively, these data

suggest that GPCRs are ubiquitinated in a regulated fashion inside cilia and subsequently

retrieved into the cell by the BBSome.

To test the functional importance of GPCR ubiquitination in BBSome-mediated exit from cilia,

we removed all ubiquitination sites from SSTR3 and GPR161 by mutating all cytoplasm-

exposed lysine residues to arginine (cK0 variants). Exit kinetics were precisely monitored by real-

time tracking of individual cilia in live cells. In both cases, the cK0 variant underwent markedly

slower signal-dependent exit from cilia than the WT allele (**Figure 2A-B**). Measurement of exit

rates revealed that exit of SSTR3cK0 was ~2-fold slower than SSTR3 and that exit of

GPR161cK0 was ~2.5-fold slower than GPR161. The residual exit rates of the cK0 mutants

indicate the existence of alternative mechanisms of exit that complement GPCR ubiquitination

or bypass BBSome-dependent retrieval (see discussion).

To rule out that the observed ciliary exit defect of the cK0 mutants was an indirect consequence

of defective endocytosis for example because of clogging of the ciliary exit path, we directly

assessed endocytosis of SSTR3 and SSTR3cK0. The surface-exposed pool of ^APSSTR3 was pulse

labeled with fluorescently labeled monovalent streptavidin (mSA) and cells were stimulated with

sst. The faint hazy signal corresponding to the plasma membrane pool of SSTR3 disappeared

after 10 min in the presence of sst in a similar fashion for both WT and cK0 variants (**Figure**

**2C**). Concurrently, the WT and cK0 variants appeared in cytoplasmic foci corresponding to

endocytic vesicles (**Figure 2C**). Counting cytoplasmic foci revealed that endocytosis proceeded

normally irrespective of the ubiquitination competence of SSTR3 (**Figure 2D**). SSTR3 thus

behaves similarly to nearly every GPCR tested to date, in that it does not require ubiquitination

for its signal-dependent internalization. Indeed, while ubiquitination of plasma membrane

proteins is a major driver of internalization in yeast, ubiquitination of signaling receptors is

generally dispensable for signal-dependent internalization in mammalian cells (Dores and Trejo,

2019; Piper et al., 2014; Skieterska et al., 2017).

These results strongly support a direct role for GPCR ubiquitination in promoting signal-

dependent exit from cilia.

**Ciliary K63 ubiquitin linkages are required for GPCR exit**

We next sought to characterize the type of ubiquitin conjugates that are attached to ciliary

proteins. The FK1 and FK2 antibodies used in our immunofluorescent studies recognize

ubiquitin either as single adduct or in chain but not free ubiquitin (Fujimuro et al., 1994;

Emmerich and Cohen, 2015). The availability of antibodies specific for K48 and K63 ubiquitin

linkages (Newton et al., 2008) enabled us to determine the types of ubiquitin conjugates that are

attached to ciliary GPCRs. While we found no detectable signal for K48-linked ubiquitin chains

(UbK48) in cilia, the signal for K63-linked ubiquitin (UbK63) linkages mirrored the ciliary

response observed with the FK1 and FK2 antibody in IMCD3-[SSTR3] cells subjected to

agonist treatment (**Figure 3A-B and S1C-D**). Given the high specificity and selectivity of the

K48 and K63 linkage antibodies, these data strongly suggest that K63Ub chains are added onto

SSTR3 inside cilia upon activation.

To confirm that K63 of Ub is the main linkage used for the elongation of Ub chains on SSTR3,

we transfected variants of ubiquitin with lysines mutated to arginines into $Arl6^{-/-}$ cells stably

expressing SSTR3. Signal-dependent ubiquitination of SSTR3 was abrogated when all seven

lysines of Ub were mutated to arginines (**Figure 3C-D**). However, when K63 was the only lysine

left intact on Ub, signal-dependent ubiquitination of SSTR3 was restored to the same extent as

with WT Ub (**Figure 3C-D**). We conclude that UbK63 chains are assembled onto SSTR3 in

response to SSTR3 activation inside cilia.

To determine the functional importance of UbK63 linkages in tagging GPCRs for exit from cilia,

we sought to specifically interfere with UbK63 linkages inside cilia (**Figure 4A**). Structural and

biochemical studies have demonstrated that the deubiquitinase AMSH possesses a near-absolute

specificity for UbK63 linkages and does not cleave other Ub chain linkages or Ub-substrates

linkages (Sato et al., 2008; McCullough et al., 2004). AMSH normally function on the surface of

late endosomes in concert with the endosomal sorting complex required for transport (ESCRT)

protein STAM (McCullough et al., 2006). AMSH was previously fused to the epidermal growth

factor receptor EGFR to demonstrate that UbK63 chains assembled on EGFR are dispensable

for internalization but required for sorting into lysosomes (Huang et al., 2013). We targeted

AMSH to cilia using the well-validated ciliary targeting signal from the ciliopathy protein

NPHP3 (Nakata et al., 2012; Wright et al., 2011) and analyzed the exit of three distinct GPCRs:

SMO, GPR161 and SSTR3. To alleviate concerns related to the interaction of AMSH with the

STAM, we deleted the STAM-interacting domain of AMSH and only expressed the catalytic

domain of AMSH. The signal-dependent exit of SSTR3 was assessed by treating IMCD3-

[SSTR3] cells with sst for 2h. In cells expressing cilia-AMSH (a fusion of NPHP3[1-200] to GFP

and the catalytic domain of AMSH), sst no longer triggered a significant reduction in the ciliary

levels of SSTR3 (**Figure 4B-C**). Care was taken to only analyze cells that expressed modest

levels of cilia-AMSH as judged by GFP fluorescence (**Figure S3**). In contrast, SSTR3 exit

proceeded normally in cells that expressed NPHP3[1-200]-GFP alone or a catalytically dead

variant of cilia-AMSH that we named cilia-ASMH[†] (**Figure 4B-C**). These controls strongly

suggest that cilia-AMSH exerts its effects on SSTR3 exit by cleaving UbK63 chains inside cilia, a

conclusion borne out by imaging ciliary UbK63 in cells transfected with cilia-AMSH (**Figure

S4**). Similarly, the exit of endogenous GPR161 from RPE cell cilia upon activation of the

Hedgehog pathway was abolished in cells expressing moderate levels of cilia-AMSH (**Figure

4D-E**). Neither NPHP3[1-200]-GFP nor cilia-AMSH[†] blocked the signal-dependent exit of

GPR161 from cilia (**Figure 4D-E**).

Unlike GPR161 and other ciliary GPCRs such as the Somatostatin receptor 3 (SSTR3) that are

removed from cilia by the BBSome only when they become activated, SMO undergoes

constitutive BBSome-dependent exit from cilia in the absence of Hedgehog pathway stimulation

(Nachury, 2018; Ye et al., 2018; Ocbina and Anderson, 2008; Zhang et al., 2011). Signal-

dependent accumulation of SMO in cilia is achieved, at least in part, by suppression of its exit

(Milenkovic et al., 2015; Nachury and Mick, 2019; Ye et al., 2018). In cells expressing cilia-

AMSH, endogenous SMO spontaneously accumulated inside cilia in the absence of Hedgehog

pathway activation (**Figure 4F-G**). Again, neither NPHP3[1-200]-GFP nor cilia-AMSH[†]

influenced the ciliary accumulation of SMO (**Figure 4F-G**).

Together, these results indicate that K63-linked ubiquitin chains inside cilia, presumably built

onto each ciliary GPCRs under a specific signaling stimulus, are required for GPCR removal

from cilia.

**β-arrestin 2 mediates signal-dependent ubiquitination of ciliary GPCRs**

As β-arrestin 2 is required for signal-dependent exit of GPR161 and SSTR3 from cilia (Green et al., 2015; Pal et al., 2016; Ye et al., 2018), we sought to determine the functional relationship between β-arrestin 2 and the BBSome. As previously described (Green et al., 2015), β-arrestin 2 is initially undetectable inside cilia and becomes recruited to cilia within minutes of SSTR3 agonist addition (**Figure 5A-C**). The ciliary β-arrestin 2 signal reaches a plateau after about 10 min and the $t_{1/2}$ for ciliary accumulation is less than 5 min (**Figure 5C**). Meanwhile the $t_{1/2}$ for BBSome recruitment to the tip of cilia was 10 min and the first signs of detectable SSTR3 exit were seen between 10 and 15 min (**Figure 5C**). These kinetics suggested that β-arrestin 2 is first recruited to activated ciliary SSTR3 before the BBSome ferries activated SSTR3 out of cilia. When ARL6 was depleted, the basal levels of ciliary β-arrestin 2 were elevated but the kinetics of β-arrestin 2 accumulation in cilia upon SSTR3 activation were indistinguishable from control-depleted cells (**Figure 5A-B**). These data suggest that a small fraction of SSTR3 that is tonically active fails to exit cilia in ARL6-depleted cells and recruits β-arrestin 2. From these data, we conclude that β-arrestin 2 functions either upstream of, or in parallel with the BBSome in signal-dependent retrieval of GPCRs. Similarly, the kinetics of GPR161 exit and β-arrestin 2 ciliary recruitment upon activation of the Hh pathway indicated that β-arrestin 2 reaches its maximal ciliary level before the first signs of GPR161 exit become evident (**Figure 5D**).

We previously proposed that β-arrestin 2 may bridge activated GPCRs to the BBSome (Ye et al., 2018; Nachury, 2018). Yet, we have thus far failed to detect a biochemical interaction between β-arrestin 2 and the BBSome. Furthermore, the continuous distribution patterns of β-arrestin 2 and SSTR3 inside cilia are similar to one another and distinct from the discontinuous pattern of the BBSome (**Figure 5A**), arguing against a direct association of β-arrestin 2 with BBSome/IFT trains inside cilia. We considered the alternative hypothesis that β-arrestin 2 function upstream of the BBSome. Given that β-arrestin 2 functions in the signal-dependent targeting of GPCRs from the plasma membrane to the lysosome by recruiting ubiquitin ligases to activated GPCRs (Henry et al., 2012; Bhandari et al., 2007; Shenoy et al., 2008; Martin et al., 2003), we posited that β-arrestin 2 recognizes activated GPCRs inside cilia to direct their ubiquitination and subsequent

selection by the BBSome for removal from cilia. A central prediction of this model is that β-
arrestin 2 is required for the ubiquitination of ciliary GPCRs in response to stimulation. To test
this prediction, we deleted the β-arrestin 2 gene *Arrb2* in *Arl6* knockout IMCD3 cells (Nager et
al., 2017). While the signal-dependent exit of SSTR3 from cilia failed in both *Arl6$^{-/-}$* and *Arl6$^{-/-}$*
*Arrb2$^{-/-}$* cells (**Figures 1A-B and 6A-B**), ubiquitin staining in cilia and by inference signal-
dependent ubiquitination of SSTR3 was only observed in *Arl6$^{-/-}$* cells (**Figures 6A-B**). In *Arl6$^{-/-}$*
*Arrb2$^{-/-}$* cells, the signal-dependent increase in ciliary ubiquitin signal was no longer observed
(**Figures 6A-B**). Similarly, the increase in ciliary ubiquitin signal seen in *Arl6$^{-/-}$* cells treated with
the Hedgehog pathway agonist SAG was no longer observed in the absence of β-arrestin 2
(**Figure 6C-D**). These data indicate that, in the absence of BBSome function, ciliary GPCRs
become ubiquitinated in response to activation in a β-arrestin 2-dependent manner. We sought
to confirm our imaging-based finding with a biochemical analysis of SSTR3 ubiquitination.
While SSTR3 ubiquitination was readily detected in *Arl6$^{-/-}$* cells and measurably increased upon
stimulation with sst, the deletion of β-arrestin 2 in *Arl6$^{-/-}$* cells drastically reduced SSTR3
ubiquitination levels and no sst-dependent increase of SSTR3 ubiquitination was detected in
*Arl6$^{-/-}$ /Arrb2$^{-/-}$* cells (**Figure 6E-F**).

Together, these data suggest the following order of action: GPCR activation, β-arrestin 2
engagement, Ub ligase recruitment, assembly of K63Ub chains on the GPCR, and finally
selection of ubiquitinated GPCRs for BBSome-mediated retrieval (**Figure 7E**).

### Constitutively retrieved BBSome cargoes are ubiquitinated prior to exit

Besides removing GPCRs from cilia in a regulated fashion, the BBSome also clears cilia of
unwanted proteins that accidentally enter cilia. In the single-cell flagellated organism
*Chlamydomonas reinhardtii*, mutations in the BBSome subunit Bbs4 cause a constitutive
accumulation of several proteins including Phospholipase D in cilia (Lechtreck et al., 2009).
When stained for ubiquitin, *Bbs4 Chlamydomonas* showed a 2-fold enrichment in ciliary signal
compared to WT (**Fig. 7A-B**). Isolation of cilia revealed the accumulation of ubiquitin

conjugates above 80 kDa in *Bbs4* mutant *Chlamydomonas* cilia (**Fig. 7C**). These results are

consistent with the hypothesis that ciliary proteins subject to constitutive retrieval are

ubiquitinated prior to their removal from cilia by the BBSome.

In the mammalian retina, proteomics studies have found over 100 proteins accumulating in the

outer segment (the equivalent of the cilium) of *Bbs* photoreceptors compared to WT

photoreceptors (Datta et al., 2015). When we stained sections of mouse retina for ubiquitin, we

found a drastic accumulation of ubiquitin signal in the outer segment of *Bbs4$^{-/-}$* photoreceptors

compared to WT photoreceptors (**Fig. 7D**). We note that the nature of ubiquitin conjugates

remains to be determined in the case of constitutive cargoes. These results suggest that a variety

of proteins that accidentally flow into the photoreceptors outer segments –possibly caught on the

tremendous flux of rhodopsin trafficking to the outer segment (Baker et al., 2008)– are recognized

as foreign to the cilium, ubiquitinated on site and selected for removal by the BBSome (**Figure**

**7E**).

**DISCUSSION**

**Recognition of K63-linked ubiquitin chains by the ciliary exit machinery**

Our results indicate that K63-linked ubiquitin chains tag activated GPCRs and likely other

unwanted proteins for BBSome-mediated removal from cilia. Prior findings in trypanosomes

suggest that the BBSome may directly recognize ubiquitin (Langousis et al., 2016), although

direct biochemical evidence and information regarding ubiquitin chain specificity is still missing.

It is also known that the BBSome directly recognizes cytoplasmic determinants in the GPCRs it

ferries out of cilia (Jin et al., 2010; Klink et al., 2017; Ye et al., 2018; Seo et al., 2011; Domire et

al., 2011). We propose that the summation of weak molecular interactions between the BBSome

and UbK63 chains on one hand and the BBSome and GPCRs cytoplasmic loops on the other

hand increases the BBSome-cargo interaction to enable sorting of ubiquitinated GPCRs out of

cilia. In support of this bivalent recognition of ubiquitinated cargoes by the BBSome, the affinity

of the BBSome for cytoplasmic determinant of GPCRs are in the µM range (Klink et al., 2017)

and affinities of ubiquitin-binding proteins for ubiquitin typically reside in the sub mM range

(Husnjak and Dikic, 2012). The nature of the biochemical entity that directly recognizes

ubiquitin, and in particular UbK63 chains remains to be determined: while the BBSome was

pulled down on Ub-agarose resin from trypanosome lysates (Langousis et al., 2016), no known

Ub-binding domain is found in the BBSome polypeptides. Furthermore, the IFT-A subunit IFT-

139 associates with ubiquitinated α-tubulin in ciliary extracts from *Chlamydomonas*, suggesting a

possible recognition of ubiquitin by IFT139 (Wang et al., 2019). Again, the absence of known

Ub-binding domain in the sequence of IFT139 presents a challenge to this hypothesis.

**Requirements for K63-linked ubiquitination in ciliary exit and sorting to the**

**lysosome**

Wang et al. (2019) found that α-tubulin is modified by UbK63 chains and not UbK48 chain

upon ciliary disassembly. The ubiquitin ligases that have been associated with ciliary clearance of

Patched 1 and of PDGFRαα specifically assemble UbK63 chains (Kim et al., 2015; Yue et al.,

2014; Schmid et al., 2018). The suppression of GPR161, SSTR3 and SMO exit by cilia-AMSH

(**Figure 4**) demonstrates that UbK63 chains are recognized inside cilia to direct trafficking out of

cilia. The milder exit defect observed when acceptor lysines are mutated on ciliary GPCRs

(**Figure 2**) than when cilia-AMSH is expressed (**Figure 4**) suggests that K63Ub chains added

onto other ciliary proteins besides GPCRs may participate in ciliary exit. A role for

ubiquitination of the ciliary transport machinery in exit is in line with the functional importance

of ubiquitination of the endolysosomal machinery in GPCR trafficking (Dores and Trejo, 2019).

β arrestin ubiquitination promotes internalization of the β2-adrenergic receptor (β2AR) (Shenoy

et al., 2001, 2007; Shenoy and Lefkowitz, 2003), ubiquitination of the ESCRT components Hrs

and STAM participates in sorting of the chemokine receptor CXCR4 into the MVB (Malik and

Marchese, 2010; Marchese et al., 2003) and ubiquitination of the ESCRT-III recruitment factor

ALIX enhances MVB sorting of the protease activated receptor PAR-1 (Dores et al., 2015).

The requirement for UbK63 chains in signal-dependent exit from the cilium contrasts with the

more limited role of ubiquitin and UbK63 chains in signal-dependent endocytosis of signaling

receptors at the plasma membrane in mammals. Although ubiquitination is a major driver of

endocytosis in yeast (Stringer and Piper, 2011; Piper et al., 2014), foundational studies on

epidermal growth factor receptor (EGFR) trafficking found that signal-dependent receptor

ubiquitination is not essential for internalization because it is redundant with other

internalization mechanisms (Goh et al., 2010; Huang et al., 2013; Fortian et al., 2015). Studies of

an EGFR-AMSH fusion found that UbK63 chains attached onto EGFR become increasingly

necessary as EGFR progresses along the degradative route and a strict requirement for K63-

linked ubiquitin chains is observed at the terminal step of EGFR sorting into the lumen of

multivesicular body (MVB) (Huang et al., 2013). A strict requirement for receptor ubiquitination

in degradative sorting but not in internalization can be generalized to nearly all GPCRs

(Skieterska et al., 2017), namely β2AR (Shenoy et al., 2001), CXCR4 (Marchese and Benovic,

2001), μ opioid receptor (Hislop et al., 2011), κ opioid receptor (Li et al., 2008), δ opioid receptor

(Henry et al., 2011), neurokinin-1 receptor (Cottrell et al., 2006), protease activated receptor 2

(PAR-2) (Jacob et al., 2005) and vasopressin receptor 2 (V2R) (Martin et al., 2003). Similarly, we

find that SSTR3 endocytosis proceeds normally in the absence of SSTR3 ubiquitination (**Figure**
**2C-D**). Studying the importance of ubiquitin K63 in mammalian cells via genetics remains
challenging because of the multiple genes encoding Ub but in yeast strains expressing UbK63R
as the sole source of Ub, the Gap1 permease is internalized normally and fails to get sorted into
the MVB (Lauwers et al., 2009).

The parallels between sorting at the late endosome and ciliary trafficking extend to β arrestin-
directed ubiquitination. While β arrestin functions inside cilia to direct the addition of ubiquitin
onto activated SSTR3 and GPR161 (**Figures 5 and 6**), β arrestin-mediated ubiquitination
selectively affects sorting at the level of endosomes rather that at the plasma membrane for
activated CXCR4 (Bhandari et al., 2007), $\beta_2AR$ (Shenoy et al., 2008) and V2R (Martin et al.,
2003) and β arrestin-mediated ubiquitination of CXCR4 was shown to take place on endosomes
(Malik and Marchese, 2010). In this context, the recognition of K63-linked ubiquitin chain inside
cilia may befit the endosomal origin of primary cilia (Sorokin, 1962; Westlake et al., 2011;
Mitchell, 2017).

The parallels between endosomal and ciliary trafficking lead us to speculate that some
ubiquitinated proteins might be accidentally sorted to cilia instead of late endosomes. These
proteins would then need to be retrieved from cilia by the BBSome. Thus, the Ub signal detected
in photoreceptor outer segment and in *Chlamydomonas* flagella might result from the accidental
import of ubiquitinated proteins into cilia rather than *in situ* ubiquitination of unwanted proteins
inside cilia.

**Potential coupling between BBSome-mediated exit and degradative sorting**

An interesting consideration is the relationship between ciliary exit and endo-lysosomal sorting. It
has been proposed that endocytosis of signaling receptors is intimately coupled to their exit from
cilia (Pedersen et al., 2016). However, single molecule imaging of GPR161 exiting cilia suggests
that activated GPR161 diffuses into the plasma membrane after it exits from the ciliary
compartment (Ye et al., 2018). Prior findings that worm mutants for BBSome or the ESCRT

machinery both fail to remove ciliary proteins fused to Ub from cilia (Hu et al., 2007; Xu et al., 2015) suggest the provocative possibility that ciliary exit may be tightly coupled to degradative sorting. Nonetheless, the hypothesis that interfering with UbK63 chain formation on ciliary GPCRs indirectly blocks ciliary exit because of a primary defect in endocytosis can be rejected because ubiquitination is not a major determinant of SSTR3 endocytosis and because cilia-AMSH potently and specifically blocks the signal-dependent exit of ciliary GPCRs.

In this context, the contrast between the dramatic increase in ubiquitinated SSTR3 levels in $Arl6^{-/-}$ cells compared to WT cells even in the absence of stimulation (**Figure 1C-D**) and the modest increase in ciliary Ub levels in $Arl6^{-/-}$ cells compared to WT cells in the absence of stimulation (**Figure 1A-B**) suggests that a considerable amount of ubiquitinated SSTR3 accumulates outside of cilia in $Arl6^{-/-}$ cells. One interpretation consistent with a direct coupling between ciliary exit and degradation is that a small fraction of ubiquitinated SSTR3 escapes cilia in $Arl6^{-/-}$ cells but fails to reach to correct degradative route. In this interpretation, the BBSome would deliver its cargoes to the ESCRT machinery for routing into the lysosomal degradative pathway.

**Ubiquitination and control of the ciliary proteome**

A role for ubiquitination in selecting unwanted proteins for removal from cilia may unify the functions of the BBSome in signal-dependent retrieval of GPCRs and the constitutive clearance of proteins that accidentally enter cilia. In both of these cases, unwanted ciliary proteins need to be recognized as 'non-self' by a ciliary surveillance machinery. For GPCRs, β arrestin orchestrate the recognition of activated GPCRs as unwanted by directing their ubiquitination. A fascinating question for future investigations is how the more than 100 proteins that accumulate in outer segments of *Bbs* photoreceptors are recognized as 'non-self' and tagged with ubiquitin.

**REFERENCES**

Anvarian, Z., K. Mykytyn, S. Mukhopadhyay, L.B. Pedersen, and S.T. Christensen. 2019.
Cellular signalling by primary cilia in development, organ function and disease. *Nat. Rev.*
*Nephrol.* doi:10.1038/s41581-019-0116-9.

Baker, S.A., M. Haeri, P. Yoo, S.M. Gospe, N.P. Skiba, B.E. Knox, and V.Y. Arshavsky. 2008.
The outer segment serves as a default destination for the trafficking of membrane proteins
in photoreceptors. *J. Cell Biol.* 183:485–498. doi:10.1083/jcb.200806009.

Bhandari, D., J. Trejo, J.L. Benovic, and A. Marchese. 2007. Arrestin-2 interacts with the
ubiquitin-protein isopeptide ligase atrophin-interacting protein 4 and mediates
endosomal sorting of the chemokine receptor CXCR4. *J. Biol. Chem.* 282:36971–36979.
doi:10.1074/jbc.M705085200.

Chiang, A.P., J.S. Beck, H. Yen, M.K. Tayeh, T.E. Scheetz, R.E. Swiderski, D.Y. Nishimura,
T.A. Braun, K.-Y.A. Kim, J. Huang, K. Elbedour, R. Carmi, D.C. Slusarski, T.L.
Casavant, E.M. Stone, and V.C. Sheffield. 2006. Homozygosity mapping with SNP
arrays identifies TRIM32, an E3 ubiquitin ligase, as a Bardet-Biedl syndrome gene
(BBS11). *Proc. Natl. Acad. Sci. U. S. A.* 103:6287–6292. doi:10.1073/pnas.0600158103.

Corbit, K.C., P. Aanstad, V. Singla, A.R. Norman, D.Y.R. Stainier, and J.F. Reiter. 2005.
Vertebrate Smoothened functions at the primary cilium. *Nature.* 437:1018–1021.
doi:10.1038/nature04117.

Cottrell, G.S., B. Padilla, S. Pikios, D. Roosterman, M. Steinhoff, D. Gehringer, E.F. Grady, and
N.W. Bunnett. 2006. Ubiquitin-dependent Down-regulation of the Neurokinin-1
Receptor. *J. Biol. Chem.* 281:27773–27783. doi:10.1074/jbc.M603369200.

Craige, B., J.M. Brown, and G.B. Witman. 2013. Isolation of Chlamydomonas Flagella. *Curr.*
*Protoc. Cell Biol.* 59:3.41.1-3.41.9. doi:10.1002/0471143030.cb0341s59.

Datta, P., C. Allamargot, J.S. Hudson, E.K. Andersen, S. Bhattarai, A.V. Drack, V.C. Sheffield,
and S. Seo. 2015. Accumulation of non-outer segment proteins in the outer segment
underlies photoreceptor degeneration in Bardet-Biedl syndrome. *Proc Natl Acad Sci U A.*
112:E4400–9. doi:10.1073/pnas.1510111112.

Domire, J.S., J.A. Green, K.G. Lee, A.D. Johnson, C.C. Askwith, and K. Mykytyn. 2011.
Dopamine receptor 1 localizes to neuronal cilia in a dynamic process that requires the
Bardet-Biedl syndrome proteins. *Cell. Mol. Life Sci. CMLS.* 68:2951–2960.
doi:10.1007/s00018-010-0603-4.

Dores, M.R., H. Lin, N. J Grimsey, F. Mendez, and J. Trejo. 2015. The α-arrestin ARRDC3
mediates ALIX ubiquitination and G protein-coupled receptor lysosomal sorting. *Mol.*
*Biol. Cell*. 26:4660–4673. doi:10.1091/mbc.E15-05-0284.

Dores, M.R., and J. Trejo. 2019. Endo-lysosomal sorting of G-protein-coupled receptors by
ubiquitin: Diverse pathways for G-protein-coupled receptor destruction and beyond.
*Traffic Cph. Den.* 20:101–109. doi:10.1111/tra.12619.

Emmerich, C.H., and P. Cohen. 2015. Optimising methods for the preservation, capture and
identification of ubiquitin chains and ubiquitylated proteins by immunoblotting. *Biochem.*
*Biophys. Res. Commun.* 466:1–14. doi:10.1016/j.bbrc.2015.08.109.

Fortian, A., L.K. Dionne, S.H. Hong, W. Kim, S.P. Gygi, S.C. Watkins, and A. Sorkin. 2015.
Endocytosis of Ubiquitylation-Deficient EGFR Mutants via Clathrin-Coated Pits is
Mediated by Ubiquitylation. *Traffic Cph. Den.* 16:1137–1154. doi:10.1111/tra.12314.

Fujimuro, M., H. Sawada, and H. Yokosawa. 1994. Production and characterization of
monoclonal antibodies specific to multi-ubiquitin chains of polyubiquitinated proteins.
*FEBS Lett.* 349:173–180. doi:10.1016/0014-5793(94)00647-4.

Gigante, E.D., and T. Caspary. 2020. Signaling in the primary cilium through the lens of the
Hedgehog pathway. *Wiley Interdiscip. Rev. Dev. Biol.* e377. doi:10.1002/wdev.377.

Goh, L.K., F. Huang, W. Kim, S. Gygi, and A. Sorkin. 2010. Multiple mechanisms collectively
regulate clathrin-mediated endocytosis of the epidermal growth factor receptor. *J. Cell*
*Biol.* 189:871–883. doi:10.1083/jcb.201001008.

Gorman, D.S., and R.P. Levine. 1965. Cytochrome f and plastocyanin: their sequence in the
photosynthetic electron transport chain of Chlamydomonas reinhardi. *Proc. Natl. Acad. Sci.*
*U. S. A.* 54:1665–1669. doi:10.1073/pnas.54.6.1665.

Green, J.A., C.L. Schmid, E. Bley, P.C. Monsma, A. Brown, L.M. Bohn, and K. Mykytyn. 2015.
Recruitment of β-Arrestin into Neuronal Cilia Modulates Somatostatin Receptor
Subtype 3 Ciliary Localization. *Mol. Cell. Biol.* 36:223–235. doi:10.1128/MCB.00765-15.

Haglund, K., S. Sigismund, S. Polo, I. Szymkiewicz, P.P. Di Fiore, and I. Dikic. 2003. Multiple
monoubiquitination of RTKs is sufficient for their endocytosis and degradation. *Nat. Cell*
*Biol.* 5:461–466. doi:10.1038/ncb983.

Henry, A.G., J.N. Hislop, J. Grove, K. Thorn, M. Marsh, and M. von Zastrow. 2012.
Regulation of endocytic clathrin dynamics by cargo ubiquitination. *Dev. Cell*. 23:519–532.
doi:10.1016/j.devcel.2012.08.003.

Henry, A.G., I.J. White, M. Marsh, M. von Zastrow, and J.N. Hislop. 2011. The Role of
Ubiquitination in Lysosomal Trafficking of δ-Opioid Receptors. *Traffic*. 12:170–184.
doi:10.1111/j.1600-0854.2010.01145.x.

Hislop, J.N., A.G. Henry, and M. von Zastrow. 2011. Ubiquitination in the first cytoplasmic loop
of μ-opioid receptors reveals a hierarchical mechanism of lysosomal down-regulation. *J.*
*Biol. Chem.* 286:40193–40204. doi:10.1074/jbc.M111.288555.

Howarth, M., and A.Y. Ting. 2008. Imaging proteins in live mammalian cells with biotin ligase
and monovalent streptavidin. *Nat. Protoc.* 3:534–545. doi:10.1038/nprot.2008.20.

Hu, J., S.G. Wittekind, and M.M. Barr. 2007. STAM and Hrs down-regulate ciliary TRP
receptors. *Mol. Biol. Cell.* 18:3277–3289. doi:10.1091/mbc.E07-03-0239.

Huang, F., X. Zeng, W. Kim, M. Balasubramani, A. Fortian, S.P. Gygi, N.A. Yates, and A.
Sorkin. 2013. Lysine 63-linked polyubiquitination is required for EGF receptor
degradation. *Proc. Natl. Acad. Sci.* 110:15722–15727. doi:10.1073/pnas.1308014110.

Huang, K., D.R. Diener, and J.L. Rosenbaum. 2009. The ubiquitin conjugation system is
involved in the disassembly of cilia and flagella. *J. Cell Biol.* 186:601–613.
doi:10.1083/jcb.200903066.

Husnjak, K., and I. Dikic. 2012. Ubiquitin-binding proteins: decoders of ubiquitin-mediated
cellular functions. *Annu. Rev. Biochem.* 81:291–322. doi:10.1146/annurev-biochem-
051810-094654.

Jacob, C., G.S. Cottrell, D. Gehringer, F. Schmidlin, E.F. Grady, and N.W. Bunnett. 2005. c-
Cbl mediates ubiquitination, degradation, and down-regulation of human protease-
activated receptor 2. *J. Biol. Chem.* 280:16076–16087. doi:10.1074/jbc.M500109200.

Jiang, W., X. Yao, Z. Shan, W. Li, Y. Gao, and Q. Zhang. 2019. E3 ligase Herc4 regulates
Hedgehog signalling through promoting Smoothened degradation. *J. Mol. Cell Biol.*
11:791–803. doi:10.1093/jmcb/mjz024.

Jin, H., S.R. White, T. Shida, S. Schulz, M. Aguiar, S.P. Gygi, J.F. Bazan, and M.V. Nachury.
2010. The conserved Bardet-Biedl syndrome proteins assemble a coat that traffics
membrane proteins to cilia. *Cell.* 141:1208–1219. doi:10.1016/j.cell.2010.05.015.

Kim, J., E.Y. C Hsia, A. Brigui, A. Plessis, P.A. Beachy, and X. Zheng. 2015. The role of ciliary
trafficking in Hedgehog receptor signaling. *Sci. Signal.* 8:ra55–ra55.
doi:10.1126/scisignal.aaa5622.

Klink, B.U., E. Zent, P. Juneja, A. Kuhlee, S. Raunser, and A. Wittinghofer. 2017. A
recombinant BBSome core complex and how it interacts with ciliary cargo. *eLife*. 6.
doi:10.7554/eLife.27434.

Langousis, G., M.M. Shimogawa, E.A. Saada, A.A. Vashisht, R. Spreafico, A.R. Nager, W.D.
Barshop, M.V. Nachury, J.A. Wohlschlegel, and K.L. Hill. 2016. Loss of the BBSome
perturbs endocytic trafficking and disrupts virulence of Trypanosoma brucei. *Proc Natl*
*Acad Sci U A*. 113:632–637. doi:10.1073/pnas.1518079113.

Lauwers, E., C. Jacob, and B. André. 2009. K63-linked ubiquitin chains as a specific signal for
protein sorting into the multivesicular body pathway. *J. Cell Biol.* 185:493–502.
doi:10.1083/jcb.200810114.

Lechtreck, K.-F., E.C. Johnson, T. Sakai, D. Cochran, B.A. Ballif, J. Rush, G.J. Pazour, M.
Ikebe, and G.B. Witman. 2009. The Chlamydomonas reinhardtii BBSome is an IFT
cargo required for export of specific signaling proteins from flagella. *J. Cell Biol.*
187:1117–1132. doi:10.1083/jcb.200909183.

Li, J.-G., D.S. Haines, and L.-Y. Liu-Chen. 2008. Agonist-Promoted Lys63-Linked
Polyubiquitination of the Human κ-Opioid Receptor Is Involved in Receptor Down-
Regulation. *Mol. Pharmacol.* 73:1319–1330. doi:10.1124/mol.107.042846.

Liew, G.M., F. Ye, A.R. Nager, J.P. Murphy, J.S. Lee, M. Aguiar, D. Breslow, S.P. Gygi, and
M.V. Nachury. 2014. The intraflagellar transport protein IFT27 promotes BBSome exit
from cilia through the GTPase ARL6/BBS3. *Dev. Cell.* 31:265–278.
doi:10.1016/j.devcel.2014.09.004.

Loktev, A.V., and P.K. Jackson. 2013. Neuropeptide Y family receptors traffic via the Bardet-
Biedl syndrome pathway to signal in neuronal primary cilia. *Cell Rep.* 5:1316–1329.
doi:10.1016/j.celrep.2013.11.011.

Malik, R., and A. Marchese. 2010. Arrestin-2 interacts with the endosomal sorting complex
required for transport machinery to modulate endosomal sorting of CXCR4. *Mol. Biol.*
*Cell.* 21:2529–2541. doi:10.1091/mbc.e10-02-0169.

Marchese, A., and J.L. Benovic. 2001. Agonist-promoted ubiquitination of the G protein-coupled
receptor CXCR4 mediates lysosomal sorting. *J. Biol. Chem.* 276:45509–45512.
doi:10.1074/jbc.C100527200.

Marchese, A., C. Raiborg, F. Santini, J.H. Keen, H. Stenmark, and J.L. Benovic. 2003. The E3
ubiquitin ligase AIP4 mediates ubiquitination and sorting of the G protein-coupled
receptor CXCR4. *Dev. Cell.* 5:709–722. doi:10.1016/s1534-5807(03)00321-6.

Martin, N.P., R.J. Lefkowitz, and S.K. Shenoy. 2003. Regulation of V2 Vasopressin Receptor
Degradation by Agonist-promoted Ubiquitination. *J. Biol. Chem.* 278:45954–45959.
doi:10.1074/jbc.M308285200.

McCullough, J., M.J. Clague, and S. Urbé. 2004. AMSH is an endosome-associated ubiquitin
isopeptidase. *J. Cell Biol.* 166:487–492. doi:10.1083/jcb.200401141.

McCullough, J., P.E. Row, Ó. Lorenzo, M. Doherty, R. Beynon, M.J. Clague, and S. Urbé.
2006. Activation of the Endosome-Associated Ubiquitin Isopeptidase AMSH by STAM,
a Component of the Multivesicular Body-Sorting Machinery. *Curr. Biol.* 16:160–165.
doi:10.1016/j.cub.2005.11.073.

Michel, M.A., P.R. Elliott, K.N. Swatek, M. Simicek, J.N. Pruneda, J.L. Wagstaff, S.M.V.
Freund, and D. Komander. 2015. Assembly and specific recognition of k29- and k33-
linked polyubiquitin. *Mol. Cell.* 58:95–109. doi:10.1016/j.molcel.2015.01.042.

Milenkovic, L., L.E. Weiss, J. Yoon, T.L. Roth, Y.S. Su, S.J. Sahl, M.P. Scott, and W.E.
Moerner. 2015. Single-molecule imaging of Hedgehog pathway protein Smoothened in
primary cilia reveals binding events regulated by Patched1. *Proc Natl Acad Sci U A*.
112:8320–8325. doi:10.1073/pnas.1510094112.

Mitchell, D.R. 2017. Evolution of Cilia. *Cold Spring Harb. Perspect. Biol.* 9:a028290.
doi:10.1101/cshperspect.a028290.

Mukhopadhyay, S., X. Wen, N. Ratti, A. Loktev, L. Rangell, S.J. Scales, and P.K. Jackson.
2013. The ciliary G-protein-coupled receptor Gpr161 negatively regulates the Sonic
hedgehog pathway via cAMP signaling. *Cell.* 152:210–223.
doi:10.1016/j.cell.2012.12.026.

Mykytyn, K., and C. Askwith. 2017. G-Protein-Coupled Receptor Signaling in Cilia. *Cold Spring*
*Harb. Perspect. Biol.* a028183. doi:10.1101/cshperspect.a028183.

Mykytyn, K., R.F. Mullins, R.F. Mullins, M. Andrews, M. Andrews, A.P. Chiang, A.P. Chiang,
R.E. Swiderski, R.E. Swiderski, B. Yang, B. Yang, T. Braun, T. Braun, T. Casavant, T.
Casavant, E.M. Stone, E.M. Stone, and V.C. Sheffield. 2004. Bardet-Biedl syndrome
type 4 (BBS4)-null mice implicate Bbs4 in flagella formation but not global cilia assembly.
*Proc. Natl. Acad. Sci. U. S. A.* 101:8664–8669. doi:10.1073/pnas.0402354101.

Nachury, M.V. 2018. The molecular machines that traffic signaling receptors into and out of
cilia. *Curr. Opin. Cell Biol.* 51:124–131. doi:10.1016/j.ceb.2018.03.004.

Nachury, M.V., and D.U. Mick. 2019. Establishing and regulating the composition of cilia for
signal transduction. *Nat. Rev. Mol. Cell Biol.* 20:389–405. doi:10.1038/s41580-019-0116-4.

Nager, A.R., J.S. Goldstein, V. Herranz-Pérez, D. Portran, F. Ye, J.M. García-Verdugo, and
M.V. Nachury. 2017. An Actin Network Dispatches Ciliary GPCRs into Extracellular
Vesicles to Modulate Signaling. *Cell*. 168:252–263.e14. doi:10.1016/j.cell.2016.11.036.

Nakata, K., D. Shiba, D. Kobayashi, and T. Yokoyama. 2012. Targeting of Nphp3 to the
primary cilia is controlled by an N-terminal myristoylation site and coiled-coil domains.
*Cytoskelet. Hoboken NJ*. 69:221–234. doi:10.1002/cm.21014.

Newton, K., M.L. Matsumoto, I.E. Wertz, D.S. Kirkpatrick, J.R. Lill, J. Tan, D. Dugger, N.
Gordon, S.S. Sidhu, F.A. Fellouse, L. Komuves, D.M. French, R.E. Ferrando, C. Lam,
D. Compaan, C. Yu, I. Bosanac, S.G. Hymowitz, R.F. Kelley, and V.M. Dixit. 2008.
Ubiquitin chain editing revealed by polyubiquitin linkage-specific antibodies. *Cell*.
134:668–678. doi:10.1016/j.cell.2008.07.039.

Ocbina, P.J.R., and K.V. Anderson. 2008. Intraflagellar transport, cilia, and mammalian
Hedgehog signaling: analysis in mouse embryonic fibroblasts. *Dev. Dyn.* 237:2030–2038.
doi:10.1002/dvdy.21551.

Pal, K., S. Hwang, B. Somatilaka, H. Badgandi, P.K. Jackson, K. DeFea, and S.
Mukhopadhyay. 2016. Smoothened determines β-arrestin-mediated removal of the G
protein-coupled receptor Gpr161 from the primary cilium. *J. Cell Biol.* 212:861–875.
doi:10.1083/jcb.201506132.

Pedersen, L.B., J.B. Mogensen, and S.T. Christensen. 2016. Endocytic Control of Cellular
Signaling at the Primary Cilium. *Trends Biochem. Sci.* 41:784–797.
doi:10.1016/j.tibs.2016.06.002.

Piper, R.C., I. Dikic, and G.L. Lukacs. 2014. Ubiquitin-Dependent Sorting in Endocytosis. *Cold
Spring Harb. Perspect. Biol.* 6:a016808–a016808. doi:10.1101/cshperspect.a016808.

Postma, M., and J. Goedhart. 2019. PlotsOfData-A web app for visualizing data together with
their summaries. *PLoS Biol.* 17:e3000202. doi:10.1371/journal.pbio.3000202.

Rohatgi, R., L. Milenkovic, and M.P. Scott. 2007. Patched1 regulates hedgehog signaling at the
primary cilium. *Sci. N. Y. NY*. 317:372–376. doi:10.1126/science.1139740.

Sato, Y., A. Yoshikawa, A. Yamagata, H. Mimura, M. Yamashita, K. Ookata, O. Nureki, K.
Iwai, M. Komada, and S. Fukai. 2008. Structural basis for specific cleavage of Lys 63-
linked polyubiquitin chains. *Nature*. 455:358–362. doi:10.1038/nature07254.

Schmid, F.M., K.B. Schou, M.J. Vilhelm, M.S. Holm, L. Breslin, P. Farinelli, L.A. Larsen, J.S.
Andersen, L.B. Pedersen, and S.T. Christensen. 2018. IFT20 modulates ciliary PDGFRα
signaling by regulating the stability of Cbl E3 ubiquitin ligases. *J. Cell Biol.* 217:151–161.
doi:10.1083/jcb.201611050.

Seo, S., Q. Zhang, K. Bugge, D. Breslow, C.C. Searby, M.V. Nachury, and V.C. Sheffield.
2011. A novel protein LZTFL1 regulates ciliary trafficking of the BBSome and
Smoothened. *PLoS Genet.* 7:e1002358. doi:10.1371/journal.pgen.1002358.

Shaner, N.C., G.G. Lambert, A. Chammas, Y. Ni, P.J. Cranfill, M.A. Baird, B.R. Sell, J.R.
Allen, R.N. Day, M. Israelsson, M.W. Davidson, and J. Wang. 2013. A bright
monomeric green fluorescent protein derived from Branchiostoma lanceolatum. *Nat.*
*Methods*. 10:407–409. doi:10.1038/nmeth.2413.

Shankar, H., H. Shankar, A. Michal, A. Michal, R.C. Kern, R.C. Kern, D.S. Kang, D.S. Kang,
V.V. Gurevich, V.V. Gurevich, and J.L. Benovic. 2010. Non-visual arrestins are
constitutively associated with the centrosome and regulate centrosome function. *J. Biol.*
*Chem.* 285:8316–8329. doi:10.1074/jbc.M109.062521.

Shearer, R.F., and D.N. Saunders. 2016. Regulation of primary cilia formation by the ubiquitin-
proteasome system. *Biochem. Soc. Trans.* 44:1265–1271. doi:10.1042/BST20160174.

Shenoy, S.K., L.S. Barak, K. Xiao, S. Ahn, M. Berthouze, A.K. Shukla, L.M. Luttrell, and R.J.
Lefkowitz. 2007. Ubiquitination of beta-arrestin links seven-transmembrane receptor
endocytosis and ERK activation. *J. Biol. Chem.* 282:29549–29562.
doi:10.1074/jbc.M700852200.

Shenoy, S.K., and R.J. Lefkowitz. 2003. Trafficking Patterns of β-Arrestin and G Protein-
coupled Receptors Determined by the Kinetics of β-Arrestin Deubiquitination. *J. Biol.*
*Chem.* 278:14498–14506. doi:10.1074/jbc.M209626200.

Shenoy, S.K., P.H. McDonald, T.A. Kohout, and R.J. Lefkowitz. 2001. Regulation of receptor
fate by ubiquitination of activated beta 2-adrenergic receptor and beta-arrestin. *Science.*
294:1307–1313. doi:10.1126/science.1063866.

Shenoy, S.K., K. Xiao, V. Venkataramanan, P.M. Snyder, N.J. Freedman, and A.M. Weissman.
2008. Nedd4 mediates agonist-dependent ubiquitination, lysosomal targeting, and
degradation of the beta2-adrenergic receptor. *J. Biol. Chem.* 283:22166–22176.
doi:10.1074/jbc.M709668200.

Skieterska, K., P. Rondou, and K. Van Craenenbroeck. 2017. Regulation of G Protein-Coupled
Receptors by Ubiquitination. *Int. J. Mol. Sci.* 18. doi:10.3390/ijms18050923.

Sorokin, S. 1962. Centrioles and the formation of rudimentary cilia by fibroblasts and smooth
muscle cells. *J. Cell Biol.* 15:363–377. doi:10.1083/jcb.15.2.363.

Stringer, D.K., and R.C. Piper. 2011. A single ubiquitin is sufficient for cargo protein entry into
MVBs in the absence of ESCRT ubiquitination. *J. Cell Biol.* 192:229–242.
doi:10.1083/jcb.201008121.

Swatek, K.N., and D. Komander. 2016. Ubiquitin modifications. *Cell Res.* 26:399–422.
625             doi:10.1038/cr.2016.39.

626    Wang, Q., Z. Peng, H. Long, X. Deng, and K. Huang. 2019. Polyubiquitylation of α-tubulin at
627             K304 is required for flagellar disassembly in Chlamydomonas. *J. Cell Sci.* 132.
628             doi:10.1242/jcs.229047.

629    Westlake, C.J., L.M. Baye, M.V. Nachury, K.J. Wright, K.E. Ervin, L. Phu, C. Chalouni, J.S.
630             Beck, D.S. Kirkpatrick, D.C. Slusarski, V.C. Sheffield, R.H. Scheller, and P.K. Jackson.
631             2011. Primary cilia membrane assembly is initiated by Rab11 and transport protein
632             particle II (TRAPPII) complex-dependent trafficking of Rabin8 to the centrosome. *Proc*
633             *Natl Acad Sci U A.* 108:2759–2764. doi:10.1073/pnas.1018823108.

634    Wingfield, J.L., K.-F. Lechtreck, and E. Lorentzen. 2018. Trafficking of ciliary membrane
635             proteins by the intraflagellar transport/BBSome machinery. *Essays Biochem.* 62:753–763.
636             doi:10.1042/EBC20180030.

637    Wright, K.J., L.M. Baye, A. Olivier-Mason, S. Mukhopadhyay, L. Sang, M. Kwong, W. Wang,
638             P.R. Pretorius, V.C. Sheffield, P. Sengupta, D.C. Slusarski, and P.K. Jackson. 2011. An
639             ARL3-UNC119-RP2 GTPase cycle targets myristoylated NPHP3 to the primary cilium.
640             *Genes Dev.* 25:2347–2360. doi:10.1101/gad.173443.111.

641    Xia, R., H. Jia, J. Fan, Y. Liu, and J. Jia. 2012. USP8 promotes smoothened signaling by
642             preventing its ubiquitination and changing its subcellular localization. *PLoS Biol.*
643             10:e1001238. doi:10.1371/journal.pbio.1001238.

644    Xu, Q., Y. Zhang, Q. Wei, Y. Huang, Y. Li, K. Ling, and J. Hu. 2015. BBS4 and BBS5 show
645             functional redundancy in the BBSome to regulate the degradative sorting of ciliary
646             sensory receptors. *Sci. Rep.* 5:11855. doi:10.1038/srep11855.

647    Yau, R., and M. Rape. 2016. The increasing complexity of the ubiquitin code. *Nat. Cell Biol.*
648             18:579–586. doi:10.1038/ncb3358.

649    Ye, F., A.R. Nager, and M.V. Nachury. 2018. BBSome trains remove activated GPCRs from
650             cilia by enabling passage through the transition zone. *J. Cell Biol.* 217:1847–1868.
651             doi:10.1083/jcb.201709041.

652    Yue, S., L.-Y. Tang, Y. Tang, Y. Tang, Q.-H. Shen, J. Ding, Y. Chen, Z. Zhang, T.-T. Yu, Y.E.
653             Zhang, and S.Y. Cheng. 2014. Requirement of Smurf-mediated endocytosis of Patched1
654             in sonic hedgehog signal reception. *eLife.* 3:128. doi:10.7554/eLife.02555.

655    Zhang, Q., D. Nishimura, S. Seo, T. Vogel, D.A. Morgan, C. Searby, K. Bugge, E.M. Stone, K.
656             Rahmouni, and V.C. Sheffield. 2011. Bardet-Biedl syndrome 3 (Bbs3) knockout mouse

657     model reveals common BBS-associated phenotypes and Bbs3 unique phenotypes. *Proc*
658     *Natl Acad Sci U A*. 108:20678–20683. doi:10.1073/pnas.1113220108.

## ACKNOWLEDGMENTS

We thank Drs. Kirk Mykytyn, Val Sheffield, Alice Ting, and Mark Scott for the gifts of cDNAs;

Kathryn Anderson, Vishva Dixit, and Felice Dunn for the gifts of antibodies; Val Sheffield for

the gift of the $Bbs4^{-/-}$ mice, George B. Witman and the Chlamydomonas Resource Center for the

gift of *Chlamydomonas* strains; Yien-Ming Kuo for help with cryosections of mouse retina and core

support; Dhivya Kumar and Tyler Picariello for the assistance with *Chlamydomonas* cultures;

Nicholas Morante and Jeremy Reiter for providing $Bbs4^{-/-}$ mouse eyes; Fan Ye for generating

some of the cell lines used in the study; Irene Ojeda Naharros for help with single cilia tracking,

the Nachury and von Zastrow labs for helpful discussions and the Ogden and Caspary labs for

comments on the preprint. This work was funded by NIGMS (R01-GM089933, M.V.N.). This

work was made possible, in part, by NEI EY002162 - Core Grant for Vision Research and by

the Research to Prevent Blindness Unrestricted Grant (M.V.N.). Authors contributions: SRS

conducted all experimental work except for the experiments in Figure 5 which were conducted

by ARN. MVN supervised research. SRS and MVN wrote the manuscript.

**Figure 1**

**Figure 1. Ubiquitinated signaling receptors accumulate inside cilia of *Arl6⁻/⁻* cells**

**A.** SSTR3 fused to the fluorescent protein mNeonGreen (NG) at its intracellular C-terminus and a biotinylation Acceptor Peptide (AP) tag at its extracellular N-terminus was expressed under the control of an attenuated EF1α promoter (pEF1α$^A$) in wild type (WT) and *Arl6⁻/⁻* IMCD3 cells. Cells were treated with or without somatostatin-14 (sst) for 2 h, then fixed and stained for acetylated tubulin (acTub, magenta) and ubiquitin (Ub, yellow). $^{AP}$SSTR3$^{NG}$ (cyan) was imaged through the intrinsic fluorescence of NG. Channels are shifted in the insets to facilitate visualization of overlapping ciliary signals. Scale bar: 5µm (main panel), 2µm (inset). In WT cells, the ciliary SSTR3 signal decreases over the experimental time course. In *Arl6⁻/⁻* cells, SSTR3 fails to exit cilia and an increase in the ciliary Ub level is detected. As a control, *Arl6⁻/⁻* cells that did not express SSTR3-NG were tested; no increase in ciliary Ub levels was observed upon addition of sst. **B.** The fluorescence intensity of the Ub channel in the cilium was measured in each condition and the data are represented as violin plots. The thick bar indicates the median and the dotted lines the first and third quartiles. An 11-fold increase in ciliary Ub signal is observed upon addition of sst to SSTR3-expressing cells. Asterisks indicate ANOVA significance value. ****, p= <0.0001. **C.** WT or *Arl6⁻/⁻* IMCD3 cells stably expressing $^{AP}$SSTR3$^{NG}$ and the biotin ligase BirA targeted to the ER lumen (BirA-ER ) were transiently transfected with HA-tagged ubiquitin (HA-Ub). 10 µM biotin was added to cells 24 h post transfection for maximal biotinylation of $^{AP}$SSTR3$^{NG}$ and, after another 18 h, cells were treated with sst (10µM) for indicated times. Cells were lysed under denaturing conditions and biotinylated SSTR3 was captured on streptavidin resin. Eluates were probed for HA via immunoblotting and for biotin via streptavidin-HRP. Two major biotinylated proteins endogenous to cells are marked by asterisks. Whole cell lysates were probed for Arl6 and, as a loading control, actin. A non-specific band cross-reacting with the anti-Arl6 antibody is marked with a dot. **D.** Quantitation of SSTR3 ubiquitination. The signals of HA-Ub conjugated to SSTR3 in the streptavidin eluates were measured. The experiment shown in **C** was repeated three times and for each experiment, Ub-SSTR3 signals were normalized to the value in *Arl6⁻/⁻* cells at t = 0 of sst stimulation and plotted as grey circles. The horizontal blue lines represent mean values. **E.** IMCD3 cells of the indicated genotypes were treated with the

Smoothened agonist SAG or the vehicle DMSO for 2h. Cells were then fixed and stained for

acTub and Ub. Channels are shifted in the insets to facilitate visualization of overlapping ciliary

signals. Scale bar 5µm (main panel), 2µm (inset). Activation of Hh signaling promotes a

detectable increase in ciliary Ub levels only in *Arl6*$^{-/-}$ cells. **F.** Violin plots of the fluorescence

intensity of the Ub channel in the cilium in each condition are shown. Asterisks indicate

ANOVA significance value. ****, p= <0.0001.

**Figure 2**

**Figure 2. Ubiquitination of ciliary GPCRs is required for signal-dependent exit but not for endocytosis.**

**A.** IMCD3-[pEF1α$^\Delta$-$^{AP}$SSTR3] cells stably expressed the ER-localized biotin ligase BirA to enables the biotinlylation of $^{AP}$SSTR3. SSTR3 was either the wild-type allele (WT) or a variant where all five cytoplasm-facing lysine residues (listed in Methods) were mutated to arginine (cK0). Ciliary $^{AP}$SSTR3 was pulse-labeled by addition of Alexa647-conjugated mSA (mSA647) to the medium for 5-10 min before addition of sst. Far red fluorescence was tracked in individual cilia at 10 min capture intervals. For each individual cilia, fluorescence intensities were normalized to the value at t = 0. A comparison of the ciliary levels of SSTR3 at t=0 is shown in **Fig. S2A**. Data were plotted and fitted to a line. Error bars: 95% CI. $\mathcal{N}$ =18-22 cilia. **B.** GPR161 fused to three tandem repeats of NG at its C-terminus was expressed under the control of the ultra-weak δ-crystallin promoter (pCrys) in IMCD3 cells. GPR161 was either the wild-type allele (WT) or a variant where all eighteen cytoplasm-facing lysine residues (listed in Methods) where mutated to arginine (cK0). IMCD3-[pCrys-GPR161$^{3NG}$] cells were treated with SAG for 2h. During the course of the experiment, NG fluorescence was tracked in individual cilia at 10 min capture intervals. Fluorescence data were acquired and analyzed as in panel **(A)**. A comparison of the ciliary levels of GPR161 at t=0 is shown in **Fig. S2B**. Error bars: 95% CI. $\mathcal{N}$ =10-19 cilia. **C.** IMCD3-[$^{AP}$SSTR3; BirA-ER] cells expressing either WT or cK0 SSTR3 were pulse-labeled by addition of Alexa647-conjugated mSA (mSA647) to the medium for 5 min before addition of sst and were imaged by far red fluorescence immediately after addition of sst ($t_0$) and 10 min later ($t_{10}$). The contrast level was adjusted to reveal plasma membrane-localized and internalized $^{AP}$SSTR3, causing the cilia-localized $^{AP}$SSTR3 signal to reach saturation. Scale bar 5μm. **D.** Internalized $^{AP}$SSTR3 foci were counted immediately after addition of sst ($t_0$) and 10 min later ($t_{10}$), n = 34 cells.

**Figure 3**

**Figure 3. Signal-dependent accumulation of K63-linked ubiquitin chains inside cilia of *Bbs* mutant cells.**

**A.** IMCD3 cells of the indicated genotypes expressing $^{AP}$SSTR3$^{NG}$ were treated with somatostatin-14 for 2 h. Cells were fixed and stained for AcTub (magenta) and with antibodies specific for the lysine 63 (UbK63) or lysine 48 (UbK48) Ub chain linkages (yellow). SSTR3$^{NG}$ (cyan) was imaged through the intrinsic fluorescence of NG. Channels are shifted in the insets to facilitate visualization of overlapping ciliary signals. Scale bar 5µm (main panel), 2µm (inset). **B.** The fluorescence intensity of the UbK48 and UbK63 channels in the cilium are represented as violin plots. A 14-fold increase in ciliary Ub abundance is detected with the K63Ub linkage-specific antibody. Asterisks indicate ANOVA significance value. ****, p= <0.0001. No ciliary signal is detected with the K48Ub linkage-specific antibody. **C.** *Arl6*$^{-/-}$ IMCD3-[$^{AP}$SSTR3; BirA-ER] cells were transfected with the HA-tagged ubiquitin variants WT, noK0 (all seven acceptor lysine residues mutated to arginine) or K63 (where all lysine residues are mutated to arginine except for K63). Biotin was included in the culture medium and cells were treated with sst for 0 or 10 min. Cells were lysed under denaturing conditions and biotinylated SSTR3 was captured on streptavidin resin. Eluates were probed for HA via immunoblotting and for biotin via streptavidin-HRP. Two major biotinylated proteins endogenous to cells are marked by asterisks. Whole cell lysates were probed for Arl6 and, as a loading control, actin. A non-specific band cross-reacting with the anti-Arl6 antibody is marked with a dot. WT IMCD3 cells were processed in parallel as a control. **D.** Quantitation of SSTR3 ubiquitination. The signals of HA-Ub conjugated to SSTR3 in the streptavidin eluates were measured. The experiment shown in **C** was repeated four times and for each Ub variant, Ub-SSTR3 signals were normalized to the value at t = 0 of sst stimulation and plotted as grey circles. The horizontal blue lines represent mean values. Asterisks indicate ANOVA significance value. *** p =< 0.001; ** p=<0.01.

**Figure 4**

**Figure 4. Ciliary K63-linked ubiquitin chains are required for GPCR exit from cilia.**

**A.** Diagram of the working model and the experimental strategy. **B.** IMCD3-[pEF1α$^Δ$-$^{AP}$SSTR3; pEF1α-BirA•ER] were transfected with plasmids expressing the ciliary targeting signal (CTS) of NPHP3 fused to GFP only (left panels), to GFP and the catalytic domain of the K63-specific deubiquitinase AMSH (middle panels), or to GFP and the catalytically inactive E280A variant of AMSH catalytic domain (AMSH$^†$, right panels). Ciliary $^{AP}$SSTR3 was pulse-labeled with mSA647 for 5-10 min and cells were then treated with or without sst for 2h, before fixation and staining for acetylated tubulin (acTub, magenta) and DNA (blue). The NPHP3$^{CTS}$ fusions were visualized through the intrinsic fluorescence of GFP (cyan) and $^{AP}$SSTR3 was visualized via mSA647 (yellow). Channels are shifted in the insets to facilitate visualization of overlapping ciliary signals. Scale bar 5µm (main panel), 2µm (inset). **C.** The fluorescence intensities of ciliary $^{mSA647-AP}$SSTR3 are represented as violin plots. Asterisks indicate Kruskal-Wallis test significance values. **\*\*\***, p= <0.001; **\*\***, p= <0.01. Addition of sst triggers the exit of SSTR3 from cilia in cells transfected with the controls NPHP3$^{CTS}$ and NPHP3$^{CTS}$-AMSH$^†$ but NPHP3$^{CTS}$-AMSH blocks ciliary exit of SSTR3. **D.** RPE1-hTERT (RPE) cells transfected with the indicated constructs were treated with SAG or vehicle (DMSO) for 2h, then fixed and stained for acetylated tubulin (magenta) and GPR161 (yellow). The NPHP3$^{CTS}$ fusions were visualized through the intrinsic fluorescence of GFP (cyan). Channels are shifted in the insets to facilitate visualization of overlapping ciliary signals. Scale bar 5µm (main panel), 2µm (inset). **E.** The fluorescence intensities of ciliary GPR161 are represented as violin plots. Asterisks indicate ANOVA significance value. **\*\*\*\***, p= <0.0001. NPHP3$^{CTS}$-AMSH specifically blocks the SAG-dependent exit of GPR161 from cilia. **F.** IMCD3 cells transfected with the indicated constructs were fixed and stained for acetylated tubulin (magenta) and SMO (yellow). The NPHP3$^{CTS}$ fusions were visualized through the intrinsic fluorescence of GFP (cyan). Channels are shifted in the insets to facilitate visualization of overlapping ciliary signals. **G.** The fluorescence intensities of ciliary SMO are represented as violin plots. Asterisks indicate ANOVA significance value. **\*\*\*\***, p=

788    <0.0001. NPHP3[CTS]-AMSH specifically blocks the constitutive exit of SMO from cilia and

789    promotes accumulation of SMO in cilia in the absence of pathway activation.

**Figure 5**

**Figure 5. β-arrestin 2 is recruited to cilia upon activation of SSTR3 or GPR161.**

**A.** β-Arrestin2 is rapidly recruited to cilia upon activation of SSTR3. siRNA-treated IMCD3-[$^{AP}$SSTR3, β-Arrestin2$^{GFP}$] cells were pulse-labeled with mSA647 before addition of sst. Channels are shifted to facilitate visualization of overlapping ciliary signals. As shown previously (Shankar et al., 2010), β-Arrestin2 is at the basal body in unstimulated cells. Scale bar: 2 µm. **B.** Kinetics of β-Arrestin2$^{GFP}$ recruitment to cilia upon sst stimulation in control- and ARL6-depleted IMCD3-[$^{AP}$SSTR3, β-Arrestin2$^{GFP}$] cells. Data were fit to a simple exponential. (n=10 cilia). **C-D.** Kinetics of early trafficking events for retrieval of SSTR3 **(C)** and GPR161 **(D)**. The top panels show removal of stably expressed $^{AP}$SSTR3$^{NG}$ or $^{AP}$GPR161$^{NG3}$ following addition of sst **(C)** or SAG **(D)**. The bottom panels show the kinetics of β-Arrestin2$^{GFP}$ entry **(C-D)**, $^{AP}$Smoothened$^{NG}$ entry **(D)**, and $^{NG3}$BBS5 tip accumulation **(C)** measured in IMCD3 cells stably expressing the indicated proteins. Fluorescence values are normalized to the initial ($t_0$) and final ($t_f$) values. Error bars: SEM. (n=6-11 cilia).

**Figure 6**

**Figure 6. β-arrestin 2 directs the signal-dependent ubiquitination of ciliary GPCRs.**

**A.** IMCD3-[pEF1α$^{\Delta}$-$^{AP}$SSTR3$^{NG}$] cells of the indicated genotypes were treated with sst for 2h before fixation and staining for acetylated tubulin (acTub, magenta) and ubiquitin (Ub, yellow). SSTR3$^{NG}$ was visualized through the intrinsic fluorescence of NG (cyan). Channels are shifted in the insets to facilitate visualization of overlapping ciliary signals. SSTR3 exit is blocked in *Arl6$^{-/-}$* cells regardless of the *β-Arrestin2* genotype but the ciliary Ub signal is only evident when β-Arrestin2 function is intact. Scale bar: 5µm (main panel), 1µm (inset). **B.** Violin plots representing the ciliary levels of Ub under the indicated conditions. Asterisks indicate ANOVA significance value. ****, p= <0.0001. **C.** IMCD3 WT, knockout for *Arl6* only and double knockout for *Arl6* and the β-Arrestin2 gene *Arrb2* were treated with SAG for 2h before fixation and staining for acetylated tubulin (acTub, magenta) and ubiquitin (Ub, yellow). Channels are shifted in the insets to facilitate visualization of overlapping ciliary signals. Scale bar: 5µm (main panel), 1µm (inset). **D.** Violin plots representing the ciliary Ub levels. Asterisks indicate ANOVA significance value. ****, p= <0.0001; **, p= <0.01. The appearance of a Ub signal in cilia in *Arl6$^{-/-}$* cells depends on β-Arrestin2. **E.** IMCD3-[$^{AP}$SSTR3; BirA-ER] cells of the indicated genotypes stably expressing $^{AP}$SSTR3$^{NG}$ and BirA-ER were transfected with HA-Ub, biotin was added to the medium and cells were treated with sst for indicated times. Biotinylated SSTR3 was captured from cell lysates under denaturing conditions on streptavidin resin. Eluates were probed for HA via immunoblotting and for biotin via streptavidin-HRP. Two major biotinylated proteins endogenous to cells are marked by asterisks. Whole cell lysates were probed for Arl6, β-Arrestin2 to verify genotypes and, as a loading control, actin. A non-specific band cross-reacting with the anti-Arl6 antibody is marked with a dot. WT IMCD3 cells were processed in parallel as a control. **F.** Quantitation of SSTR3 ubiquitination. The signals of HA-Ub conjugated to SSTR3 in the streptavidin eluates were measured. The experiment shown in **E** was repeated twice and for each experiment, Ub-SSTR3 signals were normalized to the value in *Arl6$^{-/-}$* cells at t = 0 of sst stimulation and plotted as grey circles. The horizontal blue lines represent mean values.

**Figure 7**

**Figure 7. Accumulation of ubiquitinated proteins in cilia of *Chlamydomonas* cells**

**and mammalian photoreceptors.**

**A.** *Chlamydomonas* cells of the indicated genotypes were fixed and stained for acetylated tubulin

(acTub, magenta) and ubiquitin (Ub, yellow). Scale bar: 5 µm. A weak ubiquitin signal detected

in WT cells is increased in *bbs4* mutant cells. **B.** Violin plots of the ciliary Ub levels in WT and

*bbs4 Chlamydomonas* cells. Asterisks indicate Mann-Whitney test significance values. **\*\*\*\***, p=

<0.0001. The ciliary levels of Ub are increased more than 2-fold in *bbs4* cells as compared to

WT. **C.** Cilia purified from WT and *bbs4* cells via deciliation with dibucaine and differential

centrifugation were solubilized, and the protein contents resolved by SDS-PAGE. Immunoblots

for Ub (using the P4D1 antibody) and tubulin are shown. A rise in ubiquitin conjugates is

observed in *bbs4 Chlamydomonas* cells compared to WT. **D.** Mouse retina were fixed, embedded

and sectioned before staining for ubiquitin (Ub, yellow, lower panel), cone arrestin (magenta, a

marker of the photoreceptor outer segment layer) and DNA (cyan). OS: outer segment; IS: inner

segment; ONL: outer nuclear layer; OPL: outer plexiform layer; INL: inner nuclear layer. Scale

bar: 10µm. **E.** Model for the role of ubiquitin chains in selecting cargoes for signal-dependent

and constitutive exit via BBSome-mediated transport.

**A** IMCD3-[pEF1α$^Δ$-$^{AP}$SSTR3$^{NG}$]

**B** IMCD3-[pEF1α$^Δ$-$^{AP}$SSTR3$^{NG}$]

**C** IMCD3-[pEF1α$^Δ$-$^{AP}$SSTR3$^{NG}$]

**D** IMCD3-[pEF1α$^Δ$-$^{AP}$SSTR3$^{NG}$]

**Supplementary Figure 1**

**Figure S1. Signal-dependent ubiquitination of ciliary GPCRs**

**A.** IMCD3 cells of the indicated genotypes expressing $^{AP}$SSTR3$^{NG}$ were treated with

somatostatin (sst) for 2 h. Cells were fixed and stained for ubiquitin with the FK1 monoclonal

antibody (Ub (FK1), yellow) and acetylated tubulin (acTub, magenta). $^{AP}$SSTR3$^{NG}$ (cyan) was

imaged through the intrinsic fluorescence of NG. Channels are shifted in the insets to facilitate

visualization of overlapping ciliary signals. Scale bar: 5μm (main panel), 2μm (inset). **B.** Violin

plots of the fluorescence intensity of the Ub channel in the cilium in each condition are shown. A

3-fold increase in ciliary Ub signal is observed upon addition of sst to SSTR3-expressing cells.

Asterisks indicate ANOVA significance value. ****, p= <0.0001. **C.** IMCD3 cells of the

indicated genotypes expressing $^{AP}$SSTR3$^{NG}$ were treated with sst for 2 h. Cells were fixed and

stained for AcTub (magenta) and UbK63 (yellow). $^{AP}$SSTR3$^{NG}$ (cyan) was imaged through the

intrinsic fluorescence of NG. Channels are shifted in the insets to facilitate visualization of

overlapping ciliary signals.  Scale bar 5μm (main panel), 2μm (inset). **D.** The fluorescence

intensity of the UbK63 channel in the cilium is represented as violin plots. Asterisks indicate

ANOVA significance value. ****, p= <0.0001.

**Supplementary Figure 2**

**Supplementary figure 3**

**Supplementary figure 4**

**Figure S2. Fluorescence intensities of WT and cK0 GPCR variants at t=0.**

**A.** Ciliary $^{AP}$SSTR3 was pulse-labeled by addition of Alexa647-conjugated mSA (mSA647) to the medium for 5-10 min before addition of sst. Violin plots of the ciliary fluorescence intensities for WT and cK0 SSTR3 imaged in the far-red channel at t=0 are shown. **B.** Violin plots of the fluorescence intensities of NG fluorescence corresponding to WT and cK0 GPR161$^{NG}$ at t=0 are shown.

**Figure S3. Expression levels of ciliary AMSH.**

IMCD3-[pEF1α$^{Δ}$-$^{AP}$SSTR3; pEF1α-BirA•ER] cells were transfected with the plasmid expressing NPHP3$^{CTS}$-AMSH as in Figure 4, before fixation and staining for acetylated tubulin (acTub, magenta) and DNA (blue). The NPHP3$^{CTS}$ fusions were visualized through the intrinsic fluorescence of GFP (cyan) A representative micrograph is shown. A cell expressing modest levels of cilia-AMSH is indicated by an arrow and possesses normal ciliary staining of acetylated tubulin. A cell expressing high levels of cilia-AMSH indicated by an arrowhead displays abnormal ciliary levels of acetylated tubulin. Cells with high ciliary levels of NPHP3$^{CTS}$-AMSH were excluded from the analysis and only cells with modest ciliary levels of NPHP3$^{CTS}$-AMSH were included in the experiment. Scale bar 5µm (main panel), 1µm (inset).

**Figure S4. Cilia-targeted AMSH removes UbK63 chains from ciliary substrates.**

**A.** IMCD3-[pEF1α$^{Δ}$-$^{AP}$SSTR3; pEF1α-BirA•ER] cells were transfected with control or Arl6 siRNAs. Cell lysates were immunoblotted for Arl6 or actin. A non-specific band cross-reacting with the anti-Arl6 antibody is marked with a dot. **B.** IMCD3-[$^{AP}$SSTR3; BirA•ER] cells were transfected with siRNA targeting Arl6 and plasmids expressing NPHP3$^{CTS}$-AMSH or NPHP3$^{CTS}$- AMSH$^{†}$. Surface-exposed $^{AP}$SSTR3 was pulse-labeled with mSA647 for 5-10 min and cells were then treated with sst for 2h before fixation and staining for K63-linked ubiquitin chains (UbK63, yellow). The NPHP3$^{CTS}$-AMSH fusions were visualized through the intrinsic fluorescence of GFP (cyan), $^{AP}$SSTR3 was visualized via mSA647 (magenta) and DNA is blue.

891 Channels are shifted in the insets to facilitate visualization of overlapping ciliary signals. Scale bar

892 5µm (main panel), 2µm (inset). **C.** The fluorescence intensities of the UbK63 signal inside cilia

893 are represented as violin plots. Asterisks indicate Mann Whitney test significance value. **\*\*\*\***, p=

894 <0.0001.

  **MATERIALS AND METHODS**

**Cell culture**

The mouse IMCD3 cell lines used in the study were generated from a parental IMCD3-FlpIn

cell line (gift from P.K. Jackson, Stanford University, Stanford, CA). IMCD3-FlpIn cells were

cultured in DMEM/F12 (11330-057; Gibco) supplemented with 10% FBS (100-106; Gemini

Bio-products), 100 U/ml penicillin-streptomycin (400-109; Gemini Bio-products), and 2 mM L-

glutamine (400-106; Gemini Bio-products). The RPE1-hTERT cell line (ATCC CRL-4000) was

cultured in DMEM/F12 supplemented with 10% FBS, 100 U/ml penicillin-streptomycin, 2 mM

L-glutamine and 0.26% sodium bicarbonate (25080; Gibco).

Ciliation was induced by serum starvation in media containing 0.2% FBS for 16 to 24 h.

**Plasmid construction and Generation of stable cell lines**

Stable isogenic IMCD3 cell lines were generated using the Flp-In system (ThermoFisher

Scientific). Low-expression promoters and additional expression cassettes were introduced into

pEF5/FRT plasmid as described (Nager et al., 2017; Ye et al., 2018). To reduce expression

levels, the EF1α promoter was attenuated by mutating the TATA-box (pEF1α$^\Delta$) or replaced with

a minimal chicken lens δ-crystallin promoter (pCrys).

Coding sequences were amplified from plasmids encoding mouse GPR161 (BC028163;

Mammalian Gene Collection [MGC]; GE Healthcare), mouse SSTR3 (gift from Kirk Mykytyn,

Ohio State University, Columbus, OH), and human BBS5 (gifts from V. Sheffield, University of

Iowa, Iowa City, IA), BirA-ER (gift from Alice Ting). β-arrestin 2 (a gift from Mark Scott) was

stably expressed from a pCMV-based plasmid (pEGFP-N). SSTR3 and SMO expression were

driven by pEF1α$^\Delta$, GPR161 expression by pCrys, and BBS5 expression by pEF1α. Coding

sequences were fused to GFP, mNeonGreen (NG, (Shaner et al., 2013), or an acceptor peptide

for the biotin ligase BirA (AP) (Howarth and Ting, 2008). Multiple rounds of site-directed

mutagenesis were performed to generate SSTR3-cK0, a variant with all the cytoplasm-facing

lysine residues (K233, K330, K356, K407, and K421) mutated to arginine. GPR161-cK0, a

variant with 18 cytoplasm-exposed lysine residues (K83, K84, K93, K167, K247, K250, K269,

K296, K358, K362, K455, K469, K473, K481, K486, K497, K504, and K548) mutated to

arginine, was gene synthesized (GenScript). Cilia-AMSH was generated by fusing the catalytic

domain of mouse AMSH (gift from David Komander; Addgene plasmid #66712; (Michel et al.,

2015) with NPHP3(1-200) and GFP to create NPHP3[1-200]-GFP-AMSH[243-424]. A

catalytically dead version of AMSH was generated by mutating the water-activating Glu280

residue to Ala (Sato et al., 2008; Huang et al., 2013). pRK5-HA-Ubiquitin WT, K0 (all seven

acceptor lysine residues − K6, K11, K27, K29, K33, K48, and K63, are mutated to arginine) ,

and K63 only (all lysines except K63 mutated to arginines)(gift from Ted Dawson; Addgene

plasmid numbers − 17608, 17603, and 17606 respectively).

CRISPR-edited *Arl6*$^{-/-}$ and *Arl6*$^{-/-}$/*Arrb2*$^{-/-}$ cell lines were described previously (Liew et al., 2014;

Nager et al., 2017). The genotypes are *Arl6*, NM_001347244.1:c.- 10_25del; c.3_6del; and *Arrb2*,

NM_001271358.1:c.1 12del;c112_113del.

## Transfection

For generation of all stable cell lines except for the ones expressing β-arrestin 2$^{GFP}$, FRT-based

plasmids were reverse transfected using XtremeGene9 (Roche) into IMCD3 Flp-In cells along

with a plasmid encoding the Flp recombinase (pOG44) and stable transformants were selected by

blasticidin resistance (4 µg/ml). The transfection mixture was then added to the cells. pEGFP-N•

β-arrestin 2-GFP was transfected into IMCD3 cells using Lipofectamine 2000, and clones were

selected using neomycin resistance.

## Antibodies and drug treatments

The following monoclonal antibodies were used for immunofluorescence: anti-acetylated tubulin

(mouse; clone 6-11B-1; Sigma-Aldrich; 1:500), anti-ubiquitin clone FK2  was used in all

experiments except where indicated (mouse; D058-3; Medical and Biological Laboratories;

1:500), anti-ubiquitin FK1 in **Fig. S1A-B** (mouse; D071-3; Medical and Biological Laboratories;

1:500), anti-ubiquitin K48 linkage (rabbit; clone Apu2; 05-1307; Millipore-Genentech; 1:500),

anti-ubiquitin K63 linkage (human; clone Apu3; a gift from Genentech; 1:500). The following

monoclonal antibodies were used for immunoblotting: anti-ubiquitin (mouse; clone P4D1; 3936;

Cell Signaling; 1:1000), anti-α tubulin  (mouse; clone DM1A; MS-581-P1ABX; Thermo

Scientific;1:1000), anti-HA (mouse; clone 16B12; 901501; Biolegend;1:500), streptavidin-HRP

(Pierce-21140;Thermo Scientific; 1:10000). The following polyclonal antibodies were used for

immunofluorescence: anti-GPR161 (rabbit; 13398-1-AP; Proteintech; 1:100), anti-Smoothened

(rabbit; a gift from Kathryn Anderson, Memorial Sloan Kettering Cancer Center, New York,

NY; 1:500), anti-cone-arrestin (rabbit; AB15282; Millipore; 1:500). Biotinylated SSTR3 and

GPR161 were detected using Alexa Fluor 647–labeled monovalent streptavidin (mSA647) (Ye et

al., 2018). The following reagents were used at the indicated concentrations: 200 nM SAG and

10 µM somatostatin 14. Somatostatin 14 stocks were made in DMEM/F12 media, HEPES, no

phenol red (11039-021, GIBCO). SAG was dissolved in DMSO (276855, Sigma-Aldrich).

**Imaging and microscopy**

Cells were imaged either on a DeltaVision system (Applied Precision) equipped with a PlanApo

60x/1.40 objective lens (Olympus), CoolSNAP HQ2 camera (Photometrics), and solid-state

illumination module (InsightSSI), or on a confocal LSM 700 (Zeiss) microscope equipped with

40x Plan-Apochromat 1.3 DIC oil objective. Z stacks were acquired at 0.5 µm interval (LSM

700, Figs. 1, 3, 7, and S1) or 0.2 µm interval (DeltaVision, Figs. 4, 6, S3, and S4).

For fixed imaging, 60,000 cells were seeded on acid-washed coverglass (12 mm #1.5; 12-545-81;

Thermo Fisher Scientific). Cells were grown for 24 h and then starved for 16-24 h in 0.2% FBS

media before experimental treatment. After treatment, cells were fixed with 4%

paraformaldehyde in PBS for 15 min at 37°C and extracted in -20°C methanol for 5 min. Cells

were then permeabilized in IF buffer (PBS supplemented with 0.1% Triton X-100, 5% normal

donkey serum (017-000-121; Jackson ImmunoResearch Laboratories), and 3% bovine serum

albumin (BP1605-100; Thermo Fisher Scientific)), incubated at room temperature for 1h with

primary antibodies diluted in IF buffer, washed three times in IF buffer, incubated with

secondary antibodies (Jackson ImmunoResearch Laboratories) diluted in IF buffer for 30 min,

and washed three times with IF buffer. DNA was stained with Hoechst 33258 (H1398; Molecular

Probes), cells were washed twice more with PBS, and coverglass mounted on slides using

fluoromount-G (17984-25; Electron Microscopy Sciences).

For live-cell imaging, 300,000 cells were seeded on acid-washed 25mm cover glass (Electron

Microscopy Sciences). After 24 h of growth, cells were serum starved for 16 h and transferred to

the DeltaVision stage for imaging at 37°C within an environmental chamber that encloses the

microscope and the stage. Cells were imaged in DMEM/F12 media, HEPES, no phenol red

(11039-021, GIBCO). To measure SSTR3 exit, cells expressing [AP]SSTR3 were first washed three

times with PBS and then pulse-labeled with 2μg/ml mSA647 for 5 min at 37°C. Cells were then

imaged after addition of sst (for SSTR3) or SAG (for GPR161) for 2 h with an acquisition every

10 min. Each acquisition consisted of a 3-plane Z-stack with each plane separated by 0.5 μm.

Transmittance was set to 5% and exposure time to 500 ms.

WT or *Bbs4⁻/⁻* mice (Mykytyn et al., 2004) eyes were enucleated at P21. Whole eyes were fixed in

4% paraformaldehyde/PBS overnight at 4°C and then washed three times with PBS. The eyes

were infiltrated in 30% sucrose/PBS at 4 °C overnight. Eyes were placed in OCT compound

embedding medium (4583; Tissue-Tek), frozen on dry ice, and stored at −80 °C. 14 μm sections

were cut on a cryostat (CM 1850; Leica). Sections were processed for immunofluorescence as

follows. Sections were blocked in blocking solution (50 mM Tris pH7.4, 100 mM NaCl, 0.1%

Triton-X 100, 3% normal donkey serum, and 0.1% BSA) for 1 h at room temperature, then

incubated with primary antibodies overnight at 4°C, followed by three washes in wash solution

(50 mM Tris pH7.4, 100 mM NaCl, and 0.1% TX100). After rinsing in wash solution, sections

were incubated with secondary antibodies for 1 h at room temperature. DNA was counterstained

using Hoechst 33258 dye, sections washed twice more with PBS, and mounted on slides using

Fluoromount-G. Sections were imaged on Confocal LSM 700 (Zeiss ) microscope.

1005

**Image analysis**

Files were imported from the Deltavision or LSM700 workstations into ImageJ/ Fiji (National

Institutes of Health) for analysis. For the quantification of ciliary ubiquitin signal in fixed cells,

maximum intensity projections were used. The ciliary ubiquitin intensity was measured using the

following equation:

$F_{Ub\text{-}cilia} = F_{Ub\text{-}cilia\_measured} - F_{background}$

$F_{Ub\text{-}cilia\_measured}$ is the total ciliary ubiquitin fluorescence detected, $F_{background}$ is the background

ubiquitin fluorescence measured in the adjacent area. For all measurements, the fluorescence

integrated density was used. $F_{Ub\text{-}cilia}$ were plotted as violin plots using the PlotsOfData web tool

(https://huygens.science.uva.nl/PlotsOfData/). Each violin represents the distribution of data,

which includes all the data points. Median and interquartile range is marked by solid and dotted

lines, respectively. No gamma adjustment was applied during figure preparation; all the

representative micrographs for respective panels are displayed within the same dynamic range.

For some representative micrographs, the most in-focus plane was used rather than the

maximum intensity projection (Fig.3, and Fig. 6, panel A and C).

To measure SSTR3 exit, the integrated density of Alexa Fluor 647 fluorescence in the maximum

intensity projection was measured for each time point. The cilia-adjacent fluorescence was

subtracted as the background, and a mathematical photobleaching correction was applied:

$F_{cilia} = (F_{mSA647\_measured}/F_{mSA647\_1}) + ((1 - e^{-\lambda}) * (n - 1))$, where $\lambda$ is the photobleaching decay

constant, n is the number of images taken, $F_{mSA647\_measured}$ is the integrated mSA647 fluorescence

measured for image n, $F_{mSA657\_1}$ is the measurement for the first time point, and $F_{cilia}$ is the

reported fluorescence. In this equation, $F_{cilia}$ is reported in relative fluorescence units (RFUs).

[AP]GPR161[3NG] exit was followed similarly with the difference that NG fluorescence intensity was

measured. For SSTR3 as well as GPR161, photobleaching corrected data ($F_{cilia}$) for each

condition were linearly fitted ($F_{cilia} = m * time + c$) and plotted in Excel.

<mark>Endocytosed SSTR3 foci were revealed by contrast enhancement and counted with the ImageJ</mark>
<mark>particle analysis tool.</mark>

### *Chlamydomonas* culture, isolation of flagella and immunofluorescence

*Chlamydomonas* WT- g1 (nit1, agg1, mt+) [gift from George B. Witman, University of

Massachusetts Medical School, Worcester, Massachusetts]and *Bbs4*- CC-4377 ptx5-1/bbs4-1::

NIT1 agg1 mt+ (Chlamydomonas Resource Center) strains were cultured asynchronously under

light in TAP media (Gorman and Levine, 1965) for 72 h. Flagellar fractions were prepared from

two liters of cultures of WT or *Bbs4* cells as described (Craige et al., 2013). Briefly, cells were

harvested by centrifugation for 5 min at 2,000 rpm (1,100 x *g*) at room temperature. Cells were

resuspended in 10 mM HEPES, pH 7.4 and centrifuged for 5 min at 2,000 rpm (1,100 x *g*). Next,

the cell pellet was gently resuspended in ice-cold HMDS (10 mM HEPES, pH 7.4, 5 mM

$MgSO_4$, 1 mM DTT, and 4% (w/v) sucrose). Deflagellattion was achieved by addition of 5 mM

dibucaine to each tube of cells and by pipetting up and down ~10 times using a 10-ml plastic

serological pipette. The suspension was spun for 5 min at 1,800 x *g* at 4°C. The supernatant

containing flagella was underlaid with 9 ml ice-cold HMDS-25% sucrose (10 mM HEPES, pH

7.4, 5 mM $MgSO_4$,1 mM DTT, and 25% (w/v) sucrose), followed by a centrifugation step for 10

min at 2,800 rpm (2,400 x *g*), 4°C. The supernatant above the sucrose interface was collected

using a 25-ml serological pipette and flagella were pelleted by centrifugation for 20 min at 30,000

x *g* (16,000 rpm), 4°C. The flagellar pellet was resuspended in 100 µl HMDS buffer. Protein

concentrations were measured by Bradford and 25 ug were resolved by SDS-PAGE for

immunoblotting.

*Chlamydomonas* immunofluorescence was performed as follows. Cells were fixed with 4 %

paraformaldehyde in MT buffer* (30 mM HEPES, pH7, 5 mM EGTA, 5 mM MgSO4, and

25mM KCl) for 20 minutes in suspension. The cells were centrifuged at 1000 rpm for 5 minutes,

resuspended in 100 µl of fixative and transferred onto slides coated with 1 mg/ml poly-L-lysine.

After 5 minutes, the unadhered cells were washed off by rinsing with PBS. Cells were

permeabilized in 0.5% Triton-X 100 for 20 minutes followed by blocking for 1 h in blocking

buffer (3% fish skin gelatin, 1% BSA, 0.1% Tween20 in PBS) at room temperature. Cells were

incubated at 4°C overnight with primary antibodies diluted in blocking buffer, washed five times

in PBS and incubated with secondary antibodies for 2 h at room temperature. After five washes

in PBS, coverglasses were mounting on slides using Fluoromount-G. Cells were imaged on the

LSM700 confocal microscope.

1063

**Biochemical analysis of SSTR3 ubiquitination**

IMCD3 cells stably expressing $pEF1\alpha^{\Delta\text{-AP}}SSTR3^{NG}$ and BirA-ER were transiently transfected

with HA-tagged ubiquitin plasmids. Cells were reverse transfected using XtremeGene9 and

plated into 15 cm plates. Cells were moved to medium containing 0.2% serum and

supplemented with 10 μM biotin 24 h after transfection to promote ciliation and maximize

biotinylation of SSTR3. 18h after the medium change, sst was added for 0, 10 or 20 min. The

assay was stopped by washing cells twice with ice-cold 1xPBS and scraping cells off on ice into

1ml of ice-cold RIPA buffer (50 mM Tris/HCl pH 8, 150 mM NaCl, 1% Nonidet P-40, 0.5%

sodium deoxycholate, 0.1% SDS, 100 mM iodoacetamide, 10mM EDTA) supplemented with

protease inhibitors (1 mM AEBSF, 0.8 mM Aprotinin, 15 mM E-64, 10 mg/mL Bestatin, 10

mg/mL Pepstatin A and 10 mg/mL Leupeptin). After gently rocking at 4°C for 5 minutes,

lysates were clarified by spinning in a tabletop Eppendorf centrifuge at 15,000 rpm and 4°C for

20 minutes. Each clarified lysate was incubated with 50µl of streptavidin Sepharose beads (GE

Healthcare) for 1 h at 4°C to capture biotinylated SSTR3. Beads were washed four times with

500µl of RIPA buffer and bound proteins were eluted by heating the beads in 1X LDS sample

loading buffer containing 2 mM biotin at 70°C for 10 minutes. After SDS-PAGE and transfer,

PVDF membranes were probed with anti-HA antibody and streptavidin-HRP.

Signals were acquired with a ChemiDoc imager (Bio-Rad) and analyzed with ImageLab 6.0.1

(Bio-Rad). The integrated densities of the SSTR3 bands in the streptavidin-HRP blot were

normalized to the integrated densities of the endogenous biotinylated proteins to correct for

variations in the recovery of biotinylated proteins on streptavidin Sepharose. The resulting value

is $SSTR3_{bio}$. Second, the integrated densities of ubiquitinated SSTR3 were collected by

measuring the area from 100 kDa to the top of the gel on the HA blot. The values were

background-corrected by subtracting the value of the equivalent area in the IMCD3 control lane.

The resulting value is Ub-SSTR3$_{raw}$. The relative amount of SSTR3 that is ubiquitinated, Ub-

SSTR3$_{frac}$ = Ub-SSTR3$_{raw}$/SSTR3$_{bio}$ was plotted on the graphs using PlotsOfData (Postma and

Goedhart, 2019).

1091