Polar Gini Curve: a Technique to Discover Single-cell Biomarker Using 2D Visual Information

Thanh Minh Nguyen[1], Jacob John Jeevan[1], Nuo Xu[2], Jake Chen[1*]

[1]Informatics Institute, the University of Alabama at Birmingham, AL, United States

[2]Collat School of Business, the University of Alabama at Birmingham, AL, United States

[*]Corresponding author: Jake Chen

Email: jakechen@uab.edu

Running title: *Nguyen et al / Polar Gini Curve single cell*

Authors' ORCID No

Thanh Nguyen: 0000-0002-8440-1594

Jake Chen: 0000-0001-8829-7504

Total word counts: 3446

Total figures: 10

Total tables: 0

Total supplementary figures: 0

Total supplementary tables: 0

Total supplementary files: 3

Abstract

In this work, we design the Polar Gini Curve (PGC) technique, which combines the gene expression and the 2D embedded visual information to detect biomarkers from single-cell data. Theoretically, a Polar Gini Curve characterizes the shape and 'evenness' of cell-point distribution of cell-point set. To quantify whether a gene could be a marker in a cell cluster, we can combine two Polar Gini Curves: one drawn upon the cell-points expressing the gene, and the other drawn upon all cell-points in the cluster. We hypothesize that the closers these two curves are, the more likely the gene would be cluster markers. We demonstrate the framework in several simulation case-studies. Applying our framework in analyzing neonatal mouse heart single-cell data, the detected biomarkers may characterize novel subtypes of cardiac muscle cells. The source code and data for PGC could be found at https://figshare.com/projects/Polar_Gini_Curve/76749.

KEYWORDS: Single-cell gene expression; Gini coefficient; Polar Gini Curve; Biomarker

Introduction

Discovering biomarkers from the single-cell gene expression data is an interesting yet challenging problem [1]. Compared to the well-established bulk gene expression data, the expression distribution in single-cell is significantly more heterogeneous [2-4]. Therefore, as shown in [5, 6], the bulk-analysis strategies [7, 8] achieve low sensitivity in detecting markers. In addition, as embedding [9-11] and clustering [12-14] are the essential components in many single-cell expression analytical pipelines [15, 16], the biomarker detection techniques would need to tackle the challenges and errors from embedding and clustering [17, 18].

From the statistical point of view, there are two different directions among the current state-of-the-art methods in solving the single-cell biomarker discovery problem. The first direction is using non-parametric approaches [19]. Non-parametric approaches do not attempt to construct the model characterizing the gene expression distribution [20]. They do not require too many prior assumptions about the expression data. Therefore, in theory, they could be applied in most of the heterogeneous scenarios in single-cell expression. For example, Seurat [16] and the SINCERA [21] pipelines use the Mann–Whitney test [22]. The disadvantages of non-parametric

approaches include lacking the point-estimator (for example, we could not tell how much of
fold-change when comparing the expressions of the same gene in two populations) and the lower
true positive rate [5, 6]. On the other hand, the parametric approaches model the underlying
expression distribution. For example, [23] applies Bayesian statistics, Monocle2 [11, 24] and
MAST [2] apply different linear models, and [25] applies the Poisson models to single-cell
differential expression analysis. The parametric approaches, compared to the non-parametric
ones, are significantly more sensitive [5, 6], especially in detecting markers in small cell-cluster
since they may require less number of cell-samples. However, these approaches assume that the
gene expression distribution has specific shapes; therefore, these approaches tend to have higher
false-positive rates.

In this work, we developed a new framework based on the novel idea of integrating
expression and the embedded visual information of single-cell data into one metric to identify
biomarkers. This idea has been successfully implemented in spatial single-cell data, in which the
visualization space reflects the relative position of the cells in a tissue image [26, 27]. In this
framework, we decided to take advantage of cluster shape and cell-point distribution from the 2D
visual space. Our strategy was to project the single-cell 2D cluster onto multiple angle-axes to
explore all viewing angles of the cluster. On each 'viewing angle', we captured the visual
distribution using the Gini coefficient [28]. Together, for each set of points in 2D, we constructed
a Polar Gini Curve (PGC) from the correspondent between viewing angle and Gini coefficient.
We hypothesized that for the marker gene, its expressing cell set should have its PGC close to
the PGC computed from the whole cluster cell-set. We demonstrated the framework in several
simulation case-studies. Applying our framework in analyzing neonatal mouse heart single-cell
data [29], the detected biomarkers may characterize novel subtypes of cardiac muscle cells. We
named the framework PGC-RSMD (Polar Gini Curve – Root Mean Square Deviation). The
source code and dataset, including supplemental data, used in this manuscript could be found in
https://figshare.com/projects/Polar_Gini_Curve/76749.

83 Material and Method

84 Computing PGC-RSMD for one gene in one cluster

85 **Figure 1** demonstrates the workflow to compute PGC-RSMD for one gene in a cell cluster

86 from the single-cell expression data. Our approach used the 2D embedding [9] and clustering

87 results from single-cell expression data as the input. Starting from the 2D *x-y* embedding space,

88 for an arbitrary angle θ, the pipeline projects the x-y coordinate [30] *for every cell-point* onto the

89 θ-axis (*z* score)

$$z = x\cos(\theta) + y\sin(\theta) \quad (1)$$

90 We subtracted the scores from (1) with the smallest *z* to ensure that all *z* scores are non-negative,

91 which is the requirement for computing the Gini coefficient. Then, it computed two Gini

92 coefficients g_{sub} and g_{whole} to measure the inequality among the *z* scores. The g_{sub} coefficient only

93 used the distribution of *z* scores obtained from cells expressing the gene. The g_{whole} coefficient

94 would use the distribution of all *z* scores. The Gini coefficient formula is as in [28]

$$g = \frac{1}{2n^2\bar{z}}\sum_{i=1}^{n}\sum_{j=1}^{n}\left|z_i - z_j\right| \quad (2)$$

Here, *i* and *j* are arbitrary indices in the list of *z* scores being used in the computation, $\bar{z}$ is the

average of these *z* scores, and *n* is the size of the *z* score list. Repeating (1) and (2) for multiple

angles θ spanning from 0 to 2π would yield the corresponding lists between *g* and θ, as shown in

the bottom-right table in **Figure 1**. This would lead to two polar curves for Gini coefficients, one

for the cell-points expressing the gene in the cluster, and one for all cell-points in the cluster. We

hypothesize that *the two curves would be closer in the marker-gene scenario than in the non-*

marker gene scenario. Therefore, we used the root-mean-square deviation (RSMD) metric,

which is popular in computing fitness in Bioinformatics [31], to determine whether a gene is a

marker in the cluster.

$$RSMD = \frac{\sum_{\forall\theta}\left(g_{sub}(\theta) - g_{whole}(\theta)\right)^2}{n_\theta} \quad (3)$$

Here, n_θ, also called resolution, is the number of angles θ for which we repeat (1) and (2). In this

work, we chose $n_\theta = 1000$, which makes the angle list θ = 0, π/500, 2π/500, …, 999π/500, 2π.

To compute the RSMD statistical p-value for each gene in each cluster, first, we linearly

normalized (scaled) the RSMD computed in (3) such that the normalized RSMD is between 0

and 1. This could be done by diving (3) by the largest RSMD among all genes in each cluster.

Then, we applied the estimated p-value calculation in [32] to assign a p-value for each gene in

each cluster. Briefly, from the RSMD scores in (3), we verified that the RSMD scores followed a

bell-shaped distribution. Then, we computed the mean μ and standard deviation σ of the

normalized RSMD. Then, the p-value for each gene in the cluster is

$$p - value(i) = \frac{1}{\sqrt{2\pi\sigma^2}} \int_{-\infty}^{U} e^{-\frac{(u-\mu)^2}{2\sigma^2}} du \ (4)$$

In (4), *U* stands for the normalized RSMD.

Setting up simulation

In this work, to demonstrate how the PGC-RSMD functions, we setup two simulations. In the

first simulation, the cell cluster in the *x-y* embedding space had 5000 points, which were

uniformly generated in the unit circle $x^2 + y^2 \leq 1$. In the second simulation, the 2D visualization

of cell clusters had the shape identical to the real-world cluster obtained from visualizing the

mouse fetal lung single-cell data [29] using tSNE [33]. We applied the sampling-by-rejection

technique [34] to generate these cluster points as follows. In the first simulation, we randomly

generated a point whose coordinates are between -1 and 1 using uniform sampling, then accepted

the point if it had $x^2 + y^2 \leq 1$. In the second simulation, the random point coordinates were within

the cluster coordinate range. We extracted the cluster boundary points, compute the polygon

from these boundary points, which allowed deciding whether a point was inside the polygon

using Matlab [35, 36]. In each simulation, we randomly chose *m* percentage of points (*m* = 5, 10,

15, …, 95) and assumed that they represent the cells expressing gene. For each *m* percentage, we

repeat the simulation 1000 times.

In addition, to evaluate how the performance of PGC-RSMD would change in drop-out

scenario, we modified the single-cell data simulator in [37] as follow. First, we use [37] default

parameters to synthesize 2 clusters such that each cluster has 6000 cells, 250 markers (total 500

cluster markers) and other 4500 genes. For each cluster marker, the average expression fol-

change when comparing two clusters was between 4 and 1000. We assigned the drop-out

probability for each gene from 0, 5, 10, …, to 45% such that there were 25 markers for each
drop-out probability. Then, in each cell, we randomly change the marker expression to 0
according to the markers' drop-out probability. For each of the 4500 non-cluster markers, the
expression in each cell was randomly between 0 and 500. We assigned the sparisity – defined as
the percentage of non-expressing cells (0 expression) – for each non-cluster marker from 0, 5,
10, …. to 95%. In each cell, we randomly changed the non-cluster marker to 0 according to its
sparsity. We used the AUC metric to evaluate whether the PGC-RSMD score could differentiate
the 500 cluster-markers: whether each marker is specific for the first or the second clusters.

Identifying cardiac muscle cell clusters and marker genes from the neonatal mouse
heart single-cell data

We obtained the neonatal mouse heart single-cell case-study from the Mouse Cell Atlas [29].
We processed the data as specified in [29]. After preprocessing, the dataset covered 19,494 genes
expression in 5075 cells. We use tSNE [33] (without dimensional reduction) to embed the
dataset into the 2D space. We used the density-based clustering algorithm [38] implemented in
Matlab [39] to identify 9 cell clusters. In the implementation [39], we chose the clustering
parameters epsilon = 4, minpts = 40. There were 788, 397, 2966, 156, 288, 123, 76, 125, 87 cell-
points in cluster 1, 2, …, 9, correspondingly. There were 69 cell-points for which the algorithm
is unable to assign to any clusters (Supplemental Data 3).

We computed the percentage of expressing cells (the naïve approach) and PGC-RSMD for
all genes in all clusters. We removed genes expressing in less than 10% of the cluster cells. For
comparison, in the naïve approach, in each cluster, we selected the top genes sorted by the
highest percentage of expressing cells as the cluster markers. In the PGC-RSMD approach, we
selected the smallest-RSMD genes with its p-value < 0.05 as the cluster marker. In this work, we
focused on identifying the heart muscle cell clusters and their markers. We manually examine
the distribution of cells expressing the well-known heart muscle cell markers: *Myh7*, *Actc1*, and
Tnnt2 [40-47].

Seting up the re-identifying cluster ID problem

To compare the robustness of our PGC-RSMD markers with other approaches, we setup the

re-identifying cluster ID as follow. From the visual coordinates and 9 clusters of 5075 cells in

[29], we randomly divided the dataset into the training set (4060 cells – 80%) and the test set

(1015 cells – 20%) such that set has samples of all 9 clusters. Using the training set and markers'

expression found by PGC-RSMD, in comparison with other approaches, we applied the neural

network algorithm [48] to train models that identify cluster ID. We evaluated these models in the

test set and recorded the classification accuracy and area-under-receiving characteristic curve

(AUC). Here, we hypothesized that the 'better' markers would yield higher classification

accuracy and AUC. The other approaches being compared with PGC-RSMD are:

- The baseline approach: in this approach, we would train the classification models using all

genes expression.

- The differential expression approach: in this approach, we use Fisher's exact test [49],

which computes the likelihood of a gene being expressed (raw expression > 0) in a cluster,

compare to the likelihood of the gene being expressed outside the cluster. In this work, we select

the DEG markers in each cluster according to the following criteria: odd ratio > 5 and the

percentage of expressing cell (m) $> 50\%$.

- The SpatialDE [26] approach: SpatialDE finds the gene with high variance regarding the

distribution of 'point' on the spatial 2D space. The 'null' hypothesis in this approach is the gene

distribution in the 'spatial space' follows a multivariate normal distribution. The marker is

selected if the gene expression distribution is significantly different from the null distribution,

recorded in the p-value. In this work, we select the SpatialDE marker according to the following

criteria: q-value (adjusted p-value) < 0.05 and percentage of expressing cell (m) $> 50\%$.

In both the DEG and the SpatialDE approach, we sort the markers according to the decreasing

order of m. To make a fair comparison, we use the same number of markers, ranging from 5 to

100, found by PGC-RSMD, DEG and SpatialDE to train the classification models.

Results

PGC-RSMD strongly correlates to the percentage of expressing cell in a cluster

In **Figure 2**, we show that the fitness between the cluster PGC and the sub-cluster PGC strongly correlates to the percentage of expressing cell-points as the 'sub-cluster' m in the circle-shaped simulation. In addition, as m increases, the RSMD variance decreases. We represented the fitness by the **r**oot-**m**ean-**s**quare **d**eviation (RSMD) as showed in the method section. In this figure, for each m (from 5 to 95), we randomly generate 1000 sub-clusters and their PGCs. The detailed result of this simulation could be found at the Supplemental Data 1.

In addition, we observed a similar correlation when experimenting with the mouse fetal lung single-cell data [29]. **Figure 3a** shows the dataset clusters visualization using tSNE [33] and the chosen cluster. To synthesize a 3000-point cluster with the same shape to the chosen cluster, we still applied the random-by-rejection [34] as presented in the Material and Method section. **Figure 3b** still shows a strong correlation between m and RSMD. The detailed result of this simulation could be found at the Supplemental Data 2.

On the other hand, the PGC approach has the potential to answer whether the marker could identify subpopulations of cells in a cluster. **Figure 4a** demonstrates the 30000-point cluster with ring-shape $0.25 \leq x^2 + y^2 \leq 1$, which appears to be a sub-cluster marker. In this case, $m = 0.75$. In this example, RSMD $= 0.033$ (**Figure 4b**), which is greater than the RSMD distribution computed from the random and uniformly-distributed cluster with the same m (**Figure 4c**).

Figure 5 shows a decrease of PGC-RSMD performance in the drop-out scenario. Briefly, the synthetic data has 2 clusters, 250 distinct markers for each cluster. Each gene has a specific drop-out rate as presented in the Material and Method section. Using the PGC-RSMD scores in each cluster to differentiate these 500 the cluster-specific markers, we observed that PGC-RSMD achieves very high area-under-receiver-characteristic curve (AUC) (>0.95) when the drop-out probability is small ($\leq 5\%$). However, AUC decreases significantly with the probability of drop-out (**Figure 5a**). This phenomenon further demonstrates the strong association between RSMD and the percentage of expressing-cell. When the drop-out rate increases, the percentage of expressing-cell decreases; therefore, RSMD may mischaracterize a high-dropout marker as non-marker.

Case-study: PGC identifies heart muscle cell in neonatal mouse heart single-cell

PGC-RSMD detects markers to support cell-type identification in single-cell mouse neonatal heart data

Figure 6 summarizes the neonatal mouse heart single-cell data [29] and its 9-cluster markers. **Figure 6a** visualizes these 9 clusters with tSNE. The PGC-RSMD founds 258 genes, which are the union of the smallest 100-PCG-RSMD genes found in each cluster, marking these clusters (Supplemental Data 3). **Figures 6b** and **6c** showed that the gene-cluster marker-association reflects the underlying gene expression in the single-cell data. In these heatmap figures, each row corresponds to one gene.

We identified the muscle-cell clusters 1, 4 and 9 by the expression of *Myh7*, *Actc1*, and *Tnnt2*, which strongly express in muscle cell type [40-47] (**Figure 7**). Compared to the naïve method using the percentage of expressing cell, our PGC-RSMD is significantly better by detecting *Actc1*, which are missed by the naïve approach (**Figure 8**). Furthermore, our approach identified *Mgrn1* [50, 51], *Ifitm3* [52], *Myl6b* [53] marking cluster 1, which could play important roles in cardiac muscle functionality, heart failure, and heart development. These genes are not identified using the naïve approach (**Figure 8**). On the other hand, among genes having a high percentage of expressing cell, our PGC-RSMD suggests that *Ndufa4l2*, *Mdh2*, and *Atp5g1* may not be heart muscle cell markers. However, they could suggest a subtype of heart muscle cells (**Figure 9**). The percentage of expressing cells, PGC-RSMD, statistical p-value and ranks for all genes could be found in Supplemental Data 3.

Re-identifying the cells' cluster ID from markers

We observe that the markers found by the PGC-RSMD approach achieve better performance than the similar SpatialDE [26] markers, and similar performance to the differentially-expressed-gene (DEG) when being used to re-identify cell's cluster ID. Briefly, after computing the visual coordinate and cluster ID of all cells, we randomly split the dataset [29] into the training (80%) and test (20%) sets. We only applied the baseline PGC-RSMD, SpatialDE and DEG approaches to find the markers and built machine learning models to predict the cells' cluster ID from these

markers in the training set. In this experiment, we used all genes to train the predictor in the
baseline approach. The detailed description of this experiment could be found in the method
section. Evaluating the prediction models in the test set, the PGC-RSMD approach performs
closely to the DEG; both have cluster ID prediction accuracy above 0.9 and AUC above 0.95 on
average (**Figure 10**). These two approaches significantly outperform SpatialDE, whose accuracy
is just above the baseline.

Discussion

In this work, we show that integrating the embedded information, which does not often have
a deterministic relationship with gene expression and is primarily for clustering a visualization,
could lead to new insights to biomarkers in single-cell data. In the mouse neonatal heart case-
study, our PGC-RSMD approach could recall *Actc1* as the marker characterizing heart muscle
cell. Meanwhile, the approach using the ratio of expressing cell may fail to recall because a large
percentage of cells does not capture *Actc1* transcript. Therefore, our proposed technique has the
potential to handle analytical issues due to single-cell data quality, such as short-read and low
sequencing depth [54-56]. On the other hand, for genes having high percentage of expressing
cell, the PGC approach could further show that these genes may characterize novel cardiac
muscle cell sub-types for future studies, such as in *Mdh2* and *Myl6b*. Therefore, we suggest that
the biomarker discovery problem could be divided into two sub-problems: the 'global markers'
specify cell types and the 'local markers' specify subtypes. We could solve these two sub-
problems by the right integration of gene expression and visual information.

In addition to our proposed PGC approach, we could apply several alternative strategies to
integrate the gene expression and visual information to solve the single-cell biomarker discovery
problem. For example, the fractal dimension analysis strategies [57, 58], which focus on
evaluating the uniformity of cell-point distribution, could be applied to identify markers in which
the expressing cells distribute more densely than they are in the overall cluster. In addition, we
could also customize the statistical texture analysis in image processing, such as homogeneity
and integrity [59, 60], to analyze the difference between the overall cluster cell-point and cell-
expressing gene point as the metric to determine markers. On the other hand, choosing the

appropriate visual approach depends on the nature of the data and the problem. Our experiment
with the re-identifying cluster ID shows that the well-established SpatialDE [26] does not
outperform our approach and the DEG approach. One explanation is that in our problem, a good
marker for identifying cell type usually follows a good 'default' distribution over the visual
space; meanwhile, the SpatialDE aims to find markers that express significantly different from a
default distribution.

The major limitation of our proposed PGC-RSMD approach is the long computational time,
especially when comparing to the DEG approaches. This is similar to SpatialDE, which also used
visual information to detect marker genes. The DEG approaches may only need to compute one
statistical test to determine whether a gene is a marker for all clusters. Meanwhile, to draw the
curves, PGC-RSMD would need to compute hundreds to thousands, which depends on the
desired curve resolution, to characterize one gene in one cluster. Due to the long computational
time, we were not able to create multiple simulations, which is the ideal approach, run to
compute the statistical [32] p-value for the RSMD score. Therefore, we decided to reapply the
estimation presented to compute the p-value. This approach is computationally more efficient but
may not well-reflect the statistical characteristic of the single-cell data. In addition, we have not
fully tackled the problem of choosing the right threshold to determine whether a gene expresses
in a cell. Because of the strong association between PGC-RSMD and the percentage of
expressing-cell, we expect that the result would significantly different when choosing a different
threshold to determine whether a gene expresses in a cell. In this work, choosing 0 as the
threshold still yields good performance because of the high sparsity in the real dataset.

Conclusions

In this work, we have presented Polar Gini Curve, a novel technique to detect markers from
the single-cell RNA expression data using visual information. In principle, our technique could
complement the state-of-the-art approach: the PGC technique finds markers such that the
expressing cells are evenly distributed throughout the cluster space; meanwhile, the state-of-the-
art approach finds markers assuming a multivariate normal distribution of gene expression in the
visual space. We have demonstrated that the PGC technique performs better in some tasks in
single-cell analysis.

Authors' contribution

TN designed and implemented the core polar Gini algorithm, designed the sampling strategies in the simulation, performed the neonatal heart single-cell case-study, and primarily prepared the manuscript. JJ prepared the software package, preprocessed the single-cell data, and executed the simulation designs. NX designed different simulation scenarios, interpreted the statistical outcomes, and prepared the literature review. JC originated the idea of using Gini coefficient curves to integrate gene expression, cell-point distribution, and cluster shape to solve the biomarker discovery problem, designed the performance evaluation, and supervised the overall technical development. All authors reviewed, revised and approved the manuscript.

Competing interests

The authors have declared that no competing interests exist.

Acknowledgments

This work was supported by the 'startup budget' granted to Jake Chen from the University of Alabama at Birmingham.

References

1.   Zhu, Z., et al., Single-cell transcriptome in the identification of disease biomarkers: opportunities and challenges. J Transl Med, 2014. 12: p. 212.

2.   Finak, G., et al., MAST: a flexible statistical framework for assessing transcriptional
changes and characterizing heterogeneity in single-cell RNA sequencing data. Genome
Biol, 2015. 16: p. 278.
3.   McCarthy, D.J., et al., Scater: pre-processing, quality control, normalization and
visualization of single-cell RNA-seq data in R. Bioinformatics, 2017. 33(8): p. 1179-
1186.
4.   Poirion, O.B., et al., Single-Cell Transcriptomics Bioinformatics and Computational
Challenges. Front Genet, 2016. 7: p. 163.
5.   Jaakkola, M.K., et al., Comparison of methods to detect differentially expressed genes
between single-cell populations. Brief Bioinform, 2017. 18(5): p. 735-743.
6.   Wang, T., et al., Comparative analysis of differential gene expression analysis tools for
single-cell RNA sequencing data. BMC Bioinformatics, 2019. 20(1): p. 40.
7.   Love, M.I., W. Huber, and S. Anders, Moderated estimation of fold change and
dispersion for RNA-seq data with DESeq2. Genome Biol, 2014. 15(12): p. 550.
8.   Robinson, M.D., D.J. McCarthy, and G.K. Smyth, edgeR: a Bioconductor package for
differential expression analysis of digital gene expression data. Bioinformatics, 2010.
26(1): p. 139-40.
9.   Pezzotti, N., et al., Approximated and User Steerable tSNE for Progressive Visual
Analytics. IEEE Trans Vis Comput Graph, 2017. 23(7): p. 1739-1752.
10.   Becht, E., et al., Dimensionality reduction for visualizing single-cell data using UMAP.
Nat Biotechnol, 2018.
11.   Qiu, X., et al., Reversed graph embedding resolves complex single-cell trajectories. Nat
Methods, 2017. 14(10): p. 979-982.
12.   Yang, Y., et al., SAFE-clustering: Single-cell Aggregated (from Ensemble) clustering for
single-cell RNA-seq data. Bioinformatics, 2019. 35(8): p. 1269-1277.
13.   Kiselev, V.Y., et al., SC3: consensus clustering of single-cell RNA-seq data. Nat
Methods, 2017. 14(5): p. 483-486.
14.   Aibar, S., et al., SCENIC: single-cell regulatory network inference and clustering. Nat
Methods, 2017. 14(11): p. 1083-1086.
15.   Zheng, G.X., et al., Massively parallel digital transcriptional profiling of single cells. Nat
Commun, 2017. 8: p. 14049.

16.   Satija, R., et al., Spatial reconstruction of single-cell gene expression data. Nat
Biotechnol, 2015. 33(5): p. 495-502.

17.   Kiselev, V.Y., T.S. Andrews, and M. Hemberg, Challenges in unsupervised clustering of
single-cell RNA-seq data. Nat Rev Genet, 2019. 20(5): p. 273-282.

18.   Yuan, G.C., et al., Challenges and emerging directions in single-cell analysis. Genome
Biol, 2017. 18(1): p. 84.

19.   Conover, W.J. and W.J. Conover, Practical nonparametric statistics. 1980.

20.   Hollander, M., D.A. Wolfe, and E. Chicken, Nonparametric statistical methods. Vol. 751.
2013: John Wiley & Sons.

21.   Guo, M., et al., SINCERA: A Pipeline for Single-Cell RNA-Seq Profiling Analysis.
PLoS Comput Biol, 2015. 11(11): p. e1004575.

22.   Birnbaum, Z. On a use of the Mann-Whitney statistic. in Proceedings of the Third
Berkeley Symposium on Mathematical Statistics and Probability, Volume 1:
Contributions to the Theory of Statistics. 1956. The Regents of the University of
California.

23.   Korthauer, K.D., et al., A statistical approach for identifying differential distributions in
single-cell RNA-seq experiments. Genome Biol, 2016. 17(1): p. 222.

24.   Trapnell, C., et al., The dynamics and regulators of cell fate decisions are revealed by
pseudotemporal ordering of single cells. Nat Biotechnol, 2014. 32(4): p. 381-386.

25.   Kharchenko, P.V., L. Silberstein, and D.T. Scadden, Bayesian approach to single-cell
differential expression analysis. Nat Methods, 2014. 11(7): p. 740-2.

26.   Svensson, V., S.A. Teichmann, and O. Stegle, SpatialDE: identification of spatially
variable genes. Nat Methods, 2018. 15(5): p. 343-346.

27.   Edsgard, D., P. Johnsson, and R. Sandberg, Identification of spatial expression trends in
single-cell gene expression data. Nat Methods, 2018. 15(5): p. 339-342.

28.   Gini, C., Concentration and dependency ratios. Rivista di politica economica, 1997. 87:
p. 769-792.

29.   Han, X., et al., Mapping the Mouse Cell Atlas by Microwell-Seq. Cell, 2018. 173(5): p.
1307.

30.   Strang, G., et al., Introduction to linear algebra. Vol. 3. 1993: Wellesley-Cambridge Press
Wellesley, MA.

31.   Hyndman, R.J. and A.B. Koehler, Another look at measures of forecast accuracy.
410   International journal of forecasting, 2006. 22(4): p. 679-688.

411   32.   Yue, Z., et al., WIPER: Weighted in-Path Edge Ranking for biomolecular association
412   networks. Quantitative Biology, 2019. 7(4): p. 313-326.

413   33.   Maaten, L.v.d. and G. Hinton, Visualizing data using t-SNE. Journal of machine learning
414   research, 2008. 9(Nov): p. 2579-2605.

415   34.   Bishop, C.M., Pattern recognition and machine learning. 2006: springer.

416   35.   MathWorks.   Matlab   -   inpolygon.   2019   2019/06/05];   Available   from:
417   https://www.mathworks.com/help/matlab/ref/inpolygon.html.

418   36.   MathWorks.   Matlab   -   boundary.   2019   2019/06/05];   Available   from:
419   https://www.mathworks.com/help/matlab/ref/boundary.html.

420   37.   Baruzzo, G., I. Patuzzi, and B. Di Camillo, SPARSim Single Cell: a count data simulator
421   for scRNA-seq data. Bioinformatics, 2019.

422   38.   Ester, M., et al. A density-based algorithm for discovering clusters in large spatial
423   databases with noise. in Kdd. 1996.

424   39.   MathWorks.   Matlab   -   dbscan.   2019;   Available   from:
425   https://www.mathworks.com/help/stats/dbscan.html.

426   40.   Bashyam, M.D., et al., Molecular genetics of familial hypertrophic cardiomyopathy
427   (FHC). J Hum Genet, 2003. 48(2): p. 55-64.

428   41.   Finsterer, J., C. Stollberger, and J.A. Towbin, Left ventricular noncompaction
429   cardiomyopathy: cardiac, neuromuscular, and genetic factors. Nat Rev Cardiol, 2017.
430   14(4): p. 224-237.

431   42.   Keren, A., P. Syrris, and W.J. McKenna, Hypertrophic cardiomyopathy: the genetic
432   determinants of clinical disease expression. Nat Clin Pract Cardiovasc Med, 2008. 5(3):
433   p. 158-68.

434   43.   Morita, H., et al., Shared genetic causes of cardiac hypertrophy in children and adults. N
435   Engl J Med, 2008. 358(18): p. 1899-908.

436   44.   Jiang, H.K., et al., Reduced ACTC1 expression might play a role in the onset of
437   congenital heart disease by inducing cardiomyocyte apoptosis. Circ J, 2010. 74(11): p.
438   2410-8.

45. Kwon, C., et al., A regulatory pathway involving Notch1/beta-catenin/Isl1 determines
cardiac progenitor cell fate. Nat Cell Biol, 2009. 11(8): p. 951-7.

46. Wei, B. and J.P. Jin, TNNT1, TNNT2, and TNNT3: Isoform genes, regulation, and
structure-function relationships. Gene, 2016. 582(1): p. 1-13.

47. Ju, Y., et al., Troponin T3 expression in skeletal and smooth muscle is required for
growth and postnatal survival: characterization of Tnnt3(tm2a(KOMP)Wtsi) mice.
Genesis, 2013. 51(9): p. 667-75.

48. Russell, S. and P. Norvig, Artifical Intelligence: A Modern Approach. 2003. Prentice
Hall, Upper Saddle River, New Jersey.

49. Mehta, C.R. and N.R. Patel, A network algorithm for performing Fisher's exact test in r×
c contingency tables. Journal of the American Statistical Association, 1983. 78(382): p.
427-434.

50. Mukherjee, R. and O. Chakrabarti, Regulation of Mitofusin1 by Mahogunin Ring Finger-
1 and the proteasome modulates mitochondrial fusion. Biochim Biophys Acta, 2016.
1863(12): p. 3065-3083.

51. Liu, X., et al., Differential microRNA Expression and Regulation in the Rat Model of
Post-Infarction Heart Failure. PLoS One, 2016. 11(8): p. e0160920.

52. Lau, S.L., et al., Interferons induce the expression of IFITM1 and IFITM3 and suppress
the proliferation of rat neonatal cardiomyocytes. J Cell Biochem, 2012. 113(3): p. 841-7.

53. Wang, L., et al., Mutations in myosin light chain kinase cause familial aortic dissections.
Am J Hum Genet, 2010. 87(5): p. 701-7.

54. Shalek, A.K., et al., Single-cell RNA-seq reveals dynamic paracrine control of cellular
variation. Nature, 2014. 510(7505): p. 363-9.

55. Rizzetto, S., et al., Impact of sequencing depth and read length on single cell RNA
sequencing data of T cells. Sci Rep, 2017. 7(1): p. 12781.

56. McDavid, A., et al., Data exploration, quality control and testing in single-cell qPCR-
based gene expression experiments. Bioinformatics, 2013. 29(4): p. 461-7.

57. Fortin, C., et al., Fractal dimension in the analysis of medical images. IEEE Engineering
in Medicine and Biology Magazine, 1992. 11(2): p. 65-71.

58. Davies, S. and P. Hall, Fractal analysis of surface roughness by using spatial data. Journal
of the Royal Statistical Society: Series B (Statistical Methodology), 1999. 61(1): p. 3-37.

59. Bharati, M.H., J.J. Liu, and J.F. MacGregor, Image texture analysis: methods and
comparisons. Chemometrics and intelligent laboratory systems, 2004. 72(1): p. 57-71.
60. Kunimatsu, A., et al., Comparison between glioblastoma and primary central nervous
system lymphoma using MR image-based texture analysis. Magnetic Resonance in
Medical Sciences, 2017: p. mp. 2017-0044.

Figure legends

Figure 1. Overall workflow to compute the PGC-RSMD metric for one gene in one cluster of

cells. Here, the data points, histogram, and PGC for cells expressing the gene are cyan. The ones

for the whole cells in the cluster are red.

Figure 2. Boxplot showing a strong correlation between 'subcluster' percentage (m) and cluster-

subcluster PGC fitness (RSMD) in uniformly-distributed and a circular cluster.

Figure 3. a) The selected cluster for the experiment in [29]. b) Correlation between 'subcluster'

percentage (m) and cluster-subcluster PGC fitness (RSMD) in the selected cluster.

Figure 4. The ring-shape simulation study: a) Visualization of the cluster and ring-shape sub-

cluster ($m = 0.75$); b) PGC yield RSMD = 0.033; c) Distribution of RSMD, extracted from

Figure 2 with $m = 75\%$, when the sub-cluster uniformly distributed on the cluster area.

Figure 5. PGC-RSMD performance in recalling cluster marker in drop-out simulation. a)

heatmap showing the simulation design of 500 markers and 4500 neutral genes, with drop out /

percentage of cell expressing between 5 and 100%; b) The simulation data 2D visualization; c)

the AUC drops when drop-out increases.

Figure 6. The result from mouse neonatal heart single-cell [29] analysis. a) the tSNE plot shows

9 clusters. b) gene-cluster marker relationship (from 258 genes) found by PGC-RSMD; ■ gene

is found as marker, ■ gene is found as non-marker. c) expression heatmap for these genes.

Figure 7: Heart muscle cell clusters, identified by *Myh7*, *Actc1*, and *Tnnt2*

Figure 8. PGC-RSMD highlight makers that do not have high percentage of expressing cells:

PGCs of *Actc1*, *Mgrn1*, *Ifitm3*, *Myl6b* in cluster 1. The numbers in the parenthesis are ranks of

these genes in each metric

Figure 9. PGC-RSMD shows that gene haves high percentage of expressing cells: *Ndufa4l2*,

Mdh2, and *Atp5g1*, may not be markers in cluster 1. These genes appear to highlight a local

subcluster. The numbers in the parenthesis are ranks of these genes in each metric.

Figure 10. Performance of the PGC-RSMD, SpatialDE, and DEG in re-identifying the cell's

cluster ID problem using dataset [26]. The x-axis shows the number of top-significant markers

being selected to train the prediction models. a) accuracy; b) AUC over 9 clusters.

$$\frac{1}{2N^2}\sum_{a=1}^{N}\sum_{i=1}^{N}|z_i - z_t|$$

Angle θ	E_{min}	S_{min}	$S_{min} \times (E_{min}/S_{total})^3$
0°			
30°	0.26	0.29	0.63

$$RSMD = \sum f_0 / n_0$$

a)

median RSMD = -1.65×10^{-4} m + 0.015
R-square: 0.80
p-value: 2.06×10^{-7}

a) All cluster cells / Subcluster cells

a)

b)

c)

Cluster ID