Space-time logic of liver gene expression at sublobular scale

Colas Droin[1+], Jakob El Kholtei[2+], Keren Bahar Halpern[2+], Clémence Hurni[1], Milena Rozenberg[2], Sapir Muvkadi[2], Shalev Itzkovitz[2*], Felix Naef[1*]

[1]Institute of Bioengineering, School of Life Sciences, Ecole Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland

[2]Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot 7610001, Israel

[+]These authors contributed equally to this work.

*Corresponding authors:

Shalev Itzkovitz, Department of Molecular Cell Biology, Weizmann Institute of Science, Rehovot 7610001, Israel

Email: shalev.itzkovitz@weizmann.ac.il; +972 8 9343104:

Felix Naef, Institute of Bioengineering, School of Life Sciences, Ecole Polytechnique Fédérale de Lausanne, CH-1015 Lausanne, Switzerland

Email: felix.naef@epfl.ch; phone: +41 21 6931621

Abstract

The mammalian liver performs key physiological functions for maintaining energy and metabolic homeostasis. Liver tissue is both spatially structured and temporally orchestrated. Hepatocytes operate in repeating anatomical units termed lobules and different lobule zones perform distinct functions. The liver is also subject to extensive temporal regulation, orchestrated by the interplay of the circadian clock, systemic signals and feeding rhythms. Liver zonation was previously analyzed as a static phenomenon and liver chronobiology at the tissue level. Here, we use single-cell RNA-seq to investigate the interplay between gene regulation in space and time. Categorizing mRNA expression profiles using mixed-effect models and smFISH validations, we find that many genes in the liver are both zonated and rhythmic, most of them showing multiplicative space-time

effects. Such dually regulated genes cover key hepatic functions such as lipid, carbohydrate

and amino acid metabolism, but also genes not previously associated with liver zonation

such as chaperones. Our data also suggest that rhythmic and localized expression of Wnt

targets could be explained by rhythmically expressed Wnt ligands from non-parenchymal

cells near the central vein. Core circadian clock genes are expressed in a non-zonated

manner, indicating that the liver clock is robust to zonation. Together, our comprehensive

scRNA-seq analysis revealed how liver function is compartmentalized spatio-temporally

at the sub-lobular scale.

Introduction

The liver is a vital organ maintaining body physiology and energy homeostasis. The liver

carries out a broad range of functions related to carbohydrate and lipid metabolism,

detoxification, bile acid biosynthesis and transport, cholesterol processing, xenobiotics

biotransformation, and carrier proteins secretion. Notably, the liver performs catabolic and

anabolic processing of lipids and amino acids and produces the majority of plasma

proteins[1]. Liver tissue is highly structured on the cellular scale, being heterogeneous in

both cell-type composition and microenvironment[2]. In fact, liver tissue is made up of

millions of repeating anatomical and functional subunits, called lobules, which in mice

contain hepatocytes arranged in about 15 concentric layers with a diameter of about

0.5mm[3,4]. On the portal side of the lobule, blood from the portal vein and the hepatic

arteriole enters small capillaries called sinusoids and flows to the central vein. This is

accompanied with gradients in oxygen concentration, nutrients and signaling along the

portal-central axis, with the latter notably involving the Wnt pathway[5,6]. Due to this

polarization, hepatocytes in different layers perform separate functions. This is

accompanied with gradients of gene expression along the portal-central axis, with some

genes expressed more strongly near the central vein, and vice versa for portally expressed

genes. This phenomenon is termed *liver zonation*[1,7].

Recently, Halpern et al. combined single-cell RNA-sequencing (scRNA-seq) of

dissociated hepatocytes and single-molecule RNA fluorescence in situ hybridization

(smFISH) to reconstruct spatial mRNA expression profiles along the portal-central axis[8].

This analysis revealed an unexpected breadth of spatial heterogeneity, with ~50% of genes

showing spatially non-uniform patterns. Among them, functions related to ammonia clearance, carbohydrate catabolic and anabolic processes, xenobiotics detoxification, bile acid and cholesterol synthesis, fatty acid metabolism, targets of the Wnt and Ras pathways, and hypoxia-induced genes were strongly zonated.

In addition to its spatial heterogeneity, liver physiology is also highly temporally dynamic. Chronobiology studies showed that temporally gated physiological and metabolic programs in the liver result from the complex interplay between the endogenous circadian liver oscillator, rhythmic systemic signals, and feeding/fasting cycles[9,10,11]. An intact circadian clock has repeatedly been demonstrated as key for healthy metabolism, also in humans[12]. In addition, the hepatocyte clock has specifically been shown to play a major role in the physiological coordination of nutritional signals and cell-cell communication (including non-hepatocytic cells) controlling rhythmic metabolism[13]. Temporal compartmentalization can prevent two opposite and incompatible processes from simultaneously occurring, for example, glucose is stored as glycogen following a meal and is later released into the blood circulation during fasting period to maintain homeostasis in plasma glucose levels. Functional genomics studies of the circadian liver were typically performed on bulk liver tissue[14]. In particular, several studies showed how both the circadian clock and the feeding fasting cycles pervasively drive rhythms of gene expression in bulk, impacting key sectors of liver physiology such as glucose homeostasis, lipid and steroid metabolism[15–18].

Here, we asked how these spatial and temporal regulatory programs interact on the level of individual genes and liver functions more generally. In particular, can zonated gene expression patterns be temporally modulated on a 24 h time scale? And conversely, can rhythmic gene expression patterns observed in bulk samples exhibit sub-lobular structure? More complex situations may also be envisaged, such as time-dependent zonation patterns of mRNA expression (or, equivalently, zone-dependent rhythmic patterns), or sublobular oscillations that would escape detection on the bulk level due to cancelations. On the physiological level, it is of interest to establish how hepatic functions might be compartmentalized both in space and time. To study both the spatial and temporal axes, we performed scRNA-seq of hepatocytes at four different times along the 24 h day, extending a previous approach[8,19] to reconstruct spatial profiles at each time point. The resulting

space-time patterns were statistically classified using a mixed-effect model describing both

spatial and temporal variations in mRNA levels. In total, ~5000 liver genes were classified

based on their spatio-temporal expression profiles, and a few representative profiles were

further analyzed with smFISH. Overall, this approach revealed the richness of space-time

gene expression dynamics of the liver and provides a comprehensive view on how spatio-

temporal compartmentalization is utilized at the sub-lobular scale in the mammalian liver.

Results

Single-cell RNA-seq captures spatiotemporal gene expression patterns in mouse liver

To investigate spatio-temporal gene expression patterns in mouse liver, we sequenced

mRNA from individual liver cells obtained via perfusion from 10 *ad libitum* fed mice at 4

different times of the day (ZT = 0h, 6h, 12h and 18h, two to three replicates per time point).

The interval between ZT0 and ZT12 has the light on and corresponds to the fasting period

in mice, while feeding happens predominantly between ZT12 and ZT0. We here focused

on hepatocytes by enrichment of cells according to size and *in silico* filtering, yielding a

total of 19663 cells (Methods). To validate that the obtained scRNA-seq data captured the

expected variability in both spatial and temporal mRNA levels, we generated a clustering

analysis of all cells using a standard 2-dimensional t-SNE dimensionality reduction

(Methods) and colored cells either by their positions along the central-portal axis (the a

posteriori assigned layers, see below) (Figure 1A) or time (Figure 1B). The clustering

revealed that portally and centrally expressed landmark transcripts, such as the cytochrome

P450 oxygenases *Cyp2f2* and *Cyp2e1* involved in xenobiotics metabolism, mark cells in

opposite regions of the projections (Figure 1C-D). Likewise, time-of-day gene expression

varied along an orthogonal direction (Figure 1B), as shown for the fatty acid elongase

Elovl3 peaking at ZT0 (Figure 1E).

To obtain spatial mRNA expression profiles for each gene along the central-portal

axis, we here introduced eight lobule layers, to which we assigned each individual cell. For

this, we adapted a previous method that uses expression levels of landmark zonated genes

to define a central-portal coordinate[19], with the modification that only landmark transcripts

that were sufficiently expressed and that did not vary across mice and time points were

used (27 central and 28 portal landmark genes, Methods). The resulting reconstructed

(binned) mRNA expression profiles yielded 80 (8 layers over 10 mice) data points for each

transcript. Although our resolution is lower compared with the typically 12-15 hepatocyte

layers found in the liver[3,4], these reconstructions faithfully captured reference zonated

genes, with both central, and portal, expression (Figure 1F). Two examples of such genes

are the centrally expressed glutamine synthethase (*Glul*), and the portally expressed urea

cycle gene argininosuccinate synthetase (*Ass1*), showing mutually exclusive expression

along the lobule[8]. The reconstruction also successfully identified transcripts of the core

circadian clock, such as the master transcription factor *Bmal1* (also named *Arntl*), whose

mRNA peaked between ZT18-ZT0 (Figure 1G)[20]. In addition to core clock genes,

important clock outputs such as the PAR bZip transcription factor *Dbp,* which is a direct

transcriptional target of BMAL1 regulating detoxification enzymes, peaked between ZT6-

12 (Figure 1G)[21]. Finally, genes showing both zonated and rhythmic mRNA accumulation

were found (Figure 1H), for example elongation of very long chain fatty acids 3 (*Elovl3*)

is centrally expressed and peaks near ZT0, while phosphoenolpyruvate carboxykinase 1

(*Pck1*) regulating gluconeogenesis during fasting is expressed portally and peaks shortly

before ZT12. Since most of the zonated profiles showed exponential shapes, and gene

expression changes typically occur on a log scale[22], we log-transformed the data for further

analysis (Methods, Figure 1F and Supplementary Figure 1G-H-I). Together, these

examples indicate that the obtained gene expression profiles reliably capture spatial and

temporal regulation of hepatocyte gene expression.

Figure 1. A scRNA-seq approach to space-time gene expression in mouse liver.

PC ──────────────→ PP

(A-E) Global gene expression varies in both space and time, as shown using *t*-SNE

visualizations of the scRNA-seq (19663 hepatocytes from 10 mice). Each dot represents

one cell. Individual cells are colored by the (*a posteriori* assigned) lobule layer (A),

147	Zeitgeber time (B), expression levels of the zonated genes *Cyp2f2* and *Cyp2e1* (C-D), or
148	the temporally-regulated and centrally zonated gene *Elovl3* (E).
149	**(F-H)** Reconstructed spatial profiles (lobule layers 1-8) of selected zonated genes (F, top:
150	*Glul* pericentrally (PC) expressed, bottom: *Ass1* periportally (PP) expressed); rhythmic but
151	non-zonated genes (G, top: core-clock gene *Bmal1* peaking at ZT18-0, bottom: clock-
152	controlled *Dbp*, peaking at ZT6-12); zonated and rhythmic genes (H, Top: *Elovl3*, bottom:
153	*Pck1*). Expression levels correspond to fraction of total UMI per cell in linear scale. Log-
154	transformed profiles are in Supplementary Figure 1. Dots in F-H represent data points from
155	the individual mice. Shaded areas represent one standard deviation (SD) across the mice.

Space-time mRNA expression profiles categorized according to zonation and
rhythmicity

To gain a systematic understanding of the space-time gene expression profiles, we next
investigated if zonated gene expression patterns could be dynamic along the day, or
conversely whether temporal expression patterns might be zone-dependent. To select a
reliable set of reconstructed mRNA expression profiles for subsequent analyses, we filtered
out lowly expressed genes as well as genes with significant biological variability across
replicate liver samples, although this may be at the expense of a potentially decreased
sensitivity (Methods). This yielded 5058 spatio-temporal gene expression profiles
(Supplementary Figure 2A). An exploratory analysis of variance clearly identified zonated
genes, rhythmic genes, and fewer genes showing variability along both axes, with known
zonated and rhythmic genes distributed as expected (Figure 2A).

To identify possible dependencies between spatial and temporal variations, we built
a mixed-effect linear model[23] for the space-time mRNA profiles, which extends harmonic
regression to include a spatial covariate (Figure 2B). In this model, rhythms are
parameterized with cosine and sine functions, while spatial profiles are represented with
(up to second order) polynomials. In its most complex form, the model uses nine
parameters describing spatially modulated oscillations, and one intercept per mouse
(Methods). When some of the parameters are zero, the model reduces to simpler mRNA
profiles, for example purely spatial or purely temporal expression profiles (Figure 2C). We
then used model selection[24] to identify the optimal parameterization and category for each

gene (Methods). *In fine*, we classified each mRNA profile into one of five types of patterns

(Figure 2C). If only the intercept is used, the profile will be classified as flat or noisy (F/N,

Methods). If only time-independent zonation parameters are retained, the predicted profile

will be purely zonated (Z). If only layer-independent rhythmic parameters are retained, the

predicted profile will be purely rhythmic (R). If only layer-independent rhythmic

parameters and time-independent zonation parameters are retained, the profile is classified

as independent rhythmic-zonated (Z+R). If at least one layer-dependent rhythmic

parameter is selected, the profile will be termed interacting (ZxR). This classification

revealed that, overall, about 30% of the mRNA profiles were zonated (Z, Z+R and ZxR)

and about 20% were rhythmic (R, Z+R and ZxR) (Figure 2D). The peak times of these

rhythmic transcripts were highly consistent with bulk chronobiology data[25]

(Supplementary Figure 2B). The entire analysis can be browsed as a web-app resource

along with the corresponding data (https://czviz.epfl.ch).

Interestingly, we found that 7% of the analyzed genes in the liver were both zonated and

rhythmic. Such dually regulated transcripts represent 25% of all zonated transcripts, and

36% of all rhythmic transcripts, respectively. For example, the previously shown *Elovl3*

transcript*, involved in fatty acid elongation*, and *Pck1*, a rate limiting enzyme in

gluconeogenesis, are prototypical Z+R genes (Figure 1H, Supplementary Figure 1H).

Gluconeogenesis is an energetically-demanding task[3]. As mice are in a metabolically fasted

state requiring glucose production towards the end of the light phase (~ ZT10) and oxygen

needed for ATP production is most abundant portally[26], this process is indeed both spatially

and temporally regulated. The dual regulation of zonated-rhythmic genes may therefore

ensure optimal liver function under switching metabolic conditions.

Dually regulated genes were mostly Z+R, with only a minority of ZxR patterns.

The average expression across categories showed that rhythmic genes are significantly less

expressed on average than genes in zonated categories, likely reflecting shorter half-lives

(Figure 2E and Supplementary Figure 2C). Surprisingly, we find few highly expressed flat

genes. Together, we found that mRNA expression of many zonated genes in hepatocytes

is not static, and is in fact compartmentalized both in space and time.

Figure 2. Space-time mRNA expression profiles categorized with mixed-effect models.

(A) Spatial and temporal variations for mRNA transcript profiles, calculated as standard deviations (SD) of log2 expression along spatial or temporal dimensions. Colored dots correspond to reference zonated genes (orange) and reference rhythmic genes (blue) (Methods).

(B) Extended harmonic regression model for spatio-temporal expression profiles describing a static but zonated layer-dependent mean $\mu(x)$, as well as layer-dependent harmonic coefficients ($a(x)$ and $b(x)$). All layer-dependent coefficients are modeled as second order polynomials; i denotes the biological replicates. Temporal dependency is modelled with 24h-periodic harmonic functions. μ_i are random effects needed due to the data structure hierarchy (Methods).

(C) Schema illustrating the different categories of profiles. Depending on which coefficients are non-zero (Methods), genes are assigned to: F/N (flat or noisy, not represented), Z (zonated), R (rhythmic), Z+R (additive zonation and rhythmicity), ZxR (interacting zonation and rhythmicity). Graphs emphasize either zonation (top), with the x-axis representing layers, or rhythmicity (bottom), with the x axis representing time (ZT). Right side of the panel: two examples of fits (*Elovl3* and *Cyp7a1*, respectively Z+R and ZxR).

(D) Number of transcripts in each category.

(E) Boxplot of the mean expression per category shows that zonated genes (Z, Z+R and ZxR) are more expressed than rhythmic (R) or flat/noisy (F/N). ZxR genes are the most expressed according to median expression (orange line). Box limits are lower and upper quartile, whiskers extend up to the first datum greater/lower than the upper/lower quartile plus 1.5 times the interquartile range. Remaining points are outliers. KW test stands for Kruskal-Wallis.

Properties of dually zonated and rhythmic mRNA profiles

The majority of dually regulated genes are Z+R, which denotes additive (in log) space-time effects, or dynamic patterns where slopes or shapes of spatial patterns do not change with time (Figure 2C). On the other hand, interacting patterns (ZxR) are rare. Comparing the proportions of central, mid-lobular (peaking in the middle of the portal-central axis) and portal genes among the purely zonated genes (Z), and independently zonated and rhythmic genes (Z+R), did not reveal significant differences (Figure 3A), suggesting that rhythmicity is uncoupled with the direction of zonation. Similarly, comparing the phase distribution among the purely rhythmic genes (R) and the Z+R genes did not show a significant difference (Figure 3B), indicating that zonation does not bias peak expression

time. Moreover, oscillatory amplitudes were uncorrelated with the zonation slopes in Z+R

genes (Figure 3C). Finally, for ZxR genes with potentially more complex space-time

patterns, we investigated the spreads in amplitudes and peak times across the layers (Figure

3D). For wave-like pattern (phase modulated profiles), the phase difference across the

lobule was up to 3h, which corresponds to a difference in time between neighboring

hepatocytes on the order of 10 minutes for lobules of about 15 cell layers. On the other

hand, amplitude modulated patterns showed up to 2-fold difference in oscillatory amplitude

across the lobule.

To assess the potential physiological role of dually zonated and rhythmic

transcripts, we asked if protein levels of the identified Z+R and ZxR genes accumulated

rhythmically in a previous proteomics experiment[27]. In general, proteins rhythms are fewer,

damped, and time-delayed compared to mRNA rhythms due to protein half-lives[14,27,28]

(Discussion). However, while R transcripts were twice more frequent than Z+R transcripts,

the proportions were inverted for rhythmic proteins. Indeed, we found that among 65

rhythmic proteins (with q-value<0.2 in ref.[27]), 18 corresponded to Z+R and 10 to R

transcripts. Moreover, the identified Z+R and ZxR genes with rhythmic protein

accumulation cover key hepatic and zonated functions (Supplementary Figure 3, for a

functional interpretation, see below) and include rate limiting enzymes. For example, for

Z+R transcripts (Supplementary Figure 3A), PCK1 (rate limiting for gluconeogenesis),

LPIN2 (*Lipin2*, catalyzes the conversion of phosphatidic acid to diacylglycerol during

triglyceride, phosphatidylcholine and phosphatidylethanolamine biosynthesis), POR

(cytochrome P450 oxidoreductase, required to activate P450 enzymes), DNAJA1 (HSP40

co-chaperone), ALAS1 (rate-limiting for heme biosynthesis), GNE (rate-limiting in the

sialic acid biosynthetic pathway), THRSP (biosynthesis of triglycerides from medium-

length fatty acid chains), show robust rhythms at the protein level. Similarly, for ZxR

proteins, CYP7A1 (rate limiting enzyme in bile acid synthesis), CYP2A5 (coumarin 7-

hydroxylase), SLC1A2 (high-affinity glutamate transporter), and multidrug resistance

protein ABCC2 show rhythms on the protein level (Supplementary Figure 3B). Moreover,

the protein rhythms accompanying those Z+R and ZxR transcripts peak with an expected

delay of maximally about 6 hours[28] compared to the mRNA peak times (Supplementary

Figure 3C).

Figure 3. Properties of dually zonated and rhythmic mRNA profiles.

(A) Proportions of pericentral (green) and periportal (blue) transcripts are similar in Z and
Z+R. Mid-lobular genes (orange) are rare (<2%).

(B) Peak time distributions of rhythmic transcripts are similar in R and Z+R categories.

(C-D) Effect sizes of zonation (slope) *vs.* rhythmicity (fold-change amplitude, log2 peak-
to-trough) in Z+R genes (C). Magnitude of time shifts (delta phase, in hours) *vs.* fold-
change amplitude gradient (delta amplitude, in log2) along the central-portal axis in ZxR
genes (D). Genes for which the protein is also rhythmic (p<0.05) in bulk data are indicated
in dark red with the corresponding label (represented in Supplementary Figure 3).

smFISH analysis of space-time mRNA counts

To substantiate the RNA-seq profiles, we performed RNA single molecule fluorescence *in*

situ hybridization (smFISH) experiments on a set of selected candidate genes with diverse

spatio-temporal patterns. smFISH provides a sensitive and independent, albeit low-

throughput, measurement of mRNA expression. Purely zonated genes (Z) were already

well studied with smFISH[8]. To analyze the core-clock, we measured two genes peaking at

different times, *Bmal1* and *Per1*, which were classified as R in the RNA-seq analysis.

Bmal1 (~ZT0) and *Per1* (~ZT12) phases were nearly identical in both experiments, and the

rhythms did not depend on the lobular position consistent with R genes (Figure 4A). We

analyzed three genes classified as Z+R: *Pck1* was indeed both portally biased and rhythmic

in RNA-seq and smFISH (Figure 4B); *Elovl3* is both centrally biased and rhythmic in

RNA-seq and smFISH, even though the amplitude of the oscillations was damped on the

portal side in the FISH experiment (Supplementary Figure 4A); and for *Arg1 (Arginase 1)*

the portal RNA-seq and smFISH profiles matched well (Supplementary Figure 4B).

Finally, *Acly* showed a pattern in smFISH data which validates its classification as ZxR,

with a lower amplitude on the portal side, where the transcript is more highly expressed

(Figure 4C). Thus, overall, the reconstructed scRNA-seq and smFISH profiles were

consistent, with minor discrepancies.

Figure 4. smFISH analysis of rhythmic and zonated transcripts.

308	**(A)** smFISH (RNAscope, Methods) of the core clock genes *Bmal1* and *Per1* (both assigned
309	to R) in liver slices sampled every 3 hours. **Left:** representative images at ZT0, ZT06, ZT12
310	and ZT18 for *Bmal1*. Central vein (CV) and a portal node (PN) are marked. Scale bar is
311	50μm. Endothelial cells lining the PC and cholangiocytes surrounding the PP were
312	excluded from the quantification. mRNA transcripts and nuclei were detected in PN and
313	PC zones (Methods). **Right:** temporal profiles of *Bmal1l* and *Per1* from smFISH (top,
314	quantification of the number of mRNA transcripts at 8 time points ZT0 to ZT21, every 3
315	hours, shaded area indicate SD across images), in PN and PC regions, and scRNA-seq
316	(bottom, shaded areas is SD across mice).
317	**(B-C)** smFISH (Stellaris, Methods) for *Pck1* (Z+R) and *Acly* (ZxR). smFISH
318	quantifications were made for ZT0 and ZT12 (Methods). **Left:** representative images at
319	ZT0, ZT12 for *Pck1* (B) or *Acly* (C). Central vein (CV) and a portal node (PN) are marked.
320	Scale bar - 20μm. **Right:** quantified profiles for each gene in the two time points from
321	smFISH (top, shaded area indicate SD across images), and scRNA-seq data at the four time
322	points (bottom, shaded areas is SD across mice).

Space-time logic of hepatic functions

We next used our classification to explore the spatio-temporal dynamics of hepatic

functions and signaling pathways in the liver. Given the prevalence of zonated gene

expression profiles, we first analyzed if the circadian clock is sensitive to zonation. We

found that profiles of reference core-clock genes (Supplementary Figure 5) were assigned

to the rhythmic only category (R), except for *Cry1* and *Clock* that were assigned to Z+R,

but with high probabilities also for R (Supplementary Table 2). This suggests that the

circadian clock is largely non-zonated, as also seen in the smFISH (Figure 4A), and

therefore robust to the heterogenous hepatic microenvironment.

We then systematically explored enrichment of biological functions in the zonated

category by querying the KEGG pathways database (Supplementary Table 3, Figure 5,

Methods). In addition to recapitulating well documented zonated liver functions[8,29], which

we do not discuss here, this analysis highlighted Z and Z+R functions which, to our

knowledge, had not been linked with liver zonation. For instance, we found that cytosolic

chaperones accumulate centrally, while the endoplasmic reticulum (ER) chaperones linked

with protein secretion accumulate portally (Figure 5B,D, Supplementary Figure 6). Both

groups of chaperones peak during the activity/feeding phase, probably due to body

temperature rhythms peaking during the active phase[30,31], and likely reflect increased needs

of protein folding during times of high protein synthesis. Also, we found that mRNAs of

ribosomal protein genes accumulate centrally (Z), as do proteasome components (Figure

5A), the latter also containing rhythmic members (Z+R). In the liver, ribosomal proteins

are rate-limiting for the synthesis of ribosomes, which themselves are rate-limiting for the

synthesis of proteins[32]. Therefore, the overall protein synthesis rate is likely higher in these

hepatocytes. Conversely, transcripts encoding components of the proteasome are involved

in protein degradation. Together, these observations suggest that protein turnover is higher

in centrally located hepatocytes, which are exposed to an environment with high

concentrations of xenobiotics and hypoxic stress[33].

In addition, while many mitophagy genes are expressed centrally (Z), some of those

also show robust temporal rhythms peaking during the fasting period (Z+R), in particular

two gamma-aminobutyric acid receptor-associated proteins (*Gabarap* and *Gabarapl1*)

with an important function in autophagosome mediated autophagy[34]. Consistent with this

temporal regulation, we had previously reported that the nuclear abundance of the fasting-

dependent regulators of autophagy TFEB and ZKSCAN3 peaked near ZT6[35]. Moreover,

the centrally and synchronously accumulating Ubiquitin B mRNA (*Ubb*, Z+R) may

contribute to triggering mitophagy[36]. Thus, centrally biased mitophagy may both

participate in removal of damaged mitochondria in the stressed central environment.

Similarly, genes involved in bile acid synthesis and bile secretion are known to

show zonated expression patterns[8] (Figure 5C). Here, we found that while the rate-limiting

enzyme in the bile acid biosynthetic pathway *Cyp7a1* (ZxR, Figure 2C) is known to be

clock-controlled, with its mRNA[37,38] and protein[39] (Supplementary Figure 7D) expressed

maximally early during the feeding period, the ABC transporters Sterolin-1 and 2 (ABCG5/

ABCG8, both identified as Z+R) which excrete most of the biliary cholesterol[40] peak

towards the end of the fasting period near ZT9.

Many detoxification enzymes of the cytochromes P450 (CYPs) superfamily are

known to be centrally zonated in the liver[8] and several of those are found in the Z+R

category. In particular, the Flavin-containing monooxygenases Fmo1, 2, and 5, which are

NADPH-dependent monooxygenases involved in drug and xenobiotics detoxification, exhibit Z+R mRNA patterns with peaks near ZT16 (Figure 5C). Also, the FMO5 protein accumulates rhythmically (Figure 3C, Supplementary Figure 3). Moreover, the rate limiting enzyme ALAS1 producing the P450 cofactor heme was found as a centrally zonated Z+R transcript with peak mRNA at ZT13, showing also a robust rhythm in protein expression (Supplementary Figure 3).

To substantiate the above finding on heat shock genes, we examined the space-time behavior of temperature-regulated genes. To this end, we considered targets (bound in ChIP-seq) of the heat shock transcription factor HSF1 (Chip-Atlas[41], includes our own liver data[16]). This showed that several targets, known to peak during the active phase[30,31], are also zonated (Supplementary Figure 6). Notably, the cytoplasmic *Hsp90aa1* (Hsp90A chaperone) and its interactor *Dnaja1* (HSP40), respectively mitochondrial *Hspd1* (HSP60), chaperones are expressed centrally where protein turnover is high. On the contrary, endoplasmic reticulum (ER) located chaperones: *Hspb1* (Hsp90B) and interactor *Pdia3*, as well as *Hspa5* (HSP70) and ER-resident *Dnajc3* and *Dnajb11* (DNAJ/HSP40) are expressed portally, consistent with their role in folding proteins in the secretory pathway (secretion is known to occur portally[29]). On the other hand, the analysis of cold induced genes (i.e. CIRBP and analogous, taken from ref.[42]) did not show zonated gene expression.

Finally, we note that among all KEEG pathway related to lipids, a majority show central enrichment (Figure 5A, Table S3). Inspection of the genes involved shows that this is due to the large number of genes related to peroxisomal β-oxidation, i.e. lipid catabolism, which are incidentally also listed in biosynthesis KEGG pathways (Table S3). However, fatty acid synthesis is biased portally, as supported by key portally expressed genes such as *Fasn*, *Srebf1*, *Acly*, *Acaca*, *Elovl2*, *Elovl5*, and which is consistent with the fact that oxygen needed for mitochondrial β-oxidation is most abundant portally[26]. On the other hand, *Elovl3*, which is known to be transcriptionally controlled by the peroxisome regulator PPARα and atypically regulated among Elov family fatty acid elongases[43] is expressed centrally.

Figure 5. Space-time logic of compartmentalized hepatic functions for Z+R genes.

red gradient. *: appears central due to the KEGG annotation system; however, fatty acid

elongation is biased portally (see text).

(B-C-D) KEGG analysis of Z+R genes. Representations of genes in central (B-C) and

portal (D) enriched Z+R categories (Supplementary Table 3). Polar representation: peak

expression times are arranged clockwise (ZT0 on vertical position) and amplitudes (log2,

values indicated on the radial axes) increase radially. The radius coordinate of genes with

an amplitude >0.9 is halved (indicated with a black circle around the colored dot).

Space-time logic of activity of signaling pathways

Signaling pathways that include Wnt, Ras and hypoxia have been shown to shape

hepatocyte zonation[8]. We therefore examined the space-time activities of these pathways,

extracted from the behavior of canonical target genes. We mainly focused on Wnt, as it is

often considered the master regulator of liver zonation[44]. To systematically investigate

spatio-temporal WNT/β-Catenin activity in liver, we extracted a set of Wnt targets derived

from an APC KO in mouse liver[8,45]. We found that rhythmic transcripts (in the R, Z+R and

ZxR categories) are enriched among targets of the Wnt pathway, showing a proportion that

increases with the strength of the targets, with the strongest Wnt targets containing 80% of

rhythmic transcripts (Supplementary Figure 7A). Positive Wnt targets were pericentral[8]

and peaked between ZT9 and ZT12, whereas negative Wnt targets were periportal[8] and

peaked between ZT21 and ZT3 (Figure 6A).

To obtain a temporal view of Wnt activity, we considered the top 50 Wnt pathway targets

(according to the liver APC KO data) in the liver and analyzed the temporal profiles from

high temporal resolution bulk liver mRNA and from our scRNA-seq binned in three

different zones: central (layers 1-2), mid-lobular (layers 3-4-5) and portal (layers 6-7-

8)(Figure 6B-C, Supplementary Figure 7B-C). This analysis confirmed that the peak times

of rhythmic Wnt targets is preferentially between ZT9 and ZT12, and that the bulk and

single cell data are consistent with each other, despite of the lower temporal sampling of

the scRNA-seq. In fact, among the rhythmic genes detected in bulk RNA-seq, the five

strongest Wnt targets were, in decreasing order: *Axin2*, *Glul*, *Slc1a2*, *Tuba8*, *Rnf43*, with

the latter showing the largest amplitude (Figure 6C); all but *Tuba8* peaked in the morning.

Note that *Glul*, an important marker of central zonation and canonical Wnt target (just like

Axin2 and *Rnf43*), was assigned to the Z category, but with second highest probability for

ZxR (Supplementary Table 2), peaking at ZT12. Further evidence of rhythmic WNT/β-

Catenin activity was provided by our previous proteomics data[35] showing that the potent

Wnt effector TCF4 (encoded by the *Tcf7l2* gene) has rhythmic nuclear abundance in mouse

liver with a peak phase at ZT7.5 (Figure 6D), and hence explains the accumulation of its

mRNA targets a few hours later.

In addition to the mRNA rhythms, we found that several of the Z+R or ZxR Wnt

targets showed clear rhythms in bulk proteomics with the characteristic phase delays. The

strongest five targets with rhythmic proteins (Supplementary Figure 7D) included the rate-

limiting enzyme in the bile acid biosynthetic pathway CYP7A1, the NADPH-dependent

monooxygenases involved in drug and xenobiotics detoxification FMO5, the P450

detoxification enzymes coumarin 7-hydroxylase (*Cyp2a5*), which may protect mice from

dietary coumarin-induced toxicity[46], the high-affinity glutamate transporter Slc1a2

(*Eaat2*), and the multidrug resistance protein ABCC2. All these proteins showed high

amplitude protein rhythms peaking during the feeding phase between ZT12 and ZT18.

Thus, Wnt transcription activity is clearly rhythmic in the liver, and this rhythm can

propagate to protein expression.

We next asked whether the temporal oscillations in the expression of Wnt-activated

genes might correlate with temporal oscillations in Wnt morphogens produced by

pericentral non-parenchymal liver cells. To this end, we performed smFISH experiments

and quantified the expression of the Wnt ligand *Wnt2*[5] and of *Rspo3*[6,47], a critical facilitator

of Wnt signaling, as well as the Wnt antagonist *Dkk3*[19] (Figure 6E). We found that both

Wnt2 and *Rspo3* in liver non-parenchymal cells (NPCs) exhibit non-uniform expression

around the clock, with significantly higher mRNA levels at ZT0 ($p=4\times10^{-5}$ for *Wnt2*, and

$p=2\times10^{-8}$ for *Rspo3*, Kruskal-Wallis, Figure 6E, right). Given various delays between

mRNA accumulation of ligands and expression of the Wnt targets, this timing is

compatible with the peak nuclear accumulation of the TCF4 (Tcf7l2 gene) transcription

factor observed at ZT7.5 (Figure 6D) and with the peaks in Wnt-activated genes between

ZT6-12 (Figure 6A-B). Differences in *Dkk3* expression were not significant (p=0.053).

Thus, production of Wnt morphogens by central non-parenchymal liver cells might

underlie the observed rhythmic Wnt pathway activity.

Figure 6. Rhythmic activity of Wnt signaling.

(A) Enrichment/depletion at different times (window size: 3h), of both positive and negative Wnt targets (background: all R and Z+R genes). Colormap shows p-values (two-tailed hypergeometric test): red (blue) indicates enrichment (depletion).

(B) Heatmaps representing scRNA-seq profiles of the top 50 Wnt targets (according to the Apc-KO fold change, *Lgr5* was also added) showing rhythmic mRNA in bulk (p<0.01, ref.[25]). The profiles are computed in three different zones of the central-portal axis: central (layers 1-2), mid-lobular (layers 3-4-5) and portal (layers 6-7-8). Gene profiles (log2) are mean-centered in each zone. An enrichment of the phases around ZT8-14 can be observed, in agreement with Figure 6A.

(C) mRNA profiles from bulk RNA-seq (top) and scRNA-seq (bottom). Top five targets with the highest Apc-KO fold change, and rhythmicity in the bulk data (p<0.01, harmonic regression). Rhythmicity (indicated above each panel) is computed on bulk data or in three different zones for scRNA-seq data.

(D) Nuclear protein abundance from ref.[35] of the Wnt effector TCF4 (encoded by the *Tcf7l2* gene) in mouse liver shows a rhythm (p=0.003) peaking at ZT7.5, consistent with the accumulation of mRNA targets a few hours later (panel A).

(E) Left: Representative smFISH images of *Wnt2*, *Dkk3*, and *Rspo3* expression at ZT0 (left) and ZT18 (right), shown in green. Markers of non-parenchymal cells (NPCs) are shown in red (Methods). Nuclei are stained in blue (DAPI). Scale bars, 2 μm; **Right:** Violin plots representing quantitative analysis of smFISH images (1420 cells of 189 central veins of at least two mice per time point). *Wnt2*, *Dkk3*, and *Rspo3* transcripts are quantified in NPCs lining the central vein (Methods). mRNA expression is in smFISH dots per μm^3.

Ras signalling and hypoxia are two additional pathways that have been implicated in shaping hepatocyte zonation[8]. In agreement with ref.[8], we found that the negative targets of Ras were enriched in central genes, whereas the positive Ras targets were enriched in portal genes (not shown). The rhythmic targets (R and Z+R) of hypoxia showed a pattern of temporal compartmentalization similar with those of Wnt (Supplementary Figure 7E): the negative targets were enriched around ZT0 (dark-light transition) and underrepresented around ZT14, while the positive targets were enriched around ZT10 and underrepresented around ZT3. Ras targets, positive or negative, did not exhibit significant temporal bias.

Discussion

Recent genome-wide analyses of zonated gene expression in mouse and human
liver[8,48,49] uncovered a rich organization of liver functions in space at the sublobular scale,
while chronobiology studies of bulk liver tissue revealed a complex landscape of rhythmic
regulatory layers orchestrated by a circadian clock interacting with feeding-fasting cycles
and systemic signals[35,50–52]. Here, we established how these two regulatory programs
combine to shape the daily space-time dynamics of gene expression patterns and
physiology in adult liver by extending our previous scRNA-seq approach[8]. We found that
liver uses gene expression programs with many genes exhibiting compartmentalization in
both space and time.

In this study, we chose to focus on the parenchymal cells in the liver, the
hepatocytes, for which smFISH data on landmark zonated genes was readily available,
which enabled reconstructing spatio-temporal mRNA profiles from scRNA-seq[8]. Zonation
profiles of other cell types in the liver may be obtained as well; in fact, static zonation
mRNA expression profiles have been obtained for liver endothelial cells, using a paired-
cell approach[19] in which mRNA from pairs of attached mouse cells were sequenced and
gene expression from one cell type was used to infer the pairs' tissue coordinates. In
addition, *ab initio* reconstruction methods such as diffusion pseudo time[48] or novoSpaRc[53],
in which a zonation coordinate is inferred by assuming that the major axis of variability for
a cell type reflects transcriptome-wide gene expression changes associated with zonation,
could be used for spatially sparse cell types with no available zonated marker genes, e.g.
stellate or resident immune Kupffer cells. Moreover, it was recently found that rhythmic
gene expression and metabolism in non-hepatocyte cells can be driven both by clocks in
hepatocytes via cell-cell communication as well as feeding cycles[13]. Our computational
framework for analyzing space-time logic of gene expression could be widely applicable
in such future studies.

To study whether the observed space-time expression profiles may be regulated by
either liver zonation, 24h rhythms in liver physiology, or both, we developed a mixed-
effect model, combined with model selection. This enabled classifying gene profiles into
five categories representing different modes of spatio-temporal regulation, from flat to
wave-like. To validate these, we performed smFISH in intact liver tissue, which showed

largely compatible profiles although some quantitative differences were observed. These

differences most likely reflect the lower sensitivity of RNA-seq, uncertainties in the spatial

analysis of smFISH in tissues, as well as known inter-animal variability in the physiologic

states of individual livers, notably related to the animal-specific feeding patterns[25].

Together, this temporal analysis confirms that a large proportion of gene expression

in hepatocytes is zonated[8] or rhythmic[17], and in addition reveals marked spatio-temporal

regulation of mRNA levels in mouse liver (Z+R and ZxR genes, comprising 7% of all

detected genes according to our criteria). This means that zonated gene expression patterns

can be temporally modulated on a circadian scale, or equivalently, that rhythmic gene

expression profiles can exhibit sub-lobular structure. The dominant pattern for dually

regulated gene was Z+R, which corresponds to additive effects of space and time in log, or

multiplicate effects of gene expression levels, and describes genes expression profiles that

are compartmentalized in both space and time. In other words, such patterns are

characterized by shapes (in space) that remain invariant with time, but whose magnitudes

are rhythmically rescaled in time. Or equivalently, the oscillatory amplitude (fold change)

and phases are constant along the lobular coordinate, but the mean expression is patterned

along the lobule. Such multiplicative effects could reflect the combined actions of

transcriptional regulators for the zone and rhythm on promoters and enhancers of Z+R

genes. Indeed, gene expression changes induced by several regulators combine

multiplicatively[22]. Note that though the (relative) shape of Z+R patterns is invariant in time,

threshold-dependent responses that would lie downstream of such genes would then

acquire domain boundaries which can shift in time. Similar phenomena are expected for

interacting profiles (phase and amplitude modulated) (ZxR) that we observed for a smaller

number of genes.

As shown by us and others[27,28], rhythms at the protein level are typically damped

and phase-delayed compared the cognate mRNA rhythms, depending on the protein half-

lives. Indeed, longer protein half-lives imply smaller oscillatory amplitudes and longer

delays between mRNA and protein accumulation[14]. Analysis of liver proteomes in bulk

showed that the number of cyclic proteins is significantly lower than the number of cyclic

mRNAs, with delays approaching the predicted maximum of six hours. Here, we found

genes from both Z+R and ZxR that exhibited rhythmic accumulation in bulk proteomics

experiments, including genes encoding rate-limiting enzymes, suggesting that dually space-time regulated patterns have a physiological role in the liver. Moreover, phase delays between the mRNA and protein profiles were as expected. Future studies will utilize emerging spatial proteomics approaches to reconstruct a space-time liver proteomic atlas[54].

In addition to previously discussed zonated liver functions[8], a systematic querying of KEGG pathways highlighted Z+R functions not previously associated with rhythmic liver zonation. The roles and profiles of the corresponding genes allowed us to better understand the spatiotemporal logic of the identified pathways. For instance, we found that the expression levels of both ribosome protein genes (rate-limiting for protein synthesis[32]) and proteasome components (involved in protein degradation) were higher in central hepatocytes. Since the central environment is subject to high concentrations of xenobiotics and hypoxic stress, this could indicate an elevated protein turnover in this region, which would ensure that damaged proteins are rapidly exchanged with new, undamaged proteins. This interpretation is corroborated by the observed increased levels of cytosolic and ER chaperones during the feeding phase, to assist protein synthesis and secretion, thereby counteracting such protein stress.

It was previously shown that Wnt signaling can explain the zonation of up to a third of the zonated mRNAs[7]. Wnt ligands are secreted by pericentral non-parenchymal cells, mostly endothelial cells[5,6], forming a graded spatial morphogenetic field. As a result, and as observed in our enrichment analysis, Wnt-activated genes were pericentrally-zonated. Moreover, both the scRNA-seq data and previous bulk mRNA and protein measurements showed that Wnt activity is rhythmic in the liver. Our smFISH analysis suggested that temporal fluctuations in the expression of *Wnt2* and *Rspo3*, two key Wnt ligands, secreted by pericentral non-parenchymal cells might underlie oscillatory and zonated expression of Wnt targets at times near the fasting/feeding transition.

In summary, we demonstrate how liver gene expression can be quantitatively investigated with spatial and temporal resolution and how liver functions are compartmentalized along these two axes. Our approach could be used to reconstruct spatio-temporal gene expression patterns in other zonated tissues such as the intestine and kidney[3].

Material and methods

Animals and ethics statement

All animal care and handling were approved by the Institutional Animal Care and Use Committee of WIS and by the Canton de Vaud laws for animal protection (authorization VD3197.b). Male C57bl6 mice aged of 6 weeks, housed under reverse-phase cycle and under ad libitum feeding were used to generate sc-RNA-seq data of hepatocytes and single-molecule RNA-FISH (smFISH). Male mice between 8 to 10 weeks old, housed under 12:12 light-dark cycle, and having access to food only during the night (restricted-feeding) were used for smFISH of circadian clock genes.

Hepatocytes isolation and single-cell RNA-seq

Liver cells were isolated using a modified version of the two-step collagenase perfusion method of Seglen[55]. The tissue was digested with Liberase Blendzyme 3 recombinant collagenase (Roche Diagnostics) according to the manufacturer instructions. To enrich for hepatocytes, we applied a centrifuge step at 30g for 3 min to pull down all hepatocytes while discarding most of the non-parenchymal cells that remained in the sup. We next enriched for live hepatocytes by 2 cycles of percoll gradient, hepatocytes pellet was resuspended in 25 ml of PBS, percoll was added for a final concentration of 45% and mixed with the hepatocytes. Dead cells were discarded after a centrifuge step (70g for 10min) cells were resuspended in 10x cells buffer (1x PBS, 0.04% BSA) and proceeded directly to the 10x pipeline. The cDNA library was prepared with the V2 chemistry of 10X genomics Chromium system according to manufactures instructions and sequencing was done with Illumina Nextseq 500 at estimated depth of 40,000 reads per cell.

Conceivably, the dissociation of liver tissue into individual cells and the purification of hepatocytes are relatively lengthy and may thus lead to alterations in mRNA expression. While it has been shown that mRNA levels do not change much during the purification and 24-hour cultivation of hepatocytes[56], transcription rates on the other hand can be diminished by 8-fold to nearly 100-fold during this process[57]. This difference between nascent transcript and mature mRNA levels can be explained by the relatively long half-lives of liver-specific RNAs. In our case, the time needed from the dissociation of the tissue

until the cell lysis is approximately 1 hour and the cell are not placed in culture. Since we

are measuring mature transcripts, with half-lives in the range of typically 1-5 hours

(Supplementary Figure 2C), the changes in mRNA levels due to the protocol will remain

contained. In particular, the scRNA-seq of cells carry the typical hepatocyte gene

expression signatures, for example, genes such as Alb or Apoa2 rank at 2nd and 5th

position genome-wide. As further validation, we compared our reconstructed gene

expression zonation profiles with the zonation profiles from the massively validated

zonation study of ref.[8], revealing a near-perfect agreement (Supplementary Figure 1F).

Filtering of raw scRNA-seq data

The initial data analysis was done in R v3.4.2 using Seurat v2.1.0[58]. Each expression matrix

was filtered separately to remove dead, dying and low-quality cells. We firstly only kept

genes that were expressed in at least 5 cells in any of the ten samples. We then defined a

set of valid cells with more than 500 expressed genes and between 1000 and 10000 unique

molecular identifiers (UMIs) and secondly an additional expression matrix with cells

having between 100 and 300 UMIs which was used for background estimation

(Supplementary Figure 1A). Other UMI-filters have been tried, but yielded equal or less

reliable profiles. The mean expression of each gene was then calculated for the background

dataset and subtracted from the set of valid cells. This was subsequently filtered to only

include hepatocytes by removing cells with expression of non-parenchymal liver cell

genes. Next, the cells were filtered based on the fraction of mitochondrial gene expression.

First, expression levels in each cell were normalized by the sum of all genes excluding

mitochondrial and major urinary protein (*Mup*) genes. Indeed, as mitochondria are more

abundant in periportal hepatocytes, the expression of mitochondrial genes is higher in this

area[59]; and since these genes are very highly expressed, including them would reduce the

relative expression of all other genes based on the cell's lobular location. *Mup* genes are

also highly abundant and mapping their reads to a reference sequence is unreliable due to

their high sequence homology[60]. Moreover, *Mup* genes encode for pheromones that vary

greatly between individuals to facilitate individual recognition[61].

Mitochondrial content is often used to remove non-viable cells[62]. The mitochondrial

content of sequenced hepatocytes exhibited a bi-modal behavior (Supplementary Figure

1B). To identify the range of mitochondrial fractions that included viable hepatocytes we

used an intrinsic property of hepatocytes, which is the anti-correlation of the pericentral

landmark gene *Cyp2e1* and the periportal landmark gene *Cyp2f2*[7] (Supplementary Figure

1C). We found that hepatocytes with mitochondrial fraction in the range of 9-35%

exhibited an almost perfect anticorrelation between *Cyp2e1* and *Cyp2f2* (Supplementary

Figure 1D,E), suggesting that these are the best quality, and we consequently kept

hepatocytes within this range of mitochondrial content for further analysis.

t-SNE clustering

To validate that the expected spatial and temporal axes of variation are present in the

scRNA-seq data, we generated a low-dimensional representation of all cells using the

standard t-SNE (t-distributed stochastic neighbor embedding)[63], a nonlinear dimensionality

reduction technique that embeds high-dimensional data on a 2-dimensional plane such that

points that are similar in high-dimensional space are close together on the 2-dimensional

representation. We then colored cells either by their position along the central-portal axis,

or by time of day.

Spatial reconstruction of zonation profiles from scRNA-seq data

Choice of landmark genes

The reconstruction algorithm relies on *a priori* knowledge about the zonation of a small

set of landmark genes to infer the location of the cells. Ref.[8] used smFISH to determine the

zonation pattern *in situ* of 6 such landmark genes and used them to reconstruct the spatial

profiles of all other genes at a single time point. Since we here aimed at reconstructing

zonation profiles at different time points, we could not rely on those landmark genes, which

might be subject to temporal regulation. Therefore, we used an alternative strategy where

we selected landmark zonated genes from ref.[8] (q < 0.2), with the additional constraints

that those should be highly expressed (mean expression in fraction UMI of more than

0.01% and less than 0.1%), and importantly vary little across mice and time. Specifically,

we calculated the variability in the mean expression (across all layers) between all mice for

every gene and removed genes with >= 10% variability. This yielded 27 central (*Akr1c6,*

Alad, Blvrb, C6, Car3, Ccdc107, Cml2, Cyp2c68, Cyp2d9, Cyp3a11, Entpd5, Fmo1,

Gsta3, Gstm1, Gstm6, Gstt1, Hpd, Hsd17b10, Inmt, Iqgap2, Mgst1, Nrn1, Pex11a, Pon1,

Psmd4, Slc22a1, Tex264); and 28 portal (*Afm, Aldh1l1, Asl, Ass1, Atp5a1, Atp5g1, C8a,*

C8b, Ces3b, Cyp2f2, Elovl2, Fads1, Fbp1, Ftcd, Gm2a, Hpx, Hsd17b13, Ifitm3, Igf1,

Igfals, Khk, Mug2, Pygl, Sepp1, Serpina1c, Serpina1e, Serpind1, Vtn) landmark genes.

Reconstruction algorithm

The reconstruction algorithm is based on the algorithm in ref.[8] and was used in the modified

version from ref.[19]. The procedure was applied independently on each mouse, yielding ten

spatial gene expression profiles for each gene, given as fraction of UMI per cell.

Spatiotemporal analysis of liver gene expression profiles

Data

Each profile for the 14678 genes includes 8 layers from the pericentral to the periportal

zone and 4 time points: ZT0 (n=3 biological replicates from individual mice), ZT6 (n=2),

ZT12 (n=3) and ZT18 (n=2). The expression levels (noted as x) are then log-transformed

as follows:

$$y = log_2(x + \Delta) - B \quad (i)$$

The offset $\Delta = 10^{-4}$ buffers variability in lowly expressed genes, while the shift $B =$

$-log_2(11 \times 10^{-5})$ changes the scale so that $y = 0$ corresponds to about 10 mRNA copies

per cell (we expect on the order of 1M mRNA transcripts per liver cell).

Reference genes

For ease of interpretation (Figure 2 and Supplementary Figure 2), we used a set of reference

circadian genes and a set of reference zonated genes, highlighted in several figures.

The reference core circadian clock and clock output genes are the following: *Bmal1, Clock,*

Npas2, Nr1d1, Nr1d2, Per1, Per2, Cry1, Cry2, Dbp, Tef, Hlf, Elovl3, Rora, Rorc.

The reference zonated genes are the following: *Glul, Ass1, Asl, Cyp2f2, Cyp1a2, Pck1,*

Cyp2e1, Cdh2, Cdh1, Cyp7a1, Acly, Alb, Oat, Aldob, Cps1.

Gene expression variance in space and time

To analyze variability in space and time (Figure 2A) we computed, for each gene, the

spatial variance V_x and the temporal variance V_T. Let $y_{x,t,j}$ represent the expression profile,

with j the replicate index, $t \in \{1,2,...,N_t\}$ the time index, and $x \in \{1,2,...,N_x\}$ the layer

index. Then, V_x and V_T are computed as follows:

717
$$V_X = \frac{1}{N_t}\sum_t \frac{\Sigma_x[\Sigma_j(y_{x,t,j} - \frac{1}{N_x}\Sigma_x y_{x,t,j})]^2}{N_j^2 N_x} \quad (ii)$$

$$V_T = \frac{1}{N_x}\sum_x \frac{\Sigma_t[\Sigma_j(y_{x,t,j} - \frac{1}{N_t}\Sigma_t y_{x,t,j})]^2}{N_j^2 N_t} \quad (iii)$$

Thus, the spatial variance V_x is computed along the space (and averaged over the replicates)

for each time condition, and then averaged over time. The procedure is similar,

symmetrically, for V_t.

Gene filtering

For the analyses in Figure 2, we selected transcripts that were reproducible between

replicates, as well as sufficiently highly expressed (see scatterplot in Supplementary Figure

2A). To assess reproducibility across replicates, we computed the average relative variance

of the spatiotemporal profiles over the replicates:

727
$$V_J = \frac{\frac{1}{N_x N_t}\Sigma_{x,t}\left[\frac{1}{N_j}\Sigma_j\left(y_{x,t,j} - \frac{1}{N_j}\Sigma_j y_{x,t,j}\right)^2\right]}{\frac{1}{N_x N_t N_j}\Sigma_{x,t,j}\left[(y_{x,t,j} - \frac{1}{N_x N_t N_j}\Sigma_{x,t,j} y_{x,t,j})^2\right]} \quad (iv)$$

We considered genes with values below 50% (Supplementary Figure 2). To filter lowly

expressed genes, we required the maximum expression level across layers and time points

to exceed 10^{-5} (fraction of UMIs) which corresponds to y = 0 or about 10 copies of mRNA

per cell. While this was quite more permissive than previous scRNA-seq studies it allowed

to keep most reference circadian and zonated genes. However, scRNA-seq has still limited

sensitivity and some potentially important genes may have been removed in the filtering

process. In the end, our filters kept 5085 genes (1437 were removed due to low expression,

4733 due to high variance, and 3543 due to both), which were then used for subsequent

analyses.

Mixed-effect model for spatiotemporal mRNA profiles

Since the data is longitudinal is space (8 layers measured in each animal), modelling the

space-time profiles require the use of mixed-effect models. To systematically analyze the

spatiotemporal mRNA profiles, we used a parameterized function. Specifically, the model

uses sine and cosine functions for the time, and polynomials (up to degree 2) for space.

Possible interaction between space and time are described as space-dependent oscillatory

functions, or equivalently, time-dependent polynomial parameters. Our model for the

transformed mRNA expression y reads:

$$y_{x,t,i} = \mu_i + \mu(x) + a(x)\cos(\omega t) + b(x)\sin(\omega t) + \varepsilon_{x,t,i} \qquad (v)$$

747 Here t is the time, x the spatial position along the liver layers, and $i \in \{1,2,...,10\}$ the

748 animal index. This function naturally generalizes harmonic regression, often used for

749 analysis of circadian gene expression[25], by introducing space-dependent coefficients:

$$750 \quad \begin{cases} \mu(x) &= \mu_0 + \mu_1 P_1(x) + \mu_2 P_2(x) \\ a(x) &= a_0 + a_1 P_1(x) + a_2 P_2(x) \\ b(x) &= b_0 + b_1 P_1(x) + b_2 P_2(x) \end{cases}$$

Here, P_1 and P_2 are the Legendre polynomials of degrees 1 and 2, respectively; μ_0, μ_1 and

μ_2 represent the static zonation profile, a_0 and b_0 represent the global (space-independent)

rhythmicity of the gene, while a_1, a_2, b_1, b_2 represent layer-dependent rhythmicity. $\varepsilon_{x,t,j}$

is a Gaussian noise term with standard deviation σ. In addition to the fixed-effect

parameters described so far, we also introduced a mouse-specific random-effect μ_i (with

zero mean). This parameter groups the dependent layer measurements (obtained in the

same animal) and thereby properly adjusts the biological sample size for the rhythmicity

analysis.

Phases φ (related to peak times t through $t = \varphi * 24/2\pi$) and amplitudes A, for each

profile can then be computed for any layer from the coefficients $a(x)$ and $b(x)$:

$$761 \quad \varphi(x) = arctan2\big(b(x), a(x)\big) \qquad A(x) = \sqrt{a(x)^2 + b(x)^2} \qquad (vi)$$

762 Note that peak-to-trough difference is $2A(x)$. The peak-to-trough ratio or fold change of

763 the original expression levels is then $2^{2A(x)}$. We also note that an equivalent writing of the

764 model formulates the problem in terms of time-dependent zonation parameters instead of

765 space-dependent rhythmicity:

$$766 \quad y_{x,t,i} = \mu_i + \mu_0(t) + \mu_1(t)P_1(x) + \mu_2(t)P_2(x) + \varepsilon_{x,t,i} \qquad (vii)$$

767   where:

$$\begin{cases} \mu_0(t) & = & \mu_0 + a_0 \cos(\omega t) + b_0 \sin(\omega t) \\ \mu_1(t) & = & \mu_1 + a_1 \cos(\omega t) + b_1 \sin(\omega t) \\ \mu_2(t) & = & \mu_2 + a_2 \cos(\omega t) + b_2 \sin(\omega t) \end{cases}$$

In this study, we fixed $\omega = \frac{2\pi}{24\,h}$ since the animals were entrained in a 24 h light-dark cycle

and the low time resolution would prevent us from studying ultradian rhythms.

The model parameters, including the variance of the random effects and Gaussian noise

strength σ, are estimated for each gene using the *fit* function from the Python library

StatsModels (version 0.9.0). Nelder-Mead was chosen as the optimization method, and the

use of a standard likelihood was favored over the REML likelihood to allow for model

comparison[64]. To prevent overfitting of the gene profiles, we added a noise offset $\sigma_0 = 0.15$

[log_2] to the estimated noise σ, in the expression of the likelihood function used in the

mixed-effect model optimization.

Depending on the gene, the model presented in (v) and (vii) may be simplified by setting

all or some of the (fixed) parameters to 0. For example, a non-oscillatory gene profile

would normally have non-significant a_j and b_j parameters. In practice, considering the

fixed effects, 2^9 sub-models of various complexity can be generated. However, we added

a few reasonable requirements to reduce the number of models. First, the intercept μ_0 must

be present in every model. Similarly, the parameters a_0 and b_0, providing a global rhythm,

must be present in every rhythmic model. Finally, the parameters a_j and b_j for j=0,1,2 must

be paired to ensure a proper phase definition (vi).

The models can then be classified in different categories, depending on the retained (non-

zero) parameters (Figure 2C):

• The model comprising only the intercepts μ_0 and μ_i, termed flat or noisy (F/N).

• The models comprising only the intercepts and zonation parameters: μ_1 and/or μ_2,

termed purely zonated (Z).

• The models comprising only the intercepts and rhythmic parameters: a_0 and b_0,

termed purely rhythmic (R).

• The models comprising only the intercepts, zonated parameters and rhythmic

parameters: μ_1 and/or μ_2, and a_0, b_0, termed independent (Z+R).

•   The models comprising interaction parameters: a_j and b_j for j=1,2, termed

interacting (ZxR).

Note that we only plot the fixed effects in the predicted gene profiles. The Bayesian

Information Criterion (BIC) is then used for model selection, enabling to choose the most

parsimonious model for each gene. Consequently, the F/N class also contains noisy

profiles, since genes that are not well fitted with any complex model will then be assigned

to the simplest model. Additionally, it appears that, for some profiles, several competing

models can result in close BIC values (see e.g. the discussion on *Clock* and *Cry*1 in the

Results). Therefore, when assigning hard classes, if some models have a relative difference

of less than 1% in their BIC, we systematically keep the most complex model. Moreover,

we also assigned probabilities to the different categories (F, Z, R, Z+R and ZxR), computed

as Schwartz BIC weights[65], which is useful in case of ambiguous classification

(Supplementary Table 2). All best fits with their parameter values are listed in

Supplementary Table 1.

Comparison of peak times with a bulk dataset

We compared the peak phases our rhythmically classified genes with those obtained from

ref.[25]. These data consist of bulk liver RNA-seq data sampled every 2 hours for 24 hours,

with 4 replicates per time condition. We only compared genes that for which rhythmicity

is not changing across layers, *viz.* the R and Z+R categories. Note that since our dataset has

a lower temporal resolution and fewer replicates per time point, we found overall less

rhythmic genes.

To assess gene rhythmicity from ref.[25], we used harmonic regression on the log-

transformed profiles as previously. Using the same notation as above, we define the two

following models:

821
$$\begin{cases} y_{t,i} & = \mu + \varepsilon & (ix) \\ y_{t,i} & = \mu + a\cos(\omega t) + b\sin(\omega t) + \varepsilon & (x) \end{cases}$$

We then fit eq. ($viii$) and eq. (ix) to every transcript, and, for each of them, keep the model

with the lowest BIC. We then compared the phases of transcripts classified as rhythmic in

both datasets, and computed the circular correlation coefficient[66].

KEGG pathway Enrichment analysis

Functional annotation clustering from Enrichr[67] for the categories F/N, Z and Z+R (which

is then subdivided in central and portal), Zc+R (central zonated and rhythmic), Zp+R

(portal zonated and rhythmic) and finally R was ran with standard parameters, using the

standard KEGG 2019 Mouse set of pathways. The enriched pathways (adjusted p-value <

0.1) was then further annotated to compute, e.g. the number of central/portal genes in each

category and the phase of each gene. This analysis is available Supplementary Table 3.

smFISH

Analysis of Z+R and ZxR genes (Stellaris smFIH probes)

Preparation of probe libraries, hybridization procedure and imaging conditions were

previously described[19]. smFISH probe libraries were coupled to TMR, Alexa594 or Cy5.

Cell membranes were stained with alexa fluor 488 conjugated phalloidin (Rhenium

A12379) that was added to GLOX buffer[68]. Portal node was identified morphologically on

DAPI images based on bile ductile, central vein was identified using smFISH for Glul in

TMR, included in all hybridizations. Images were taken as scans spanning the portal node

to the central vein. Images were analyzed using ImageM[68]. Quantification of zonation

profiles in different circadian time point were generated by counting dots and dividing the

number of dots in radial layers spanning the portal-central axis by the layer volume.

Temporal analysis of circadian genes (RNA scope smFISH probes)

smFISH of R genes were done on fresh-frozen liver cryosections (8μm) embedded in

O.C.T Compound (Tissue-Tek; Sakura-Finetek USA), sampled every three hours (ZT0 to

ZT21). RNAscope® probes for *Bma1l* mRNA (Mm-Arntl, catalog #: 438748-C3) and *Per1*

mRNA (Mm-Per1, catalog #: 438751) were used, according to the manufacturer's

instructions for the RNAscope Fluorescent Multiplex V1 Assay (Advanced Cell

Diagnostics). To detect the central vein, an immunofluorescence of Glutamine Synthetase

(ab49873, Abcam, diluted 1:2000 in PBS/BSA 0.5%/Triton-X0.01%) was done together

with smFISH. Nuclei were counterstained with DAPI and sections were mounted with

ProLong™ Gold Antifade Mountant. Liver sections were imaged with a Leica DM5500

widefield microscope and an oil-immersion x63 objective. Z-stacks were acquired (0.2μm

between each Z position) and mRNA transcripts were quantified using ImageJ, as

described previously in ref.[50]. Pericentral (PC) and Periportal (PP) veins were manually

detected based on Glutamine Synthetase IF or on bile ducts (DAPI staining). The Euclidean

distance between two veins and the distance from the vein of each mRNA transcript were

calculated. mRNA transcripts were assigned to a PP or PC zone if the distance from the

corresponding vein was smaller than one-third of the distance between the PP and PC veins

(ranging from 50 to 130μm).

Wnt2, Rspo3 and Dkk3 expression in non-parenchymal cells (NPCs)

Preparation of probe libraries, hybridization protocol and imaging conditions were

previously described[19]. The *Aqp1, Igfbp7 and Ptprb* probe libraries were coupled to TMR,

the *Wnt2* library was coupled to *Alexa594* and the *Dkk3* or *Rspo3* library were coupled to

Cy5. Cell membranes were stained with alexa fluor 488 coupled to phalloidin (Rhenium

A12379) that was added to GLOX buffer[68]. The central vein was identified based on

morphological features inspected in the DAPI and Phalloidin channels and presence of

Wnt2-mRNA (detected by smFISH). Central vein niche NPCs were identified by co-

staining of *Aqp1*, *Igfbp7* and *Ptprb*. The central vein area was imaged and the images were

analyzed using ImageM[68]. We counted dots of *Wnt2, Rspo3* and *Dkk3* expression

(corresponding to single mRNA molecules) in NPCs lining the central vein and removed

background dots larger than 25 pixels. We then divided the dot count by the segmented

cell volume. In total 1420 endothelial cells from 189 central veins of at least 2 mice per

time point. In total 1420 NPCs from 189 central veins of at least 2 mice per time point

(ZT0,6,12,18) were imaged and a Kruskal-Wallis test based on the mean mRNA dot

concentration in each cell was performed to compare the ZT0 and ZT18 time points.

Data availability

scRNA-seq data

All scRNA-seq data is deposited in GEO with accession code GSE145197 (reviewer token

xxx).

Reconstructed gene profiles

Reconstructed spatio-temporal gene profiles are available as Matlab files at
https://c4science.ch/diffusion/10261/

Web-application

The whole dataset of gene profiles along with the analysis is available online as a web-
application at the URL https://czviz.epfl.ch/. The application was built in Python using the
library *Dash by Plotly* (version 1.0).

Code availability

The code for fitting the mixed-effects models and generating the main figures is available
at https://c4science.ch/diffusion/10261/

References

1.   Gebhardt, R. Metabolic zonation of the liver: Regulation and implications for liver
function. *Pharmacol. Ther.* **53**, 275–354 (1992).

2.   Hoehme, S. *et al.* Prediction and validation of cell alignment along microvessels as
order principle to restore tissue architecture in liver regeneration. *Proc. Natl. Acad. Sci.*
107, 10371–10376 (2010).

3.   Ben-Moshe, S. & Itzkovitz, S. Spatial heterogeneity in the mammalian liver. *Nat.*
Rev. Gastroenterol. Hepatol. **16**, 395–410 (2019).

4.   Colnot, S. & Perret, C. Liver Zonation. in *Molecular Pathology of Liver Diseases*
(ed. Monga, S. P. S.) vol. 5 7–16 (Springer US, 2011).

5.   Wang, B., Zhao, L., Fish, M., Logan, C. Y. & Nusse, R. Self-renewing diploid
Axin2+ cells fuel homeostatic renewal of the liver. *Nature* **524**, 180–185 (2015).

6.   Planas-Paz, L. *et al.* The RSPO–LGR4/5–ZNRF3/RNF43 module controls liver
zonation and size. *Nat. Cell Biol.* **18**, 467–479 (2016).

7.   Jungermann, K. & Kietzmann, T. Zonation of parenchymal and nonparenchymal
metabolism in liver. *Annu. Rev. Nutr.* **16**, 179–203 (1996).

8.   Halpern, K. B. *et al.* Single-cell spatial reconstruction reveals global division of
labour in the mammalian liver. *Nature* **542**, 352–356 (2017).

9.   Dibner, C., Schibler, U. & Albrecht, U. The Mammalian Circadian Timing System:
Organization and Coordination of Central and Peripheral Clocks. *Annu. Rev. Physiol.* **72**,
517–549 (2010).

10.   Green, C. B., Takahashi, J. S. & Bass, J. The Meter of Metabolism. *Cell* **134**, 728–
742 (2008).

11.   Yeung, J. & Naef, F. Rhythms of the Genome: Circadian Dynamics from
Chromatin Topology, Tissue-Specific Gene Expression, to Behavior. *Trends Genet. TIG*
34, 915–926 (2018).

12.   Maury, E., Ramsey, K. M. & Bass, J. Circadian Rhythms and Metabolic Syndrome:
From Experimental Genetics to Human Disease. *Circ. Res.* **106**, 447–462 (2010).

13.   Guan, D. *et al.* The hepatocyte clock and feeding control chronophysiology of
multiple liver cell types. *Science* **369**, 1388–1394 (2020).

14.   Mermet, J., Yeung, J. & Naef, F. Systems Chronobiology: Global Analysis of Gene
Regulation in a 24-Hour Periodic World. *Cold Spring Harb. Perspect. Biol.* **9**, a028720
(2017).

15.   Zhang, R., Lahens, N. F., Ballance, H. I., Hughes, M. E. & Hogenesch, J. B. A
circadian gene expression atlas in mammals: implications for biology and medicine. *Proc.
Natl. Acad. Sci. U. S. A.* **111**, 16219–16224 (2014).

16.   Sobel, J. A. *et al.* Transcriptional regulatory logic of the diurnal cycle in the mouse
liver. *PLOS Biol.* **15**, e2001069 (2017).

17.   Vollmers, C. *et al.* Time of feeding and the intrinsic circadian clock drive rhythms
in hepatic gene expression. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 21453–21458 (2009).

18.   Yeung, J. *et al.* Transcription factor activity rhythms and tissue-specific chromatin
interactions explain circadian gene expression across organs. *Genome Res.* **28**, 182–191
(2018).

19.   Halpern, K. B. *et al.* Paired-cell sequencing enables spatial gene expression
mapping of liver endothelial cells. *Nat. Biotechnol.* **36**, 962–970 (2018).

20.   Bunger, M. K. *et al.* Mop3 Is an Essential Component of the Master Circadian
Pacemaker in Mammals. *Cell* **103**, 1009–1017 (2000).

21.   Ripperger, J. A., Shearman, L. P., Reppert, S. M. & Schibler, U. CLOCK, an
essential pacemaker component, controls expression of the circadian transcription factor
DBP. *Genes Dev.* **14**, 679–689 (2000).

22.   Beal, J. Biochemical complexity drives log-normal variation in genetic expression.
Eng. Biol. **1**, 55–60 (2017).

23.   McLean, R. A., Sanders, W. L. & Stroup, W. W. A Unified Approach to Mixed
Linear Models. *Am. Stat.* **45**, 54 (1991).

24.   Wagenmakers, E.-J. & Farrell, S. AIC model selection using Akaike weights.
Psychon. Bull. Rev. **11**, 192–196 (2004).

25.   Atger, F. *et al.* Circadian and feeding rhythms differentially affect rhythmic mRNA
transcription and translation in mouse liver. *Proc. Natl. Acad. Sci.* **112**, E6579–E6588
(2015).

26.   Kietzmann, T. Metabolic zonation of the liver: The oxygen gradient revisited.
Redox Biol. **11**, 622–630 (2017).

27.   Robles, M. S., Cox, J. & Mann, M. In-Vivo Quantitative Proteomics Reveals a Key
Contribution of Post-Transcriptional Mechanisms to the Circadian Regulation of Liver
Metabolism. *PLoS Genet.* **10**, e1004047 (2014).

28.   Mauvoisin, D. *et al.* Circadian clock-dependent and -independent rhythmic
proteomes implement distinct diurnal functions in mouse liver. *Proc. Natl. Acad. Sci.* **111**,
167–172 (2014).

29.   Gebhardt, R. Liver zonation: Novel aspects of its regulation and its impact on
homeostasis. *World J. Gastroenterol.* **20**, 8491 (2014).

30.   Kornmann, B., Schaad, O., Bujard, H., Takahashi, J. S. & Schibler, U. System-
Driven and Oscillator-Dependent Circadian Transcription in Mice with a Conditionally
Active Liver Clock. *PLoS Biol.* **5**, e34 (2007).

31.   Reinke, H. *et al.* Differential display of DNA-binding proteins reveals heat-shock
factor 1 as a circadian transcription factor. *Genes Dev.* **22**, 331–345 (2008).

32.   Sinturel, F. *et al.* Diurnal Oscillations in Liver Mass and Cell Size Accompany
Ribosome Assembly Cycles. *Cell* **169**, 651-663.e14 (2017).

33.   Kietzmann, T., Cornesse, Y., Brechtel, K., Modaressi, S. & Jungermann, K.
Perivenous expression of the mRNA of the three hypoxia-inducible factor alpha-subunits,
HIF1alpha, HIF2alpha and HIF3alpha, in rat liver. *Biochem. J.* **354**, 531–537 (2001).

34.   Ma, D., Panda, S. & Lin, J. D. Temporal orchestration of circadian autophagy
rhythm by C/EBPβ: C/EBPβ regulates circadian autophagy rhythm. *EMBO J.* **30**, 4642–
4651 (2011).

35.   Wang, J. *et al.* Nuclear Proteomics Uncovers Diurnal Regulatory Landscapes in

Mouse Liver. *Cell Metab.* **25**, 102–117 (2017).

36.   Harper, J. W., Ordureau, A. & Heo, J.-M. Building and decoding ubiquitin chains

for mitophagy. *Nat. Rev. Mol. Cell Biol.* **19**, 93–108 (2018).

37.   Le Martelot, G. *et al.* REV-ERBα Participates in Circadian SREBP Signaling and

Bile Acid Homeostasis. *PLoS Biol.* **7**, e1000181 (2009).

38.   Ferrell, J. M. & Chiang, J. Y. L. Short-Term Circadian Disruption Impairs Bile

Acid and Lipid Homeostasis in Mice. *Cell. Mol. Gastroenterol. Hepatol.* **1**, 664–677

(2015).

39.   Ma, K. *et al.* Circadian Dysregulation Disrupts Bile Acid Homeostasis. *PLoS ONE*

4, e6843 (2009).

40.   Yu, L. *et al.* Disruption of Abcg5 and Abcg8 in mice reveals their crucial role in

biliary cholesterol secretion. *Proc. Natl. Acad. Sci.* **99**, 16237–16242 (2002).

41.   Oki, S. *et al.* Ch IP -Atlas: a data-mining suite powered by full integration of public

Ch IP -seq data. *EMBO Rep.* **19**, (2018).

42.   Gotic, I. *et al.* Temperature regulates splicing efficiency of the cold-inducible

RNA-binding protein gene *Cirbp. Genes Dev.* **30**, 2005–2017 (2016).

43.   Jakobsson, A., Jörgensen, J. A. & Jacobsson, A. Differential regulation of fatty acid

elongation enzymes in brown adipocytes implies a unique role for *Elovl3* during increased

fatty acid oxidation. *Am. J. Physiol.-Endocrinol. Metab.* **289**, E517–E526 (2005).

44.   Burke, Z. D. & Tosh, D. The Wnt/β-catenin pathway: master regulator of liver

zonation? *BioEssays* **28**, 1072–1077 (2006).

45.   Gougelet, A. *et al.* T-cell factor 4 and β-catenin chromatin occupancies pattern

zonal liver metabolism in mice. *Hepatology* **59**, 2344–2357 (2014).

46.   Lavery, D. J. *et al.* Circadian Expression of the Steroid 15 α-Hydroxylase (Cyp2a4)

and Coumarin 7-Hydroxylase (Cyp2a5) Genes in Mouse Liver Is Regulated by the PAR

Leucine Zipper Transcription Factor DBP. *Mol. Cell. Biol.* **19**, 6488–6499 (1999).

47.   Carmon, K. S., Gong, X., Lin, Q., Thomas, A. & Liu, Q. R-spondins function as

ligands of the orphan receptors LGR4 and LGR5 to regulate Wnt/beta-catenin signaling.

Proc. Natl. Acad. Sci. U. S. A. **108**, 11452–11457 (2011).

48.   Aizarani, N. *et al.* A human liver cell atlas reveals heterogeneity and epithelial

progenitors. *Nature* **572**, 199–204 (2019).

49.   Brosch, M. *et al.* Epigenomic map of human liver reveals principles of zonated
morphogenic and metabolic control. *Nat. Commun.* **9**, (2018).

50.   Mermet, J. *et al.* Clock-dependent chromatin topology modulates circadian
transcription and behavior. *Genes Dev.* **32**, 347–358 (2018).

51.   Le Martelot, G. *et al.* Genome-Wide RNA Polymerase II Profiles and RNA
Accumulation Reveal Kinetics of Transcription and Associated Epigenetic Changes
During Diurnal Cycles. *PLoS Biol.* **10**, e1001442 (2012).

52.   Wang, J. *et al.* Circadian clock-dependent and -independent posttranscriptional
regulation underlies temporal mRNA accumulation in mouse liver. *Proc. Natl. Acad. Sci.*
115, E1916–E1925 (2018).

53.   Nitzan, M., Karaiskos, N., Friedman, N. & Rajewsky, N. Gene expression
cartography. *Nature* (2019) doi:10.1038/s41586-019-1773-3.

54.   Ben-Moshe, S. *et al.* Spatial sorting enables comprehensive characterization of
liver zonation. *Nat. Metab.* **1**, 899–911 (2019).

55.   Seglen, P. O. Chapter 4 Preparation of Isolated Rat Liver Cells. in *Methods in Cell*
Biology vol. 13 29–83 (Elsevier, 1976).

56.   Darnell, A. M., Subramaniam, A. R. & O'Shea, E. K. Translational Control through
Differential Ribosome Pausing during Amino Acid Limitation in Mammalian Cells. *Mol.*
Cell **71**, 229-243.e11 (2018).

57.   Clayton, D. F., Harrelson, A. L. & Darnell, J. E. Dependence of liver-specific
transcription on tissue organization. *Mol. Cell. Biol.* **5**, 2623–2632 (1985).

58.   Satija, R., Farrell, J. A., Gennert, D., Schier, A. F. & Regev, A. Spatial
reconstruction of single-cell gene expression data. *Nat. Biotechnol.* **33**, 495–502 (2015).

59.   Loud, A. V. A QUANTITATIVE STEREOLOGICAL DESCRIPTION OF THE
ULTRASTRUCTURE OF NORMAL RAT LIVER PARENCHYMAL CELLS. *J. Cell*
Biol. **37**, 27–46 (1968).

60.   Kuhn, N. J., Woodworth-Gutai, M., Gross, K. W. & Held, W. A. Subfamilies of
the mouse major urinary protein (MUP) multi-gene family: sequence analysis of cDNA
clones and differential regulation in the liver. *Nucleic Acids Res.* **12**, 6073–6090 (1984).

61.   Hurst, J. L. *et al.* Individual recognition in mice mediated by major urinary proteins.
Nature **414**, 631–634 (2001).

62. Ilicic, T. *et al.* Classification of low quality cells from single-cell RNA-seq data.
Genome Biol. **17**, 29 (2016).

63. Maaten, L. van der & Hinton, G. Visualizing Data using t-SNE. *J. Mach. Learn.*
Res. **9**, 2579–2605 (2008).

64. Müller, S., Scealy, J. L. & Welsh, A. H. Model Selection in Linear Mixed Models.
Stat. Sci. **28**, 135–167 (2013).

65. Schwarz, G. Estimating the Dimension of a Model. *Ann. Stat.* **6**, 461–464 (1978).

66. Jammalamadaka, S. R. & SenGupta, A. *Topics in Circular Statistics*. vol. 5
(WORLD SCIENTIFIC, 2001).

67. Chen, E. Y. *et al.* Enrichr: interactive and collaborative HTML5 gene list
enrichment analysis tool. *BMC Bioinformatics* **14**, 128 (2013).

68. Lyubimova, A. *et al.* Single-molecule mRNA detection and counting in
mammalian tissue. *Nat. Protoc.* **8**, 1743–1758 (2013).
1054

Acknowledgments

This work was supported by the Rothschild Caesarea Foundation' fund managed by Weizmann Institute and EPFL, a Swiss National Science Foundation Grant 310030_173079 (to F.N.), and the EPFL. S.I. is supported by the Henry Chanoch Krenter Institute for Biomedical Imaging and Genomics, The Leir Charitable Foundations, Richard Jakubskind Laboratory of Systems Biology, Cymerman-Jakubskind Prize, The Lord Sieff of Brimpton Memorial Fund, the Wolfson Foundation SCG, the Wolfson Family Charitable Trust, Edmond de Rothschild Foundations, the I-CORE program of the Planning and Budgeting Committee and the Israel Science Foundation (grants 1902/12 and 1796/12, the Israel Science Foundation grant no. 1486/16, the Chan Zuckerberg Initiative grant no. CZF2019-002434, the Broad Institute-Israel Science Foundation grant no. 2615/18, the European Research Council (ERC) under the European Union's Horizon 2020 research and innovation program (grant agreement no. 768956, the Bert L. and N. Kuggie Vallee Foundation, and the Howard Hughes Medical Institute (HHMI) international research scholar award.

Author contributions

F.N. and S.I. conceived the study. K.B.H., J.E.K. and C.H. prepared all the samples and performed the experiments. C.D. and F.N designed the modeling. K.B.H., J.E.K., C.H. and C.D. analyzed the data. M.R. and S.M. assisted with the scRNA-seq and smFISH experiments. F.N and S.I. supervised the study. C.D., J.E.K. and F.N. wrote the manuscript. All authors reviewed the manuscript and provided input.

Ethics statement

All animal care and handling were approved by the Institutional Animal Care and Use Committee of Weizmann Institute of Science and by the Canton de Vaud laws for animal protection (authorization VD3197.b).

Competing interests

The authors declare no competing interests.