Modification of forests by people means only 40% of remaining forests have high ecosystem integrity

Authors: Grantham, H.S.[1*], Duncan, A.[1], Evans, T. D.[1], Jones, K.[1], Beyer, H.[2], Schuster, R.[3],

Walston, J.[1], Ray, J.[1], Robinson, J.[1], Callow, M.[1], Clements, T.[1], Costa, H.M.[1], DeGemmis, A.[1],

Elsen, P.R.[1], Ervin, J.[4], Franco, P.[1], Goldman, E.[5], Goetz, S.[6], Hansen, A.[7], Hofsvang, E.[8], Jantz,

P. [6], Jupiter, S. [1], Kang, A.[1], Langhammer, P.[9,10], Laurance, W.F.[11], Lieberman, S., Linkie, M.[1],

Malhi, Y.[12], Maxwell, S.[13], Mendez, M.[1], Mittermeier, R.[9], Murray, N.[11], Possingham, H.[2,14],

Radachowsky, J.[1], Samper, C.[1], Silverman, J.[1], Shapiro, A.[15], Strassburg, B.[17], Stevens, T.[1],

Stokes, E.[1], Taylor, R.[5], Tear, T.[1], Tizard, R.[1], Venter, O. [16], Visconti, P.[18], Wang, S.[1], Watson,

J.E.M.[1,13]

Affiliations

[1]Wildlife Conservation Society, Global Conservation Program, Bronx, New York, 10460 USA.

[2]School of Biological Sciences, University of Queensland, St. Lucia, Queensland, Australia

[3]Department of Biology, 1125 Colonel By Drive, Carleton University, Ottawa ON, K1S 5B6

Canada.

[4]United Nations Development Programme, One United Nations Plaza, New York, NY, 10017,

USA

[5]World Resources Institute, Washington, DC, USA.

[6]Global Earth Observation & Dynamics of Ecosystems Lab, School of Informatics, Computing,

and Cyber Systems, Northern Arizona University, Flagstaff, AZ, 86011, USA

[7]Landscape Biodiversity Lab, Ecology Department, Montana State University, Bozeman, MT, 59717, USA

[8] Rainforest Foundation Norway, Mariboes gate 8, 0183 Oslo

[9]Global Wildlife Conservation, P.O. Box 129, Austin, Texas 78767, USA

[10]School of Life Sciences, Arizona State University, P.O. Box 874501, Tempe, Arizona 85287, USA

[11]Centre for Tropical Environmental and Sustainability Science, College of Science and Engineering, James Cook University, Cairns, QLD 4878, Australia

[12]Environmental Change Institute, School of Geography and the Environment, University of Oxford, Oxford, United Kingdom

[13]School of Earth and Environmental Sciences, University of Queensland, Brisbane, Australia

[14]The Nature Conservancy, Arlington, VA, USA

[15]World Wide Fund for Nature Germany, Space+Science

[16]Natural Resource and Environmental Studies Institute, University of Northern British Columbia, Prince George, Canada

[17]International Institute of Sustainability, Rio de Janeiro, 22460-320, Brazil

[18]International Institute for Applied Systems Analysis, Laxenburg, Austria

*hgrantham@wcs.org

Abstract: Many global environmental agendas, including halting biodiversity loss, reversing land degradation, and limiting climate change, depend upon retaining forests with high ecological integrity, yet the scale and degree of forest modification remains poorly quantified and mapped. By integrating data on observed and inferred human pressures and an index of lost connectivity, we generate the first globally-consistent, continuous index of forest condition as determined by

degree of anthropogenic modification. Globally, only 17.4 million km^2 of forest (40.5%) have

high landscape level integrity (mostly found in Canada, Russia, the Amazon, Central Africa and

New Guinea) and only 27% of this area is found in nationally-designated protected areas. Of the

forest in protected areas, only 56% has high landscape level integrity. Ambitious policies that

prioritize the retention of forest integrity, especially in the most intact areas, are now urgently

needed alongside current efforts aimed at halting deforestation and restoring the integrity of

forests globally.

MAIN TEXT

Introduction

Deforestation is a major environmental issue [1], but far less attention has been given to the degree

of anthropogenic modification of remaining forests, which reduces ecosystem integrity and

diminishes many of the benefits that these forests provide [2,3]. This is worrying since modification

is potentially as significant as outright forest loss in determining overall environmental outcomes

[4]. There is increasing recognition of this issue, for forests and other ecosystems, in synthesis

reports by global science bodies e.g. [5], and it is now essential that the scientific community

develop improved tools and data to facilitate the consideration of levels of integrity in decision-

making. Mapping and monitoring this globally will provide essential information for coordinated

global, national and local policy-making, planning and action, to help nations and other

70 stakeholders achieve the Sustainable Development Goals (SDGs) and implement other shared

71 commitments such as the United Nations Convention on Biological Diversity (CBD), Convention

72 to Combat Desertification (UNCCD), and Framework Convention on Climate Change

73 (UNFCCC).

Ecosystem integrity is foundational to all three of the Rio Conventions (UNFCCC, UNCCD,

CBD). As defined by Parrish et al. [6], it is essentially the degree to which a system is free from

anthropogenic modification of its structure, composition and function. Such modification causes

the reduction of many ecosystem benefits, and is often also a precursor to outright deforestation

[7,8]. Forests largely free of significant modification (i.e. forests having high ecosystem integrity),

typically provide higher levels of many forest benefits than modified forests of the same type [9],

including; carbon sequestration and storage [10], healthy watersheds [11], traditional homelands for

imperiled cultures [12], contribution to local and regional climate processes [13], and forest-dependent

biodiversity [14-17]. Industrial-scale logging, fragmentation by infrastructure, farming (including

cropping and ranching) and urbanization, as well as less visible forms of modification such as

over-hunting, wood fuel extraction and changed fire or hydrological regimes [18,19], all degrade the

degree to which forests still support these benefits, as well as their long-term resilience to climate

change [9]. There can be trade-offs however, between the benefits best provided by less-modified

forests (e.g., regulatory functions such as carbon sequestration) and those production services that

require some modification (e.g., timber production). These trade-offs can, at times, result in

disagreement among stakeholders as to which forest benefits should be prioritized [20].

In recent years, easily accessible satellite imagery and new analytical approaches have

dramatically improved our ability to map and monitor forest extent globally [21-23]. However, while

progress has been made in developing tools for assessment of global forest losses and gains,

consistent monitoring of the degree of forest modification has proved elusive [24,25].

Technical challenges include the detection of low intensity and unevenly distributed forest

modification, the wide diversity of changes that comprise forest modification, and the fact that

many changes are concealed by the forest canopy [24]. New approaches are emerging on relevant

forest indicators, such as canopy height, canopy cover and fragmentation, and maps of different human pressures, which are used as proxies for impacts on forests e.g., [26,27,28]. Some binary measures of forest modification, such as Intact Forest Landscapes [29] and wilderness areas [30], have also been mapped at the global scale and used to inform policy, but do not resolve the degree of modification within remaining forests, which we aimed to do with this assessment.

Human activities influence the integrity of forests at multiple spatial scales, including intense, localized modifications such as road-building and canopy loss, more diffuse forms of change that are often spatially associated with these localized pressures (e.g. increased accessibility for hunting, other exploitation, and selective logging), and changes in spatial configuration that alter landscape-level connectivity. Previous studies have quantified several of these aspects individually e.g. [26,27,28], but there is a need to integrate them to measure and map the overall degree of modification considering these landscape-level anthropogenic influences at each site. Here, we integrate data on forest extent defined as all woody vegetation taller than 5 m, following [22], 'observed' human pressures (e.g. infrastructure) which can be directly mapped using current datasets, other 'inferred' human pressures (e.g. collection of forest materials) that occur in association with those that are observed but cannot be mapped directly, and alterations in forest connectivity, to create the "Forest Landscape Integrity Index" (FLII), that describes the degree of forest modification for the beginning of 2019 (Fig. 1). The result is the first globally applicable, continuous-measure map of landscape level forest integrity (hereafter, integrity), which offers a timely indicator of the status and management needs of Earth's remaining forests, as well as a flexible methodological framework (Fig. 1) for measuring changes in forest integrity that can be adapted for more detailed analysis at national or subnational scales.

Results

Forest modification caused by human activity is both highly pervasive and highly variable across

the globe (Fig. 2). We found 31.2% of forests worldwide are experiencing some form of

'observed' human pressure. Our models also inferred the likely occurrence of other pressures, and

the impacts of lost connectivity, in almost every forest location (91.2% of forests), albeit

sometimes at very low levels. Diverse, recognizable patterns of forest integrity can be observed in

our maps at a range of scales, depending on the principal forms and general intensity of human

activity in an area. Broad regional trends can be readily observed, for example the overall gradient

of decreasing human impact moving northwards through eastern North America (Fig. 2), and

finer patterns of impact are also clearly evident, down to the scale of individual protected areas,

forest concessions, settlements and roads (Fig. S2).

FLII scores range from 0 (lowest integrity) to 10 (highest). We discretized this range to define

three broad illustrative categories: low (≤6.0); medium (>6.0 and <9.6); and high integrity (≥9.6)

by benchmarking against reference locations worldwide (see Methods). Only 40.5% (17.4

million km^2) of forest was classified as having high integrity (Fig. 3; Table 1). Moreover, even in

this category of high integrity (36%) still showed at least a small degree of human modification.

The remaining 59% (25.6 million km^2) of forest was classified as having low or medium integrity,

including 25.6% (11 million km^2) with low integrity (Fig. 3; Table 1). When we analyzed across

biogeographical realms defined by [31] not a single biogeographical realm of the world had more

than half of its forests in the high category (Fig. 3; Table 1).

The biogeographical realms with the largest area of forest with high integrity are the Paleartic,

particularly northern Russia, and the Neartic, in northern Canada, and Alaska. There are also large

areas of forest with high integrity in the Neotropics, concentrated in the Amazon region, including

within the Guianas (Fig. 3, Table 1). The Afrotropic realm has significant areas with high

integrity, particularly within the humid forests of central Africa (e.g., in Republic of Congo and

Gabon) and in some of the surrounding drier forest/woodland belts (e.g., in South Sudan, Angola

and Mozambique) (Fig. 3). In tropical Asia, the largest tracts of forest with high integrity are in

New Guinea. Smaller but still very significant tracts of forest with high integrity are also scattered

elsewhere in each of the main forested regions, including parts of Sumatra, Borneo, Myanmar and

other parts of the greater Mekong subregion, Madagascar, West Africa, Mesoamerica, the

Atlantic forests of Brazil, southern Chile, the Rocky Mountains, northern Assam, the Pacific

forests of Colombia, the Caucasus, and the Russian Far East (Fig. 3).

Concentrations of forest with low integrity are found in many regions including west and central

Europe, the south-eastern USA, island and mainland South-East Asia west of New Guinea, the

Andes, much of China and India, the Albertine Rift, West Africa, Mesoamerica and the Atlantic

Forests of Brazil (Fig. 3). The overall extent of forests with low integrity is greatest in the

Paleartic realm, followed by the Neotropics, which are also those biogeographic realms with the

largest forest cover (Table 1). The Indo-Malayan realm has the highest percentage with low

integrity, followed by the Afrotropics (Fig. 3; Table 1).

These patterns result in variation with forest integrity scores in ways that allow objective

comparisons to be made between locations and at a resolution relevant for policy and

management planning, such as at national and sub-national scales. The global average FLII score

for all countries is 5.48, representing generally low forest integrity, and a quarter of forested

countries have a national average score < 4. National mean scores vary widely, ranging from >9

in Guyana, French Guiana, Gabon, Sudan and South Sudan to <3 in Sierra Leone and many west

European countries (see Fig 4. and Table S5 for full list of countries). Provinces and other sub-

national units vary even more widely (see Fig. S2 and Table S6)

Over one-quarter (26.1%) of all forests with high integrity fall within protected areas, compared

to just 13.1% of low and 18.5% of medium integrity forests respectively. For all forests that are

found within nationally designated protected areas (around 20% of all forests globally), we found

the proportions of low, medium and high integrity forests were 16.8%, 30.3%, and 52.8%

respectively (Table 2). Within the different protected area categories, we typically found that

there was more area within the high integrity category versus the medium and low except for

Category V (protected landscape/seascape) (Table 2). However, with 47.1% of forests within

protected areas having low to medium integrity overall, it is clear that forests considered

'protected' are already often fairly modified (Table 2). Even though they are quite modified, some

of these forests might still have high conservation importance, such as containing endangered

species.

Discussion

By providing a transparent and defensible methodological framework, and by taking advantage of

global data on forest extent, human drivers of forest modification, and changes in forest

connectivity, our analysis paints a new, sobering picture of the extent of human impacts on the

world's forests. This analysis enables the changes that degrade many forest values *(8)* to be

visualized in a new and compelling way and for policy makers and decision makers to see where

forests that survive in good condition are found. By integrating data on multiple pressures that are

known to modify forests, our analysis is the first to move global quantification beyond the use of

simple categories, or solely using pressure indicators as proxies for integrity, to a more nuanced

depiction of this issue as a continuum, recognising that not all existing forests are in the same

condition. Our analysis reveals that severe and extensive forest modification has occurred across all biogeographic regions of the world. Consequently, indices only using forest extent may inadequately capture the true impact of human activities on forests, and are insensitive to many drivers of forest modification and the resulting losses of forest benefits.

A plan is clearly needed to put in place retention strategies for the remaining forests with high integrity, tailored towards the context in each country or jurisdiction and its different forest types [32,33], because such areas are known to hold exceptional value. Avoiding the loss of integrity is a better strategy than aiming to restore forest condition after it is lost, because restoration is more costly, has a risk of failure , and is unlikely to lead to full recovery of benefits [5]. For the forests with highest integrity to be retained they should ideally be mapped using nationally appropriate criteria by the countries that hold them, formally recognized, prioritized in spatial plans, and placed under effective management (e.g. protected areas and other effective conservation areas, lands under Indigenous control etc.). These forests must be protected from industrial development impacts that degrade them through sensible public and private sector policy that is effective at relevant scales [12,34]. Our global assessment reveals where these places are found, and can be refined at more local scales where better data are available.

Around a third of global forests had already been cleared by 2000 [35], and we show that at least 59% of what remains has low to medium integrity, with > 50% falling in these two broad categories in every biogeographical realm. These levels of human modification result partly from the large areas affected by relatively diffuse anthropogenic pressures whose presence is inferred near forest edges, and by lost connectivity. We also map a surprising level of more localized, observed pressures, such as infrastructure and recent forest loss, which are seen in nearly a third of forested pixels worldwide.

Conservation strategies in these more heavily human-modified forests should focus on securing

any remaining fragments of forests in good condition, proactively protecting those forests most

vulnerable to further modification [7] and planning where restoration efforts might be most

effective [36-38]. In addition, effective management of production forests is needed to sustain yields

without further worsening their ecological integrity [39]. More research is required on how to

prioritize, manage, and restore forests with low to medium integrity [38,40], and the FLII presented

here might prove useful for this, for example, by helping prioritize where the best returns on

investment are, in combination with other sources of data [41].

Loss of forest integrity severely compromises many benefits of forests that are central to

achieving multiple Sustainable Development Goals and other societal targets [42,43]. Therefore,

governments must adopt policies and strategies to retain and restore the ecological integrity of

their forests, whilst ensuring that the solutions are also economically viable, socially equitable,

and politically acceptable within complex and highly diverse local contexts. This is an enormous

challenge and our efforts to map the degree of forest modification are designed both to raise

awareness of the importance of the issue, and to support implementation through target setting,

evidence-based planning, and enhanced monitoring efforts.

Whilst policy targets for halting deforestation are generally precise and ambitious, only vague

targets are typically stipulated around reducing levels of forest modification [9,44]. We urgently

need SMART (specific, measurable, achievable, realistic, and time-bound) goals and targets for

maintaining and restoring forest integrity that directly feed into higher-level biodiversity, climate,

land degradation, and sustainable development goals [45]. These types of targets should be included

within an over-arching target on ecosystems within the post-2020 Global Biodiversity

Framework, which is currently being negotiated among Parties to the CBD [46]. This target should

be outcome-focused and address both the extent and the integrity of ecosystems (e.g. using FLII

for forests), in a way that enables quantitative, measurable goals to be set but allows flexibility for

implementation between Parties.

In addition to broader goals in global frameworks, the retention and restoration of forest integrity

should also be addressed in nationally-defined goals embodied in, and aligned between,

Nationally Determined Contributions under the UNFCCC, efforts to stop land degradation and

achieve land degradation neutrality under the UNCCD, and National Biodiversity Strategy and

Action Plans under the CBD. Since no single metric can capture all aspects of a country's

environmental values, efforts to conserve high levels of forest integrity should be complemented

by consideration of areas support important values according to other measures (e.g. Key

Biodiversity Areas [47] and notable socio-cultural landscapes).

The overall level and pervasiveness of impacts on Earth's remaining forests is likely even more

severe than our findings suggest, because some input data layers, despite being the most

comprehensive available, are still incomplete as there are lags between increases in human

pressures and our ability to capture them in spatial datasets e.g., infrastructure, [48,49, see also Fig. S1 and

text S5]. For example, roads and seismic lines used for natural resource exploration and extraction in

British Columbia, Canada, are not yet fully reflected in global geospatial datasets Fig. S1; see also

[50]. Furthermore, because natural fires are such an important part of the ecology of many forest

systems (e.g. boreal forests) and because we cannot consistently identify anthropogenic fires from

natural fires at a global scales [51] we have taken a strongly conservative approach to fire in our

calculations, treating all tree cover loss in 10 km pixels where fire was the dominant driver *(23)* as

temporary, and not treating such canopy loss as evidence of observed human pressure. Varying

these assumptions where human activity is shown to be causing permanent tree cover losses,

increasing fire return frequencies, or causing fire in previously fire-free systems would result in

lower forest extent and/or lower forest integrity scores in some regions than we report.

We map forest integrity based on quantifiable processes over the recent past (since 2000). In

some areas modification that occurred prior to this (e.g. historical logging) is not detectable by

our methods but may have influenced the present-day integrity of the forest so, in such cases, we

may overestimate forest integrity. This is another reason why our index should be considered as

conservative, and we therefore recommend that the index be used alongside other lines of

evidence to determine the absolute level of ecological integrity of a given area. Moreover, the

definition of forest in this study is all woody vegetation taller than 5 m, following [22] and hence

includes not only naturally regenerated forests but also tree crops, planted forests, wooded

agroforests and urban tree cover in some cases. Users should be mindful of this when interpreting

the results, especially when observing areas with low forest integrity scores. Inspection of the

results for selected countries with reliable plantation maps [52] shows that the great majority of

planted forests have low forest integrity scores, because they are invariably associated with dense

infrastructure, frequent canopy replacement and patches of farmland.

We note our measure of forest integrity does not address past, current and future climate change.

As climate change affects forest condition both directly and indirectly, this is a clear shortfall and

needs research attention. The same is true for invasive species, as there is no globally coherent

data on the range of those invasive species that degrade forest ecosystems, although this issue is

indirectly addressed since the presence of many invasive species are likely spatially correlated

with the human pressures that we use as drivers in our model [53]. If global data became available it

would also be valuable to incorporate governance effectiveness into our model, because there are

potentially contexts (e.g. well-managed protected areas and community lands, production forests under 'sustainable forest management') where the impacts associated with the human pressures we base our map on are at least partially ameliorated [39], and enhanced governance is also likely to be a significant component of some future strategies to maintain and enhance forest integrity.

The framework we present has great potential to be tailored for use at smaller scales, ranging from regional to national and sub-national scales, and even to individual management units. Forest definitions and the relative weights of the global parameters we use can be adjusted to fit local contexts and, in many cases, better local data could be substituted, or additional variables incorporated. This would increase the precision of the index in representing local realities, and increase the degree of ownership amongst national and local stakeholders whose decisions are so important in determining forest management trajectories.

Methods

To produce our global Forest Landscape Integrity Index (FLII), we combined four sets of spatially explicit datasets representing: (i) forest extent [22]; (ii) 'observed' pressure from high impact, localized human activities for which spatial datasets exist, specifically: infrastructure, agriculture, and recent deforestation [53]; (iii) 'inferred' pressure associated with edge effects [54], and other diffuse processes, (e.g. activities such as hunting and selective logging) [55] modelled using proximity to observed pressures; and iv) anthropogenic changes in forest connectivity due to forest loss [56 see Table S1 for data sources]. These datasets were combined to produce an index score for each forest pixel (300m), with the highest scores reflecting the highest forest integrity (Fig 1), and applied to forest extent for the start of 2019. We use globally consistent parameters for all

elements (i.e. parameters do not vary geographically). All calculations were conducted in Google

Earth Engine (GEE) [57].

Forest extent

We derived a global forest extent map for 2019 by subtracting from the Global Tree Cover

product for 2000 [22] annual Tree Cover Loss 2001-2018, except for losses categorized by Curtis

and colleagues [23] as those likely to be temporary in nature (i.e. those due to fire, shifting

cultivation and rotational forestry). We applied a canopy threshold of 20% based on related

studies e.g. [29,58] and resampled to 300m resolution and used this resolution as the basis for the rest

of the analysis (see text S1 for further mapping methods).

Observed human pressures

We quantify observed human pressures (P) within a pixel as the weighted sum of impact of

infrastructure (I; representing the combined effect of 41 types of infrastructure weighted by their

estimated general relative impact on forests (Table S3), agriculture (A) weighted by crop intensity

(indicated by irrigation levels), and recent deforestation over the past 18 years (H; excluding

deforestation from fire, see Discussion). Specifically, for pixel i:

$$P_i = \exp(-\beta_1 I_i) + \exp(-\beta_2 A_i) + \exp(-\beta_3 H_i)$$

whereby the values of β were selected so that the median of the non-zero values for each

component was 0.75. This use of exponents is a way of scaling variables with non-commensurate

units so that they can be combined numerically, while also ensuring that the measure of observed

pressure is sensitive to change (increase or decrease) in the magnitude of any of the three

components, even at large values of I, A or H. This is an adaptation of the 'Human Footprint'

methodology [53]. See text S3 for further details.

Inferred human pressures

Inferred pressures are the diffuse effects of a set of processes for which directly observed datasets

do not exist, that include microclimate and species interactions relating to the creation of forest

edges [59] and a variety of intermittent or transient anthropogenic pressures such as: selective

logging, fuelwood collection, hunting; spread of fires and invasive species, pollution, and

livestock grazing [55,60,61]. We modelled the collective, cumulative impacts of these inferred effects

through their spatial association with observed human pressure in nearby pixels, including a

decline in effect intensity according to distance, and a partitioning into stronger short-range and

weaker long-range effects. The inferred pressure (P') on pixel *i* from source pixel *j* is:

$$P'_{i,j} = P_j\,(w_{i,j} + v_{i,j})$$

where $w_{i,j}$ is the weighting given to the modification arising from short-range pressure, as a

function of distance from the source pixel, and $v_{i,j}$ is the weighting given to the modification

arising from long-range pressures.

Short-range effects include most of the processes listed above, which together potentially affect

most biophysical features of a forest, and predominate over shorter distances. In our model they

decline exponentially, approach zero at 3 km, and are truncated to zero at 5 km (see text S4).

$$w_{i,j} = \alpha \exp(-\lambda \times d_{i,j}) \qquad \text{[for } d_{i,j} \leq 5 \text{ km]}$$

$$w_{i,j} = 0 \qquad \text{[for } d_{i,j} > 5 \text{ km]}$$

where α is a constant set to ensure that the sum of the weights across all pixels in range is 1.85

(see below), λ is a decay constant set to a value of 1 (see [62] and other references in text S4) and $d_{i,j}$

is the Euclidean distance between the centres of pixels i and j expressed in units of km.

Long-range effects include over-exploitation of high socio-economic value animals and plants,

changes to migration and ranging patterns, and scattered fire and pollution events. We modelled

long-range effects at a uniform level at all distances below 6 km and they then decline linearly

with distance, conservatively reaching zero at a radius of 12 km [55,63 and other references in text S4]:

$$v_{i,j} = \gamma \qquad \text{[for } d_{i,j} \leq 6 \text{ km]}$$

$$v_{i,j} = \gamma \times (12 - d_{i,j})/6 \qquad \text{[for } 6 \text{ km} < d_{i,j} \leq 12 \text{ km]}$$

$$v_{i,j} = 0 \qquad \text{[for } d_{i,j} > 12 \text{ km]}$$

Where γ is a constant set to ensure that the sum of the weights across all pixels in range is 0.15

and $d_{i,j}$ is the Euclidean distance between the centres of pixels i and j, expressed in kilometres.

The form of the weighting functions for short- and long-range effects and the sum of the weights

$(\alpha+\gamma)$ were specified based on a hypothetical reference scenario where a straight forest edge is

adjacent to a large area with uniform human pressure, and ensuring that in this case total inferred

pressure immediately inside the forest edge is equal to the pressure immediately outside, before

declining with distance. γ is set to 0.15 to ensure that the long-range effects conservatively

contribute no more than 5% to the final index in the same scenario, based on expert opinion and

supported e.g. Berzaghi *et al.* [64] regarding the approximate level of impact on values that would

be affected by severe defaunation and other long-range effects.

The aggregate effect from inferred pressures (P') on pixel *i* from all *n* pixels within range (*j=1* to

j=n) is then the sum of these individual, normalized, distance-weighted pressures, i.e.

$$P'_i = \Sigma_{[j=1....n]}\, P'_{i,j}$$

Loss of forest connectivity

Average connectivity of forest around a pixel was quantified using a method adapted from Beyer

et al. [56]. The connectivity C_i around pixel *i* surrounded by n other pixels within the maximum

radius (numbered *j*=1, 2…n) is given by:

$$C_i = \Sigma_{[j=1...n]}\, (F_j G_{i,j})$$

where F*j* is the forest extent is a binary variable indicating if forested (1) or not (0) and G*i,j* is the

weight assigned to the distance between pixels *i* and *j*. G*i,j* uses a normalized Gaussian curve,

with σ = 20km and distribution truncated to zero at 4σ for computational convenience (see text

S3). The large value of σ captures landscape connectivity patterns operating at a broader scale

than processes captured by other data layers. C_i ranges from 0 to 1 ($C_i \in [0,1]$).

Current Configuration (CC_i) of forest extent in pixel i was calculated using the final forest extent

map and compared to the Potential Configuration (PC) of forest extent without extensive human

modification, so that areas with naturally low connectivity, e.g. coasts and natural vegetation

mosaics, are not penalized. PC was calculated from a modified version of the map of Laestadius

et al. [35] and resampled to 300 m resolution (see text S2 for details). Using these two measures, we

calculated Lost Forest Configuration (LFC) for every pixel as:

$$LFC_i = 1 - (CC_i/PC_i)$$

Values of $CC_i/PC_i > 1$ are assigned a value of 1 to ensure that LFC is not sensitive to apparent

increases in forest connectivity due to inaccuracy in estimated potential forest extent – low values

represent least loss, high values greatest loss ($LFC_i \in [0,1]$).

Calculating the Forest Landscape Integrity Index

The three constituent metrics, LFC, P and P', all represent increasingly modified conditions the

larger their values become. To calculate a forest integrity index in which larger values represent

less degraded conditions we therefore subtract the sum of those components from a fixed large

value (here, 3). Three was selected as our assessment indicates that values of LFC + P + P' of 3 or

more correspond to the most severely degraded areas. The metric is also rescaled to a convenient

scale (0-10) by multiplying by an arbitrary constant (10/3). The FLII for forest pixel *i* is thus

calculated as:

$$FLII_i = [10/3] * (3 - \min(3, [P_i + P'_i + LFC_i]))$$

where $FLII_i$ ranges from 0 - 10, forest areas with no modification detectable using our methods

scoring 10 and those with the most scoring 0.

Illustrative forest integrity classes

Whilst a key strength of the index is its continuous nature, the results can also be categorized for a

range of purposes. In this paper three illustrative classes were defined, mapped and summarized

to give an overview of broad patterns of integrity in the world's forests. The three categories were

defined as follows.

High Forest Integrity (scores ≥9.6) Interiors and natural edges of more or less unmodified

naturally-regenerated (i.e. non-planted) forest ecosystems, comprised entirely or almost entirely

of native species, occurring over large areas either as continuous blocks or natural mosaics with

non-forest vegetation; typically little human use other than low intensity recreation or spiritual

uses and/or low intensity extraction of plant and animal products and/or very sparse presence of

infrastructure; key ecosystem functions such as carbon storage, biodiversity and watershed

protection and resilience expected to be very close to natural levels (excluding any effects from

climate change) although some declines possible in the most sensitive elements (e.g. some high

value hunted species).

Medium Forest Integrity (scores >6.0 but <9.6) Interiors and natural edges of naturally-

regenerated forest ecosystems in blocks smaller than their natural extent but large enough to have

some core areas free from strong anthropogenic edge effects (e.g. set asides within forestry areas,

fragmented protected areas), dominated by native species but substantially modified by humans

through a diversity of processes that could include fragmentation, creation of edges and proximity

to infrastructure, moderate or high levels of extraction of plant and animal products, significant

timber removals, scattered stand-replacement events such as swidden and/or moderate changes to

fire and hydrological regimes; key ecosystem functions such as carbon storage, biodiversity,

watershed protection and resilience expected to be somewhat below natural levels (excluding any

effects from climate change).

Low Forest Integrity (score ≤6.0): Diverse range of heavily modified and often internally

fragmented ecosystems dominated by trees, including (i) naturally regenerated forests, either in

the interior of blocks or at edges, that have experienced multiple strong human pressures, which

may include frequent stand-replacing events, sufficient to greatly simplify the structure and

species composition and possibly result in significant presence of non-native species, (ii) tree

plantations and, (iii) agroforests; in all cases key ecosystem functions such as carbon storage,

biodiversity, watershed protection and resilience expected to be well below natural levels

(excluding any effects from climate change).

The numerical category boundaries were derived by inspecting FLII scores for a wide selection of

benchmark locations whose forest integrity according to the category definitions was known to

the authors, see text S6 and Table S4.

Protected areas analysis

Data on protected area location, boundary, and year of inscription were obtained from the

February 2018 World Database on Protected Areas [65]. Following similar global studies e.g. [66], we

extracted protected areas from the WDPA database by selecting those areas that have a status of

"designated", "inscribed", or "established", and were not designated as UNESCO Man and

Biosphere Reserves. We included only protected areas with detailed geographic information in

the database, excluding those represented as a point only. To assess integrity of protected forest,

we extracted all 300m forest pixels that were at least 50% covered by a formal protected area and

measured the average FLII score.

References and Notes

Seymour, F. & Harris, N. L. Reducing tropical deforestation. *Science* **365**, 756-757, doi:10.1126/science.aax8546 (2019).
Pearson, T. R. H., Brown, S., Murray, L. & Sidman, G. Greenhouse gas emissions from tropical forest degradation: an underestimated source. *Carbon Balance and Management* **12**, 3, doi:10.1186/s13021-017-0072-2 (2017).
Moen, J. *et al.* Eye on the Taiga: Removing Global Policy Impediments to Safeguard the Boreal Forest. *Conservation Letters* **7**, 408-418, doi:10.1111/conl.12098 (2014).
Erb, K.-H. *et al.* Unexpectedly large impact of forest management and grazing on global vegetation biomass. *Nature* **553**, 73-76, doi:10.1038/nature25138 (2018).
Scholes, R. *et al.* IPBES (2018): Summary for policymakers of the assessment report on land degradation and restoration of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. (2018).
Parrish, J. D., Braun, D. P. & Unnasch, R. S. Are we conserving what we say we are? Measuring ecological integrity within protected areas. *Bioscience* **53**, 851-860 (2003).
Cochrane, M. A. *et al.* Positive Feedbacks in the Fire Dynamic of Closed Canopy Tropical Forests. *Science* **284**, 1832-1835, doi:10.1126/science.284.5421.1832 (1999).
Erdozain, M. *et al.* Demand for provisioning ecosystem services as a driver of change in the Canadian boreal zone1. *Environmental Reviews* **27**, 166-184, doi:10.1139/er-2018-0064 (2018).
Watson, J. E. *et al.* The exceptional value of intact forest ecosystems. *Nature ecology & evolution*, 1 (2018).
Lewis, S. L. *et al.* Increasing carbon storage in intact African tropical forests. *Nature* **457**, 1003, doi:10.1038/nature07771
https://www.nature.com/articles/nature07771#supplementary-information (2009).
11   Mello, K. d., Valente, R. A., Randhir, T. O. & Vettorazzi, C. A. Impacts of tropical forest cover on water quality in agricultural watersheds in southeastern Brazil. *Ecological Indicators* **93**, 1293-1301, doi:https://doi.org/10.1016/j.ecolind.2018.06.030 (2018).
12   Garnett, S. T. *et al.* A spatial overview of the global importance of Indigenous lands for conservation. *Nature Sustainability* **1**, 369-374, doi:10.1038/s41893-018-0100-6 (2018).
13   Bonan, G. B. Forests and Climate Change: Forcings, Feedbacks, and the Climate Benefits of Forests. *Science* **320**, 1444-1449, doi:10.1126/science.1155121 (2008).
14   Barlow, J. *et al.* Anthropogenic disturbance in tropical forests can double biodiversity loss from deforestation. *Nature* **535**, 144, doi:10.1038/nature18326 (2016).
15   Betts, M. G. *et al.* Global forest loss disproportionately erodes biodiversity in intact landscapes. *Nature* **547**, 441, doi:10.1038/nature23285
https://www.nature.com/articles/nature23285#supplementary-information (2017).
Gibson, L. Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* **478**, doi:10.1038/nature10425 (2011).
Di Marco, M., Ferrier, S., Harwood, T. D., Hoskins, A. J. & Watson, J. E. M. Wilderness areas halve the extinction risk of terrestrial biodiversity. *Nature* **573**, 582-585, doi:10.1038/s41586-019-1567-7 (2019).

Laurance, W. F. & Peres, C. A. *Emerging threats to tropical forests*. (University of Chicago Press, 2006).

Gauthier, S., Bernier, P., Kuuluvainen, T., Shvidenko, A. & Schepaschenko, D. Boreal forest health and global change. *Science* **349**, 819-822 (2015).

FAO. Assessing forest degradation. Towards the development of globally applicable guidelines. . (FAO, Rome, 2011).

FAO. Global Forest Land-Use Change 1990-2005. (Food and Agriculture Organisation of the United Nations, Rome, 2012).

Hansen, M. C. *et al.* High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science* **342**, 850-853, doi:10.1126/science.1244693 (2013).

Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A. & Hansen, M. C. Classifying drivers of global forest loss. *Science* **361**, 1108-1111, doi:10.1126/science.aau3445 (2018).

Mitchell, A. L., Rosenqvist, A. & Mora, B. Current remote sensing approaches to monitoring forest degradation in support of countries measurement, reporting and verification (MRV) systems for REDD+. *Carbon Balance and Management* **12**, 9, doi:10.1186/s13021-017-0078-9 (2017).

Wang, Y. *et al.* Mapping tropical disturbed forests using multi-decadal 30 m optical satellite imagery. *Remote Sensing of Environment* **221**, 474-488, doi:https://doi.org/10.1016/j.rse.2018.11.028 (2019).

Venter, O. *et al.* Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nature Communications* **7**, 12558 (2016).

Shapiro, A. C., Aguilar-Amuchastegui, N., Hostert, P. & Bastin, J.-F. Using fragmentation to assess degradation of forest edges in Democratic Republic of Congo. *Carbon Balance and Management* **11**, 11, doi:10.1186/s13021-016-0054-9 (2016).

Hansen, A. *et al.* Global humid tropics forest structural condition and forest structural integrity maps. *Scientific Data* **6**, 232, doi:10.1038/s41597-019-0214-3 (2019).

Potapov, P. *et al.* The last frontiers of wilderness: Tracking loss of intact forest landscapes from 2000 to 2013. *Science Advances* **3**, doi:10.1126/sciadv.1600821 (2017).

Watson, James E. M. *et al.* Catastrophic Declines in Wilderness Areas Undermine Global Environment Targets. *Current Biology* **26**, 2929-2934, doi:https://doi.org/10.1016/j.cub.2016.08.049 (2016).

Olson, D. M. & Dinnerstein, E. G. The Global 200: a representation approach to conserving the Earth's most biologically valuable ecoregions. *Conserv. Biol.* **12**, 502-515 (1998).

Pressey, R. L., Watts, M. E. & Barret, T. W. Is maximising protection the same as minimizing loss? Efficiency and retention as alternative measures of the effectiveness of proposed reserves. *Ecol Letters* **7**, 1035-1046 (2004).

Maron, M., Simmonds, J. S. & Watson, J. E. M. Bold nature retention targets are essential for the global environment agenda. *Nature Ecology & Evolution* **2**, 1194-1195, doi:10.1038/s41559-018-0595-2 (2018).

Dudley, N. *et al.* The essential role of other effective area-based conservation measures in achieving big bold conservation targets. *Global Ecology and Conservation* **15**, e00424, doi:https://doi.org/10.1016/j.gecco.2018.e00424 (2018).

Laestadius, L. *et al.* Opportunities for forest landscape restoration. *Unasylva* **62**, 238 (2011).

Edwards, D. P., Tobias, J. A., Sheil, D., Meijaard, E. & Laurance, W. F. Maintaining ecosystem function and services in logged tropical forests. *Trends Ecol. Evol.* **29**, doi:10.1016/j.tree.2014.07.003 (2014).

Runting, R. K. *et al.* Larger gains from improved management over sparing–sharing for tropical forests. *Nature Sustainability* **2**, 53-61, doi:10.1038/s41893-018-0203-0 (2019).

Chazdon, R. L. *et al.* A Policy-Driven Knowledge Agenda for Global Forest and Landscape Restoration. *Conservation Letters* **10**, 125-132, doi:doi:10.1111/conl.12220 (2017).

MacDicken, K. G. *et al.* Global progress toward sustainable forest management. *For. Ecol. Manage.* **352**, 47-56, doi:https://doi.org/10.1016/j.foreco.2015.02.005 (2015).

Ghazoul, J. & Chazdon, R. Degradation and Recovery in Changing Forest Landscapes: A Multiscale Conceptual Framework. *Annual Review of Environment and Resources* **42**, 161-188, doi:10.1146/annurev-environ-102016-060736 (2017).

Strassburg, B. B. N. *et al.* Strategic approaches to restoring ecosystems can triple conservation gains and halve costs. *Nature Ecology & Evolution* **3**, 62-70, doi:10.1038/s41559-018-0743-8 (2019).

Pandit, R. *et al.* Summary for policymakers of the assessment report on land degradation and restoration of the Intergovernmental Science-Policy Platform on Biodiversity and Ecosystem Services. (2018).

Osuri, A. M. *et al.* Contrasting effects of defaunation on aboveground carbon storage across the global tropics. *Nature Communications* **7**, 11351, doi:10.1038/ncomms11351 https://www.nature.com/articles/ncomms11351#supplementary-information (2016).

Mackey, B. Policy options for the world's primary forests in multilateral environmental agreements. *Conserv. Lett.* **8**, doi:10.1111/conl.12120 (2015).

Butchart, S. H. M., Di Marco, M. & Watson, J. E. M. Formulating Smart Commitments on Biodiversity: Lessons from the Aichi Targets. *Conservation Letters* **9**, 457-468, doi:10.1111/conl.12278 (2016).

Watson, J. E. *et al.*   (Nature Publishing Group, 2020).

Stuart, S. N. *et al.* Clarifying the key biodiversity areas partnership and programme. *Biodivers. Conserv.* **27**, 791-793, doi:10.1007/s10531-017-1490-x (2018).

Barrington-Leigh, C. & Millard-Ball, A. The world's user-generated road map is more than 80% complete. *PLOS ONE* **12**, e0180698, doi:10.1371/journal.pone.0180698 (2017).

Hughes, A. C. Have Indo-Malaysian forests reached the end of the road? *Biol. Conserv.* **223**, 129-137, doi:https://doi.org/10.1016/j.biocon.2018.04.029 (2018).

Pasher, J., Seed, E. & Duffe, J. Development of boreal ecosystem anthropogenic disturbance layers for Canada based on 2008 to 2010 Landsat imagery. *Canadian Journal of Remote Sensing* **39**, 42-58, doi:10.5589/m13-007 (2013).

Erb, K.-H. *et al.* Land management: data availability and process understanding for global change studies. *Global Change Biol.* **23**, 512-533, doi:10.1111/gcb.13443 (2016).

Harris, N., E. , Goldman, L. & Gibbes, S. Spatial Database of Planted Trees (SDPT) Version 1.0. (World Resources Institute, Washington, DC, 2018).

Venter, O. Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nat. Commun.* **7**, doi:10.1038/ncomms12558 (2016).

Haddad, N. M. *et al.* Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* **1**, e1500052, doi:10.1126/sciadv.1500052 (2015).

Peres, C. A., Emilio, T., Schietti, J., Desmoulière, S. J. M. & Levi, T. Dispersal limitation induces long-term biomass collapse in overhunted Amazonian forests. *Proceedings of the National Academy of Sciences* **113**, 892-897, doi:10.1073/pnas.1516525113 (2016).

Beyer, H. L., Venter, O., Grantham, H. S. & Watson, J. E. M. Substantial losses in ecoregion intactness highlight urgency of globally coordinated action. *Conservation Letters* **n/a**, e12592, doi:10.1111/conl.12692.

Gorelick, N. *et al.* Google Earth Engine: Planetary-scale geospatial analysis for everyone. *Remote Sensing of Environment* **202**, 18-27, doi:https://doi.org/10.1016/j.rse.2017.06.031 (2017).

Heino, M. *et al.* Forest Loss in Protected Areas and Intact Forest Landscapes: A Global Analysis. *PLOS ONE* **10**, e0138918, doi:10.1371/journal.pone.0138918 (2015).

Laurance, W. F. *et al.* Ecosystem decay of Amazonian forest fragments: a 22-year investigation. *Conserv. Biol.* **16**, 605-618 (2002).

Cochrane, M. A. & Laurance, W. F. Fire as a large-scale edge effect in Amazonian forests. *Journal of Tropical Ecology* **18**, 311-325, doi:10.1017/S0266467402002237 (2002).

Zimmerman, B. L. & Kormos, C. F. Prospects for Sustainable Logging in Tropical Forests. *Bioscience* **62**, 479-487, doi:10.1525/bio.2012.62.5.9 (2012).

Chaplin-Kramer, R. *et al.* Degradation in carbon stocks near tropical forest edges. *Nature Communications* **6**, 10158, doi:10.1038/ncomms10158 https://www.nature.com/articles/ncomms10158#supplementary-information (2015).

Maisels, F. *et al.* Devastating Decline of Forest Elephants in Central Africa. *PLOS ONE* **8**, e59469, doi:10.1371/journal.pone.0059469 (2013).

Berzaghi, F. *et al.* Carbon stocks in central African forests enhanced by elephant disturbance. *Nature Geoscience* **12**, 725-729, doi:10.1038/s41561-019-0395-6 (2019).

IUCN, U.-W. a. Protected Planet: The World Database on Protected Areas (WDPA). (UNEP-WCMC, Cambridge, UK, 2018).

Butchart, S. H. M. *et al.* Shortfalls and Solutions for Meeting National and Global Conservation Area Targets. *Conservation Letters* **8**, 329-337, doi:10.1111/conl.12158 (2015).

Gibson, L. *et al.* Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* **478**, 378-381 (2011).

ESA. Land Cover CCI Product User Guide Version 2. Tech. Rep., (2017).

Theobald, D. M. A general model to quantify ecological integrity for landscape assessments and US application. *Landscape Ecol.* **28**, 1859-1874 (2013).

Sanderson, E. W. *et al.* The human footprint and the last of the wild. *Bioscience* **52**, 891-904 (2002).

Kennedy, C. M., Oakleaf, J. R., Theobald, D. M., Baruch-Mordo, S. & Kiesecker, J. Managing the middle: A shift in conservation priorities based on the global human modification gradient. *Global Change Biol.* **25**, 811-826 (2019).

Thompson, I. D. *et al.* An Operational Framework for Defining and Monitoring Forest Degradation. *Ecology and Society* **18** (2013).

Contributors.   (ed OpenStreetMap) (2015).

Teluguntla, P. *et al.*   (ed NASA EOSDIS Land Processes DAAC) (2016).

Tulloch, A. I., Gordon, A., Runge, C. A. & Rhodes, J. R. Integrating spatially realistic infrastructure impacts into conservation planning to inform strategic environmental assessment. *Conservation Letters* **12**, e12648 (2019).

Benítez-López, A., Alkemade, R. & Verweij, P. A. The impacts of roads and other infrastructure on mammal and bird populations: a meta-analysis. *Biol. Conserv.* **143**, 1307-1316 (2010).

Broadbent, E. N. *et al.* Forest fragmentation and edge effects from deforestation and selective logging in the Brazilian Amazon. *Biol. Conserv.* **141**, 1745-1757 (2008).

Benítez-López, A., Santini, L., Schipper, A. M., Busana, M. & Huijbregts, M. A. J. Intact but empty forests? Patterns of hunting-induced mammal defaunation in the tropics. *PLOS Biology* **17**, e3000247, doi:10.1371/journal.pbio.3000247 (2019).

Lamarche, C. *et al.* Compilation and validation of SAR and optical data products for a complete and global map of inland/ocean water tailored to the climate modeling community. *Remote Sensing* **9**, 36 (2017).

Gumma, M. *et al.* NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Global Food Security-support Analysis Data (GFSAD) Cropland Extent 2015 South Asia, Afghanistan, Iran 30 m V001. (2017).

Massey, R. *et al.*   (ed NASA EOSDIS Land Processes DAAC) (2017).

Oliphant, A. *et al.* NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Global Food Security-support Analysis Data (GFSAD) Cropland Extent 2015 Southeast Asia 30 m V001. (2017).

Phalke, A. *et al.*   (ed NASA EOSDIS Land Processes DAAC) (2017).

Teluguntla, P. *et al.* NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Global Food Security-support Analysis Data (GFSAD) Cropland Extent 2015 Australia, New Zealand, China, Mongolia 30 m V001. (2017).

Xiong, J. *et al.* NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Global Food Security-support Analysis Data (GFSAD) Cropland Extent 2015 Africa 30 m V001. (2017).

Zhong, Y. *et al.*   (ed NASA EOSDIS Land Processes DAAC) (2017).

Pekel, J.-F., Cottam, A., Gorelick, N. & Belward, A. S. High-resolution mapping of global surface water and its long-term changes. *Nature* **540**, 418-422 (2016).

Acknowledgments: We thank Peter Potapov, Dmitry Aksenov, and Matthew Hansen for

comments and advice.

Funding: The research for this paper was in part funded by the John D. and Catherine T.

MacArthur Foundation, Natasha and Dirk Ziff, Trillion Trees (a joint venture between

BirdLife International, Wildlife Conservation Society, and WWF-UK), and other generous

donors.

Author contributions: Conceived and design the study: HG, TE and JEW, collected data and

developed the model: AD, HG, TE, HB, RS, analyzed and interpreted the results: AD,

HG, TE, HB, RS, KJ, JEW, wrote draft manuscript: HG, TE and JEW, contributed to the

writing of the manuscript: all co-authors.

Competing interests: There are no competing interests

Data and materials availability: Data will be available for download when published.

Figures and Tables

Table 1. A summary of the Forest Landscape Integrity Index scores for each biogeographic realm

globally, measuring the mean score, in addition to the area and proportion of realm for each

category of integrity. Scores are divided into three categories of integrity: high, medium and low.

Table 2. A summary of the Forest Landscape Integrity Index scores for each type of protected

area designation based on the IUCN Protected Areas categories measuring mean score, in

addition to the area and proportion of realm for each category of integrity. Scores are divided into

three categories of integrity: high, medium and low.

Figure 1. The Forest Landscape Integrity Index was constructed based on three main data inputs:

1) observed pressures (infrastructure, agriculture, tree cover loss), 2) inferred pressure modelled

based on proximity to the observed pressures, and 3) change in forest connectivity.

Figure 2. A global map of Forest Landscape Integrity for 2019. Three regions are highlighted

including A) USA, B) Equatorial Guinea C) Myanmar. For a) shows the edge of Smoky

Mountains National Park in Tennessee b) shows a logging truck passing through some partially

degraded forest along a newly constructed highway in Shan State, c) shows an intact mangrove

forest within Reserva Natural del Estuario del Muni, near the border with Gabon. The stars

indicate approximately where the photos were taken (A2, B2 and C2).

Figure 3. The Forest Landscape Integrity Index for 2019 categorized into three broad, illustrative

classes and mapped for across each biogeographic realm (A – G). The size of the pie charts

indicates the relative size of the forests within each realm (A - G), and H shows all the world's

forest combined.

50 **Figure 4.** The Forest Landscape Integrity Index for 2019 categorized into three broad, illustrative

51 classes for each major forested country in the world. (A) countries with a forest extent larger than

52 1 million km^2, and (B) countries with forest extent between 1 million km^2 and 100,000 km^2 of

53 forest. The size of the bar represents the area of a country's forests.

56
57

58   Table 1
59

Biogeographic Realm	Total forest	FLII	High (9.6 - 10)		Medium (6 – 9.6)		Low (0 - 6)	
	Km²	*Mean*	*Km²*	*% of realm*	*Km²*	*% of realm*	*Km²*	*% of realm*
Afrotropic	7,362,740	7.34	2,450,953	33.3	2,903,483	39.4	2,008,304	27.3
Australasia	1,711,684	8.05	656,701	38.4	753,188	44.0	301,796	17.6
Indo-malayan	3,596,249	5.9	420,977	11.7	1,599,049	44.5	1,576,223	43.8
Neotropic	10,271,519	7.81	4,579,406	44.6	3,122,706	30.4	2,569,407	25.0
Oceania	23,389	7.66	5,279	22.6	14,331	61.3	3,780	16.2
Palearctic	12,172,668	8	5,571,997	45.8	3,910,629	32.1	2,690,042	22.1
Nearctic	7,794,117	7.84	3,716,855	47.7	2,257,518	29.0	1,819,744	23.3
Total	42,932,367	7.76	17,402,170		14,560,903		10,969,294	

771   Table 2.
772
773
774

Protected Area Category	Total forest	FLII	High (score 9.6 - 10)		Medium (score 6 – 9.6)		Low (score 0 - 6)	
	Km²	Mean	Km²	% of protected area	Km²	% of protected area	Km²	% of protected area
Ia (strict nature reserve)	439,082	9.27	304,329	69.31	106,703	24.3	28,049	6.39
Ib (wilderness area)	367,330	9.22	240,453	65.46	102,096	27.79	24,780	6.75
II (national park)	1,900	9.14	1,223,138	64.38	540,805	28.46	136,056	7.16
III (natural monument or feature)	113,805	8.49	54,476	47.87	40,021	35.17	19,308	16.97
IV (habitat/species management area)	838,707	8.69	432,828	51.61	268,027	31.96	137,850	16.44
V (protected landscape/seascape)	840,919	6.4	224,491	26.7	295,769	35.17	320,658	38.13
VI (Protected area with sustainable use of natural resources)	1,472,278	9.21	1,026,169	69.7	344,617	23.41	101,491	6.89
Not Applicable / Not Assigned / Not Reported	2,613,541	8.29	1,030,430	39.42	906,745	34.69	676,365	25.88
All Protected Areas	8,585,661	8.55	4,536,314	52.83	2,694,784	30.34	1,444,562	16.82

75   Fig 1

76
77
78

Forest Landscape Integrity Index

Low Integrity (0 - 6)	Medium Integrity (6 – 9.6)	High Integrity (>9.6)

Fig 4.

Supplementary Materials

Text S1. Mapping forest extent

We generated a preliminary base map of global forest extent for the start of 2019 at 30 m resolution by subtracting annual Tree Cover Loss 2001-2018 (with exceptions noted in the next paragraph) from the Global Tree Cover 2000 product [22] using a canopy cover threshold of 20%. This is one of the most widely used tree cover datasets globally, so it has been widely tested in many settings and its strengths and constraints are well understood. It has many advantages, including its high resolution, high accuracy, global coverage, annual time series and good prospects of sustainability in the coming years. The definition of forest in the source dataset is all woody vegetation taller than 5 m and hence includes naturally regenerated forests as well as tree crops, planted forests, wooded agroforests and urban tree cover. No globally consistent dataset was available that allowed natural and planted tree cover to be consistently distinguished in this study. Therefore, we should be mindful of the many differences between planted and natural tree cover (e.g.[67]).

More than 70% of the tree cover loss shown by the Hansen *et al.* [22] products has been found to be in 10 km pixels where the dominant loss driver is temporary and so tree cover is expected to return above the forest definition threshold within a short period [23]. It is important to take account of this issue as treating all such areas as permanent loss would severely under-estimate current forest cover in many regions. However, no global map of forest cover gain exists for the study period other than the 2000-2012 gain product from Hansen *et al.* [22], so we developed an alternative approach. When removing annual loss shown by the Global Tree Cover Loss product cited above we elected not to remove any loss that was in a 10 km pixel categorized by Curtis *et*

al. [23] as dominated by temporary loss under the categories of fire, shifting cultivation or rotational

forestry. This resulted in the adjusted preliminary forest base map. The balance of evidence is that

the great majority of such areas would have begun to regenerate and hence qualify as forest by

our definition again by 2019 or soon after [23]. The anthropogenically disturbed nature of many of

these areas of temporary tree cover loss and recovery is reflected in scoring within the index,

because temporary tree cover loss in the categories of shifting cultivation or rotational forestry is

treated as an observed pressure. We do not treat tree cover loss through fire as an observed

pressure, because fires are often part of natural processes, especially in the boreal zone. This

makes our global index conservative as a measure of degradation in these zones, because in some

locations fires are anthropogenic in nature.

The adjusted preliminary base map was then resampled to a final base map for 2019 at 300m

resolution using a pyramid-by-mode decision rule, with the resulting pixels simply classified as

forest or non-forest based on a majority rule. The FLII was calculated for every forest pixel but

not for non-forest pixels. GEE performs calculations in WGS84. Supplementary analyses outside

GEE were applied using a Mollweide equal-area projection.

Text S2. Mapping potential forest configuration

Potential connectivity (PC) is calculated from an estimate of the potential extent of the forest zone

taken from Laestadius *et al.* [35], treating areas below 25% crown cover (this was the nearest class

to the threshold used in our tree cover dataset of 20%) as non-forest and resampling to 300 m

resolution. To minimize false instances of lost connectivity and ensure measures of forest

modification are conservative we masked from this data layer areas which we believe to include a

significant proportion of naturally unforested land using selected land-cover categories in ESA

([68]; see Table S1). Because these natural non-forest patches are shown in the Hansen *et al.* [22]

dataset but not Laestadius *et al.* [35], not excluding such classes would result in an inflated estimate

of the loss of connectivity and hence the level of degradation. We have elected to remain

conservative in our estimate of modification.

Text S3. Mapping observed human pressure

Several recent analyses have developed composite, multi-criteria indices of human pressure to

provide assessments of ecosystem condition for the USA [69] or globally [26,70,71]. Thompson *et al.* [72]

set out a framework specific to forest ecosystems that could indicate modification through a

balanced mix of available pressure and state variables. We adapted the methodology of Venter et

al. [26], informed by the other studies cited, to generate measures of (i) the modification of forest

associated with observed human pressure from infrastructure, agriculture and deforestation and

(ii) the more diffuse inferred modification effects (e.g. edge effects) whose presence is inferred

from proximity to these focal areas of human activity. Edge effects resulting entirely from natural

processes are excluded, because they do not represent modification by our definition, although,

like many other natural factors, they do also have a role in determining ecosystem benefits.

50 *Infrastructure*

52 We generated the infrastructure (I') data layer by rasterizing the OpenStreetMap data [73] from Feb

53 2018, using weights for each type of infrastructure as noted in Table S3. The weights were

54 derived from authors' expert opinion and experimentation with weights according to their relative

55 impact on forest condition.

Agriculture

For agriculture (A') we made a global binary composite of the croplands datasets produced by the USGS (Table 1) at 30 m resolution, and weighted each cropped pixel at this resolution by the likely intensity of cropping using the global irrigation dataset at 1km resolution (Teluguntla et al, [74]), with values of Irrigation Major = 2, Irrigation Minor = 1.5, Rainfed = 1. The average cropping intensity (including uncropped areas, which score zero) was then calculated across the whole of each 300 m pixel of our final basemap.

Deforestation

For deforestation (H') we made a binary composite of tree cover loss 2001-2018 at 30 m resolution [22], masked out 30 m pixels already classified as agriculture in the preceding step to avoid double-counting, and excluded loss predicted by Curtis *et al*. [23] to be most likely caused by fires, to give a conservative data layer of recent permanent and temporary tree cover loss indicative of human activity in the immediate vicinity. We excluded small clusters of 6 or fewer pixels (0.54 ha) because they may have been natural tree cover loss (e.g. small windthrows) or classification errors. Each 30 m pixel was then weighted by its year of loss, giving higher weight to the most recent loss (2001 = 1, 2002 = 2, etc.). The average 'recentness' of deforestation (including areas not deforested, which score zero) was then calculated across the whole of each 300 m pixel of our base map.

Transformations

The exponential transformations described in the main text were used to convert I', A' and H' to the variables I, A and H respectively.

Text S4. Modelling inferred pressures using proximity to observed pressures

Each cell also experiences modification as a result of pressures originating from nearby cells that have observed human pressures, largely through the family of processes known as 'edge effects' [54]. Edge effects are partly a result of the changes relating to biophysical factors (such as humidity, wind, temperature and the increased presence of non-forest species) that accompany the creation of new edges in formerly continuous forest (as exemplified by the carefully controlled studies in tropical forests summarized by Laurance *et al.* [59]). They also result in part from the increased pressure associated with human activities within tropical forest near to edges such as logging [61], anthropogenic fire [60], hunting [55], livestock grazing, pollution, visual and auditory disturbances, etc. These multiple factors are synergistic and so we model them together, notwithstanding regional and local variations in the relative intensity of each one.

We model the inferred effect caused by each nearby source cell as a function of (a) the observed human pressure observed in that source cell and (b) a decline in the intensity of edge effects with distance from the source cell, based on a review of the literature. We then determine the total inferred effect on a given cell by summing the individual effects from all source cells within a certain range.

Two complementary types of inferred effect are modelled and added together. One relates to the diverse, strong, relatively short-range edge effects which decay to near zero over a few kilometers and have the potential to affect most biophysical features of a forest to a greater or lesser extent.

The other relates to weaker, longer-range effects such as over-hunting of high-value animals that

affect fewer biophysical features of a forest (and so have a much smaller maximum effect on

overall integrity) but can nonetheless have detectable effects in locations more than 10 km from

the nearest permanent human presence.

The literature on the spatial influence of short-term effects uses a variety of mathematical

descriptors, in two broad categories – continuous variables and distance belts. As we wish to

model edge effects as a continuous variable we concentrated on studies that have taken a similar

approach, and used distance-belt studies as ancillary data.

Chaplin-Kramer *et al.* [62] is a good example of a continuous variable approach, estimating detailed

biomass loss curves near tropical forest edges. Because they analyze a key forest condition

variable with a very large pantropical dataset we hypothesize that the exponential declines in

degradation with distance that they find are likely to be a common pattern and so we use a similar

framework for our more general model of degradation. We consider that a model of exponential

decay is also a sufficient approximation to the evidence presented by some authors as graphs

without an associated mathematical model (e.g., [60,75]) or analyzed using logistic regression (e.g.,

[76]). In our model we set the exponential decay constant to be broadly consistent with these four

studies, resulting in degradation at 1 km inside a forest that is approximately 37% of that at the

forest edge, declining to 14% at 2 km and near zero at 3 km. We truncate the distribution at 5 km

to minimize computational demands.

Distance-belt studies define the width of a belt within which edge effects are considered to occur,

and beyond which forests are considered to be free of edge effect. Belts of 1 km are commonly

used (e.g., [54]) but smaller distances may be used for specific parameters (e.g. 300 m for biomass

reduction near edges in DRC's primary forests; [27]). Our continuous variable approach is broadly

consistent with these studies, with the majority of our modelled degradation within a 1 km belt

and little extending beyond 2 km. While most individual edge effects reported in the literature

penetrate less than 100-300 m (e.g., [59,77]) most of the effects reported on in these studies relate to

the changed natural factors mentioned in an earlier paragraph, and are likely to be dwarfed in both

intensity and extent by edge effects relating to spillovers of human activity, so our model

emphasizes the spatial distribution of the latter (e.g., [60]). We consider our model of the levels of

modification to be conservative.

For the weaker, more widespread long-range effects we use recent large-scale studies of

defaunation, which is one of the key long-range pressures and also acts as a proxy for other

threats including harvest of high value plants (such as eaglewood *Aquilaria* spp. in tropical Asia),

occasional remote fires, pollution associated with artisanal mining, etc. We adopt a simplified

version of the distribution used by Peres *et al.* [55] to model hunting around settlements in the

Amazon, which sets $2\sigma=12$ km; this is likely conservative compared to evidence for hunting-

related declines in forest elephants in central Africa up to 60 km from roads [63] and the extensive

declines in large-bodied quarry species in remote areas in many regions modelled by Benitez-

Lopez *et al.* [78].

Text S5. Limitations in data: example with infrastructure data in British Columbia, Canada

OpenStreetMap (OSM) represents the most detailed publicly available relevant global dataset but

is nonetheless noted to be incomplete, even for one of the most heavily used categories of

infrastructure, paved roads [48]. No global assessment is available for the completeness of other

categories in the dataset. One of the key categories for forest integrity, unpaved roads used for

resource extraction, has been shown to be incomplete over much of insular South-east Asia [49]. In

Canada, for example, roads and other linear corridors used to explore, access and extract natural

resources (e.g., logging, oil and gas, and minerals) are sometimes missing. Government data for

the province of British Columbia (available at https://catalogue.data.gov.bc.ca/dataset/digital-

road-atlas-dra-master-partially-attributed-roads) demonstrates, for example, the larger extent and

density of regional roads as compared to OSM (Fig S1).

Text S6. Classification of Forest Landscape Integrity Index scores

In this paper, three illustrative classes were defined, mapped and summarized to give an overview

of broad patterns of degradation in the world's forests. Three categories were defined as set out in

the Materials and Methods. To determine the approximate levels of the FLII associated with these

three categories, benchmark locations were selected in sites that could unambiguously be assigned

to one of the categories using the authors' personal knowledge. At each site a single example

pixel was selected within a part of the area with relatively uniform scores. The sample points are

summarized in Table S4; they are widely spread across the world to ensure that the results are not

only applicable to a limited region. The scores at these points suggest the following category

boundaries:

- High FLII – 9.6-10
- Medium FLII – 6-9.6
- Low FLII – 0-6

Table S1. The datasets used to develop the Forest Ecosystem Integrity Index. The factor column indicates the component of the index the dataset was used in.

Dataset	Factor	Sources
Tree cover and tree cover loss	Forest extent, connectivity, observed and inferred pressures	Global Forest Cover datasets; Hansen *et al.* [22]; updates to 2018 available on-line from: http://earthenginepartners.appspot.com/science-2013-global-forest.
Major tree cover loss driver	Forest extent, observed and inferred pressures, connectivity	Curtis *et al.* [23]
Landover and ocean extent	Forest extent	Lamarche *et al.* [79]
Potential forest cover	Connectivity	Laestadius *et al.* [35]
Natural non-forest areas within extent of potential forest	Connectivity	ESA-CCI Land Cover dataset; ESA [68]
Infrastructure	Observed and inferred pressures	Open Street Map (selected elements) as of 2018; OpenStreetMap contributors [73]
Cropland	Observed and inferred pressures	GFSAD 2015 Cropland Extent; Gumma *et al.* [80], Massey *et al.* [81], Oliphant *et al.* [82], Phalke *et al.* [83], Teluguntla *et al.* [84], Xiong *et al.* [85] and Zhong *et al.* [86]
Cropping intensity (irrigation)	Observed and inferred pressures	GFSAD 2010 Cropland Mask; Teluguntla *et al.* [74]
Water surface	Observed and inferred pressures	JRC Global Surface Water Occurrence (all classes with >75% occurrence); Pekel *et al.* [87]

Table S2. Classes in ESA-CCI dataset excluded from our potential forest cover layer because they overlap extensively with potential forest cover mapped by Laestadius *et al.* [35] but contain significant areas of natural non forest

Legend code	Class name
60	Treecover, broadleaved, deciduous, closed to open, >15%
100	Mosaic tree and shrub (>50%]/ Herbaceous cover (<50%)
120	Shrubland
121	Evergreen shrubland
122	Deciduous shrubland
130	Grassland
140	Lichens and mosses
150	Sparse vegetation (tree, shrub, herbaceous cover) (<15%)
152	Sparse shrub (<15%)
180	Shrub or herbaceous cover, flooded, fresh/saline/brackish water
200	Bare areas
201	Consolidated bare areas
202	Unconsolidated bare areas
220	Permanent snow and ice

Table S3. Weightings used for Open Street Map (OSM) to combine into the Infrastructure data layer.

OSM Category	OSM Subcategory	Weighting applied for FPI
Aeroway	Apron / Helipad / Runway / Taxiway	8
	Hangar / Terminal	4
	Aerodrome / Heliport / Spaceport	3
Amenity / Landuse / Man-made object	Fuel station / Gasometer / Petroleum well / Pipeline / Adit / Mineshaft / Quarry / Landfill / Sanitary dump station / Wastewater plant	15
	Chimney	10
	Industrial	8
	Basin / Covered Reservoir / Pumping station / Water tower / Water well / Water works / Watermill	7
	Silo / Storage tank / Works	6
	Aerialway / Beacon / Lighthouse / Breakwater / Dyke / Embankment / Groyne / Pier / Communications tower / Mast / Observatory / Tower / Telescope	5
	Salt pond	4
	Alpine hut / Beach resort / Camp site / Cemetery / Golf course / Marina / Pitch / Village green / Wilderness hut	3
Barrier	City wall / Retaining wall / Wall	5
	Ditch / Snow fence / Snow net	3
	Hedge	2
Road	Motorway / Motorway link / Raceway	15
	Trunk / Trunk link	11
	Primary / Primary link	9
	Secondary / Secondary link	7
	Tertiary / Tertiary link	6
	Bus guideway / Service	5
	Living street / Mini roundabout / Residential / Turning circle / Unclassified / Unknown/ Elevator / Rest area	4
	Escape / Track	3
	Bridleway / Cycleway/ Footway / Path / Pedestrian / Steps	2
Military	Nuclear explosion site	30
	Danger area / Range / Trench	15
	Ammunition / Barracks / Bunker / Checkpoint	7
	Airfield / Military-owned land / Naval base / Training area	3
Power	Plant/generator - coal	20
	Plant/generator - oil	15
	Plant/generator – gas/ Plant/generator - bio / waste	10
	Plant/generator – hydro; nuclear; other / Line, Substation	7
	Plant/generator - solar / Heliostat / wind / Windmill	5
	Cable	3
Railway	Funicular / Preserved / Rail / Monorail / Subway	10
	Light rail / Miniature / Narrow gauge/ Tram	7
	Station	5
	Halt / Platform	4
	Abandoned / Disused	2
Waterway	Dam / Lock gate	20
	Canal	13
	Ditch/ Drain / Weir	3

Table S4. Points assessed to determine category boundaries for classifying the FHI into high, medium and low classes.

Category	Code	Point description	Country	Point Score
High	103	Interior of Lopé National Park	Gabon	10.000
High	106	Interior of Taï National Park	Cote d'Ivoire	10.000
High	108	Interior of Pacaya-Samiria National Reserve	Peru	10.000
High	109	Interior of Central Suriname Nature Reserve	Suriname	10.000
High	116	Interior of Liard River area	Canada	10.000
High	101	Interior of Okapi Faunal Reserve	DRC	9.997
High	104	Interior of Nyungwe National Park	Rwanda	9.992
High	111	Interior of Rio Platano Biosphere Reserve	Honduras	9.990
High	102	Interior of Odzala National Park	RoC	9.974
High	117	Interior of Wells Gray Provincial Park	Canada	9.972
High	119	Interior of Øvre Pasvik National Park	Norway	9.944
High	115	Interior of Tasmania Wilderness World Heritage Area	Australia	9.918
High	107	Interior of Marojejy National Park	Madagascar	9.910
High	112	Interior of Khao Yai National Park	Thailand	9.908
High	105	Interior of Niassa Special Reserve	Mozambiuque	9.819
High	110	Interior of Maya Biosphere Reserve	Guatemala	9.798
High	114	Interior of Batang Ai National Park	Malaysia	9.756
High	118	Interior of Quetico Provincial Park	Canada	9.750
High	113	Interior of Sundarbans National Park	Bangladesh	9.606
Medium	215	Interior of Bialowieża National Park	Poland	9.086
Medium	208	Interior of Mabira Central Forest Reserve	Uganda	9.067
Medium	211	Area of selective logging	Gabon	8.840
Medium	219	Near main tourism corridor, Mt Myohyang National Park	DPR Korea	8.762
Medium	203	Interior of Phnom Kulen Wildlife Sanctuary	Cambodia	8.710
Medium	210	Area of selective logging	Guyana	8.364
Medium	202	Interior of Dong Hua Sao National Protected Area	Lao PDR	8.078
Medium	212	Area of selective logging	DRC	7.981
Medium	206	Interior of Manga Forest Reserve	Tanzania	7.960
Medium	207	Near margin of Nyungwe National Park	Rwanda	7.938
Medium	204	South part of Nagarahole National Park	India	7.759
Medium	213	Area of selective logging	Cameroon	7.379
Medium	201	Tat Leuk, Phou Khaokhoay National Protected Area	Lao PDR	7.251
Medium	216	Interior of Loch Garten Nature Reserve	UK	7.146
Medium	209	Area of selective logging	Congo	6.734
Medium	217	Tourism area, Lamington National Park	Australia	6.729
Medium	214	Lowlands of Guanacaste National Park	Costa Rica	6.719
Medium	218	Near margin of Sepilok Forest Reserve	Malaysia	6.353
Medium	205	Interior of Similajau National Park	Malaysia	6.130
Low	305	Dong Nathat	Lao PDR	5.638
Low	317	Foothills of Mt Makiling	Philippines	5.395
Low	310	Suburban woodlot, Dobbs Ferry	USA	4.710
Low	309	Jozani Forest Reserve	Tanzania	4.680
Low	316	Foothills of Mt Canlaon	Philippines	4.597

Low	320	Forest fragment near Paramaribo	Suriname	4.566
Low	302	Central Park, New York	USA	3.575
Low	301	Bagley Wood, Oxford	UK	3.525
Low	307	Boeng Yeak Lom Protected Area	Cambodia	3.323
Low	304	Angkor Thom	Cambodia	3.122
Low	315	Forest in rural complex, Mambasa area	DRC	2.689
Low	312	Woodland in Beaumont area	USA	2.581
Low	318	Swidden near Andoung Kraloeng village	Cambodia	2.304
Low	319	Forest mosaic near Kaev Seima village	Cambodia	2.187
Low	303	Thetford Forest	UK	2.082
Low	313	Woodland in Augusta area	USA	0.686
Low	314	Woodland in Emporia area	USA	0.589
Low	311	River Park, Chicago	USA	0.566
Low	306	Houei Nhang Forest Reserve	Lao PDR	0.000
Low	308	Pugu Forest Reserve	Tanzania	0.000

16 **Table S5**. Mean Forest Landscape Integrity Index scores and areas for forest integrity categories
17 by country.
18

Country	Mean FHI	Low health (km²)	Medium health (km²)	High health (km²)	Total forest area (km²)
Afghanistan	8.85	90	1,475	977	2,542
Albania	6.77	2,426	5,256	122	7,805
Algeria	5.22	7,418	6,044	81	13,543
Andorra	4.45	170	49	0	219
Angola	8.35	105,487	284,054	315,895	705,436
Antigua and Barbuda	4.72	114	92	0	206
Argentina	7.21	98,249	189,966	72,557	360,772
Armenia	5.46	1,894	1,681	3	3,577
Australia	7.22	117,672	239,624	103,852	461,148
Austria	3.55	36,666	12,422	21	49,109
Azerbaijan	6.55	4,820	7,189	1,534	13,543
Bahamas	7.35	741	1,935	399	3,075
Bangladesh	5.45	10,013	7,251	1,947	19,211
Belarus	3.63	77,870	20,847	91	98,808
Belgium	1.36	8,803	297	0	9,099
Belize	6.15	7,004	7,957	2,744	17,705
Benin	5.86	4,724	3,698	1,769	10,191
Bhutan	8.85	1,620	16,769	10,140	28,529
Bolivia	8.47	78,745	280,532	272,007	631,284
Bosnia and Herzegovina	5.99	13,387	17,031	574	30,993
Botswana	9.13	13	187	372	572
Brazil	7.52	1,374,902	1,354,961	2,338,101	5,067,963
Brunei Darussalam	7.71	1,102	2,842	1,498	5,442
Bulgaria	6.09	18,884	26,325	847	46,057
Burundi	4.5	6,882	3,841	46	10,769
Cabo Verde	6.37	27	38	0	65
Cambodia	6.31	30,143	31,939	16,349	78,431
Cameroon	8	66,191	181,336	119,263	366,789
Canada	8.99	480,206	1,027,386	2,968,268	4,475,860
Central African Republic	9.28	30,161	139,350	379,097	548,608
Chad	6.18	5,261	6,016	1,910	13,187
Chile	7.37	56,849	41,971	93,537	192,357
China	7.14	533,800	974,431	301,051	1,809,282
Colombia	8.26	150,737	272,442	428,320	851,499
Comoros	7.69	284	1,149	82	1,515
Congo	8.89	24,512	124,215	158,184	306,911
Congo DRC	7.56	533,118	935,508	727,983	2,196,608

Costa Rica	4.65	27,164	12,838	4,164	44,167
Cote d'Ivoire	3.64	158,010	41,005	7,288	206,303
Croatia	4.92	15,732	10,522	379	26,633
Cuba	5.4	22,605	18,460	1,632	42,697
Cyprus	7.06	388	1,026	18	1,432
Czechia	1.71	32,161	1,611	0	33,772
Denmark	0.5	5,756	31	0	5,787
Dominica	1.06	531	2	0	533
Dominican Republic	4.19	19,890	9,364	518	29,772
Ecuador	7.66	48,822	77,585	73,492	199,900
Egypt	0.56	4,772	218	69	5,059
El Salvador	4.05	8,837	2,947	0	11,784
Equatorial Guinea	7.99	3,982	17,595	5,007	26,585
Estonia	3.05	24,473	4,832	52	29,358
Ethiopia	7.16	52,652	84,430	44,397	181,479
Fiji	8.35	1,753	10,802	3,594	16,148
Finland	5.08	144,310	83,572	9,294	237,176
France	4.52	161,987	49,496	74,121	285,604
Gabon	9.07	11,780	118,348	120,852	250,979
Gambia	4.56	181	85	0	266
Georgia	7.79	6,982	17,803	9,784	34,570
Germany	2.28	122,168	11,307	0	133,475
Ghana	4.53	57,519	28,901	2,160	88,580
Greece	6.6	14,548	27,833	1,078	43,459
Grenada	4.22	221	86	0	308
Guatemala	3.85	58,572	18,764	5,592	82,928
Guinea	4.9	81,702	54,877	2,895	139,475
Guinea-Bissau	5.7	9,274	8,702	855	18,831
Guyana	9.58	4,162	40,817	147,413	192,391
Haiti	4.01	7,116	2,831	12	9,959
Honduras	4.48	57,899	23,802	3,692	85,392
Hungary	2.25	18,729	2,047	0	20,776
India	7.09	117,992	254,792	54,428	427,211
Indonesia	6.6	535,370	509,018	431,973	1,476,361
Iran	7.67	3,361	12,930	2,162	18,453
Iraq	3.59	104	9	0	113
Ireland	0.92	5,283	96	0	5,378
Israel	4.14	170	85	0	255
Italy	3.65	79,403	26,858	25	106,286
Jamaica	5.01	5,362	3,249	158	8,770
Japan	5.8	135,783	133,480	16,005	285,268
Jordan	2.79	12	0	0	12
Kazakhstan	8.23	6,068	18,926	15,294	40,288

Country					
Kenya	4.2	28,427	13,558	4,702	46,686
Kosovo	5.19	2,628	1,775	47	4,450
Kyrgyzstan	8.86	329	2,819	2,761	5,909
Laos	5.59	92,986	80,564	19,252	192,801
Latvia	2.09	38,164	2,137	0	40,301
Lebanon	3.76	541	115	0	656
Lesotho	7.4	1	4	0	5
Liberia	4.79	51,975	31,162	11,025	94,163
Libya	4.85	15	2	0	17
Liechtenstein	4.5	59	42	0	101
Lithuania	1.62	24,554	930	0	25,484
Luxembourg	1.12	1,170	0	0	1,170
Macedonia	7.42	2,034	7,090	459	9,583
Madagascar	4.63	120,340	66,584	11,922	198,846
Malawi	5.74	12,514	12,167	2,396	27,078
Malaysia	5.01	130,825	91,957	21,499	244,281
Mali	7.16	451	996	140	1,586
Mauritius	5.46	567	478	0	1,045
Mexico	6.82	193,908	280,445	121,842	596,195
Micronesia	7.55	8	35	0	43
Moldova	2.2	3,113	202	0	3,315
Mongolia	9.36	520	11,915	27,407	39,841
Montenegro	6.41	2,949	4,778	82	7,809
Morocco	6.74	2,260	4,076	451	6,787
Mozambique	6.93	150,665	189,362	115,379	455,406
Myanmar	7.18	129,745	220,188	96,924	446,857
Namibia	8.43	5	13	17	36
Nepal	7.23	13,785	41,992	3,760	59,538
Netherlands	0.6	5,250	72	0	5,322
New Zealand	7.12	34,503	44,155	35,334	113,992
Nicaragua	3.63	65,356	17,646	4,858	87,860
Nigeria	6.2	64,621	65,355	24,307	154,283
North Korea	8.02	8,374	40,156	8,410	56,939
Norway	6.98	39,343	67,383	16,627	123,352
Pakistan	7.42	2,090	7,859	1,139	11,088
Palau	8.09	45	333	9	387
Panama	6.37	25,420	21,310	14,605	61,336
Papua New Guinea	8.84	37,294	183,415	216,355	437,064
Paraguay	6.39	78,538	102,626	29,877	211,041
Peru	8.86	85,793	190,547	509,720	786,061
Philippines	5.91	91,820	100,831	8,393	201,044
Poland	2.24	101,886	7,103	0	108,989
Portugal	8.82	25,966	553	0	26,519

Romania	5.95	38,395	48,394	607	87,395
Russian Federation	9.02	739,484	2,245,281	5,137,079	8,121,843
Rwanda	3.85	5,665	2,170	619	8,454
Saint Kitts and Nevis	4.55	95	50	0	145
Saint Lucia	6.17	235	316	0	551
Saint Vincent and the Grenadines	6.95	91	221	0	312
San Marino	0.01	7	0	0	7
Sao Tome and Principe	6.64	31	140	0	171
Senegal	7.11	847	2,456	162	3,465
Serbia	5.29	17,513	14,112	516	32,141
Seychelles	10	0	0	68	68
Sierra Leone	2.76	52,512	11,858	640	65,010
Singapore	1.11	170	2	0	172
Slovakia	4.34	17,615	8,165	0	25,781
Slovenia	3.78	11,065	3,791	0	14,856
Solomon Islands	7.19	6,871	15,310	3,149	25,329
Somalia	7.16	347	1,384	46	1,777
South Africa	4.94	45,489	34,968	3,196	83,653
South Korea	6.02	25,060	32,009	888	57,956
South Sudan	9.45	5,083	59,389	146,218	210,691
Spain	4.23	82,770	46,013	133	128,916
Sri Lanka	5.83	20,541	22,390	1,613	44,544
Sudan	9.8	1	72	495	569
Suriname	9.39	6,796	25,031	107,954	139,781
Swaziland	4.21	5,054	2,501	14	7,569
Sweden	5.35	174,415	109,779	23,494	307,687
Switzerland	3.53	13,636	4,412	10	18,058
Syria	3.64	841	282	0	1,123
Taiwan	6.38	8,786	14,547	1,453	24,786
Tajikistan	8.65	34	137	130	301
Tanzania	7.13	123,997	159,712	122,812	406,521
Thailand	6	86,276	89,326	33,612	209,214
Timor-Leste	7.11	1,783	7,008	47	8,838
Togo	5.88	5,064	4,522	1,076	10,662
Trinidad and Tobago	6.62	1,478	2,176	418	4,072
Tunisia	5.14	1,354	987	0	2,340
Turkey	6.39	43,043	68,243	3,516	114,801
Turkmenistan	6.31	5	33	0	37
Uganda	4.36	77,303	36,381	7,507	121,190
Ukraine	3.3	89,540	20,183	176	109,900
United Kingdom	1.65	29,149	2,917	35	32,101
United States	6.65	1,328,079	1,144,693	658,645	3,131,417

Uruguay	3.61	11,793	3,998	0	15,791
Uzbekistan	6.77	214	227	199	640
Vanuatu	8.82	734	5,322	4,448	10,504
Venezuela	8.78	64,650	170,792	351,112	586,554
Vietnam	5.35	82,551	75,353	9,588	167,492
Zambia	7.5	96,969	164,376	110,822	372,167
Zimbabwe	6.31	9,450	14,417	1,644	25,511

Table S6. Mean Forest Landscape Integrity Index scores for provinces of Democratic Republic of

Congo (DRC), Indonesia and Canada.

DRC		Indonesia		Canada	
Province	**Mean FHI**	**Province**	**Mean FHI**	**Province**	**Mean FHI**
Lualaba	8.57	Papua	9.34	Northwest Territories	9.90
Tshuapa	8.55	West Papua	9.00	Yukon	9.86
Tshopo	8.39	Kalimantan Utara	8.52	Newfoundland and Labrador	9.66
Bas-Uélé	8.38	Maluku	8.03	Nunavut	9.65
Équateur	8.37	Maluku Utara	7.41	Manitoba	9.58
Haut-Lomami	8.29	Nusa Tenggara Barat	6.86	Saskatchewan	9.40
Tanganyika	8.24	Aceh	6.83	Ontario	8.94
Nord-Ubangi	8.19	Nusa Tenggara Timur	6.80	Québec	8.80
Haut-Katanga	8.05	Gorontalo	6.60	Alberta	8.46
Kwango	7.83	Sulawesi Utara	6.58	British Columbia	8.22
Maï-Ndombe	7.58	Sulawesi Tengah	6.54	Nova Scotia	6.07
Haut-Uélé	7.46	Kalimantan Timur	6.42	New Brunswick	5.15
Maniema	7.44	Sulawesi Barat	6.31	Prince Edward Island	2.74
Sankuru	7.34	Sumatera Barat	6.20		
Lomami	7.20	Sulawesi Tenggara	5.99		
Kasaï	7.11	Kalimantan Tengah	5.84		
Ituri	6.70	Sulawesi Selatan	5.63		
Mongala	6.23	Banten	4.97		

Nord-Kivu	6.22	**Bengkulu**	4.94		
Sud-Kivu	6.20	**Sumatera Utara**	4.89		
Kasaï-Central	5.95	**Kalimantan Barat**	4.87		
Sud-Ubangi	5.93	**Kepulauan Riau**	4.86		
Kwilu	5.65	**Jawa Barat**	4.76		
Kinshasa	4.75	**Lampung**	4.73		
Kasaï-Oriental	4.13	**Jawa Tengah**	4.59		
Kongo-Central	3.95	**Bali**	4.43		
		Jawa Timur	4.40		
		Jambi	4.01		
		Riau	3.92		
		Kalimantan Selatan	3.24		
		Sumatera Selatan	2.86		
		Yogyakarta	2.83		

24
25

Figure S1. A map overlaying the Open Street Maps data (blue) and provincial government data

(green) for roads and other linear infrastructure associated with resource access.

Figure S2. A global map of Forest Landscape Integrity for 2019. Highlighted regions show **A.** A

remote road in Russia, **B.** Clearcut logging in Canada, **C.** Selective logging in Borneo, **D.**

Swidden agriculture in Madagascar, **E.** Forest fragmentation in Western Australia, **F.** Remote

settlements in the Brazilian Amazon.

References

Hansen, M. C. *et al.* High-Resolution Global Maps of 21st-Century Forest Cover Change. *Science* **342**, 850-853, doi:10.1126/science.1244693 (2013).
Gibson, L. *et al.* Primary forests are irreplaceable for sustaining tropical biodiversity. *Nature* **478**, 378-381 (2011).
Curtis, P. G., Slay, C. M., Harris, N. L., Tyukavina, A. & Hansen, M. C. Classifying drivers of global forest loss. *Science* **361**, 1108-1111, doi:10.1126/science.aau3445 (2018).
Laestadius, L. *et al.* Opportunities for forest landscape restoration. *Unasylva* **62**, 238 (2011).
ESA. Land Cover CCI Product User Guide Version 2. Tech. Rep., (2017).
Theobald, D. M. A general model to quantify ecological integrity for landscape assessments and US application. *Landscape Ecol.* **28**, 1859-1874 (2013).
Sanderson, E. W. *et al.* The human footprint and the last of the wild. *Bioscience* **52**, 891-904 (2002).
Venter, O. *et al.* Sixteen years of change in the global terrestrial human footprint and implications for biodiversity conservation. *Nature Communications* **7**, 12558 (2016).
Kennedy, C. M., Oakleaf, J. R., Theobald, D. M., Baruch-Mordo, S. & Kiesecker, J. Managing the middle: A shift in conservation priorities based on the global human modification gradient. *Global Change Biol.* **25**, 811-826 (2019).
Thompson, I. D. *et al.* An Operational Framework for Defining and Monitoring Forest Degradation. *Ecology and Society* **18** (2013).
Contributors.   (ed OpenStreetMap) (2015).
Teluguntla, P. *et al.*   (ed NASA EOSDIS Land Processes DAAC) (2016).
Haddad, N. M. *et al.* Habitat fragmentation and its lasting impact on Earth's ecosystems. *Science Advances* **1**, e1500052, doi:10.1126/sciadv.1500052 (2015).
Laurance, W. F. *et al.* Ecosystem decay of Amazonian forest fragments: a 22-year investigation. *Conserv. Biol.* **16**, 605-618 (2002).
Zimmerman, B. L. & Kormos, C. F. Prospects for Sustainable Logging in Tropical Forests. *Bioscience* **62**, 479-487, doi:10.1525/bio.2012.62.5.9 (2012).
Cochrane, M. A. & Laurance, W. F. Fire as a large-scale edge effect in Amazonian forests. *Journal of Tropical Ecology* **18**, 311-325, doi:10.1017/S0266467402002237 (2002).
Peres, C. A., Emilio, T., Schietti, J., Desmoulière, S. J. M. & Levi, T. Dispersal limitation induces long-term biomass collapse in overhunted Amazonian forests. *Proceedings of the National Academy of Sciences* **113**, 892-897, doi:10.1073/pnas.1516525113 (2016).
Chaplin-Kramer, R. *et al.* Degradation in carbon stocks near tropical forest edges. *Nature Communications* **6**, 10158, doi:10.1038/ncomms10158 https://www.nature.com/articles/ncomms10158#supplementary-information (2015).
Tulloch, A. I., Gordon, A., Runge, C. A. & Rhodes, J. R. Integrating spatially realistic infrastructure impacts into conservation planning to inform strategic environmental assessment. *Conservation Letters* **12**, e12648 (2019).
Benítez-López, A., Alkemade, R. & Verweij, P. A. The impacts of roads and other infrastructure on mammal and bird populations: a meta-analysis. *Biol. Conserv.* **143**, 1307-1316 (2010).

Shapiro, A. C., Aguilar-Amuchastegui, N., Hostert, P. & Bastin, J.-F. Using fragmentation to assess degradation of forest edges in Democratic Republic of Congo. *Carbon Balance and Management* **11**, 11, doi:10.1186/s13021-016-0054-9 (2016).

Broadbent, E. N. *et al.* Forest fragmentation and edge effects from deforestation and selective logging in the Brazilian Amazon. *Biol. Conserv.* **141**, 1745-1757 (2008).

Maisels, F. *et al.* Devastating Decline of Forest Elephants in Central Africa. *PLOS ONE* **8**, e59469, doi:10.1371/journal.pone.0059469 (2013).

Benítez-López, A., Santini, L., Schipper, A. M., Busana, M. & Huijbregts, M. A. J. Intact but empty forests? Patterns of hunting-induced mammal defaunation in the tropics. *PLOS Biology* **17**, e3000247, doi:10.1371/journal.pbio.3000247 (2019).

Barrington-Leigh, C. & Millard-Ball, A. The world's user-generated road map is more than 80% complete. *PLOS ONE* **12**, e0180698, doi:10.1371/journal.pone.0180698 (2017).

Hughes, A. C. Have Indo-Malaysian forests reached the end of the road? *Biol. Conserv.* **223**, 129-137, doi:https://doi.org/10.1016/j.biocon.2018.04.029 (2018).

Lamarche, C. *et al.* Compilation and validation of SAR and optical data products for a complete and global map of inland/ocean water tailored to the climate modeling community. *Remote Sensing* **9**, 36 (2017).

Gumma, M. *et al.* NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Global Food Security-support Analysis Data (GFSAD) Cropland Extent 2015 South Asia, Afghanistan, Iran 30 m V001. (2017).

Massey, R. *et al.*   (ed NASA EOSDIS Land Processes DAAC) (2017).

Oliphant, A. *et al.* NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Global Food Security-support Analysis Data (GFSAD) Cropland Extent 2015 Southeast Asia 30 m V001. (2017).

Phalke, A. *et al.*   (ed NASA EOSDIS Land Processes DAAC) (2017).

Teluguntla, P. *et al.* NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Global Food Security-support Analysis Data (GFSAD) Cropland Extent 2015 Australia, New Zealand, China, Mongolia 30 m V001. (2017).

Xiong, J. *et al.* NASA Making Earth System Data Records for Use in Research Environments (MEaSUREs) Global Food Security-support Analysis Data (GFSAD) Cropland Extent 2015 Africa 30 m V001. (2017).

Zhong, Y. *et al.*   (ed NASA EOSDIS Land Processes DAAC) (2017).

Pekel, J.-F., Cottam, A., Gorelick, N. & Belward, A. S. High-resolution mapping of global surface water and its long-term changes. *Nature* **540**, 418-422 (2016).