Bayesian Genome-wide TWAS method to leverage both cis- and trans- eQTL information through summary statistics

Justin M. Luningham[1,5], Junyu Chen[5], Shizhen Tang[2,5], Philip L. De Jager[3], David A. Bennett[4], Aron S. Buchman[4], Jingjing Yang[5*]

1. Department of Population Health Sciences, Georgia State University School of Public Health, Atlanta, GA, 30303, USA
2. Department of Biostatistics and Bioinformatics, Emory University School of Public Health, Atlanta, GA, 30322, USA
3. Center for Translational and Computational Neuroimmunology, Department of Neurology and Taub Institute for Research on Alzheimer's Disease and the Aging Brain, Columbia University Irving Medical Center, New York, NY10032, USA
4. Rush Alzheimer's Disease Center, Rush University Medical Center, Chicago, IL, 60612, USA
5. Center for Computational and Quantitative Genetics, Department of Human Genetics, Emory University School of Medicine, Atlanta, GA, 30322, USA

* Correspondence author: jingjing.yang@emory.edu

Keywords: Transcriptome-wide association study; TWAS; Bayesian variable selection model; Cis- eQTL; Trans- eQTL; Summary statistics; Alzheimer's dementia

Abstract

Transcriptome-wide association studies (TWAS) have been widely used to integrate gene expression and genetic data for studying complex traits. Due to the computational burden, existing TWAS methods do not assess distant trans- expression quantitative trait loci (eQTL) that are known to explain important expression variation for most genes. We propose a Bayesian Genome-wide TWAS (BGW-TWAS) method which leverages both cis- and trans-eQTL information for TWAS. Our BGW-TWAS method is based on Bayesian variable selection regression, which not only accounts for cis- and trans- eQTL of the target gene but also enables efficient computation by using summary statistics from standard eQTL analyses. Our simulation studies illustrated that BGW-TWAS achieved higher power compared to existing TWAS methods that do not assess trans-eQTL information. We further applied BWG-TWAS to individual-level GWAS data (N=~3.3K), which identified significant associations between the genetically regulated gene expression (GReX) of gene *ZC3H12B* and Alzheimer's dementia (AD) (p-value= 5.42×10^{-13}), neurofibrillary tangle density (p-value= 1.89×10^{-6}), and global measure of AD pathology (p-value=9.59×10^{-7}). These associations for gene *ZC3H12B* were completely driven by trans-eQTL. Additionally, the GReX of gene *KCTD12* was found to be significantly associated with β-amyloid (p-value= 3.44×10^{-8}) which was driven by both cis- and trans- eQTL. Four of the top driven trans-eQTL of *ZC3H12B* are located within gene *APOC1*, a known major risk gene of AD and blood lipids. Additionally, by applying BGW-TWAS with summary-level GWAS data of AD (N=~54K), we identified 13 significant genes including known GWAS risk genes *HLA-DRB1* and *APOC1*, as well as *ZC3H12B.*

Introduction

Although genome-wide association studies (GWAS) have identified thousands of variants associated with complex traits over the past decades[1-5], most of these associations are located within noncoding regions and the underlying biological mechanisms by which these

variants impact a phenotype are unknown[6; 7]. Recent studies have shown that GWAS

associations were enriched for regulatory elements such as expression quantitative trait loci

(eQTL)[8-10], suggesting that integrating transcriptomic and genetic data could help identify key

molecular mechanisms underlying complex traits.

One such integrative method is transcriptome-wide association study (TWAS)[11-13], which

takes advantage of a reference panel with profiled transcriptomic and genetic data from the

same individuals. TWAS first utilizes such reference data to fit an imputation regression model

for the expression quantitative trait of a target gene with nearby genotypes (e.g., cis-SNPs

within 1MB region of transcription starting site) as predictors, and then examines the gene-

based association between the imputed genetically regulated gene expression (GReX) and the

phenotype of interest. With fitted gene expression imputation models from reference data,

TWAS can be conducted with test samples that have either individual-level or summary-level

GWAS data[12-14]. The SNPs with non-zero effect sizes on reference transcriptome in the fitted

imputation models are referred to as broad sense "eQTL" in TWAS. Examples of publicly

available reference data include the Genotype-Tissue Expression (GTEx) project with

transcriptomic data for 54 human tissues[8], Genetic European Variation in Health and Disease

(GEUVADIS) for lymphoblastoid cell lines[15], and North American Brain Expression Consortium

(NABEC) for cortex tissues[16].

Essentially, TWAS is equivalent to a burden type gene-based test taking "cis-eQTL

effect sizes" that are non-zero coefficients of cis-SNPs from the fitted GReX imputation model

as their corresponding burden weights[11-13]. By weighting genetic variants using cis-eQTL effect

sizes, TWAS assumes the effects of risk genes on the phenotype of interest are potentially

mediated through their transcriptome variations. Recent studies of a wide range of complex

traits such as schizophrenia, breast cancer, and Alzheimer's dementia (AD)[17-21] using TWAS

have identified additional risk genes besides known GWAS risk loci, demonstrating that

additional significant associations can be identified by TWAS.

However, existing TWAS methods only use genetic data of cis-SNPs of the target gene

as predictors to fit the GReX imputation model[11-13]. As shown by recent studies, trans-SNPs

(e.g., outside of the 1MB region) of the target gene not only explain a significant amount of

variation for most expression quantitative traits, but also often contain significant trans-eQTL

that are likely to inform molecular mechanisms[22; 23]. Thus, using both cis- and trans- SNPs is

likely to increase the imputation accuracy of GReX and the power of TWAS. Nonetheless, the

enormous computational cost required to fit ~20K GReX imputation models for genome-wide

genes and genotypes per tissue type makes the routine use of existing TWAS methods

impractical.

We propose a Bayesian Genome-Wide TWAS (BGW-TWAS) method that accounts for

both cis- and trans- SNPs based on a Bayesian variable selection regression (BVSR) model[24]

for imputing GReX. Our BGW-TWAS method circumvents the current computational burden

impeding TWAS by enabling efficient computation via the scalable EM-MCMC algorithm[25] and

the summary statistics of standard eQTL analyses based on single variant tests. First, we

demonstrate the feasibility of this Bayesian approach by simulation studies with varying

proportions of true causal cis- and trans- eQTL for expression quantitative traits. We compared

BGW-TWAS with several existing TWAS methods including PrediXcan[11] and TIGAR[13] that

assess only cis-SNPs. Then we applied BGW-TWAS to clinical and postmortem data from older

adults with individual-level GWAS data (N=~3.3K)[26], to study several clinical and pathological

AD related phenotypes including clinical diagnosis of AD, neurofibrillary tangle density, β-

amyloid load, and a global summary measure of AD pathology. Further, we compared BGW-

TWAS with alternative methods by using GWAS summary statistics for AD available from the

International Genomics of Alzheimer's Project (IGAP)[27] (N=~54K).

Our simulation studies revealed that BGW-TWAS achieved higher TWAS power by

considering both cis- and trans- SNPs when trans-eQTL accounted for a non-negligible

proportion of transcriptome variance. Our studies of human AD GWAS datasets identified

several risk genes associated with AD phenotypes that were driven by trans-eQTL and thus not

identified by alternative methods. The software for implementing BGW-TWAS is available freely

on Github.

Method

TWAS Procedure

The first step in TWAS is to train an imputation model for profiled gene expression levels

using genotype data as predictors on a per-gene basis[11-13]. The general imputation model based

on linear regression is given by

$$\mathcal{E}_g = Xw + \epsilon, \qquad\qquad \text{(Equation 1)}$$

where $\mathcal{E}_g$ denotes the expression quantitative trait of the target gene, centered and adjusted for

non-genetic covariates; X denotes centered genotype data; w denotes the corresponding

"eQTL" effect sizes for the target gene; and ϵ is an error term following a $N(0, \sigma_\epsilon^2 I)$ distribution.

The intercept term is dropped for centering both response ($\mathcal{E}_g$) and explanatory (X) variables.

With $\hat{w}$ estimated from the training data (i.e., reference data) that have both transcriptomic and

genetic data profiled for the same subjects, TWAS will test the association between the

phenotype of interest and the imputed GReX obtained from individual-level genotype data $\tilde{X}$ of

the test cohort as follows

$$\widehat{GReX} = \tilde{X}\hat{w}.$$

Bayesian Variable Selection Regression

Existing TWAS methods only consider SNPs within 1MB of the flanking 5' and 3' ends

(cis-SNPs) in the gene expression imputation model (Equation 1)[11-13]. In order to leverage

additional information provided by trans-eQTL that are located outside the 1MB flanking region

of the target gene, we utilize the Bayesian Variable Selection Regression (BVSR)[24] model to

account for both cis- and trans- SNPs as follows:

$$\mathcal{E}_g = X_{cis}w_{cis} + X_{trans}w_{trans} + \epsilon, \ \ \epsilon_i \sim N(0, \sigma_\epsilon^2).$$ (Equation 2)

128   The BVSR model assumes a spike-and-slab prior distribution for w_i. That is, the prior on w_i is a

129   mixture distribution of a normal distribution with zero mean and a point-mass density function at

130   0. In order to model potentially different distributions of the effect sizes for cis- and trans- SNPs,

131   we assume the following respective priors,

$$w_{cis,i} \sim \pi_{cis} N\left(0, \sigma_{cis}^2 \sigma_\epsilon^2\right) + (1 - \pi_{cis})\delta_0\left(w_{cis,i}\right);$$ (Equation 3)

$$w_{trans,i} \sim \pi_{trans} N(0, \sigma_{trans}^2 \sigma_\epsilon^2) + (1 - \pi_{trans})\delta_0\left(w_{cis,i}\right);$$

132   where $(\pi_{cis}, \ \pi_{trans})$ denote the respective probability that the coefficient is normally distributed,

133   and $\delta_0(w_i)$ is the point mass density function that takes value 0 when $w_i \neq 0$ and 1 when

134    $w_i = 0$. Further, the following conjugate hyper prior distributions are respectively assumed for

135   the cis- and trans- specific parameters,

$$\pi_{cis} \sim Beta(a_{cis}, b_{cis}); \ \sigma_{cis}^2 \sim IG(k_1, k_2);$$ (Equation 4)

$$\pi_{trans} \sim Beta(a_{trans}, b_{trans}); \ \sigma_{trans}^2 \sim IG(k_3, k_4);$$

where *IG* indicates the Inverse Gamma distribution and hyper parameters

$(a_{cis}, b_{cis}, a_{trans}, b_{trans}, k_1, k_2, k_3, k_4)$ will be chosen to enable non-informative hyper prior

distributions (see Supplemental Methods for model details).

To facilitate computation, a latent indicator γ_i is assumed such that $w_i = 0$ if $\gamma_i = 0$, and

w_i follows a normal distribution if $\gamma_i = 1$. Then the expected value of this indicator, $\mathrm{E}[\gamma_i]$,

represents the posterior probability (PP_i) for each individual SNP to have a non-zero effect size

(i.e., to be an eQTL of the target gene)[24]. Moreover, we propose a Bayesian approach to

estimate GReX for test samples that can account for the uncertainty for each SNP to be an

eQTL:

$$\widehat{GReX} = \sum_{i=1}^{p} \tilde{X}_l(\widehat{PP}_l \, \widehat{w}_i) \qquad \text{(Equation 5)}$$

where $\tilde{X}_l$ represents the genotype data of variant i for test samples and $(\widehat{w}_i, \widehat{PP}_l)$ denote the

estimate of effect size and posterior probability (PP) of having a non-zero effect size from the

BVSR model (Equations 2-4) (see Supplemental Methods for detailed Bayesian inference

procedure). This Bayesian GReX estimate can then be used to conduct TWAS with individual-

level GWAS data by testing the association between the imputed GReX and the phenotype of

interest.

<u>BGW-TWAS with Summary-level GWAS Data</u>

With summary-level GWAS data that were generated by single variant tests, we

employed the S-PrediXcan[14] approach to obtain a burden TWAS Z-score test statistic, including

not only cis- but also trans- eQTL in the test. Let $\widehat{\beta}_l$ denote the SNP effect size of SNP l from

GWAS, $SE(\widehat{\beta}_l)$ denote the standard error of $\widehat{\beta}_l$, Z_l denote the Z-score statistic value by single

variant test, $\widehat{\sigma}_l$ denote the estimated standard deviation of the genotype data of SNP l from

reference panel, and $\widehat{\sigma}_g$ denote the estimated standard deviation of the imputed expression of

gene g from reference panel. The burden TWAS Z-score test statistic for gene g is given by

$$Z_g = \sum_{l \in Model_g} \widehat{w}_{lg} \frac{\widehat{\sigma}_l}{\widehat{\sigma}_g} \frac{\widehat{\beta}_l}{SE(\widehat{\beta}_l)} = \sum_{l \in Model_g} \widehat{w}_{lg} \frac{\widehat{\sigma}_l}{\widehat{\sigma}_g} Z_l = \sum_{l \in Model_g} \frac{(\widehat{w}_{lg} \widehat{\sigma}_l) Z_l}{\sqrt{\widehat{w}' \boldsymbol{V} \widehat{w}}},$$

$$\widehat{\sigma_l^2} = Var(x_l), \qquad \widehat{\sigma_g^2} = \widehat{w}' \boldsymbol{V} \widehat{w}, \qquad \boldsymbol{V} = Cov(\boldsymbol{X})$$

where $\widehat{w}_{lg} = \widehat{PP}_l \widehat{w}_i$ is the product of posterior probability for SNP l to have non-zero eQTL effect

size from the BVSR model (Equation 2). Here, $\boldsymbol{X}$ denotes the genotype matrix of analyzed

SNPs from reference panels of the same ethnicity and V denotes the corresponding genotype

covariance matrix.

<u>Efficient Computation Techniques</u>

In theory, the estimates of eQTL effect sizes and corresponding posterior probabilities

$(\widehat{w}_i, \widehat{PP}_i)$ can be obtained by using a standard Markov Chain Monte Carlo (MCMC)[28] algorithm.

However, in practice, the computation burden for modeling genome-wide genotype data is

nearly impossible because of enormous required memory capacity and slow convergence rate

for MCMC. To circumvent these practical limitations, we employ several techniques to enable

computational efficiency such that BGW-TWAS method can be deployed to leverage both cis-

and trans- eQTL information in practice. In particular, we adapt a previously developed scalable

expectation-maximization Markov Chain Monte Carlo (EM-MCMC) algorithm[25]. Unlike the

original EM-MCMC algorithm requiring individual-level GWAS data, we can reduce up to 90% of

the computation time by adapting the EM-MCMC algorithm to utilize only summary statistics,

including the pre-calculated linkage disequilibrium (LD) coefficients and score statistics from

standard eQTL analyses by single variant tests. Additionally, we prune genome-wide genotypes

into a subset of genome regions that are approximately independent and contain either at least

one cis-SNP or one trans-SNP with p-value $< 1 \times 10^{-5}$ by standard eQTL analyses.

<u>Simulation Study Design</u>

We conducted simulation studies to validate the performance of our proposed BGW-

TWAS method through comparing with the alternative existing methods, e.g., PrediXcan,

TIGAR, as well as BVSR using only cis-eQTL. To mimic real studies, we used real genotype

data from the ROS/MAP study to simulate gene expression and phenotype data. We took 499

samples as our training data and 1,209 samples as our test data. GReX imputation models

were fitted using the training data where "eQTL" effect sizes and the corresponding posterior

probabilities were estimated. Given these fitted GReX imputation models, GReX data were

imputed for follow-up TWAS with the test data.

We arbitrarily selected five approximately independent genome blocks, including one

"cis-" and four "trans-" genotype blocks (variants were filtered with minor allele frequency (MAF)

$> 5\%$ and Hardy-Weinberg p-value $> 10^{-5}$). With genotype matrix X_g of the randomly selected

causal eQTL, we generated effect-sizes w_i to target a selected gene expression heritability h_e^2

and that all causal eQTL explain equal expression heritability. Gene expression levels were

generated by $\mathcal{E}_g = X_g w + \epsilon_e$, with $\epsilon_e \sim N(0, (1 - h_e^2))$. Then we simulated phenotypes by

$Y = \beta \mathcal{E}_g + \epsilon_p$, where β was selected with respect to a selected phenotype heritability h_p^2 and

$\epsilon_p \sim N(0, (1 - h_p^2))$.

To mimic the complex genomic architecture of gene expression in practice, we

considered two scenarios, one with 5 true causal eQTL representing the scenario with relatively

large effect sizes and the other one with 22 true causal eQTL representing the scenario with

relatively small effect sizes. For the scenario with 5 true causal eQTL, we considered three sub-

scenarios with respect to how these true causal eQTL distributed over considered genome

blocks: i) all causal eQTL are from the cis-block; ii) two causal eQTL are from the cis-block

explaining 70% of the specified h_e^2 while the other three causal eQTL are from the trans-blocks

explaining the other 30% of h_e^2; iii) all causal QTL are from the trans-block. Similarly, For the

scenario with 22 true causal eQTL, we considered three scenarios where 30%, 50%, and 70%

of the causal eQTL were from *cis-* genome blocks. We also varied the total expression trait

heritability and phenotype heritability in both scenarios, i.e., $\left(h_e^2, h_p^2\right) = \left((5\%, 90\%), (10\%, 45\%),\right.$

$(20\%, 20\%), (50\%, 6\%)\big)$ for   the   scenario   with   5   true   causal   eQTL   and

$\left(h_e^2, h_p^2\right) = \left((5\%, 99\%), (10\%, 80\%), (20\%, 35\%), (50\%, 8\%)\right)$ for the scenario with 22 true causal

eQTL. Here, different levels of phenotype heritability were arbitrarily selected to achieve similar

levels of TWAS power across all scenarios.

In each simulation, with training data, we first fitted GReX imputation models by BVSR

(BGW-TWAS) with both *cis-* and *trans-* genome blocks, as well as by Elastic-Net (PrediXcan)

and nonparametric Bayesian Dirichlet process regression (TIGAR) with only *cis-* genome block.

Then we conducted TWAS with imputed *GReX* by respective method. We also compared BGW-

TWAS with using only cis-eQTL estimates from the same BVSR model. The performance was

compared in terms of R^2 of the imputed GReX and TWAS power in test samples. Test R^2 was

calculated as the squared correlation between imputed GReX and simulated gene expression

values of the test samples. TWAS power was calculated as the proportion of 1,000 repeated

simulations of each scenario with *p*-value $< 2.5 \times 10^{-6}$ (genome-wide significance threshold for

gene-based association studies).

ROS/MAP and Mayo Clinic GWAS data of AD

Following simulation studies, we applied BGW-TWAS method to individual-level

genomic and AD related phenotype data from older adults available from several studies. We

used transcriptomic data, GWAS data, clinical diagnosis of AD and postmortem indices of AD

pathology from the Religious Orders Study (ROS) and Rush Memory and Aging Project (MAP)

[29-31] and GWAS data from the Mayo Clinic Alzheimer's Disease Genetics Studies (MCADGS)[32-

34]. All participants from ROS/MAP sign an informed consent, an Anatomic Gift Act, and a

consent for their data to be deposited in the Rush Alzheimer's Disease Center (RADC)

repository. ROS/MAP studies were approved by the Institutional Review Board of Rush

University Medical Center, Chicago, IL. MCADGS contains samples from two clinical AD Case-

Control series (Mayo Clinic Jacksonville and Mayo Clinic Rochester) as well as a

neuropathological series of autopsy-confirmed subjects from the Mayo Clinic Brain Bank.

Microarray genotype data generated for 2,093 European-decent subjects from

ROS/MAP[35] and 2,099 European-decent subjects from MCADGS were further imputed to the

1000 Genome Project Phase 3[36] in our analysis[37].

Post-mortem brain samples from the dorsal lateral prefrontal cortex from ~30% of these

ROS/MAP participants with assayed genotype data were profiled for transcriptomic data by

next-generation RNA-seqencing[38]. These data were used as reference data to train GReX

prediction models in this study. We conducted TWAS for both clinical and pathological AD

phenotypes. The clinical diagnosis of late-onsite Alzheimer's dementia was available for both

ROS/MAP and MCADGS. Postmortem pathology indices of AD were only available for

ROS/MAP and included PHFtau tangle density, β-amyloid load, and a global measure of AD

pathology based on measures of neuritic and diffuse plaques and neurofibrillary tangles[29-31].

Additional details about the ongoing ROS/MAP cohort studies and how postmortem indices of

tangles and β-amyloid load were quantified are included in prior publications[29-31] and

summarized in the Supplemental Text.

Results

<u>Simulation Results</u>

For the scenario with 5 true causal eQTL and various expression heritability, our

simulation studies showed that BGW-TWAS obtained the highest test R^2 for GReX and TWAS

power than PrediXcan and TIGAR when any portion of the true causal eQTL are distributed

over trans- genome blocks (Figure 1 (A, B)). This is because BGW-TWAS leverages both cis-

and trans- eQTL information while the alternative methods fail to account for trans- eQTL.

Especially, when all true causal eQTL are from trans- genome regions, the alternative methods

barely have any power to identify the TWAS association with nearly zero test R^2. As expected,

BGW-TWAS and PrediXcan performed comparably when all causal eQTL were from the cis-

genome block, while TIGAR performed slightly worse with sparse true causal eQTL (Figure 1A).

For the scenario with 22 mixed cis- and trans- eQTL, the performance comparison

became more complicated with respect to various true expression heritability levels (Figure 1

(C, D)). Particularly, when $h_e^2 = 0.05$, all methods had difficulties accurately estimating eQTL

effect sizes and resulted in nearly zero test R^2. As expression heritability increased, the

advantage of modeling both cis- and trans- genotype data by BGW-TWAS arisen and led to

higher test R^2 and TWAS power. When $h_e^2 = (0.1, 0.2)$ and 70% of the true causal eQTL were

cis-, BGW-TWAS was less effective than PrediXcan and TIGAR while TIGAR achieved the best

performance. This is likely due to the fact that the nonparametric Bayesian Dirichlet process

regression model used by TIGAR is preferred when true causal eQTL manifest relatively small

effect sizes, which is consistent with previous findings[13].

In contrast, when true causal eQTL signals have relatively large effect sizes and are

distributed outside the cis-region of the target gene, BGW-TWAS method is preferred due to the

improved accuracy for GReX prediction by leveraging trans- SNP data. By comparing with using

only BVSR estimates of cis-eQTL, we showed that a significant proportion of transcriptome

variation due to trans-eQTL was missed and the follow-up TWAS was underpowered.

<u>TWAS of AD Related Phenotypes with Individual-level GWAS Data</u>

Next, we applied BGW-TWAS to conduct TWAS using the individual-level GWAS data

from ROS/MAP[26; 31] and MCADGS[32]. First, we trained the BVSR GReX imputation models using

samples (N=499) from the ROS/MAP cohort that contained both profiled genotype data and

transcriptomic data obtained from the dorsal lateral prefrontal cortex. All expression quantitative

traits were normalized and corrected for age at death, sex, postmortem interval (PMI), study

(ROS or MAP), batch effects, RNA integrity number (RIN), top three principal components

derived from genome-wide genotype data, and cell type proportions (oligodendrocytes,

astrocytes, microglia, neurons). The cell type proportions were derived by using CIBERSORT

pipeline[39] with single-cell RNA-seq transcriptome profiles from human brain tissues as the

reference[40] to de-convolute bulk RNA-seq data[41].

When we applied BGW-TWAS, we obtained GReX imputation models for 14,156 genes,

compared to respective 6,011 genes and 14,214 genes by PrediXcan and TIGAR that have at

least one cis-eQTL with nonzero effect size on expression quantitative trait. Across the 6,011

genes with GReX imputation models by PrediXcan, our BGW-TWAS approach had a smaller

train R^2 (squared correlation between fitted and profiled gene expression values) value for

expression quantitative traits for only 855 genes (Figure S1A). While TIGAR and BGW-TWAS

yielded a similarly number of GReX imputation models, BGW method is expected to result in

higher imputation accuracy when trans-eQTL play an important role in affecting gene

expression levels as shown by our simulation results. Of 13,142 genes that had imputation

models fitted by both TIGAR and BGW-TWAS, BGW-TWAS had smaller train R^2 for only 3,304

genes. That is, BGW-TWAS would be preferred for genes that have sparse eQTL, especially

trans-eQTL, while TIGAR would be preferred for genes that have less sparse eQTL that are

mostly cis-eQTL (Figure S 1B).

We imputed Bayesian GReX values for all remaining individuals with genotype data in

ROS/MAP and MCADGS by using Equation 4. We then conducted TWAS by testing the

association between the standardized GReX values (with unit variance) and both clinical and

pathological AD phenotypes. TWAS for these phenotypes controlled for age at death, sex,

smoking, ROS or MAP study, education level, and top three principal components derived from

genome-wide genotype data.

For the dichotomous phenotype of clinical diagnosis of AD, the case/control status was

determined by different rules and the available confounding variables were different for

ROS/MAP and MCADGS. Cognitive status at death for individuals from the ROS/MAP cohort is

based on the review of all longitudinal clinical data available at the time of death blinded to all

pathologic data. Individuals were classified as having no cognitive impairment, mild cognitive

impairment or AD. In this study, samples from individuals with AD were taken as cases and

samples from individuals without dementia i.e., either with no cognitive impairment or mild

cognitive impairment were taken as controls. For the MCADGS samples, cases were

determined for samples with a medical history of late onsite AD diagnosis, and available

confounding variables only included age, sex, and top three principal components derived from

GWAS data. Therefore, we meta-analyzed these two cohorts for AD clinical diagnosis by

applying the inverse-variance weighting method[42] to summary statistics obtained by TWAS per

cohort. We compared the meta-TWAS results obtained with BGW-TWAS to alternative TWAS

methods.

BGW-TWAS identified gene *ZC3H12B* (located on chromosome X) whose GReX values

were associated with AD with effect size $\beta = 0.265$, p-value 5.42×10^{-13}, and FDR $= 3.07 \times$

10^{-8} (Figure 2A; Table 1). Both within-cohort TWAS obtained positive effect sizes ($\beta =$

$0.22, 0.29$) for ROS/MAP and MCADGS, with respective p-value $= 2 \times 10^{-4}, 4.12 \times 10^{-10}$. On

the other hand, this gene was not identified by either PrediXcan or TIGAR because the

association of this gene is completely drive by trans-eQTL (Figures S2 and S4).

TWAS of pathological AD phenotypes were restricted to ROS/MAP from whom

postmortem AD indices were collected. TWAS were conducted for individuals with GWAS

genotype data and AD pathology indices — tangles (N=1,121), β-amyloid (N=1,114), and global

AD pathology (1,139). These results are shown in the Manhattan plots in Figures 2B, 3(A, B)

and Figures S2-S3 and S5-S7.

Using BGW-TWAS, *ZC3H12B* was identified to be associated with global AD pathology

with p-value$= 9.59 \times 10^{-7}$ (Figure 2B; Table 1) as well as neurofibrillary tangle density with p-

value$= 1.89 \times 10^{-6}$ (Figure 3A; Table 1). Gene *KCTD12* located on chromosome 12 was

identified to be significantly associated with β-amyloid load with p-value $= 3.44 \times 10^{-8}$ (Figure

3B; Table 1).

We show the BVSR posterior probabilities for considered SNPs to be eQTL for

ZC3H12B and *KCTD12* in Figure 4, and the standard eQTL analyses results for these two

genes in Figure S4. These data suggest that that the association between the imputed GReX

values of *ZC3H12B* and AD phenotypes is completely driven by trans-eQTL, while the

association between the GReX values of *KCTD12* and β-amyloid load is driven by both cis- and

trans- eQTL. Four of the top driven trans-eQTL (*rs12721051, rs4420638, rs56131196, rs157592;*

Table 2) for *ZC3H12B* are located in gene *APOC1* that is a known risk gene for AD[43] and blood

lipids[44-46], which is <12KB away from the well-known AD risk gene *APOE*[47]. Particularly,

rs12721051 located in the 3' UTR region of *APOC1* was identified as a GWAS signal of total

cholesterol levels[46]; *rs4420638* located in the downstream of *APOC1* is in linkage disequilibrium

(LD) with the *APOE-E4* allele (*rs429358*) and was identified to be a GWAS signal of various

blood lipids measurements (i.e., low density lipoprotein cholesterol measurement, C-reactive

protein measurement, triglyceride measurement, and total cholesterol measurement)[44] and AD[48];

rs56131196 located in the downstream and *rs157592* located in the regulatory region of *APOC1*

were identified as GWAS signals of AD and independent of *APOE-E4*[49]. Additionally, gene

ZC3H12B was found to regulate pro-inflammatory activation of macrophages[50] and has higher

expression in brain, spinal cord and thymus tissue types compared to other tissues[51]. These

results showed that the effects of these known GWAS signals (*rs4420638, rs56131196,*

rs157592) of AD could be mediated through the gene expression levels of *ZC3H12B*.

<u>TWAS of AD with Summary-level GWAS Data</u>

To validate our findings using individual-level GWAS data from ROS/MAP and

MCADGS, we conducted TWAS of AD using the publicly available IGAP GWAS summary

statistics[27]. Specifically, we used the GWAS summary statistics that were generated by meta-

analysis of four consortia (~17K cases and ~37K controls, Europeans): the Alzheimer's Disease

Genetic Consortium (ADGC), the Cohorts for Heart and Aging Research in Genomic

Epidemiology (CHARGE) Consortium, the European Alzheimer's Disease Initiative (EADI) and

the Genetic and Environmental Risk in Alzheimer's Disease (GERAD) Consortium.

BGW-TWAS identified 13 significant genes located in chromosome 3, 6, 7, 10, 11, 19,

and X, including known GWAS risk genes *HLA-DRB1*[52] and *APOC1,* and gene *ZC3H12B* that

was identified using individual-level GWAS data from ROS/MAP and MCADGS (Table 3).

Moreover, 7 of these genes (including *HLA-DRB1* and *APOC1*) were also identified when we

only considered cis-eQTL estimates by BVSR in TWAS. Gene *CEACAM19* near the well-known

GWAS risk gene *APOE* was also identified by S-PrediXcan and TIGAR. Known GWAS risk

gene *HLA-DRB1*[52] was also identified by S-PrediXcan (Table 3; Tables S1-S3).

Our results showed that by using BVSR estimates of cis- and trans- eQTL (BGW-

TWAS), most independent risk loci were identified including loci driven by trans-eQTL. For those

significant genes driven mainly by cis-eQTL, TWAS using BVSR estimates of cis-eQTL still

identified more independent significant risk loci (distributed over chromosomes 2, 3, 6, 11, and

19) than S-PrediXcan and TIGAR, including all 4 significant genes (*HLA-DRB1, SLC39A13,*

PVR, CEACAM19) identified by S-PrediXcan and 4 out of 21 significant genes (*ZNF227,*

ZFP112, PVR, CEACAM19) identified by TIGAR (Tables S1-3S). Although TIGAR identified

most significant TWAS genes (21), these genes are from chromosomes 11 and 19, which are

likely to be driven by the same cis-eQTL from two independent loci.

These TWAS results using summary-level GWAS data with a much larger sample size

validated our findings obtained with BGW-TWAS using individual-level GWAS data from

ROS/MAP and MCADGS.

Insights about eQTL Genetic Architecture

In addition to imputing Bayesian GReX values (Equation 5), the posterior probabilities of

having non-zero eQTL effect sizes estimated by BVSR also provide insights into the genetic

architecture of eQTL, especially about how potential eQTL are distributed across the genome.

Note that the posterior probability obtained from the BVSR model (Equations 2-4) is essentially

the expected probability for a SNP to be an eQTL. Therefore, the sum of posterior probabilities

of having non-zero eQTL effect sizes represents the expected number of eQTL.

From our simulation studies, we observed that the expected proportions of cis-eQTL

were consistent with the true proportions of causal cis-eQTL. The expected number of eQTL

obtained across simulation scenarios is presented in Table 4, where two out of five (40%)

causal eQTL and 11 out of 22 (50%) causal eQTL are from cis- genome regions. We can see

that, with higher true expression heritability, the expected number of eQTL is closer to the true

number of causal eQTL. We can also see that the expected number of eQTL is more accurate

for the scenario with 5 true causal eQTL than with 22 true causal eQTL, which is due to the fact

that BVSR model prefers relatively larger eQTL effect sizes. These simulation results

demonstrated the validity of our BGW-TWAS method based on the BVSR model as well as the

usefulness of the sum of posterior probabilities of having non-zero eQTL effect sizes.

For 14,156 genes with fitted GReX prediction models by BVSR using the ROS/MAP

data, after excluding 19 outlier genes with >100 expected eQTL, we obtained the average

number of expected eQTL as 2.44 (SD = 5.70) across genome-wide regions, 0.25 (SD = 1.24)

for cis-eQTL, and 2.48 (SD = 5.49) for trans-eQTL. That is, on overage, 88% of eQTL were from

trans- genome regions with respect to the target gene. We can see that ~90% genes with train

$R^2 > 0.05$ have ~2-3 average expected eQTL, and ~10% genes with train R^2 < 0.05 have >5

average expected eQTL (Table 4). By linking these findings with our simulation studies where

train R^2 is likely to be >0.05 when true expression heritability is >0.1, we can conclude that

~90% genes are likely to have true expression heritability >0.1.

Additionally, from our Bayesian estimates of the cis- and trans- specific posterior

probabilities of having non-zero eQTL effect sizes (i.e., π_{cis}, π_{trans} in Equation 3) for genome-

wide genes using ROS/MAP data (Figure S11), we can see that π_{cis} and π_{trans} clearly follow

different distributions. This also validates our assumptions of respective prior distribution for cis-

and trans- hyper parameters.

Discussion

In this paper, we proposed and validated a Bayesian Genome-wide TWAS (BGW-TWAS) method based on the BVSR[24] model to leverage the information of both cis- and trans- eQTL. We derived an efficient computational approach to fit the BVSR model with large-scale genomic data, by pruning genome regions that contain either at least one cis-SNP or one potential trans-eQTL and adapting the previously developed scalable EM-MCMC algorithm[25] with pre-calculated LD coefficients and summary statistics from standard eQTL analyses. BGW-TWAS extends previous TWAS methods[11-13] that only utilize partial genotype information from a small window of cis- SNPs to train the GReX imputation model.

Genotype data of trans-eQTL have been shown to explain a significant amount of variation of expression quantitative traits and provide important molecular mechanisms underlying known GWAS loci of complex diseases[22; 23]. The results from our simulation and application studies demonstrated that BGW-TWAS improves the yield of TWAS by levering both cis- and trans- eQTL information. For example, higher precision of GReX prediction and power of TWAS were obtained in our simulation studies when true causal trans-eQTL existed. These results showed that BGW-TWAS has a greater advantage for scenarios where eQTL have relatively large effect sizes for the expression quantitative traits (e.g., 5 vs. 22 true causal eQTL with the same expression heritability). This is because variable selection by the BVSR model is designed to select sparse signals with relatively large effect sizes as shown in previous GWAS[24; 25].

By applying our Bayesian approach to several human AD datasets, we identified a risk gene (*ZC3H12B*) with GReX values that were significantly associated with both clinical diagnosis of AD and postmortem AD pathology indices (neurofibrillary tangle density and global measure of AD pathology). This association was not identified by existing TWAS methods because this gene is shown to be completely driven by trans-eQTL. Importantly, a potential

biological mechanism was revealed by showing that the top driven trans-eQTL of *ZC3H12B* are

known GWAS signals of AD[43] and blood lipids[44-46] and <12KB away from the famous well-

known AD risk gene (*APOE*)[47]. Thus, we expect BGW-TWAS leveraging both cis- and trans-

eQTL has potential for making a large impact on advancing our understanding of complex

human diseases and traits.

By fitting BVSR models using both cis- and trans- eQTL, we can not only account for the

uncertainty for a SNP to be an eQTL to predict GReX (Equation 5), but also use the sum of

posterior probabilities of having non-zero eQTL effect sizes to estimate the expected number of

eQTL[24; 25]. The distribution of expected eQTL can also help characterize the underlying genetic

architecture of expression quantitative traits.

The current study has several limitations. First, while BGW-TWAS reduces the

computational burden for modeling both cis- and trans- eQTL, its computing costs are still

substantial to train GReX prediction models for genome-wide genes (~20K) per tissue type. It

requires approximately 30 minutes of computation time and 3GB memory per gene (with parallel

computation implemented in 4 CPU cores). Parallel computation can be employed to make use

of high-performance computation clusters with multiple cores to reduce computation time.

Second, our current method is designed to use pre-calculated in-sample LD coefficients and

summary statistics from single variant eQTL analyses, further work is required to expand this

approach to use approximate LD coefficients generated from reference samples of the same

ethnicity. Third, our simulation studies showed that the non-parametric Bayesian method TIGAR

performed best when all causal eQTL are cis- with relatively small effect sizes (e.g., 22 true

causal cis-eQTL). Our TWAS results of AD using the IGAP summary statistics demonstrated

that TIGAR and BGW-TWAS yield complementary findings. These results highlight the potential

utility of leveraging both methods especially for studies in which the true distributions of cis- and

trans- eQTL of the test genes are generally unknown.

467   In conclusion, the BGW-TWAS method presented herein provides a framework for

468 leveraging information from both cis- and trans- eQTL to conduct gene-based association

469 studies. Because trans- QTL are common for other quantitative omics traits, e.g., epigenetic,

470 proteomic, and metabonomic, our proposed computational procedure would be to investigate

471 other quantitative omics traits in gene-based association studies. Integrative method

472 developments will stand to benefit from our BGW-TWAS method, especially the perspectives of

473 leveraging information from trans- QTL and efficient computation techniques derived from this

474 paper. In addition, BGW-TWAS can be applied to study other complex human phenotypes to

475 identify potential risk genes that could be targeted in further drug discovery.

478 **Tables**

479 *Table 1. Significant risk genes identified by BGW-TWAS using individual-level GWAS data from*
480 *ROS/MAP and MCADGS cohorts.*

Gene	CHR	Position	Train R^2	p-value	Effect size (SD)	Phenotype
ZC3H12B	X	64,708,614	0.24	5.42 x 10^{-13}	0.265 (0.037)	AD
ZC3H12B	X	64,708,614	0.24	9.59 x 10^{-7}	0.142 (0.029)	Global AD pathology
ZC3H12B	X	64,708,614	0.24	1.89 x 10^{-6}	0.138 (0.029)	Tangles
KCTD12	13	77,454,311	0.09	3.44 x 10^{-8}	0.143 (0.026)	β-Amyloid

481
482
483 *Table 2. Trans-SNPs with top five posterior probability (PP)* > 0.003 *of having non-zero eQTL*
484 *effect sizes for gene ZC3H12B.*

CHR	POS	rsID	Function	MAF	PP	w	p-value
1	159,135,282	*rs3026946*	Intergenic	0.213	0.0147	-0.071	6.25×10^{-7}
19	45,422,160	*rs12721051*	3' UTR (APOC1)	0.161	0.0031	0.071	3.94×10^{-6}
19	45,422,846	*rs56131196*	Downstream (APOC1)	0.173	0.0048	0.069	1.75×10^{-6}
19	45,422,946	*rs4420638*	Downstream (APOC1)	0.173	0.0051	0.068	1.77×10^{-6}
19	45,424,514	*rs157592*	Regulatory Region (APOC1)	0.181	0.0056	0.075	1.43×10^{-6}

485
486

Table 3. Significant genes identified by BGW-TWAS using IGAP GWAS summary
statistics of AD. TWAS p-values by alternative methods, i.e., using BVSR cis-eQTL estimates
only, PrediXcan, and TIGAR are also listed. P-VALUEs for genes that were missed by TWAS
were indicated as "-". Significant gene *ZC3H12B* that was identified by BGW-TWAS using
individual-level GWAS data from ROS/MAP and MCADGS was also identified by BGW-TWAS
using IGAP summary-level GWAS statistics. Genes that were also identified as significant by
using BVSR cis-eQTL estimates only are denoted by *, identified by PrediXcan are denoted by
†; and identified by TIGAR are denoted by ⊗. In particular, gene *CEACAM19* from chromosome
19 was identified by all TWAS methods, and gene *HLA-DRB1* from chromosome 6 is a known
GWAS risk locus.
497

Gene	CHR	Position	TWAS P-VALUE			
			BGW-TWAS	BVSR cis-eQTL	PrediXcan	TIGAR
GPX1*	3	49,394,608	2.45×10^{-98}	2.45×10^{-98}	-	3.15×10^{-1}
FAM86DP	3	75,484,261	1.55×10^{-13}	4.81×10^{-1}	5.38×10^{-1}	9.63×10^{-1}
BTN3A2*	6	26,378,546	1.59×10^{-26}	1.56×10^{-26}	3.17×10^{-1}	5.04×10^{-1}
ZNF192*	6	28,124,089	1.26×10^{-32}	1.25×10^{-32}	8.56×10^{-2}	2.07×10^{-1}
AL022393.7*	6	28,144,452	3.25×10^{-178}	2.24×10^{-178}	1.50×10^{-1}	8.36×10^{-2}
***HLA-DRB1**†**	**6**	**32,557,625**	$\mathbf{1.02 \times 10^{-12}}$	$\mathbf{8.99 \times 10^{-13}}$	$\mathbf{2.06 \times 10^{-6}}$	-
AEBP1	7	44,154,161	5.55×10^{-220}	8.62×10^{-1}	6.69×10^{-1}	4.19×10^{-1}
BUB3	10	124,924,886	6.64×10^{-18}	1.05×10^{-2}	-	4.76×10^{-1}
FBXO3	11	33,796,089	1.48×10^{-9}	6.88×10^{-1}	-	1.13×10^{-1}
***CEACAM19**†⊗**	**19**	**45,187,631**	$\mathbf{4.7 \times 10^{-13}}$	$\mathbf{2.54 \times 10^{-13}}$	$\mathbf{3.60 \times 10^{-12}}$	$\mathbf{2.83 \times 10^{-16}}$
APOC1*	19	45,422,606	8.9×10^{-11}	1.11×10^{-10}	3.18×10^{-6}	7.2×10^{-3}
ZC3H12B	**X**	**64,727,767**	$\mathbf{2.08 \times 10^{-37}}$	-	-	-
CXorf56	X	118,699,397	6.02×10^{-07}	-	-	-

499 *Table 4. Average sums of posterior probabilities of having non-zero eQTL effect sizes that are*
500 *stratified based on gene expression heritability (either true simulated heritability in*
501 *simulation studies or the range of train R^2 of the fitted BVSR models with ROS/MAP data). The*
502 *simulation scenarios presented here are those with 2 of 5 and 11 of 22 true causal eQTL from*
503 *the cis- regions.*

Gene Expression Heritability		Sum of Posterior Probabilities		
		Whole Genome	Cis-Region	Trans-Region
5 True Causal eQTL	0.05	0.79	0.46	0.33
	0.1	2.28	1.13	1.15
	0.2	3.72	1.44	2.28
	0.5	4.91	1.56	3.35
22 True Causal eQTL	0.05	0.05	0.02	0.03
	0.1	0.21	0.11	0.10
	0.2	1.43	0.87	0.56
	0.5	6.46	3.89	2.57
ROS/MAP	(0, 0.05) 1,504 genes	6.63	0.60	6.23
	(0.05, 0.1) 1,964 genes	1.45	0.13	1.32
	(0.1, 0.25) 6,617 genes	2.00	0.17	1.83
	(0.25, 0.5) 3,224 genes	2.66	0.22	2.44
	(0.5, 1) 474 genes	3.04	0.31	2.73

Figure Legends

Figure 1. Simulation TWAS Studies Comparing BGW-TWAS, BVSR with cis-eQTL only, PrediXcan and TIGAR Methods. Simulation studies used various gene expression heritability $h_e^2 = (0.05, 0.1, 0.2, 0.5)$ and various true causal cis-eQTL proportions. (A, B) Test R² and TWAS power comparison with 5 true causal eQTL. BGW-TWAS was found to out-perform the alternative methods when a non-negligible proportion of true causal eQTL were from trans- genome regions. (C, D) Test R² and TWAS power comparison with 22 true causal eQTL. BGW-TWAS was found to out-perform alternative method when >50% of true causal eQTL were from trans- genome regions and $h_e^2 > 0.1$.

Figure 2. Manhattan plots of BGW-TWAS results of AD clinical diagnosis (A) and global AD pathology (B). Red lines denote genome-wide significant threshold (2.5×10^{-6}) for gene-based association studies. Gene ZC3H12B was found to be significantly associated with both AD clinical diagnosis and global AD pathology.

Figure 3. Manhattan plots of BGW-TWAS results of neurofibrillary tangle density (A) and β-amyloid load (B). Red lines denote genome-wide significant threshold (2.5×10^{-6}) for gene-based association studies. Gene ZC3H12B was found to be significantly associated with neurofibrillary tangle density. Gene KCTD12 was found to be significantly associated with and β-amyloid load.

Figure 4. BVSR posterior probabilities (PP) of having non-zero eQTL effect sizes for analyzed cis- and trans- SNPs, with target genes ZC3H12B (A) and KCTD12 (B). Gene ZC3H12B located on chromosome X has top trans-eQTL from chromosomes 1, 6, and 19, where all eQTL are of trans-eQTL. Gene KCTD12 located on chromosome 12 has top cis-eQTL from chromosome 12 and trans-eQTL from chromosomes 4 and 6.

Supplemental Data

Supplemental Figures, Tables, Methods, and References.

Data and Code Availability

ROS/MAP data can be requested through Rush Alzheimer's Disease Center (http://www.radc.rush.edu/) and synapse (https://www.synapse.org/#!Synapse:syn3219045). MCADGS data can be requested through synapse (https://www.synapse.org/#!Synapse:syn2910256). IGAP summary statistics are available from http://web.pasteur-lille.fr/en/recherche/u744/igap/igap_download.php. Summary statistics

547   generated from our BGW-TWAS methods for studying AD are publicly available through

548   synapse (XXX). Source code of BGW-TWAS is available through Github with link

549   https://github.com/yanglab-emory/BGW-TWAS.

551   **Declaration of Interests**

552   The authors declare no competing interests.
553
554   **Acknowledgement**
555

J.Y. was supported by the startup funding from Department of Human Genetics at Emory

University School of Medicine. ROS/MAP study data were provided by the Rush Alzheimer's

Disease Center, Rush University Medical Center, Chicago, IL. Data collection was supported

through funding by NIA grants P30AG10161, R01AG15819, R01AG17917, R01AG30146,

R01AG36836, U01AG32984, U01AG46152, U01AG61356, the Illinois Department of Public

Health, and the Translational Genomics Research Institute. The MCADGC led by Dr. Nilüfer

Ertekin-Taner and Dr. Steven G. Younkin, Mayo Clinic, Jacksonville, FL uses samples from the

Mayo Clinic Study of Aging, the Mayo Clinic Alzheimer's Disease Research Center, and the

Mayo Clinic Brain Bank. MCADGC data collection was supported through funding by NIA grants

P50 AG016574, R01 AG032990, U01 AG046139, R01 AG018023, U01 AG006576, U01

AG006786, R01 AG025711, R01 AG017216, R01 AG003949, NINDS grant R01 NS080820,

CurePSP Foundation, and support from Mayo Foundation.

Web Resources

BGW-TWAS, https://github.com/yanglab-emory/BGW-TWAS

TIGAR, https://github.com/yanglab-emory/TIGAR

PrediXcan, https://github.com/hakyim/PrediXcan
573

References
1. Hirschhorn, J.N., and Daly, M.J. (2005). Genome-wide association studies for common
diseases and complex traits. Nature reviews Genetics 6, 95-108.
2. Wellcome Trust Case Control, C. (2007). Genome-wide association study of 14,000 cases
of seven common diseases and 3,000 shared controls. Nature 447, 661-678.
3. McCarthy, M.I., Abecasis, G.R., Cardon, L.R., Goldstein, D.B., Little, J., Ioannidis, J.P., and
Hirschhorn, J.N. (2008). Genome-wide association studies for complex traits:
consensus, uncertainty and challenges. Nature reviews Genetics 9, 356-369.
4. Nikpay, M., Goel, A., Won, H.H., Hall, L.M., Willenborg, C., Kanoni, S., Saleheen, D.,
Kyriakou, T., Nelson, C.P., Hopewell, J.C., et al. (2015). A comprehensive 1,000
Genomes-based genome-wide association meta-analysis of coronary artery disease.
Nature genetics 47, 1121-1130.
5. Visscher, P.M., Wray, N.R., Zhang, Q., Sklar, P., McCarthy, M.I., Brown, M.A., and Yang, J.
(2017). 10 Years of GWAS Discovery: Biology, Function, and Translation. American
journal of human genetics 101, 5-22.
6. Huang, Q. (2015). Genetic study of complex diseases in the post-GWAS era. J Genet
Genomics 42, 87-98.
7. Gallagher, M.D., and Chen-Plotkin, A.S. (2018). The Post-GWAS Era: From Association to
Function. American journal of human genetics 102, 717-730.
8. GTEx Consortium. (2017). Genetic effects on gene expression across human tissues.
Nature 550, 204-213.
9. Nicolae, D.L., Gamazon, E., Zhang, W., Duan, S., Dolan, M.E., and Cox, N.J. (2010). Trait-
associated SNPs are more likely to be eQTLs: annotation to enhance discovery from
GWAS. PLoS genetics 6, e1000888.
10. Pickrell, J.K., Marioni, J.C., Pai, A.A., Degner, J.F., Engelhardt, B.E., Nkadori, E., Veyrieras,
J.B., Stephens, M., Gilad, Y., and Pritchard, J.K. (2010). Understanding mechanisms
underlying human gene expression variation with RNA sequencing. Nature 464,
768-772.
11. Gamazon, E.R., Wheeler, H.E., Shah, K.P., Mozaffari, S.V., Aquino-Michaels, K., Carroll, R.J.,
Eyler, A.E., Denny, J.C., GTEx Consortium, Nicolae, D.L., et al. (2015). A gene-based
association method for mapping traits using reference transcriptome data. Nature
genetics 47, 1091-1098.
12. Gusev, A., Ko, A., Shi, H., Bhatia, G., Chung, W., Penninx, B.W., Jansen, R., de Geus, E.J.,
Boomsma, D.I., Wright, F.A., et al. (2016). Integrative approaches for large-scale
transcriptome-wide association studies. Nature genetics 48, 245-252.
13. Nagpal, S., Meng, X., Epstein, M.P., Tsoi, L.C., Patrick, M., Gibson, G., De Jager, P.L.,
Bennett, D.A., Wingo, A.P., Wingo, T.S., et al. (2019). TIGAR: An Improved Bayesian
Tool for Transcriptomic Data Imputation Enhances Gene Mapping of Complex
Traits. American journal of human genetics 105, 258-266.
14. Barbeira, A.N., Dickinson, S.P., Bonazzola, R., Zheng, J., Wheeler, H.E., Torres, J.M.,
Torstenson, E.S., Shah, K.P., Garcia, T., Edwards, T.L., et al. (2018). Exploring the
phenotypic consequences of tissue specific gene expression variation inferred from
GWAS summary statistics. Nat Commun 9, 1825.
15. Lappalainen, T., Sammeth, M., Friedlander, M.R., t Hoen, P.A., Monlong, J., Rivas, M.A.,
Gonzalez-Porta, M., Kurbatova, N., Griebel, T., Ferreira, P.G., et al. (2013).

Transcriptome and genome sequencing uncovers functional variation in humans.
Nature 501, 506-511.
16. Gibbs, J.R., van der Brug, M.P., Hernandez, D.G., Traynor, B.J., Nalls, M.A., Lai, S.L.,
Arepalli, S., Dillman, A., Rafferty, I.P., Troncoso, J., et al. (2010). Abundant
quantitative trait loci exist for DNA methylation and gene expression in human
brain. PLoS genetics 6, e1000952.
17. Mancuso, N., Shi, H., Goddard, P., Kichaev, G., Gusev, A., and Pasaniuc, B. (2017).
Integrating Gene Expression with Summary Association Statistics to Identify Genes
Associated with 30 Complex Traits. American journal of human genetics 100, 473-
487.
18. Gusev, A., Mancuso, N., Won, H., Kousi, M., Finucane, H.K., Reshef, Y., Song, L., Safi, A.,
Schizophrenia Working Group of the Psychiatric Genomics, C., McCarroll, S., et al.
(2018). Transcriptome-wide association study of schizophrenia and chromatin
activity yields mechanistic disease insights. Nature genetics 50, 538-548.
19. Wu, L., Shi, W., Long, J., Guo, X., Michailidou, K., Beesley, J., Bolla, M.K., Shu, X.O., Lu, Y.,
Cai, Q., et al. (2018). A transcriptome-wide association study of 229,000 women
identifies new candidate susceptibility genes for breast cancer. Nature genetics 50,
968-978.
20. Raj, T., Li, Y.I., Wong, G., Humphrey, J., Wang, M., Ramdhani, S., Wang, Y.C., Ng, B., Gupta,
I., Haroutunian, V., et al. (2018). Integrative transcriptome analyses of the aging
brain implicate altered splicing in Alzheimer's disease susceptibility. Nature
genetics 50, 1584-1592.
21. Wainberg, M., Sinnott-Armstrong, N., Mancuso, N., Barbeira, A.N., Knowles, D.A., Golan,
D., Ermel, R., Ruusalepp, A., Quertermous, T., Hao, K., et al. (2019). Opportunities and
challenges for transcriptome-wide association studies. Nature genetics 51, 592-599.
22. Lloyd-Jones, L.R., Holloway, A., McRae, A., Yang, J., Small, K., Zhao, J., Zeng, B., Bakshi, A.,
Metspalu, A., Dermitzakis, M., et al. (2017). The Genetic Architecture of Gene
Expression in Peripheral Blood. American journal of human genetics 100, 371.
23. Võsa, U., Claringbould, A., Westra, H.-J., Bonder, M.J., Deelen, P., Zeng, B., Kirsten, H.,
Saha, A., Kreuzhuber, R., Kasela, S., et al. (2018). Unraveling the polygenic
architecture of complex traits using blood eQTL metaanalysis. 447367.
24. Guan, Y.T., and Stephens, M. (2011). Bayesian Variable Selection Regression for
Genome-Wide Association Studies and Other Large-Scale Problems. Annals of
Applied Statistics 5, 1780-1815.
25. Yang, J., Fritsche, L.G., Zhou, X., Abecasis, G., and International Age-Related Macular
Degeneration Genomics, C. (2017). A Scalable Bayesian Method for Integrating
Functional Information in Genome-wide Association Studies. American journal of
human genetics 101, 404-416.
26. De Jager, P.L., Ma, Y., McCabe, C., Xu, J., Vardarajan, B.N., Felsky, D., Klein, H.U., White,
C.C., Peters, M.A., Lodgson, B., et al. (2018). A multi-omic atlas of the human frontal
cortex for aging and Alzheimer's disease research. Sci Data 5, 180142.
27. Lambert, J.C., Ibrahim-Verbaas, C.A., Harold, D., Naj, A.C., Sims, R., Bellenguez, C.,
DeStafano, A.L., Bis, J.C., Beecham, G.W., Grenier-Boley, B., et al. (2013). Meta-
analysis of 74,046 individuals identifies 11 new susceptibility loci for Alzheimer's
disease. Nature genetics 45, 1452-1458.
28. Casella, G. (2001). Empirical Bayes Gibbs sampling. Biostatistics 2, 485-500.

29. Bennett, D.A., Schneider, J.A., Arvanitakis, Z., and Wilson, R.S. (2012). Overview and
findings from the religious orders study. Curr Alzheimer Res 9, 628-645.
30. Bennett, D.A., Schneider, J.A., Buchman, A.S., Barnes, L.L., Boyle, P.A., and Wilson, R.S.
(2012). Overview and findings from the rush Memory and Aging Project. Curr
Alzheimer Res 9, 646-663.
31. Bennett, D.A., Buchman, A.S., Boyle, P.A., Barnes, L.L., Wilson, R.S., and Schneider, J.A.
(2018). Religious Orders Study and Rush Memory and Aging Project. J Alzheimers
Dis 64, S161-S189.
32. Carrasquillo, M.M., Zou, F., Pankratz, V.S., Wilcox, S.L., Ma, L., Walker, L.P., Younkin, S.G.,
Younkin, C.S., Younkin, L.H., Bisceglio, G.D., et al. (2009). Genetic variation in
PCDH11X is associated with susceptibility to late-onset Alzheimer's disease. Nature
genetics 41, 192-198.
33. Zou, F., Chai, H.S., Younkin, C.S., Allen, M., Crook, J., Pankratz, V.S., Carrasquillo, M.M.,
Rowley, C.N., Nair, A.A., Middha, S., et al. (2012). Brain expression genome-wide
association study (eGWAS) identifies human disease-associated variants. PLoS
genetics 8, e1002707.
34. Allen, M., Carrasquillo, M.M., Funk, C., Heavner, B.D., Zou, F., Younkin, C.S., Burgess, J.D.,
Chai, H.S., Crook, J., Eddy, J.A., et al. (2016). Human whole genome genotype and
transcriptome data for Alzheimer's and other neurodegenerative diseases. Sci Data
3, 160089.
35. De Jager, P.L., Shulman, J.M., Chibnik, L.B., Keenan, B.T., Raj, T., Wilson, R.S., Yu, L.,
Leurgans, S.E., Tran, D., Aubin, C., et al. (2012). A genome-wide scan for common
variants affecting the rate of age-related cognitive decline. Neurobiol Aging 33, 1017
e1011-1015.
36. 1000 Genomes Project Consortium. (2012). An integrated map of genetic variation from
1,092 human genomes. Nature 491, 56-65.
37. Das, S., Forer, L., Schonherr, S., Sidore, C., Locke, A.E., Kwong, A., Vrieze, S.I., Chew, E.Y.,
Levy, S., McGue, M., et al. (2016). Next-generation genotype imputation service and
methods. Nature genetics 48, 1284-1287.
38. De Jager, P.L., Srivastava, G., Lunnon, K., Burgess, J., Schalkwyk, L.C., Yu, L., Eaton, M.L.,
Keenan, B.T., Ernst, J., McCabe, C., et al. (2014). Alzheimer's disease: early alterations
in brain DNA methylation at ANK1, BIN1, RHBDF2 and other loci. Nat Neurosci 17,
1156-1163.
39. Newman, A.M., Liu, C.L., Green, M.R., Gentles, A.J., Feng, W., Xu, Y., Hoang, C.D., Diehn, M.,
and Alizadeh, A.A. (2015). Robust enumeration of cell subsets from tissue
expression profiles. Nature methods 12, 453-457.
40. Darmanis, S., Sloan, S.A., Zhang, Y., Enge, M., Caneda, C., Shuer, L.M., Hayden Gephart,
M.G., Barres, B.A., and Quake, S.R. (2015). A survey of human brain transcriptome
diversity at the single cell level. Proceedings of the National Academy of Sciences
112, 7285-7290.
41. Wingo, T.S., Yang, J., Fan, W., Min Canon, S., Gerasimov, E.S., Lori, A., Logsdon, B., Yao, B.,
Seyfried, N.T., Lah, J.J., et al. (2020). Brain microRNAs associated with late-life
depressive symptoms are also associated with cognitive trajectory and dementia.
npj Genomic Medicine 5, 6.
42. Hartung, J., Knapp, G., and Sinha, B.K. (2011). Statistical meta-analysis with
applications.(John Wiley & Sons).

43. Marioni, R.E., Harris, S.E., Zhang, Q., McRae, A.F., Hagenaars, S.P., Hill, W.D., Davies, G.,
Ritchie, C.W., Gale, C.R., Starr, J.M., et al. (2018). GWAS on family history of
Alzheimer's disease. Transl Psychiatry 8, 99.
44. Ligthart, S., Vaez, A., Hsu, Y.H., Inflammation Working Group of the, C.C., Pmi Wg, X.C.P.,
LifeLines Cohort, S., Stolk, R., Uitterlinden, A.G., Hofman, A., Alizadeh, B.Z., et al.
(2016). Bivariate genome-wide association study identifies novel pleiotropic loci for
lipids and inflammation. BMC genomics 17, 443.
45. Ligthart, S., Vaez, A., Vosa, U., Stathopoulou, M.G., de Vries, P.S., Prins, B.P., Van der Most,
P.J., Tanaka, T., Naderi, E., Rose, L.M., et al. (2018). Genome Analyses of >200,000
Individuals Identify 58 Loci for Chronic Inflammation and Highlight Pathways that
Link Inflammation and Complex Disorders. American journal of human genetics
103, 691-706.
46. Klarin, D., Damrauer, S.M., Cho, K., Sun, Y.V., Teslovich, T.M., Honerlaw, J., Gagnon, D.R.,
DuVall, S.L., Li, J., Peloso, G.M., et al. (2018). Genetics of blood lipids among ~300,000
multi-ethnic participants of the Million Veteran Program. Nature genetics 50, 1514-
1523.
47. Genin, E., Hannequin, D., Wallon, D., Sleegers, K., Hiltunen, M., Combarros, O., Bullido,
M.J., Engelborghs, S., De Deyn, P., Berr, C., et al. (2011). APOE and Alzheimer disease:
a major gene with semi-dominant inheritance. Mol Psychiatry 16, 903-907.
48. Kamboh, M.I., Demirci, F.Y., Wang, X., Minster, R.L., Carrasquillo, M.M., Pankratz, V.S.,
Younkin, S.G., Saykin, A.J., Alzheimer's Disease Neuroimaging, I., Jun, G., et al. (2012).
Genome-wide association study of Alzheimer's disease. Transl Psychiatry 2, e117.
49. Zhou, X., Chen, Y., Mok, K.Y., Kwok, T.C.Y., Mok, V.C.T., Guo, Q., Ip, F.C., Chen, Y.,
Mullapudi, N., Alzheimer's Disease Neuroimaging, I., et al. (2019). Non-coding
variability at the APOE locus contributes to the Alzheimer's risk. Nat Commun 10,
3310.
50. Liang, J., Wang, J., Azfer, A., Song, W., Tromp, G., Kolattukudy, P.E., and Fu, M. (2008). A
novel CCCH-zinc finger protein family regulates proinflammatory activation of
macrophages. J Biol Chem 283, 6337-6346.
51. Fagerberg, L., Hallstrom, B.M., Oksvold, P., Kampf, C., Djureinovic, D., Odeberg, J.,
Habuka, M., Tahmasebpoor, S., Danielsson, A., Edlund, K., et al. (2014). Analysis of
the human tissue-specific expression by genome-wide integration of
transcriptomics and antibody-based proteomics. Mol Cell Proteomics 13, 397-406.
52. Jansen, I.E., Savage, J.E., Watanabe, K., Bryois, J., Williams, D.M., Steinberg, S., Sealock, J.,
Karlsson, I.K., Hagg, S., Athanasiu, L., et al. (2019). Genome-wide meta-analysis
identifies new loci and functional pathways influencing Alzheimer's disease risk.
Nature genetics 51, 404-413.
748

A)

BGW TWAS of AD

B)

BGW TWAS of Global AD Pathology