1

Age-related decrements in cortical gyrification:

Evidence from an accelerated longitudinal dataset

Christopher R. Madan

School of Psychology, University of Nottingham, Nottingham, UK

† Corresponding author.

Christopher R. Madan
School of Psychology, University Park
University of Nottingham
Nottingham, NG7 2RD, UK.
Email: christopher.madan@nottingham.ac.uk.

29   **Abstract**

Cortical gyrification has been found to decrease due to aging, but thus far this has only been

examined in cross-sectional samples. Interestingly, the topography of these age-related

differences in gyrification follow a distinct gradient along the cortex relative to age effects on

cortical thickness, likely suggesting a different underlying neurobiological mechanism. Here

I examined several aspects of gyrification in an accelerated longitudinal dataset of 280

healthy adults aged 45-92 with an interval between first and last MRI session of up to 10

years (total of 815 MRI sessions). Results suggest that age changes in sulcal morphology

underlie these changes in gyrification.

Keywords: structural MRI; cortical folding; brain morphology; sulcal morphology

Introduction

Over the past decade, brain imaging studies have demonstrated several gradients in activation related to functional networks of regions (e.g., Margulies et al., 2016; Murphy et al., 2019; Sormaz et al., 2018). Distinct cortical gradients in structural properties of the brain also exist, such as in cortical thickness (e.g., Hogstrom et al., 2013; Madan & Kensinger, 2018; Salat et al., 2004; Sowell et al., 2003) and myelination (e.g., Carradus et al., 2020; Glasser & Van Essen, 2011; Grydekand et al., 2013; Mangeat et al., 2015; Shafee et al., 2015). However, one of the most identifiable characteristics of the human brain is its folding structure. Despite macro-scale consistencies between individuals, everyone's cortical folding pattern is unique. While these folding patterns change due to aging, the underlying process of this change is not well understood.

Elias and Schwartz (1969) developed a procedure to quantify cortical gyrification as a ratio between the area of the total cortical surface relative to the 'exposed' surface. Here a stereological approach was used, though was limited by available technologies. This methodology was later refined to use the pial and outer contour lengths estimated from coronal sections by Zilles et al. (1988). The outlines shown in Figure 1A display this visually, with the pial length shown in blue and the outer contour length shown in green. The ratio of these lengths provide a measure of cortical folding, referred to as the gyrification index. A brain with a lower gyrification index will be smoother and less folded; a rodent's brain has a much lower gyrification index than a human brain. Extending this procedure to 3D surfaces, Schaer et al. (2008, 2012) developed an automated approach for calculating cortical gyrification from a surface mesh, through the generation of an outer surface that encloses the gray matter surface generated by FreeSurfer (as shown in Figure 1B) along with the corresponding surface measure of gyrification (Figure 1C).

65
66 **Figure 1. Illustration of the calculation procedure for global gyrification.** (A) From the
67 original T1-weighted volume, the outer contour of the gray matter is traced, and then
68 surrounded by a smooth enclosing surface. (B) When done as a 3D surface, this results in the
69 pial (blue) and enclosing (green) surfaces. (C) The gyrification index (GI) is the ratio of the
70 cortical surface area divided by the surface area of the enclosing surface.
71

As initially shown by Zilles et al. (1988) and replicated in more recent studies (e.g.,

Hogstrom et al., 2013; Madan & Kensinger, 2016), gyrification is relatively highest in the

parietal and temporal lobes. Several studies have shown that global cortical gyrification

decreases with age in several cross-sectional samples (Cao et al., 2017; Hogstrom et al.,

2013; Lamballais et al., 2020; Madan & Kensinger, 2016, 2018; Madan, 2018). It has also

been shown that the topography of these changes is distinct from cortical thickness, where

gyrification primarily decreases in the parietal lobe (e.g., Hogstrom et al., 2013; Madan &

Kensinger, 2016, 2018). Cortical thickness, in contrast, primarily decreases in frontal and

temporal regions—with age related decreases associated with changes in cortical myelination

affecting gray/white matter contrast as well as decreases in the dendritic density of pyramidal

neurons (Dickstein et al., 2007; Duan et al., 2003; Hao et al., 2007; Peters, 2002),

mechanisms distinct from age changes in gyrification. Further demonstrating the utility of

gyrification measurements, studies of patient populations have demonstrated differences in

gyrification relative to healthy individuals in relation to Alzheimer's disease (King et al.,

2010; Liu et al., 2012), schizophrenia (Cao et al., 2017; Palaniyappan & Liddle, 2012;

Palaniyappan et al., 2015), autism (Schaer et al., 2013), and major depression disorder (Cao

et al., 2017), among other psychiatric disorders. Within healthy samples, age-related

differences in cortical folding structure have also been associated with individual functional

differences (e.g., see Lamballais et al., 2020; McDonough & Madan, in press).

Using cross-sectional data, global gyrification has previously been estimated to

decrease by approximately 0.035 GI/decade (Madan & Kensinger, 2016; Madan, 2018),

though there is also a consideration of measurement validity. For example, in an analysis of

test-retest reliability, where ten sessions were conducted just 2-3 days apart (each), a mean

within-participant deviation of 0.04 was found (Madan & Kensinger, 2017). Moreover, plots

of cross-sectional data reveal a large amount of off-axis variability (i.e., age-unrelated)

indicating that, though reproducible, the link between gyrification and aging is relatively

weak and influenced by many other factors. Based on these considerations, a longitudinal

dataset with a larger interval between scans would be prudent for evaluating the influence of

aging on cortical gyrification.

Here I examine changes in cortical gyrification in an accelerated longitudinal sample

to directly examine cortical folding changes with age at the individual level, rather than in

cross-sectional datasets. While accelerated longitudinal samples have previously been used to

examine age changes in other morphological measures, such as cortical thickness (e.g.,

Storsve et al., 2014), this has yet to be done with gyrification. It is currently unclear how

gyrification changes with aging—for instance, is the decrease in gyrification in the parietal

lobe more evident because the parietal lobe has the highest gyrification? Moreover, does the

distribution of gyrification change with age?

111   **Methods**

Dataset

Data from 280 healthy adults (aged 45-92) from the OASIS-3 (Open Access Series of

Imaging Studies 3) dataset (LaMontagne et al., 2019) were included in the analyses presented

here. From the full dataset, participants were only included in the present analyses if four

conditions were met. (1) There must be at least two MRI sessions available with T1-weighted

volumes collected. (2) MRI data had to be collected using either the Siemens TIM Trio 3 T

scanner or Siemens BioGraph mMR PET-MR 3 T scanner (less than 5% of the otherwise

available data were collected using one of three other MRI scanners). (3) There had to be an

interval from first to last MRI session spanning at least three years. (4) From the clinical

sessions, there had a Clinical Dementia Rating (CDR) score of zero at every assessment. All

raw data is available from https://www.oasis-brains.org.

123   Across the 280 included adults, the Mini-Mental State Exam (MMSE) was

administered in a total of 1904 clinical sessions, with between 2 and 20 clinical sessions per

participant ($M \pm SD = 6.70 \pm 3.21$ administrations). Of the 1904 administrations, only four

yielded an MMSE score below 25. For two individuals, these sub-25 MMSE scores were

later followed by higher MMSE scores (here, scoring at least 29) on multiple subsequent

administrations. For the remaining two individuals, these were the last MMSE scores

recorded (for these specific individuals, there were 5 and 9 MMSE administrations,

respectively).

The interval between the first and last MRI session acquired for each participant

ranged from 3 to 10 years ($M\pm SD = 5.47\pm 1.91$ years). Of the 280 participants, 117 had two

MRI sessions, 96 had three sessions, 45 had four sessions, and 22 had five or more sessions

(to a maximum of seven). In total, 1138 T1-weighted volumes from 815 MRI sessions were

examined. If more than one T1-weighted volume was acquired during an MRI session, all

volumes were processed (see procedure below) and the gyrification estimates for the scans

were averaged. The same protocol was used for the other derived measures.

Structural data was collected using two different Siemens 3 T MRI scanners with

MPRAGE sequences: (1) 581 of the 815 MRI sessions were collected with a Siemens TIM

Trio 3 T scanner, with parameters: TR=2.4 ms; TE=3.2 ms; flip angle=8°; voxel size=1×1×1

mm. (2) The remaining 234 MRI sessions were collected with a Siemens BioGraph mMR

PET-MR 3 T scanner, with parameters: TR=2.3 ms; TE=3.0 ms; flip angle=9°; voxel

size=1×1×1 mm.

MRI Processing

All T1-weighted volumes were processed using FreeSurfer v6.0 on a machine running

RedHat 4.8.5-16 (Fischl, 2012; Fischl & Dale, 2000; Fischl et al., 2002). (Note, these are not

the same FreeSurfer outputs as publicly distributed from OASIS, which were estimated using

older versions of FreeSurfer.) All T1-weighted volumes were processed independently with

the standard FreeSurfer pipeline (i.e., `recon-all`), i.e., not using the longitudinal processing

pipeline, to allow the individual scan derived measures to be comparable in reliability to

previous cross-sectional studies.

Gyrification index was calculated using FreeSurfer, as described in Schaer et al.

(2008, 2012) and illustrated in Figure 1. This process involves generating an enclosing

surface which serves the same purpose as the outer contour used by Zilles et al. (1988) but

exists in 3D space. This enclosing surface involves a parameter of how 'tight' to bridge

across gyri, as opposed to falling into sulci. The default settings were used, based on

validation work conducted by Schaer and colleagues.

Sulcal morphology, width and depth, was estimated for 16 major sulci, using the

calcSulc toolbox (Madan, 2019a). The sulci are the central, post-central, superior frontal,

inferior frontal, parieto-occipital, occipito-temporal, middle occipital and lunate, and

marginal part of the cingulate sulci, in both the left and right hemispheres. Analyses here are

only conducted on the mean sulcal width and depth, averaging across the 16 sulci.

Anterior-posterior gradient in gyrification. The standard FreeSurfer pipeline results in

variation in the number of surface mesh vertices for each participant. To adjust for this,

gyrification measurements for each MRI session were resampled to the common surface

space using FreeSurfer's spherical surface co-registration (mris_preproc; Fischl et al., 1999).

This registration allows for vertex-wise comparisons in gyrification across participants, as

shown in Figure 2A.

This common surface space was then divided into 200 coronal sections, analogous to

the contour procedure used by Zilles et al. (1988). This allows for mean gyrification to be

simplified into an anterior-posterior plot, as demonstrated in Figure 2B. Note that, however,

vertex-wise gyrification measures were first estimated based on the Schaer et al. (2012)

procedure that relies on 25-mm radius local region of interest. This results in a gyrification

index value for each vertex of the surface mesh, referred to as a local gyrification index.

Gyrification for each section is spatially autocorrelated with adjacent sections. To-date there

do not appear to be any publications that have examined the anterior-posterior gyrification

pattern using the FreeSurfer gyrification calculation implementation (i.e., the Schaer et al.

[2012] approach).

Figure 2. Illustration of the calculation of the anterior-posterior gradient. (A) Individual brain pial surfaces are used to generate local gyrification index topography (based on Schaer et al., 2012), these are then resampled to the common space, through registration of the individual pial surface to the FreeSurfer standard surface. (B) The local gyrification index is then averaged across vertices for 200 coronal sections, shown for the same two example surfaces as in panel A.

Statistics

Age-related changes in global gyrification were examined as the slope of decline in gyrification with age. In subsequent analyses, the relative contribution of sulcal width and depth, in explaining the age decrements in gyrification are evaluated using a mediation analysis.

196   **Results**

Global gyrification

Before examining the topography of gyrification, I evaluated age changes in global

gyrification, which itself is yet to be examined in a longitudinal dataset. The decline in

gyrification was estimated using a linear mixed effects model to estimate the slope (allowing

for random intercepts for each participant; i.e., different starting points, but a common slope).

As shown in Figure 3, the fixed effect of age was statistically significant [$t(813)=14.41$,

$p<.001$] with a decreasing slope of 0.04291 GI/decade [95% C.I. = 0.03499—0.04603].

For comparison, a similar analysis was conducted on fractal dimensionality, another

measures of cortical structure. The results of this analysis are reported in the Appendix.

207
Figure 3. Age changes in global gyrification. Each coloured line represents an individual
participant, with each dot corresponding to an MRI session. The plot has been divided into
panels for each set of timepoints to improve readability. Dashed line present in all panels
show the decrease in global gyrification based on all data, estimated using the linear mixed
effects model.

Anterior-posterior gradient in gyrification

The results of this analysis are shown in Figure 4. Looking from anterior to posterior, the

gradient gradually rises to a peak mid-way through the frontal lobe (approx. percentile 75),

followed by a relative plateau through the section that subtends the temporal lobe, with a

trough as the anterior-posterior section is increasingly represented by the parietal lobe

(percentile 37). A higher plateau peak subtends the parietal lobe, and then gradually declines

as gradient transitions into the occipital lobe (beginning from around percentile 15).

220   The middle and lower rows of Figure 4 show that while gyrification decreases

221   globally, they are most pronounced in the parietal lobe. Further examination of the surface

222   topography can better differentiate gyrification in parietal and temporal regions. These aging

223   results demonstrate that the overall distribution of gyrification does not change with age, it

224   merely diminishes in magnitude.

**Figure 4. Gyrification index across posterior (percentile 0; caudal) to anterior
(percentile 100; rostral).** The upper row illustrates the average gyrification across the
common surface space. Coloured labels denoting each of the four lobes (frontal: red; parietal:
green; temporal: blue; occipital: orange), insula (purple), and central sulcus (yellow) are
included to aid in interpretation. An inflated cortical surface is shown on the right to help
visualise the relative extents of these regions, though a folded brain was used for the actual
analyses. The middle row shows the mean anterior-posterior gyrification for each age
between 50 and 80, inclusive; brighter colours denote regions of higher gyrification index.
This subset of ages was selected based on having sufficient sessions per age to use in the
estimation, as shown in the right panel. A total of 746 of the available 815 MRI sessions were
in this age range. The lower row shows the mean anterior-posterior gyrification for
individuals aged 50, 60, 70, and 80. Aging is associated with overall decreases in
gyrification, but these appear to be most pronounced in the parietal lobe.

Topography of gyrification

As gyrification is calculated as the ratio of areas of the cortical surface to an enclosing

surface, the gyrification index is highest at the insula—as shown Figure 5A. Here I found a

similar pattern in the present accelerated longitudinal dataset, as shown in Figure 5B. The

decline in gyrification is most pronounced in the parietal lobe and posterior aspects of the

frontal lobe. However, on individual cortical surfaces, even examining changes over nearly a

decade, changes in the surfaces are only barely visible (see Figure 6).

A. Mean Gyrification Index

B. Variance Explained by Age

250
251 **Figure 5. Topography of gyrification.** (A) Mean gyrification, with the highest values
252 corresponding to the insula. (B) Age changes in gyrification, with the highest values
253 corresponding to the parietal lobe and posterior aspects of the frontal lobe (i.e., primary
254 motor and somatosensory regions). Values determined based on vertex-wise regression of
255 local gyrification index with age.

Participant A [8-year interval]

57 years old
GI=2.993

60 years old
GI=2.979

63 years old
GI=2.955

65 years old
GI=2.936

Participant B [9-year interval]

83 years old
GI=2.721

87 years old
GI=2.688

92 years old
GI=2.644

256
257 **Figure 6. Longitudinal age changes in cortical folding.** Cortical surface reconstructions for
258 two participants, illustrating changes in folding over nearly a decade each. GI denotes
259 gyrification index. See Madan (2015) for details related to rendering the cortical surfaces.
260

Sulcal prominence

Figure 6 shows that the overall pattern of cortical folding remains relatively consistent,

despite small reductions in the gyrification index. Visible from the MRIs themselves,

however, does indicate changes in the sulcal width and depth. Here I examined how well

global measures of sulcal morphology, both width and depth, can alternatively explain

variability in gyrification. To test this, I conducted a multi-level mediation analysis on the

longitudinal global gyrification index measurements, with random intercepts for each

participant and based on all available timepoints.

269    Standardised parameter estimates for the model are shown in Figure 7, along with all

270 associated standard error (SE) estimates in parentheses. Age was significantly related to

271 increases in sulcal width [$a_1 = 0.463$ (0.028); $t(813)=16.41$, $p<.001$] as well as decreases in

272 sulcal depth [$a_2 = -0.362$ (0.022); $t(813)=16.76$, $p<.001$]. Gyrification index was significantly

273 related to both mediators, i.e., sulcal width and depth, and a significant direct effect of age

274 remained after the mediators were modelled [sulcal width: $b_1 = -0.105$ (0.029); $t(813)=3.69$,

275 $p<.001$; sulcal depth: $b_2 = 0.676$ (0.027); $t(813)=25.12$, $p<.001$; remaining direct effect of

276 age: $c' = -0.071$ (0.020); $t(813)=3.52$, $p<.001$].

277    Mediation analyses indicated that both sulcal width and depth mediated the effects of

278 age on gyrification index [sulcal width: $a_1b_1 = -0.049$ (0.014); Sobel's $Z=3.59$, $p<.001$; sulcal

279 depth: $a_2b_2 = -0.245$ (0.018); Sobel's $Z=13.94$, $p<.001$]. The entire mediation model

280 accounted for 49.9% of the variability in gyrification (i.e., R^2). Proportion-mediated effect

281 size estimates demonstrated that most of this variability was accounted for by sulcal depth

282 (36.2%), with lesser proportions being explained by sulcal width (7.2%) and the remaining

283 direct effect of age (10.6%). Note that these summed values slightly exceed the total amount

284 of variability explained, as sulcal width and depth are not orthogonal. These results indicate

285 that age related decreases in sulcal depth largely account for the apparent changes in

286 gyrification, and both sulcal width and depth are more sensitive to effects of aging than the

287 gyrification index measure.

288
289 **Figure 7. Path diagram for age effects on gyrification index, considering the mediators**
290 **of sulcal width and depth.** Standardised parameter estimates for the model are shown for
291 each effect, along with standard error (SE) estimates in parentheses.

Discussion

From the global gyrification and anterior-posterior analyses, it is clear that the age decreases

in gyrification are gradual. Moreover, the anterior-posterior analysis was designed to evaluate

if these age changes in gyrification were related to shifts in the underlying distribution of

cortical folding, but this does not appear to be the case. Instead, there is a global decrease,

with some regions more affected than others, as highlighted in the topography analyses.

Converging with prior cross-sectional studies, regions most affected by age related

gyrification changes are distinct from those regions affected by changes in cortical thickness.

While studies examining cross-sectional data suggest that global gyrification involves

a large degree of age-independent variability, the results of the longitudinal analyses here

show a relatively consistent age decline, distinct from differences in the y-intercept of the

gyrification index (i.e., Figure 3). Though changes in gyrification due to aging have not

previously been investigated using an anterior-posterior gradient approach, the overall

gradient (i.e., the upper row of Figure 4) is relatively consistent with Zilles et al. (1988, Fig.

6). The overall topography, as shown in Figure 5, appears consistent with previously

published results (e.g., Schaer et al., 2008; Cao et al., 2017; Lamballais et al., 2020). As

discussed in the Introduction section, previous studies have demonstrated age differences in

gyrification, but as of yet, this has only been in cross-sectional samples (Cao et al., 2017;

Hogstrom et al., 2013; Lamballais et al., 2020; Madan & Kensinger, 2016, 2018; Madan,

2018).

Despite numerous prior studies reporting decreases in gyrification with age, these

studies provide little towards explaining the underlying mechanism. While a mechanism is

not presented here either, the current results provide some insight into a more specific

characterisation of how aging influences cortical folding. Many prior studies of brain

morphology have indicated that cortical surface area is not affected by aging (e.g., Hogstrom

et al., 2013; McKay et al., 2014; Storsve et al., 2014). This lack of influence of age on

cortical surface area measurements also rules out any systematic changes in the minor

deformations in the folds along the gyri. More specifically, these minor folds could be

measured using other approaches, such as indices of the spatial frequency/power spectra of

cortical folding as providing information distinct from gyrification itself (as in Madan,

2019b). Age-related changes in cortical thickness (and volume) are often found, but follow

from a distinct topography than have been reported for gyrification (e.g., Hogstrom et al.,

2013; Madan & Kensinger, 2016, 2018; McKay et al., 2014). However, long before

neuroimagers began to examine cortical thickness as an index of cortical atrophy, radiologists

have qualitatively assessed sulcal prominence—alternatively referred to as widening,

enlargement, or dilation—as a key measure of age-related atrophy (Coffey et al., 1992;

Drayer, 1988; Huckman et al., 1975; Jacoby et al., 1980; Laffey et al., 1984; LeMay, 1984;

Pasquier et al., 1996; Roberts & Caird, 1976; Scheltens et al., 1997; Tomlinson et al., 1968;

Turkheimer et al., 1984; Yue et al., 1997). Indeed, by visual inspection, cortical atrophy is

much more readily assessed from sulcal features than from cortical thickness (see Figure 6).

Using current automated methods, sulcal morphology can also be quantitatively

measured, where the width and depth of major sulci can be identified and estimated (e.g.,

Kochunov et al., 2005; Madan, 2019a). Across a number of studies and samples (albeit

generally with much smaller samples), sulcal morphology has been reliably associated with

age-related differences (Jin et al., 2018; Kochunov et al., 2005; Li et al., 2011; Liu et al.,

2010, 2013; Madan, 2019a; Rettmann et al., 2006; Shen et al., 2018). The work presented

here provides additional specificity in how gyrification appears to decrease with age, a step

towards understanding the underlying mechanism. Here I show that gyrification changes can

be considered strongly related to changes in the underlying sulcal morphology, particularly

depth. However, despite this well-established relationship between sulcal morphology and

aging, the neurobiological mechanism underlying this change in the fundamental organisation

of cortical folding remains unclear and a topic for further study.

More generally, this work adds to the growing literature demonstrating that the

availability of open-access MRI data can facilitate advances in our understanding of brain

morphology well beyond the goals of the researchers that originally collected the data (see

Madan, 2017, for an overview of benefits and considerations). It is worth acknowledging that

the results presented here likely underestimate the extent that gyrification decreases with age.

Older adults that are interested and able to participate in a research study in their 80s and 90s

are very likely to be a biased sample of individuals for their age cohort, demonstrating better

physical and mental health than many of their peers (i.e., an issue of external validity; see

Pearl & Bareinboim, 2014, and Keyes & Westreich, 2019, for more nuanced discussions).

Abbreviations

355	CDR	Clinical Dementia Rating
356	FD	fractal dimensionality
357	fMRI	functional magnetic resonance imaging
358	GI	gyrification index
359	MMSE	Mini-Mental State Exam
360	MRI	magnetic resonance imaging
361	OASIS	Open Access Series of Imaging Studies

Conflict of interest

365 The authors have no conflict of interests to disclose.

Data accessibility

369 All raw data is available from https://www.oasis-brains.org.

Acknowledgments

I would like to thank Jonathan Reardon for feedback on an earlier version of this manuscript.

This work would not have been possible without data from OASIS-3 and the funding that

supported it (Principal Investigators: T. Benzinger, D. Marcus, J. Morris; NIH P50AG00561,

P30NS09857781, P01AG026276, P01AG003991, R01AG043434, UL1TR000448,

R01EB009352). I would also like to acknowledge the support of NVIDIA Corporation with

the donation of a Titan Xp GPU that was used in this research.

References

Cao, B., Mwangi, B., Passos, I.C., Wu, M.-J., Keser, Z., Zunta-Soares, G.B., Xu, D., Hasan, K.M. & Soares, J.C. (2017) Lifespan gyrification trajectories of human brain in healthy individuals and patients with major psychiatric disorders. *Sci. Rep.*, **7**, 511. doi:10.1038/s41598-017-00582-1

Carradus, A. J., Mougin, O., Hunt, B. A. E., Tewarie, P. K., Gaedas, N., Morris, P. G., Brookes, M. J., Gowland, P. A., & Madan, C. R. (2020) Age-related differences in myeloarchitecture at 7 T. *Neurobiol. Aging,* **96**, 246-254. doi:10.1016/j.neurobiolaging.2020.08.009

Coffey, C. E., Wilkinson, W. E., Parashos, L., Soady, S., Sullivan, R. J., Patterson, L. J., ... Djang, W. T. (1992) Quantitative cerebral anatomy of the aging human brain: A cross-sectional study using magnetic resonance imaging. *Neurology,* **42***,* 527–527. doi:10.1212/wnl.42.3.527372

Drayer, B. P. (1988) Imaging of the aging brain. Part I. normal findings. *Radiology*, **166**, 785–796. doi: 10.1148/radiology.166.3.3277247

Dickstein, D.L., Weaver, C.M., Luebke, J.I. & Hof, P.R. (2013) Dendritic spine changes associated with normal aging. *Neuroscience*, **251**, 21–32. doi:10.1016/j.neuroscience.2012.09.077

Duan, H., Wearne, S.L., Rocher, A.B., Macedo, A., Morrison, J.H. & Hof, P.R. (2003) Age-related dendritic and spine changes in corticocortically projecting neurons in macaque monkeys. *Cereb. Cortex*, **13**, 950–961.

Elias, H., & Schwartz, D. (1969) Surface areas of the cerebral cortex of mammals determined by sterological methods. *Science,* **166**(3901), 111-113. doi:10.1126/science.166.3901.111

Fischl, B. (2012) FreeSurfer. *NeuroImage*, *62*, 774–781. doi:10.1016/j.neuroimage.2012.01.021

Fischl, B., & Dale, A. M. (2000) Measuring the thickness of the human cerebral cortex from magnetic resonance images. *Proc. Natl. Acad. Sci. USA*, **97**, 11050–11055. doi:10.1073/pnas.200033797

Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., … Dale, A. M. (2002) Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron*, **33**, 341–355. doi:10.1016/s0896-6273(02)00569-x

Fischl, B., Sereno, M. I., Tootell, R. B. H., & Dale, A. M. (1999) High-resolution intersubject averaging and a coordinate system for the cortical surface. *Hum. Brain Mapp.,* **8**, 272–284. doi:10.1002/(sici)1097-0193(1999)8:4<272::aid-hbm10>3.0.co;2-4

Glasser, M. F., & Van Essen, D. C. (2011) Mapping human cortical areas in vivo based on myelin content as revealed by T1- and T2-weighted MRI. *J. Neurosci.,* **31**, 11597–11616. doi:10.1523/jneurosci.2180-11.2011

Grydeland, H., Walhovd, K. B., Tamnes, C. K., Westlye, L. T., & Fjell, A. M. (2013) Intracortical myelin links with performance variability across the human lifespan: Results from T1- and T2-weighted MRI myelin mapping and diffusion tensor imaging. *J. Neurosci.,* **33**, 18618–18630. doi:10.1523/jneurosci.2811-13.2013

Hao, J., Rapp, P.R., Janssen, W.G.M., Lou, W., Lasley, B.L., Hof, P.R. & Morrison, J.H. (2007) Interactive effects of age and estrogen on cognition and pyramidal neurons in monkey prefrontal cortex. *Proc. Natl. Acad. Sci. USA*, *104*, 11465–11470. doi:10.1073/pnas.0704757104

Hogstrom, L.J., Westlye, L.T., Walhovd, K.B. & Fjell, A.M. (2013) The structure of the cerebral cortex across adult life: Age-related patterns of surface area, thickness, and gyrification. *Cereb. Cortex*, **23**, 2521–2530. doi:10.1093/cercor/bhs231

Huckman, M. S., Fox, J., & Topel, J. (1975) The validity of criteria for the evaluation of cerebral atrophy by computed tomography. *Radiology,* **116**, 85–92. doi:10.1148/116.1.85

Jin, K., Zhang, T., Shaw, M., Sachdev, P., & Cherbuin, N. (2018) Relationship between sulcal characteristics and brain aging. *Front.Aging Neurosci.,* **10**, 339. doi:10.3389/fnagi.2018.00339

Keyes, K. M., & Westreich, D. (2019) UK Biobank, big data, and the consequences of non-representativeness. *Lancet,* **393***(10178)*, 1297. doi:10.1016/s0140-6736(18)33067-8

King, R. D., Brown, B., Hwang, M., Jeon, T., & George, A. T. (2010) Fractal dimension analysis of the cortical ribbon in mild Alzheimer's disease. *NeuroImage,* **53**, 471–479. doi:10.1016/j.neuroimage.2010.06.050

Laffey, P. A., Peyster, R. G., Nathan, R., Haskin, M. E., & McGinley, J. A. (1984) Computed tomography and aging: Results in a normal elderly population. *Neuroradiology,* **26**, 273–278. doi: 10.1007/BF00339770

Lamballais, S., Vinke, E. J., Vernooij, M. W., Ikram, M. A., & Muetzel, R. L. (2020) Cortical gyrification in relation to age and cognition in older adults. *NeuroImage*, **212**, 116637. doi:10.1016/j.neuroimage.2020.116637

LaMontagne, P. J., Benzinger, T. L. S., Morris, J. C., Keefe, S., Hornbeck, R., Xiong, C., … Marcus, D. (2019) OASIS-3: Longitudinal neuroimaging, clinical, and cognitive dataset for normal aging and Alzheimer's disease. *medRxiv*. doi:10.1101/2019.12.13.19014902

LeMay, M. (1984) Radiologic changes of the aging brain and skull. *Am. J. Roentgenol.,* **143**, 383–389. doi:10.2214/ajr.143.2.383

Li, S., Xia, M., Pu, F., Li, D., Fan, Y., Niu, H., … He, Y. (2011) Age-related changes in the surface morphology of the central sulcus. *NeuroImage,* **58**, 381–390. doi:10.1016/j.neuroimage.2011.06.041

Liu, T., Lipnicki, D. M., Zhu, W., Tao, D., Zhang, C., Cui, Y., Jin, J. S., Sachdev, P. S., & Wen, W. (2012) Cortical gyrification and sulcal spans in early stage Alzheimer's disease. *PLOS ONE,* **7**, e31083. doi:10.1371/journal.pone.0031083

Liu, T., Sachdev, P. S., Lipnicki, D. M., Jiang, J., Geng, G., Zhu, W., Reppermund, S., Tao, D., Trollor, J. N., Brodaty, H., & Wen, W. (2013) Limited relationships between two-year changes in sulcal morphology and other common neuroimaging indices in the elderly. *NeuroImage,* **83**, 12–17. doi:10.1016/j.neuroimage.2013.06.058

Liu, T., Wen, W., Zhu, W., Trollor, J., Reppermund, S., Crawford, J., Jin, J. S., Luo, S., Brodaty, H., & Sachdev, P. (2010) The effects of age and sex on cortical sulci in the elderly. *NeuroImage,* **51**, 19–27. doi:10.1016/j.neuroimage.2010.02.016

Madan, C. R. (2015) Creating 3D visualizations of MRI data: A brief guide. *F1000Research,* **4**, 466. doi:10.12688/f1000research.6838.1

Madan, C. R. (2017) Advances in studying brain morphology: The benefits of open-access data. *Front. Hum. Neurosci.,* **11**, 405. doi:10.3389/fnhum.2017.00405

Madan, C. R. (2018) Age differences in head motion and estimates of cortical morphology. *PeerJ,* **6**, e5176. doi:10.7717/peerj.5176

Madan, C. R. (2019a) Robust estimation of sulcal morphology. *Brain Informatics,* **6**, 5. doi:10.1186/s40708-019-0098-1

Madan, C. R. (2019b) Evaluating the angular power spectrum of cortical folding. *Proc. Cogn. Comput. Neurosci.,* **3**, 496-499. doi:10.32470/CCN.2019.1177-0

Madan, C. R., & Kensinger, E. A. (2016) Cortical complexity as a measure of age-related brain atrophy. *NeuroImage,* **134**, 617-629. doi:10.1016/j.neuroimage.2016.04.029

Madan, C. R., & Kensinger, E. A. (2017) Test-retest reliability of brain morphology estimates. *Brain Informatics,* **4**, 107-121. doi:10.1007/s40708-016-0060-4

Madan, C. R., & Kensinger, E. A. (2018) Predicting age from cortical structure across the
lifespan. *Eur. J. Neurosci.,* **47**, 399-416. doi:10.1111/ejn.13835
Mangeat, G., Govindarajan, S. T., Mainero, C., & Cohen-Adad, J. (2015) Multivariate
combination of magnetization transfer, T2* and B0 orientation to study the myelo-
architecture of the in vivo human cortex. *NeuroImage,* **119**, 89–102.
doi:10.1016/j.neuroimage.2015.06.033
Margulies, D. S., Ghosh, S. S., Goulas, A., Falkiewicz, M., Huntenburg, J. M., Langs, G.,
Bezgin, G., Eickhoff, S. B., Castellanos, F. X., Petrides, M., Jefferies, E., &
Smallwood, J. (2016) Situating the default-mode network along a principal gradient of
macroscale cortical organization. *Proc. Natl. Acad. Sci. USA,* **113**, 12574–12579.
doi:10.1073/pnas.1608282113
McDonough, I. M., & Madan, C. R. (in press) Structural complexity is negatively associated
with brain activity: A novel multimodal test of compensation theories of aging.
Neurobiol. Aging.
McKay, D. R., Knowles, E. E. M., Winkler, A. A. M., Sprooten, E., Kochunov, P., Olvera, R.
L., … Glahn, D. C. (2014) Influence of age, sex and genetic factors on the human
brain. *Brain Imag. Behav.,* **8**, 143–152. doi:10.1007/s11682-013-9277-5
Murphy, C., Wang, H.-T., Konu, D., Lowndes, R., Margulies, D. S., Jefferies, E., &
Smallwood, J. (2019) Modes of operation: A topographic neural gradient supporting
stimulus dependent and independent cognition. *NeuroImage,* **186**, 487–496.
doi:10.1016/j.neuroimage.2018.11.009
Palaniyappan, L. & Liddle, P.F. (2012) Differential effects of surface area, gyrification and
cortical thickness on voxel based morphometric deficits in schizophrenia. *NeuroImage*,
60, 693–699. doi:10.1016/j.neuroimage.2011.12.058
Palaniyappan, L., Park, B., Balain, V., Dangi, R., & Liddle, P. (2015) Abnormalities in
structural covariance of cortical gyrification in schizophrenia. *Brain Struct. Funct.,* **220**,
2059–2071. https://doi.org/10.1007/s00429-014-0772-2
Pasquier, F., Leys, D., Weerts, J. G., Mounier-Vehier, F., Barkhof, F., & Scheltens, P. (1996)
Inter-and intraobserver reproducibility of cerebral atrophy assessment on MRI scans
with hemispheric infarcts. *Eur. Neurol.,* **36**, 268–272. doi:10.1159/000117270571
Pearl, J., & Bareinboim, E. (2014) External validity: From do-calculus to transportability
across populations. *Stat. Sci.*, **29**, 579-595. doi:10.1214/14-STS486
Peters, A. (2002) Structural changes that occur during normal aging of primate cerebral
hemispheres. *Neurosci. Biobehavl. Rev.*, *26*, 733–741. doi: 10.1016/s0149-
7634(02)00060-x
Rettmann, M. E., Kraut, M. A., Prince, J. L., & Resnick, S. M. (2006) Cross-sectional and
longitudinal analyses of anatomical sulcal changes associated with aging. *Cereb.
Cortex,* **16**, 1584–1594. doi:10.1093/cercor/bhj095
Roberts, M. A., & Caird, F. I. (1976) Computerised tomography and intellectual impairment
in the elderly. *J. Neurol. Neurosurg. Psychiatry,* **39**, 986–989.
doi:10.1136/jnnp.39.10.986
Salat, D.H., Buckner, R.L., Snyder, A.Z., Greve, D.N., Desikan, R.S.R., Busa, E., Morris,
J.C., Dale, A.M. & Fischl, B. (2004) Thinning of the cerebral cortex in aging. *Cereb.
Cortex*, **14**, 721–730. doi:10.1093/cercor/bhh032
Schaer, M., Cuadra, M. B., Tamarit, L., Lazeyras, F., Eliez, S., & Thiran, J.-P. (2008) A
surface-based approach to quantify local cortical gyrification. *IEEE Trans. Med.
Imaging,* **27**, 161–170. doi:10.1109/tmi.2007.903576
Schaer, M., Cuadra, M.B., Schmansky, N., Fischl, B., Thiran, J.-P. & Eliez, S. (2012) How to
measure cortical folding from MR images: A step-by-step tutorial to compute local
gyrification index. *J. Vis. Exp.,* **59***, e3417. doi:10.3791/3417

Schaer, M., Ottet, M.-C., Scariati, E., Dukes, D., Franchini, M., Eliez, S., & Glaser, B. (2013)
Decreased frontal gyrification correlates with altered connectivity in children with
autism. *Front. Hum. Neurosci., 7*, 750. doi:10.3389/fnhum.2013.00750
Scheltens, P., Pasquier, F., Weerts, J. G., Barkhof, F., & Leys, D. (1997) Qualitative
assessment of cerebral atrophy on MRI: Inter- and intra-observer reproducibility in
dementia and normal aging. *Eur. Neurol., 37*, 95–99. doi:10.1159/000117417
Shafee, R., Buckner, R. L., & Fischl, B. (2015) Gray matter myelination of 1555 human
brains using partial volume corrected MRI images. *NeuroImage, 105*, 473–485.
doi:10.1016/j.neuroimage.2014.10.054
Shen, X., Liu, T., Tao, D., Fan, Y., Zhang, J., Li, S., … Wen, W. (2018) Variation in
longitudinal trajectories of cortical sulci in normal elderly. *NeuroImage, 166*, 1–9.
doi:10.1016/j.neuroimage.2017.10.010
Sormaz, M., Murphy, C., Wang, H., Hymers, M., Karapanagiotidis, T., Poerio, G., Margulies,
D. S., Jefferies, E., & Smallwood, J. (2018) Default mode network can support the level
of detail in experience during active task states. *Proc. Natl. Acad. Sci. USA, 115*, 9318–
9323. doi:10.1073/pnas.1721259115
Sowell, E.R., Peterson, B.S., Thompson, P.M., Welcome, S.E., Henkenius, A.L. & Toga,
A.W. (2003) Mapping cortical change across the human life span. *Nat. Neurosci., 6*,
309–315. doi:10.1038/nn1008
Storsve, A. B., Fjell, A. M., Tamnes, C. K., Westlye, L. T., Overbye, K., Aasland, H. W., &
Walhovd, K. B. (2014) Differential longitudinal changes in cortical thickness, surface
area and volume across the adult life span: Regions of accelerating and decelerating
change. *J. Neurosci., 34*, 8488–8498. doi:10.1523/jneurosci.0391-14.2014
Tomlinson, B., Blessed, G., & Roth, M. (1968) Observations on the brains of non-demented
old people. *J. Neurol. Sci., 7*, 331–356. doi:10.1016/0022-510x(68)90154-8
Turkheimer, E., Cullum, C. M., Hubler, D. W., Paver, S. W., Yeo, R. A., & Bigler, E. D.
(1984) Quantifying cortical atrophy. *J. Neurol. Neurosurg. Psychiatry, 47,* 1314–1318.
doi:10.1136/jnnp.47.12.1314
Yue, N. C., Arnold, A. M., Longstreth, W. T., Elster, A. D., Jungreis, C. A., O'Leary, D. H.,
... Bryan, R. N. (1997) Sulcal, ventricular, and white matter changes at MR imaging in
the aging brain: Data from the cardiovascular health study. *Radiology, 202*, 33–39. doi:
10.1148/radiology.202.1.8988189
Zilles, K., Armstrong, E., Schleicher, A. & Kretschmann, H.-J. (1988) The human pattern of
gyrification in the cerebral cortex. *Anat. Embryol., 179*, 173–179.
doi:10.1007/BF00304699
562
563

Appendix: Longitudinal changes in global cortical fractal dimensionality

Madan and Kensinger (2016) demonstrated that fractal dimensionality can be a more

sensitive measure of age-related differences in cortical structure than conventional measures,

including gyrification. To provide a complementary analysis, although tangential to the focus

of the current study, here I present the age-related longitudinal changes in fractal

dimensionality, from the same data as in the other analyses. To-date, longitudinal analyses of

fractal dimensionality have not yet been conducted, and these results serve as an initial

comparison between the gyrification results (e.g., participants in each panel are in the same

line colour as in Figure 3). As expected from prior work, Figure A1 shows that there is less

off-axis variability (i.e., random y-intercepts) with fractal dimensionality than are

consistently found with gyrification. As with gyrification, a linear mixed effects model with

random slopes for each participant was calculated, with a significant age fixed effect

[$t(813)=25.55$, $p<.001$] with a decreasing slope of 0.01325 FD/decade [95% C.I. = 0.01223—

0.01426].

As fractal dimensionality is a summary statistic of the complexity of a structure, a

topographical analysis cannot be equivalently carried out as for gyrification—though fractal

dimensionality can be calculated for parcellated cortical regions.

582
583 **Figure A1. Age changes in global cortical fractal dimensionality.** Each coloured line
584 represents an individual participant, with each dot corresponding to an MRI session. The plot
585 has been divided into panels for each set of timepoints to improve readability. Dashed line
586 present in all panels shows the decrease in global fractal dimensionality based on all data,
587 estimated using the linear mixed effects model. Compare with Figure 3.