Dynamics of microbial competition, commensalism and cooperation and its implications for coculture and microbiome engineering

Peng Xu*

Department of Chemical, Biochemical and Environmental Engineering, University of Maryland

Baltimore County, Baltimore, MD 21250

* Corresponding author, Tel: +1(410)-455-2474; fax: +1(410)-455-1049. E-mail address:

pengxu@umbc.edu (PX)

Abstract:

Microbial consortium is a complex adaptive system with higher order dynamic characteristics that are not present by individual members. To accurately predict the social interactions, we formulate a set of unified unstructural kinetic models to quantitatively understand the dynamic interactions of multiple microbial species. By introducing an interaction coefficient to the growth fitness function, we analytically derived the steady state solutions for the interacting species and the substrate profile in the chemostat. We analyzed the stability of the possible co-existing states defined by competition, parasitism (predation), amensalism, commensalism and cooperation. Our model predicts that only parasitism, commensalism and cooperation could lead to stable co-existing state. We also analyzed the design constraints and operational conditions of microbial coculture with sequential metabolic reactions compartmentalized into two distinct species. Coupled with Luedeking–Piret and Michaelis-Menten equation, accumulation of metabolic precursors in one species and formation of end-product in another species could be derived and assessed. We discovered that parasitism consortia disfavor the bioconversion of intermediate to final product; and commensalism consortia could efficiently

convert metabolic intermediate to final product and maintain metabolic homeostasis with a broad range of operational conditions (i.e., dilution rates); whereas cooperative consortia leads to highly nonlinear pattern of precursor accumulation and end-product formation. The underlying dynamics and emergent properties of microbial consortia may provide critical knowledge for us to engineer efficient bioconversion process, deliver effective gut therapeutics as well as elucidate probiotic-pathogen interactions in general.

Key words: Microbial consortia, competition, commensalism, cooperation, modeling, dynamic analysis

1. Introduction

Microbes in nature form diverse social interactions and dynamically respond to metabolic and environmental variations at community level. The interacting species in a microbial community might compete for the same resource, exchange for metabolites, communicate each other *via* metabolic or genetic signals [1]. The unique interactions in a microbial community define the collective biological functions that are robust to harsh conditions than individual cells could sustain [2, 3]. Compared to monocultures, co-cultures exhibit a number of advantages, including division of labor, compartmentalization of reaction, and robustness to perturbations *et al* [4, 5]. Recently, microbial coculture or consortia has been increasingly applied to configurate different sections of metabolic pathways with improved catalytic performance. The co-cultivation of various species has led to the production of important biofuels [6], traditional food [7] and nutraceuticals [8-10]. Novel synthetic biology tools, including cell signaling translator [11] and transcriptional factor-based biosensors [12, 13] have been recently developed to autonomously regulate culture composition and eliminate

metabolic heterogeneity. In particular, microbial social interaction could define unique spatial patterns that are important for us to fabricate advanced biomaterials [14, 15], understand biofilm formation [16] and disarm antibiotic resistant superbugs [17].

Kinetic models have been increasingly important to understand microbial social interactions at the consortia-level [18-20]. Most of these kinetic equations are developed by correlating cell growth fitness with the nutrient or environmental conditions of the interacting species. In particular, the canonical Monod equation, which describes the quantitative relation between cell growth and a limiting nutrient [21], has been widely adopted and revised to incorporate multiple inhibitory terms for describing cell, substrate, product inhibition *et al* [22-24]. At ecological level, biochemical engineers have formulated a set of coupled Monod equations to describe the predator-prey relationship between *Dictyostelium discoideum* and *Escherichia coli* in Chemostat [25], where an oscillatory relationship between the two populations were predicted and experimentally validated. On the other hand, ecologists preferred to use Logistic equation due to the simplicity, existence of analytical solution and the rich dynamics of the Logistic model. For example, the popular Lotka–Volterra predator-prey model was developed and analyzed as early as 1926 by biophysicists to understand dynamic species interaction in a closed ecological niche [26, 27]. The intriguing solutions of the discrete Logistic model were discovered to have complex dynamic behaviors, ranging from stable points, to bifurcating stable cycles, to chaotic fluctuations, depending on the initial parameter conditions [28]. Both Monod and Logistic equations benefit us to quantitatively understand microbial interaction kinetics [29].

Mathematical models of microbial consortia [30] has been studied and analyzed in 1980s, which lay the foundation for us to quantitatively define microbial social interactions. However, the utilization of microbial consortia is significantly hindered by our limited understanding of the emerging dynamics arising from the interacting species [20]. Here we developed a set of unified unstructural kinetic models to quantitatively understand the dynamic interactions of multiple microbial species. We analytically derived the steady state solutions for the two interacting species and the substrate profile in the chemostat. By analyzing the stability of the possible co-existing states, we can divide microbial social interaction into eight categories: competition, coexisting parasitism, extinctive parasitism, cooperation, bistable amensalism, extinctive amensalism, coexisting commensalism and extinctive commensalism. By analyzing the solutions for microbial consortia with sequential metabolic reactions compartmentalized into distinct species, we revealed the design criteria of microbial coculture engineering in chemostat. We discovered that commensalism consortia could efficiently convert metabolic intermediate to final product and maintain metabolic homeostasis (i.e., constant final product formation) with a broad range of operational conditions (dilution rates). The simplicity and the rich dynamics of the consortia model highlight the importance to incorporate social interaction parameters into the unstructural kinetic models. The dynamics of microbial competition and cooperation may facilitate us to assemble diverse microbial species with defined social interactions for important biotechnological and biomedical applications.

2. Computational methods

2.1 Matlab computation environment

Matlab R2017b was used as the computational platform and installed on a Windows 7 professional operation system with Intel Core i3-6100 CPU processor at speed of 3.70 GHz. The installed memory (RAM) is 4.0 GHz. Matlab symbolic language package coupled with LaTex makeup language is used to derive and output the symbolic equations and solutions (Supplementary files). Analytical solutions were derived for Fig. 2 and Fig. 4 with Matlab Symbolic Language. All trajectories in Fig. 2 and Fig. 4, and phase portraits in Fig. 3 were computed by numerical ODE45 solvers. Numerical solutions for the compartmentalized sequential metabolic reactions in Fig. 5 and Fig. 6 were computed by numerical ODE45 solver. Matlab code has been compiled into the supplementary file.

2.2 Stability analysis of microbial social interactions

For the generalized systems ODE equations listed in Eqn. 3, Eqn. 4 and Eqn. 5, we have computed the parameterized Jacobian matrix with Matlab symbolic language. This parameterized Jacobina matrix represents all the nine social interactions summarized in Fig. 1. One can simply substitute specific numbers (-1, 0, or 1) into the interaction coefficients (γ_{AB} or γ_{BA}) in this generalized Jacobian matrix to retrieve the detailed matrix. Then specific eigenvalues cold be computed to evaluate the stability criteria of each steady states. A complete long form of the Jacobian Matrix is provided as supplementary equations. Some representative eigenvalues for specific parameter conditions are compiled to the supplementary table. Here is the generalized Jacobian Matrix that corresponds to systems ODEs Eqn.3 to Eqn. 5.

$$J = \begin{pmatrix} A - D & C & AF \\ E & B - D & BG \\ -\dfrac{A}{Y_{AS}} - \dfrac{E}{Y_{BS}} & -\dfrac{B}{Y_{BS}} - \dfrac{C}{Y_{AS}} & -\dfrac{AF}{Y_{AS}} - \dfrac{BG}{Y_{BS}} - D \end{pmatrix}$$

Where *A*, *B*, *C*, *D*, *E*, *F* and *G* represents

$$A = \frac{S\,\mu_{\text{Amax}}}{K_{\text{SA}} + S}\left(\frac{X_B\,\gamma_{\text{BA}}}{S_0\,Y_{\text{BS}}} + 1\right)$$

$$B = \frac{S\,\mu_{\text{Bmax}}}{K_{\text{SB}} + S}\left(\frac{X_A\,\gamma_{\text{AB}}}{S_0\,Y_{\text{AS}}} + 1\right)$$

$$C = \frac{S\,X_A\,\gamma_{\text{BA}}\,\mu_{\text{Amax}}}{S_0\,Y_{\text{BS}}\,(K_{\text{SA}} + S)}$$

$$E = \frac{S\,X_B\,\gamma_{\text{AB}}\,\mu_{\text{Bmax}}}{S_0\,Y_{\text{AS}}\,(K_{\text{SB}} + S)}$$

$$F = \frac{K_{\text{SA}}\,X_A}{(K_{\text{SA}} + S)\,S}$$

$$G = \frac{K_{\text{SB}}\,X_B}{(K_{\text{SB}} + S)\,S}$$

2.3 Parameter settings

All the solutions were derived either analytically or numerically with the following parameter settings: μ_{Amax}= 1.6 (1/h); μ_{Bmax} = 1.2 (1/h); K_{SA} = 1.0 (g/L); K_{SB} = 0.8 (g/L); S_0 = 50 (g/L); Y_{AS} = 0.5 (g/g); Y_{BS}= 0.8 (g/g); Y_{BA} = 0.8 (g/g); Y_{PS} = 0.4 (g/g); α = 0.5 (dimensionless); β = 0.5 (1/h); γ_{AB} = -1, 0, or 1 (dimensionless), depending on the social interactions as specified in Fig. 1; γ_{BA} = -1, 0, or 1 (dimensionless), depending on the social interactions as specified in Fig. 1; k = 0.8 (1/h); K_m= 1.0 (g/L). Dilution rate *D* could be varied from 0 to 1.8 (1/h). All biophysical parameters were taken from biochemical parameter database (BioNumbers) or biochemical engineering textbooks (Shuler & Kargi, Clark & Blanch, Bailey *et al.*) with a physiologically interpretable range. It should be noted that, we assume that species A has a larger growth fitness (maximal specific growth rate) than species B in all our simulations.

3. Results and discussions

3.1 A unified unstructural kinetic model to define microbial social interactions

Based on the beneficial or detrimental relationship between two species, we can define six social interactions (**Fig. 1**): competition, amensalism, parasitism, neutralism, commensalism and mutualism (cooperation). To dynamically understand the nature of these social interactions, we modified the growth fitness function of the interacting species (**Eqn.1** and **Eqn. 2**), and introduced an interaction coefficient (γ_{AB} or γ_{BA}) to describe the species-level population effect. An interaction factor ($\gamma_{BA} \frac{X_B}{S_0 Y_{BS}}$ or $\gamma_{AB} \frac{X_A}{S_0 Y_{AS}}$), which is defined as the product of the interaction coefficient and relative population of the interacting species, is incorporated to the growth fitness equation as specified by Eqn.1 and Eqn.2.

The benefits of introducing this interaction coefficient are multifold. The growth fitness equation (**Eqn.1** or **Eqn. 2**) converges to the canonical Monod equation, when the population of the interacting species is negligible ($X_A \rightarrow 0$ or $X_B \rightarrow 0$). As the population of the interacting species approaches to the capacity of the system ($X_A \rightarrow S_0 Y_{AS}$ or $X_B \rightarrow S_0 Y_{BS}$), the interaction factor ($\gamma_{BA} \frac{X_B}{S_0 Y_{BS}}$ or $\gamma_{AB} \frac{X_A}{S_0 Y_{AS}}$) reaches its maximal or minimal value (γ_{BA} or γ_{AB}) that corresponds to either beneficial, neutral or detrimental effects, depending on the sign (+1, 0, or -1) of the interaction coefficient (γ_{AB} or γ_{BA}). As the cell grows, the population starts having beneficial or detrimental effect on the interacting species; the strength of this interaction is proportional to the relative ratio of the cell populations. The sign (negative or positive) of the interaction coefficient (γ_{AB} or γ_{BA}) determines the nature (detrimental or beneficial) of the social interactions (**Fig. 1**). With the simplified microbial consortia models (Eqn. 1 to Eqn. 8), we will derive the analytical solutions and evaluate the stability of the co-existing solutions. The insight obtained from this model will be critical for us to understand the dynamic nature of microbial species interaction and the importance to maintain microbial homeostasis, which may

facilitate us to design efficient microbial process for metabolite production or gut therapeutics for human health applications.

3.2 Dynamics of microbial competition, parasitism and cooperation with resource limitations

We next sought to understand the dynamics of microbial consortia with strong interactions, namely competition, parasitism and cooperation. One important question we would like to answer is whether the two interacting species could stably co-exist. This is critical, because there would be no microbial consortia if the two species could not exist together. With Matlab Symbolic computation package, we analytically derived the steady state solutions for Eqns. 1 to Eqns. 5 (a supplementary Matlab code was provided as SI file). By varying the interaction coefficients (γ_{AB} or γ_{BA}), we could use the set of parameterized equations (Eqn. 1 to Eqn. 5) to describe the various social interactions of the microbial consortia.

When species A and species B mutually exclude each other and compete for the same substrate, this condition ($\gamma_{AB} = -1$ and $\gamma_{BA} = -1$) will lead to an unstable co-exist state (**Fig. 2a**), as evidenced that one of the eigenvalues of Jacobian Matrix is positive (Supplementary Eigenvalue tables). Indeed, any perturbation from the co-exist state (light orange line in **Fig. 2a**) will lead to a single species survival state (**Fig. 3a**), which excludes the presence of the other species. The two species mutually exclude each other, therefor they could not exist together. For example, as we increase the dilution rate (D = 0.2 for red trajectory, D = 0.6 for green trajectory and D = 1.0 for blue trajectory in **Fig. 2a**), the unstable coexisting state will shift toward either species A or species B existing state (**Fig.2a** and **Fig. 3a**). The steady state solution of species A or B decreases as we increase the dilution rate. This trend is consistent with the population concentration as observed when a single species is cultivated in the chemostat [29].

When species A benefits species B, but species B harms species A, this condition ($\gamma_{AB} = +1 \; and \; \gamma_{BA} = -1$) will lead to a coexisting parasitism regime (**Fig. 2b**). At relatively low dilution rate ($D < 1.185$ h^{-1}) , species B will exist alone regardless of the initial concentration of species A (**Fig. 2b** and **Fig. 3b**). For example, the trajectory of the solution will move to species B alone state, as we increase the dilution rate (D = 0.6 for green trajectory and D = 1.0 for blue trajectory in **Fig. 2a**). Interestingly, at relatively large dilution rate ($D \geq 1.185$ h^{-1}), the system will move to a parasitism coexisting state (dark orange line in **Fig. 2b**). Since this dilution rate is approaching to the maximal growth rate of species B (μ_{Bmax} = 1.2 h^{-1}), species B must depend on the existence of species A to survive: species A benefits the growth of species B. As a result, we observed an oscillatory trajectory (a limited cycle, **Fig. 2b** and **Fig. 3b**) where species A and species B eventually lead to a balanced distribution (D = 1.365), as specified by the dark orange line in **Fig. 2b**. On the contrary, if we flip the sign of the interaction coefficient ($\gamma_{AB} = -1 \; and \; \gamma_{BA} = +1$), the system will lead to an extinctive parasitism: where species A will always outcompete the growth of species B (**Fig. 2c** and **Fig. 3c**), due to the fact that species A has a large growth fitness than species B. This analysis indicates that parasitism interaction could either lead to co-existing or extinction sates. The exploitative relation between species A and species B may reach a coexisting state, only if the dilution rate is harmful to the exploiter (i.e., species B is an exploiter in Fig. **2b**).

When species A and B mutually benefit each other, this condition ($\gamma_{AB} = +1 \; and \; \gamma_{BA} = +1$) will lead to a cooperative state (**Fig. 2d**). A stable coexisting state (dark orange line) is possible due to the mutualistic interactions between species A and B. Interestingly, when the dilution rate is approaching to the maximal growth rate of species A (μ_{Amax} = 1.6 h^{-1}), the

system will move to a unstable co-existing state (the light orange line in Fig. **2d**). Any infinitesimal perturbation from this unstable coexisting solution (light orange line) will move the system to an extinction state (i.e., solution falls to the origin, pink trajectory for $D = 1.7$) or to the stable co-existing state (i.e., solution falls to the dark orange line, blue trajectory for $D =$ 1.7). This intriguing behavior is also exemplified in the 2-D phase portrait as D is above 1.58 (Fig. **3d**). This bistability at large dilution rate ($D > 1.58$) suggests that the relative population of species A and B is critical to maintain the co-existing state. Obviously, the cooperation allows the system to operate at a dilution rate that is larger than both species could sustain, indicating the robust nature of the mutualistic interactions.

3.3 Dynamics of microbial commensalism and amensalism with resource limitations

We next sought to understand the dynamics of microbial consortia with weak interactions, namely amensalism and commensalism. Under these conditions, we are also interested in understanding whether the two interacting species could possibly co-exist. Following the same stability criteria, we evaluated the eigenvalues of Jacobian matrix. According to the system dynamics behavior, we could categorize the weak interactions into four groups: bistable amensalism (**Fig. 4a**), extinctive amensalism (**Fig. 4b**), coexisting commensalism (**Fig. 4c**) and extinctive commensalism (**Fig. 4d**).

When species B is harmful to species A, but species A is neutral to species B, the system ($\gamma_{AB} = 0 \ and \ \gamma_{BA} = -1$) will lead to an unstable coexisting state (light orange line of **Fig. 4a**). For example, any perturbation from this coexisting state will lead to the survival of a single species (either species A or species B). As we increase the dilution rate ($D = 0.2$ for red trajectory, $D = 0.8$ for green trajectory and $D = 1.0$ for blue trajectory in **Fig. 4a**), the unstable

coexisting state will shift toward either species A or species B existing state (**Fig. 4a**). We named this as "Bistable amensalism". On the contrary, an extinctive amensalism will arise when species A is harmful to species B, but species B is neutral to A ($\gamma_{AB} = -1 \; and \; \gamma_{BA} = 0$). Under any dilution condition, species A will outcompete species B to exist alone in the system (**Fig. 4b**).

When species A benefits species B, but species B is neutral to species A, the system ($\gamma_{AB} = +1 \; and \; \gamma_{BA} = 0$) will lead to a stable coexisting state (dark orange line of **Fig. 4c**). For example, the trajectory of any initial conditions will move the system toward the co-existing state (the dark orange line of **Fig. 4c**). Under this condition, the species with larger growth fitness (i.e., species A) will promote the growth of the species with smaller growth fitness (i.e., species B). On the contrary, an extinctive commensalism will arise if species A is neutral to species B, but species B benefits species A ($\gamma_{AB} = 0 \; and \; \gamma_{BA} = +1$). Under any dilution condition, species A will outcompete species B to exist alone in the system (**Fig. 4d**).

In summary, the social interactions of two species with resource limitations could result in rich dynamics ranging from bistable competition (**Fig. 2a**), coexisting parasitism (**Fig. 2b**), extinctive parasitism (**Fig. 2c**), cooperation (**Fig. 2d**), bistable amensalism (**Fig. 4a**), extinctive amensalism (**Fig. 4b**), coexisting commensalism (**Fig. 4c**) and extinctive commensalism (**Fig. 4d**). This complex interaction was captured by a set of unstructural kinetic models developed in this work, simply with the introduction of the interaction coefficients ($\gamma_{AB} \; or \; \gamma_{BA}$) to the classical Monod equations. The parameterized system equations (**Eqn.1** to **Eqn. 5**) provides a quantitative appraoch to analyze the system dynamics, which may help us design more efficient consortia and leverage coculture engineering for biotechnological and biomedical applications.

It should be noted that neutralism (where two species grow independently) was not explored in this study, but neutralism was also captured in our model ($\gamma_{AB} = 0 \ and \ \gamma_{BA} = 0$, **Fig. 1**)

3.4 Implications for microbial co-culture engineering and microbiome engineering

Co-cultures or microbial consortia are emerging strategies to improve metabolic pathway efficiency. They exhibit a number of advantages over monoculture, including division of labor, compartmentalization of reaction, and robustness to perturbations [4, 8, 9]. Microbial consortia define the functional assembly and social interaction of multiple species. To design efficient bioconversion process, one must consider the social interactions of the individual members in the microbial community. In this section, we will determine the optimal operation conditions for microbial coculture when a 2-step sequential metabolic reaction is compartmentalized into two co-cultivating species. Namely, intermediate A (P_A) is secreted from species A and was later converted to final product B (P_B) by species B. Two additional equations (**Eqn. 6** and **Eqn. 7**) and a modified substrate consumption equation (**Eqn. 8**) were introduced to describe the system dynamics. We assume there is no bottleneck of metabolites transportation across the two species and the consortia is devoid of metabolic burden due to accumulation of intermediate or final product. The formation rate for intermediate A (P_A) was assumed to follow the Luedeking-Piret equation [31], and the formation rate for product B (P_A) should follow a Michaelis-Mention type kinetics with the turnover rate constant proportional to the concentration of species B. With biomass yield ($Y_{AS} \ and \ Y_{BS}$), intermediate A yield from substrate (Y_{PS}), and product B yield from intermediate A (Y_{BA}), we can derive the mass balance equation for substrate consumption as specified by **Eqn. 8.**

To simplify the discussion, we focus on the steady state solutions of coexisting parasitism (**Fig. 5a** and **Fig. 6a**), extinctive parasitism (**Fig. 5b** and **Fig. 6b**), coexisting commensalism (**Fig. 5c** and **Fig. 6c**), and cooperation (**Fig. 5d** and **Fig. 6d**). The distribution of the steady state solution with varying dilution rate at 3-D space is presented in **Fig. 5**, and the exact steady state solution with varying dilution rate at a 2-D panel is presented in **Fig. 6**.

The solution distribution for biomass (X_A and X_B) and metabolite (P_A and P_B) displays highly nonlinear behavior (**Fig. 5** and **Fig. 6**), especially for the case when the two species coexist. Consistent with the species distribution (**Fig. 2b**), coexisting parasitism ($\gamma_{AB} = +1 \; and \; \gamma_{BA} = -1$) allow species B to exist alone and species A is suppressed by species B at low dilution rate ($D < 1.185$ h^{-1}). Therefore, intermediate A (P_A) was maintained at minimal value (~0) and there is no B produced at low dilution rate (**Fig. 5a** and **Fig. 6a**). When the dilution rate reaches a critical value ($D \geq 1.185$ h^{-1}) in proximate to the maximal growth rate of species B ($\mu_{Bmax} = 1.2$ h^{-1}), species B must rely on the beneficial effects (public goods or welfare) of species A to survive (**Fig. 5a** and **Fig. 6a**). With such disadvantageous conditions for species B, species A will be encouraged to proliferate, and the two species will gradually reach a balanced distribution with coexisting. Under high dilution rate ($D \geq 1.185$ h^{-1}), intermediate A (P_A) secreted from species A will be converted to product B (P_B) by species B (**Fig. 5a** and **Fig. 6a**). As predicted in the simulation, we can arrive an optimal dilution rate to maximize the product B formation (P_B), despite the fact that there is a tipping (discontinuous) point for species A (X_A) and intermediate A (P_A) (**Fig. 6a**) as we increase the dilution rate. This optimal dilution rate could be analytically derived should we have enough computational power. On the contrary, an extinctive parasitism ($\gamma_{AB} = -1 \; and \; \gamma_{BA} = +1$) will lead to the existence of species A, hence

the accumulation of the intermediate A (P_A) but without formation of product B (**Fig. 5b** and **Fig. 6b**). The solution will eventually fall to the origin (washout state) when we further increase the dilution rate for all the scenario discussed here (**Fig. 5** and **Fig. 6**). This analysis indicates that a parasitism relation may allow the compartmentalization of two sequential metabolic reactions in two distinct species, and there exists an optimal dilution rate to maximize the metabolite production (P_B). In reality, this metabolite P_B might be related with some signaling molecules that are associated with antibiotic resistance in biofilm formation or a metabolic intermediate that is associated with the dysbiosis of gut microbes in living organisms.

The most complicated dynamics were displayed when the two species are commensal (**Fig. 5c**) or cooperative (**Fig. 5d**). In both scenario, coexisting is possible and product B will be formed when the two species harbor distinct sections of metabolic reactions (**Fig. 6c** and **Fig. 6d**). In particular, intermediate A (P_A) secreted from species A will be efficiently converted to product B (P_B) when the two specie form a commensalism consortium (**Fig. 5c** and **Fig. 6c**). As a result, the intermediate A (P_A) was almost kept at minimal level with a very large window of operational conditions (i.e, 0.16 < D < 1.32), as evidenced that there is almost constant product B (P_B) formed in the system (**Fig. 6c**). We can also arrive an optimal dilution rate to maximize biomass for species B (**Fig. 6c**). In addition, the simulation indicates drastic changes of species A biomass (X_A) and intermediate (P_A) at a tipping point when the dilution rate is about 1.5 h^{-1}. Beyond this tipping point, the system rapidly falls to the washout states. (**Fig. 5c** and **Fig. 6c**). As a comparison, cooperation between species A and B instead leads to the accumulation of intermediate A (P_A in **Fig. 5d** and **Fig. 6d**), which possibly due to the mutualistic beneficial interaction between species A and species B: increase in the biomass of species A, which is the

source for intermediate P_A, will benefit the fitness of species B, which is the sink for intermediate P_A. Under this scenario, the activity of the metabolic source pathway and metabolic sink pathway is proportionally increased. Due to the intrinsic parameter settings (i.e., species A has a larger fitness than species B), intermediate A (P_A) from the metabolic source strain might not be efficiently converted to product B by the metabolic sink strain. This might possibly explain the inefficient conversion of intermediate A (P_A) to product B (P_B). Biomass (X_A) and intermediate (P_A) in species A will keep increasing before the system reaches the tipping dilution rate ($D \approx 1.5$ h^{-1}) (**Fig. 6d**). Product B will keep decreasing within the operational dilution window (0 h^{-1} < D < 1.6 h^{-1}). This analysis indicates that designing a cooperative consortium will be more challenging than designing a commensal consortium due to the bidirectional interaction of the two species. Importantly, commensal consortia allow the stable existence of two species and the efficient conversion of metabolic intermediate (P_A) to final product (P_B). In particular, metabolite concentration (P_B) was almost kept constant at a very large dilution window (i.e, 0.16 < D < 1.32), which may explain the phenomenon why most of the species in gut microbiota maintain a commensal consortium. Equivalently to say, this commensal consortium is critical to maintain metabolite homeostasis (i.e. constant P_B), which could resist large perturbations of environmental condition change (i.e., the dilution rate discussed in this study).

4. Conclusions

Microbial consortium is a complex adaptive system with higher order dynamic characteristics that are not present by individual members. To accurately predict the social interactions, we formulate a set of unified unstructural kinetic models to quantitatively

understand the dynamic interactions of multiple microbial species. With the generalized social interaction model (**Eqn.1** to **Eqn. 5**), we analytically derived the steady state solutions for the two interacting species and the substrate in the continuous stirred tan reactor (CSTR or chemostat). By computing the Jacobian matrix and evaluating the eigenvalues, we analyzed the stability of the possible co-existing states and divide microbial social interaction into eight categories: competition, coexisting parasitism, extinctive parasitism, cooperation, bistable amensalism, extinctive amensalism, coexisting commensalism and extinctive commensalism. Our model predicts that only parasitism, commensalism and cooperation could lead to stable co-existing state. We then move forward to understand the dynamics of microbial consortia with sequential metabolic reactions compartmentalized into distinct species. Coupled with Luedeking–Piret equation and Michaelis-Menten equation, accumulation of metabolic intermediate in one species and formation of final product in another species could be derived and assessed. We then conclude that there is inefficient conversion of metabolic intermediate to the final product if the two species form parasitism consortia. Our simulation indicates that commensalism consortia could efficiently convert metabolic intermediate to final product and maintain metabolic homeostasis (i.e., constant final product formation) with a broad range of dilution rates. Instead, cooperative consortia may not maintain this metabolic homeostasis due to the mutualistic relation between the two species. In this study, we discovered the underlying dynamics and emergent properties of microbial consortia, which may provide critical knowledge for us to engineering them. In particular, the simplicity and the rich dynamics of the consortia model highlight the importance to integrate unstructural kinetic models and social interaction parameters to systematically improve our prediction power.

Acknowledgments

Dr. Xu would like to acknowledge the Bill and Melinda Gates Foundation (OPP1188443) and National Science Foundation under grant number 1805139 for financially supporting this project.

Conflicts of interests

None declared.

Reference

1.   Fredrickson, A.G. and G. Stephanopoulos, *Microbial Competition.* Science, 1981. **213**(4511): p. 972-979.
2.   Brenner, K., L. You, and F.H. Arnold, *Engineering microbial consortia: a new frontier in synthetic biology.* Trends in Biotechnology, 2008. **26**(9): p. 483-489.
3.   Tsoi, R., Z. Dai, and L. You, *Emerging strategies for engineering microbial communities.* Biotechnology Advances, 2019. **37**(6): p. 107372.
4.   McCarty, N.S. and R. Ledesma-Amaro, *Synthetic Biology Tools to Engineer Microbial Communities for Biotechnology.* Trends in Biotechnology, 2018.
5.   Jawed, K., S.S. Yazdani, and M.A.G. Koffas, *Advances in the development and application of microbial consortia for metabolic engineering.* Metabolic Engineering Communications, 2019. **9**: p. e00095.
6.   Minty, J.J., et al., *Design and characterization of synthetic fungal-bacterial consortia for direct production of isobutanol from cellulosic biomass.* Proceedings of the National Academy of Sciences, 2013. **110**(36): p. 14592.
7.   Lu, Z.-M., et al., *Microbial ecology of cereal vinegar fermentation: insights for driving the ecosystem function.* Current Opinion in Biotechnology, 2018. **49**: p. 88-93.
8.   Wang, R., et al., *Recent advances in modular co-culture engineering for synthesis of natural products.* Current Opinion in Biotechnology, 2020. **62**: p. 65-71.
9.   Zhang, H. and X. Wang, *Modular co-culture engineering, a new approach for metabolic engineering.* Metab Eng, 2016. **37**.
10.   Arora, D., et al., *Expanding the chemical diversity through microorganisms co-culture: Current status and outlook.* Biotechnology Advances, 2020: p. 107521.
11.   Stephens, K., et al., *Bacterial co-culture with cell signaling translator and growth controller modules for autonomously regulated culture composition.* Nature Communications, 2019. **10**(1): p. 4129.
12.   Lv, Y., et al., *Coupling feedback genetic circuits with growth phenotype for dynamic population control and intelligent bioproduction.* Metabolic Engineering, 2019. **54**: p. 109-116.
13.   Rugbjerg, P., et al., *Synthetic addiction extends the productive life time of engineered Escherichia coli populations.* Proceedings of the National Academy of Sciences, 2018.
14.   Dai, Z., et al., *Versatile biomanufacturing through stimulus-responsive cell–material feedback.* Nature Chemical Biology, 2019. **15**(10): p. 1017-1024.

15.   Ben Said, S., et al., *The engineering of spatially linked microbial consortia – potential and perspectives.* Current Opinion in Biotechnology, 2020. **62**: p. 137-145.

16.   Beaudoin, T., et al., *Staphylococcus aureus interaction with Pseudomonas aeruginosa biofilm enhances tobramycin resistance.* npj Biofilms and Microbiomes, 2017. **3**(1): p. 25.

17.   Davies, J. and D. Davies, *Origins and Evolution of Antibiotic Resistance.* Microbiology and Molecular Biology Reviews, 2010. **74**(3): p. 417.

18.   Song, H.-S., et al., *Mathematical Modeling of Microbial Community Dynamics: A Methodological Review.* Processes, 2014. **2**: p. 711-752.

19.   Succurro, A. and O. Ebenhöh, *Review and perspective on mathematical modeling of microbial ecosystems.* Biochemical Society transactions, 2018. **46**(2): p. 403-412.

20.   Kong, W., et al., *Designing microbial consortia with defined social interactions.* Nature Chemical Biology, 2018. **14**(8): p. 821-829.

21.   Monod, J., *THE GROWTH OF BACTERIAL CULTURES.* Annual Review of Microbiology, 1949. **3**(1): p. 371-394.

22.   Han, K. and O. Levenspiel, *Extended monod kinetics for substrate, product, and cell inhibition.* Biotechnology and Bioengineering, 1988. **32**(4): p. 430-447.

23.   Levenspiel, O., *The monod equation: A revisit and a generalization to product inhibition situations.* Biotechnology and Bioengineering, 1980. **22**(8): p. 1671-1687.

24.   Luong, J.H., *Generalization of monod kinetics for analysis of growth data with substrate inhibition.* Biotechnol Bioeng, 1987. **29**(2): p. 242-8.

25.   Tsuchiya, H.M., et al., *Predator-prey interactions of Dictyostelium discoideum and Escherichia coli in continuous culture.* J Bacteriol, 1972. **110**(3): p. 1147-53.

26.   Lotka, A.J., *ELEMENTS OF PHYSICAL BIOLOGY.* Science Progress in the Twentieth Century (1919-1933), 1926. **21**(82): p. 341-343.

27.   Volterra, V., *Fluctuations in the Abundance of a Species considered Mathematically1.* Nature, 1926. **118**(2972): p. 558-560.

28.   May, R.M., *Simple mathematical models with very complicated dynamics.* Nature, 1976. **261**(5560): p. 459-467.

29.   Xu, P., *Analytical solution for a hybrid Logistic-Monod cell growth model in batch and continuous stirred tank reactor culture.* Biotechnology and Bioengineering, 2019. **n/a**(n/a).

30.   Stephanopoulos, G., *The dynamics of commensalism.* Biotechnology and Bioengineering, 1981. **23**(10): p. 2243-2255.

31.   Robert Luedeking, E.L.P., *A kinetic study of the lactic acid fermentation. Batch process at controlled pH.* Journal of Biochemical and Microbiological Technology and Engineering, 1959. **1**(4): p. 393-412.

Symbols and biophysical explanations with SI units

μ_{Amax} maximal specific growth rate for species A, (1/h)

μ_A specific growth rate for species A, (1/h)

μ_{Bmax} maximal specific growth rate for species B, (1/h)

μ_B specific growth rate for species B, (1/h)

K_{SA} substrate saturation constant for species A, (g/L)

K_{SB} substrate saturation constant for species B, (g/L)

Y_{AS} Species A biomass yield from substrate S, (g/g)

Y_{BS} Species B biomass yield from substrate S, (g/g)

Y_{BA} Product B (P_B) yield from intermediate A (P_A), (g/g)

Y_{PS} Intermediate A (P_A) yield from substrate S, (g/g)

α Growth-associated intermediate A (P_A) formation coefficient, dimensionless

β Growth-unassociated intermediate A (P_A) formation rate, (1/h)

γ_{AB} Interaction coefficient of species A imposes on species B, dimensionless

γ_{BA} Interaction coefficient of species B imposes on species A, dimensionless

k Rate constant of intermediate A (P_A) converted to product B (P_B), (1/h)

K_m Intermediate A saturation constant for species B, (g/L)

X_A Species A biomass in the CSTR, (g/L)

X_B Species B biomass in the CSTR, (g/L)

P_A Intermediate A concentration in the CSTR, (g/L)

P_B Product B concentration in the CSTR, (g/L)

S Substrate concentration in the CSTR, (g/L)

S_0 Substrate concentration in the feeding stream, (g/L)

D Dilution rate in the CSTR, (1/h)

Tables

Table 1. A unified mathematical model to describe the mass balance equations that govern microbial social interactions in chemostat.

Equations	Equation No.
$$\mu_A = \frac{\mu_{Amax}S}{K_{SA} + S}\left(1 + \gamma_{BA}\frac{X_B}{S_0\,Y_{BS}}\right)$$	Eqn. 1
$$\mu_B = \frac{\mu_{Bmax}S}{K_{SB} + S}\left(1 + \gamma_{AB}\frac{X_A}{S_0\,Y_{AS}}\right)$$	Eqn. 2
$$\frac{d}{dt}X_A(t) = \mu_A\,X_A(t) - D\,X_A(t)$$	Eqn. 3
$$\frac{d}{dt}X_B(t) = \mu_B\,X_B(t) - D\,X_B(t)$$	Eqn. 4
$$\frac{d}{dt}S(t) = D\left(S_0 - S(t)\right) - \frac{\mu_A\,X_A(t)}{Y_{AS}} - \frac{\mu_B\,X_B(t)}{Y_{BS}}$$	Eqn. 5
$$\frac{d}{dt}P_A(t) = (\alpha\,\mu_A + \beta)X_A(t) - \frac{k\,X_B(t)\,P_A(t)}{Y_{BA}\left(K_m + P_A(t)\right)} - D\,P_A(t)$$	Eqn. 6
$$\frac{d}{dt}P_B(t) = \frac{k\,X_B(t)\,P_A(t)}{K_m + P_A(t)} - D\,P_B(t)$$	Eqn. 7
$$\frac{d}{dt}S(t) = D\left(S_0 - S(t)\right) - \frac{\mu_A\,X_A(t)}{Y_{AS}} - \frac{\mu_B\,X_B(t)}{Y_{BS}} - \frac{(\alpha\,\mu_A + \beta)X_A(t)}{Y_{PS}}$$	Eqn. 8

Figure captions

Fig. 1 Categories of microbial social interactions: competition, amensalism, parasitism, neutralism, commensalism and cooperation. The interaction coefficient was defined by a dimensionless factor (γ_{AB} or γ_{BA}) that describe the beneficial or detrimental interactions between species A and species B. A green arrow indicates beneficial relation, a blunt-ended orange arrow indicates harmful relation.

Fig. 2 Dynamics of microbial competition, parasitism and cooperation at different dilution rate in chemostat. Green solid line: species B exist alone; Blue solid line: species A exists alone; Light orange solid line: *unstable* co-exist solution; Dark orange solid line: *stable* co-exist solution; Dash line: trajectory of steady state solution due to a small perturbation from any random state with fixed dilution rate. In all the simulation, we assume that species A has a larger growth fitness (maximal specific growth rate) than species B. (**a**) Competition. Species A and species B mutually exclude each other, leading to an unstable co-exist solution. Any perturbation from the coexist state (light orange line) will result in the survival of a single species (either A or B). (**b**) Coexisting parasitism: Species A benefits species B, but species B is harmful to species A. Stable coexisting is possible at relatively large dilution rate (equivalently to harsh conditions). (**c**) Extinctive parasitism: Species B benefits species A, but species A is harmful to species B. Stable coexisting is impossible and B will extinct. (**d**) Cooperation: Species A and species B mutually benefit each other. A stable coexisting solution (dark orange line) is possible due to the mutualistic interactions between species A and B. Any infinitesimal perturbation from the unstable coexisting solution (light orange line) will move the system to a washout state (solution falls to the origin) or to the stable co-existing state (dark orange line).

Fig. 3 Two-dimensional phase portrait of steady state solutions of simple microbial interaction models. Phase portrait for bistable competition (**a**), coexisting parasitism (**b**), extinctive parasitism (**c**), and cooperation (**d**). The 2-D phase portrait corresponds to the trajectory of the solution from any random state at a fixed dilution rate in Fig. 2. In all the simulation, we assume that A has a larger growth fitness than species B.

Fig. 4 Dynamics of microbial amensalism and commensalism at different dilution rate in chemostat. Purple solid line: species B exist alone; Blue solid line: species A exists alone; Light orange solid line: *unstable* co-exist solution; Dark orange solid line: *stable* co-exist solution; Dash line: trajectory of steady state solution due to a small perturbation from any random state with fixed dilution rate. In all the simulation, we assume that species A has a larger growth fitness than species B. (**a**) Bistable amensalism: Species B is harmful to species A, but species A is neutral to species B. Coexistence is unstable and any perturbation will lead to the survival of a single species (either species A or species B). (**b**) Extinctive amensalism: Species A is harmful to species B, but species B is neutral to species A. The system eventually leads to the existence of species A alone (species B will extinct). (**c**) Coexisting commensalism: Species A benefits species B, but species B is neutral to species A. Species A and B will exist together. (**d**) Extinctive commensalism: Species B benefits species A, but species A is neutral to species B. The system eventually leads to the existence of species A alone (species B will extinct).

Fig. 5 Steady state solutions of microbial co-culture with sequential metabolic reactions compartmentalized in two species. Red dots: biomass; blue dots: intermediate P_A or product P_B. In all the simulation, we assume that species A has a larger growth fitness than species B. Intermediate P_A secreted from species A is converted to product P_B by species B. (**a**) Coexisting

parasitism: Species A benefits species B, but species B is harmful to species A. (**b**) Extinctive parasitism: Species B benefits species A, but species A is harmful to species B. The system will move to species A existing state with only intermediate P_A accumulation (no product P_B formation). (**c**) Coexisting commensalism: Species A benefits species B, but B is neutral to A. (**d**). Cooperation: Species A and B mutually benefit each other.

Fig. 6 Operational conditions of microbial co-culture with sequential metabolic reactions compartmentalized in two species. (**a**) Coexisting parasitism leads to the inefficient conversion of intermediate P_A to product P_B, as the dilution rate increases. Optimal dilution rate is possible to maximize P_B production. (**b**) Extinctive parasitism only leads to the presence of species A and the accumulation of intermediate PA. Optimal dilution rate is possible to maximize intermediate P_A production. (**c**) Coexisting commensalism leads to the coexistence of species A and B and the efficient conversion of intermediate P_A to product P_B. (**d**) Mutualistic cooperation leads to the accumulation of intermediate P_A and rapid declining of product P_B as the dilution rate increases.

Figures

(1) A ⊣══════⊢ B $\gamma_{AB} = -1, \gamma_{BA} = -1$ Competition

(2) A ⊢────── B $\gamma_{AB} = 0, \gamma_{BA} = -1$ Amensalism

(3) A ══════→ B $\gamma_{AB} = 1, \gamma_{BA} = -1$ Parasitism

(4) A ──────⊢ B $\gamma_{AB} = -1, \gamma_{BA} = 0$ Amensalism

(5) A　　　　B $\gamma_{AB} = 0, \gamma_{BA} = 0$ Neutralism

(6) A ──────→ B $\gamma_{AB} = 1, \gamma_{BA} = 0$ Commensalism

(7) A ←══════⊢ B $\gamma_{AB} = -1, \gamma_{BA} = 1$ Parasitism

(8) A ←────── B $\gamma_{AB} = 0, \gamma_{BA} = 1$ Commensalism

(9) A ⇄══════⇄ B $\gamma_{AB} = 1, \gamma_{BA} = 1$ Cooperation

(Fig. 1)

(Fig. 2)

(Fig. 3)

(Fig. 4)

(Fig. 5)

(Fig. 6)