Bridging the gap: Using reservoir ecology and human serosurveys to estimate Lassa virus spillover in West Africa

Andrew J. Basinski[1], Elisabeth Fichet-Calvet[2], Anna R. Sjodin[3],

Tanner J. Varrelman[4], Christopher H. Remien[1], Nathan C. Layman[3],

Brian H. Bird[5], David J. Wolking[5], Corina Monagin[5],

Bruno M. Ghersi[5], Peter A. Barry[6], Michael A. Jarvis[7],

Paul E. Gessler[8], Scott L. Nuismer[3]

1 Department of Mathematics, University of Idaho, Moscow, Idaho, USA

2 Department of Virology, Bernhard-Nocht Institute of Tropical Medicine, Hamburg, Germany

3 Department of Biological Sciences, University of Idaho, Moscow, Idaho, USA

4 Bioinformatics and Computational Biology, University of Idaho, Moscow, Idaho, USA

5 One Health Institute, School of Veterinary Medicine, University of California, Davis, California, USA

6 Center for Comparative Medicine, California National Primate Research Center, Department of Pathology and Laboratory Medicine, University of California, Davis, California, USA

7 School of Biomedical and Healthcare Sciences, University of Plymouth, Plymouth, UK

8 College of Natural Resources, University of Idaho, Moscow, Idaho, USA

* abasinski@uidaho.edu

Abstract

Forecasting the risk of pathogen spillover from reservoir populations of wild or domestic animals is essential for the effective deployment of interventions such as wildlife vaccination or culling. Due to the sporadic nature of spillover events and limited availability of data, developing and validating robust, spatially explicit, predictions is challenging. Recent efforts have begun to make progress in this direction by capitalizing on machine learning methodologies. An important weakness of existing approaches, however, is that they generally rely on combining human and reservoir infection data during the training process and thus conflate risk attributable to the prevalence of the pathogen in the reservoir population with the risk attributed to the realized rate of spillover into the human population. Because effective planning of interventions requires that these components of risk be disentangled, we developed a multi-layer machine learning framework that separates these processes. Our approach begins by training models to predict the geographic range of the primary reservoir and the subset of this range in which the pathogen occurs. The spillover risk predicted by the product of these reservoir specific models is then fit to data on realized patterns of historical spillover into the human population. The result is a geographically specific spillover risk forecast that can be easily decomposed and used to guide effective intervention. Applying our method to Lassa virus, a zoonotic pathogen that regularly spills over into the human population across West Africa, results in a model that explains a modest but statistically significant portion of geographic variation in historical patterns of spillover. When combined with a mechanistic mathematical model of infection dynamics, our spillover risk model predicts that 897,700 humans are infected by Lassa virus each year across West Africa, with Nigeria accounting for more than half of these human infections.

Author Summary

The 2019 emergence of SARS-CoV-2 is a grim reminder of the threat animal-borne pathogens pose to human health. Even prior to SARS-CoV-2, the spillover of pathogens from animal reservoirs was a persistent problem, with pathogens such as Ebola, Nipah, and Lassa regularly but unpredictably causing outbreaks. Machine-learning models that

anticipate when and where pathogen transmission from animals to humans is likely to occur would help guide surveillance efforts and preemptive countermeasures like information campaigns or vaccination programs. We develop a novel machine learning framework that uses datasets describing the distribution of a virus within its host and the range of its animal host, along with data on spatial patterns of human immunity, to infer rates of animal-to-human transmission across a region. By training the model on data from the animal host alone, our framework allows rigorous validation of spillover predictions using human data. We apply our framework to Lassa fever, a viral disease of West Africa that is spread to humans by rodents, and use the predictions to update estimates of Lassa virus infections in humans. Our results suggest that Nigeria is most at risk for the emergence of Lassa virus, and should be prioritized for outbreak-surveillance.

Introduction

Emerging infectious diseases (EIDs) pose a persistent threat to public health. Approximately 60% of EIDs are caused by pathogens that normally circulate in wild or domestic animal reservoirs (i.e., zoonotic pathogens) [1]. Prior to full scale emergence, interaction between humans and wildlife creates opportunities for the occasional transfer, or spillover, of the zoonotic pathogen into human populations [2]. These initial spillover infections, in turn, represent newly established pathogen populations in human hosts that are subject to evolutionary pressures and may potentially lead to increased transmission among humans [2,3]. Consequently, a key step in preempting the threat of EIDs is careful monitoring of when and where spillover into the human population occurs. However, because the majority of EIDs from wildlife originate in low and middle income regions with limited disease surveillance, accurately estimating the rate and geographical range of pathogen spillover, and therefore the risk of new EIDs, is a major challenge [1].

Machine learning techniques have shown promise at predicting the geographical range of spillover risk for several zoonotic diseases including Lassa fever [4–6], Ebola [7,8], and Leishmaniases [9]. Generally, these models are trained to associate environmental features with the presence or absence of case reports in humans or the

associated reservoir. Once inferred from the training process, the learned relationships 19
between disease presence and the environment can be extended across a region of 20
interest. Using these techniques, previous studies of Lassa fever (LF) have derived risk 21
maps that assess the likelihood of human LF cases being present in different regions of 22
West Africa [4, 5]. However, because these forecasts combine case-reports from both 23
rodents and humans in the training process, they conflate attributes of the human and 24
reservoir populations that increase spillover risk. Consequently, these approaches shed 25
little light on aspects of reservoir or human populations that determine the magnitude 26
of spillover at a location and thus miss opportunities to identify effective interventions. 27

We develop a multi-layer machine learning framework that accounts for the 28
differences between how data involving a wildlife reservoir, and data from human 29
serosurveys, can simultaneously inform spillover risk in people and rigorously assess 30
whether predicted risk quantifies the rate of new infections in humans. Our approach 31
uses machine learning algorithms that, when trained on data from the wildlife reservoir 32
alone, estimate the likelihood that the reservoir and the zoonotic pathogen are present 33
in an area. These predictions are then combined into a composite estimate of spillover 34
risk to humans. Next, our framework uses estimates of human pathogen seroprevalence, 35
as well as estimates of human population density, to translate the composite risk 36
estimate into a prediction of the realized rate of zoonotic spillover into humans. 37
Omitting human seroprevalence data from the training process of the risk-layer has 38
several advantages. First, in the case of LF, due to modern transportation and the 39
longevity of Lassa virus antibodies in humans, a general concern is that the reported 40
location of individual cases of human disease or Lassa virus antibody detection is not 41
the site at which the infection occurred [10–12]. If the dispersal ability of the reservoir 42
is small, training the risk layer on reservoir infections alone helps the model avoid these 43
biases when learning the spatial variation of spillover risk. Secondly, in our framework, 44
human seroprevalence estimates provide an ultimate test of the risk layer's ability to 45
correlate with spatial variation in the cumulative human exposure to the pathogen. The 46
seroprevalence data, in turn, stem from population-based surveys at a site and are 47
therefore much less likely to be influenced by the movement of individuals. 48

We apply our framework to Lassa virus (formally *Lassa mammarenavirus* [LASV]), a 49
bi-segmented, single-stranded ambisense RNA virus in the *Arenaviridae* family and the 50

causative agent of LF in West Africa [11,13]. Though LASV can transmit directly between humans and often does so in hospital settings [14], rodent-to-human transmission accounts for the majority of new LASV infections [11,15]. Specifically, the multimammate rat *Mastomys natalensis* is believed to be responsible for most of the transmission into the human population, either through consumption of food contaminated by rodent feces and urine or through hunting and consumption of the rodent reservoir itself [16]. What remains largely unknown, however, is the extent to which spatial patterns of spillover are driven by spatial variation in the abundance of *M. natalensis* and viral prevalence within *M. natalensis* relative to spatial variation in other contributing factors such as human behavior, housing materials, or other rodent reservoirs. An additional unknown is the true magnitude of spillover into the human population outside of the few areas in Sierra Leone and Nigeria where hospitals with Lassa diagnostic capacity exist. As a consequence, most estimates for the magnitude of Lassa virus spillover rely on longitudinal serosurveys conducted in the 1980s in Sierra Leone [17], yielding estimates of between 100,000 and 300,000 LASV infections each year across West Africa. Here, we use our framework to fill these important gaps in our current understanding of Lassa virus spillover within West Africa.

Data and Study Region

We used online data repositories and literature sources to collect three types of data in West Africa spanning the time-range 1970 - 2017: 1) capture-locations of *M. natalensis*, as well as occurrence locations of non-Mastomys murids; 2) locations and outcomes of LASV surveys conducted in *M. natalensis*; and 3) locations and measured seroprevalence of human LASV serosurveys. The focal region from which our data originate, shown in Fig 1, was chosen as the intersection of West Africa and the International Union for Conservation of Nature (IUCN) range map for *Mastomys natalensis* [18,19]. Though *M. natalensis* is widely distributed across all of Africa, the species consists of multiple clades that likely differ in their ability to serve as hosts to LASV [20,21]. By limiting the spatial extent of the study region to West Africa we focus on the region occupied by the A-I clade of *M. natalensis* that is believed to transmit LASV [22]. Our *M. natalensis* capture data, as well as all of the LASV survey

data, originate from within this region, thus providing a discrete bound on the area of Africa in which the learned relationships of the model apply. For these analyses, this study region was divided into 0.05°x0.05° pixels (approximately 5 km by 5 km at the equator).

Fig 1. Map of the study region. The dashed blue line indicates the study region from which rodent and human data originate. Dots indicate locations at which Lassa virus or arenavirus antibodies have been sampled in rodents or humans. Each rodent point shows the outcome of a serological or PCR test. Each human population point shows the location of a serosurvey.

The first two datasets generate response variables for the model layers that predict LASV risk. The human seroprevalence data are used to evaluate the combined LASV risk layer for its ability to predict LASV spillover in humans and are also used to calibrate the stage of the model that predicts human LASV spillover. Our full dataset and the script files used to fit the models are available in a github repository [23].

Mastomys natalensis presence data and background

We collected data on historical captures of *M. natalensis* from various sources. First, several sources were used to identify all countries of West Africa that contain *M. natalensis* [24–26]. Next, rodent and mammal databases, as well as literature sources, were cross-referenced to fill in details regarding the year of capture, latitude/longitude

coordinates, and the method of identification for each location at which *M. natalensis* was documented [17, 20, 27–42]. Because *M. natalensis* is morphologically similar to other rodents in the study region (e.g., *Mastomys erythroleucus*), we only include those presences that have been confirmed with genetic methods or skull morphology. We found 167 locations with confirmed *M. natalensis* captures. All *M. natalensis* captures occurred in the time-range 1977 - 2017.

Fitting the model requires supplementing the presence-only data with background points, also called pseudo-absences [43, 44]. Background points serve as an estimate of the distribution of sampling effort for the organism being modeled [45]. We used background points chosen from locations where rodents in the family Muridae had been captured in West Africa from the Global Biodiversity Information Facility (GBIF) website [46]. We filtered the original dataset to reduce the likelihood of including *M. natalensis* rodents that were misidentified as *M. erythroleucus* and vice versa. Namely, we omit from the collection all Murid occurrences that are within the genus *Mastomys*. In addition, to ensure that the GBIF captures are concurrent with captures of *M. natalensis*, we only retained captures that occurred in the time-frame of the *M. natalensis* captures. Finally, we only included records that are within the study region depicted in Fig 1 and that fall outside of any pixel that contains a documented *M. natalensis*. The resulting GBIF dataset spans the years 1977 - 2015.

These data were used to categorize the subset of the pixels that contained one or more captures into two exclusive categories: those in which at least one *M. natalensis* had been captured (termed presences), and those with only non-Mastomys rodent occurrences (termed pseudo-absences). In total, our dataset classified 155 unique pixels as capture-positive for *M. natalensis*, and 252 pixels as background (Table 1).

Table 1. Summary of rodent captures used in the reservoir layer.

Country	Year	# Pseudoabsences	# Presences
Benin	2001-2017	12	7
Burkina Faso	1977-2008	3	15
Ghana	1999-2011	13	9
Guinea	1996-2012	71	12
Guinea-Bissau	2013	1	0
Ivory Coast	1978-2010	21	8
Liberia	1980-2013	18	0
Mali	1979-2012	58	47
Niger	1977-2007	16	14
Nigeria	1977-2015	7	13
Senegal	1990-2005	0	13
Sierra Leone	1977-2014	31	17
Togo	1982	1	0
Aggregate	1977-2017	252	155

Pseudoabsences shows the number of unique $0.05 \times 0.05°$ pixels in the GBIF dataset for which only non-Mastomys rodents were captured. # Presences indicates the number of pixels in which one or more *M. natalensis* was captured.

Surveys of *Mastomys natalensis* for Lassa virus

We compiled a dataset that contains occurrences of LASV in rodents or humans. The dataset was established by an extensive review of LASV literature. Primary sources were found by PubMed and GenBank searches of the terms "Lassa", "Lassa fever", "Lassa virus", "Lassa arenavirus", and "Lassa mammarenavirus" [47]. Data from these primary sources was organized into an Excel workbook.

From the full LASV dataset, we collected published studies that sampled *M. natalensis* rodents for indicators of LASV. For each study, we found the sampling location for each tested rodent (either latitude/longitude or a locality name for which coordinates could be obtained). In total, we compiled thirteen rodent studies [17, 30, 34, 36, 39, 41, 42, 48–53] that tested *M. natalensis* for LASV and contained latitude/longitude coordinates. The resulting test locations originate from six countries and span the years 1972 – 2014.

Because the prevalence of LASV in rodents varies seasonally [54], and because of the sparsity of time-series data that might otherwise allow the average LASV prevalence in rodents to be estimated, we used the collected data to broadly classify pixels into the categories "Lassa positive" or "Lassa negative". Specifically, a pixel was defined as Lassa positive if, at some point, a *M. natalensis* rodent was captured within the pixel,

and the rodent tested positive for LASV using a RT-PCR assay or viral isolation. 137

Because arenavirus antibodies cross-react, a positive LASV antibody test in an 138

individual rodent only indicates past infection with an arenavirus, and not necessarily 139

LASV. In an effort to reduce the frequency of false positives in the training data, pixels 140

that only contain LASV seropositive tests of rodents, and no positive LASV viral 141

detection, were not used as training data. These criteria led to the omission of eight 142

pixels from the training data. Fitting the model with these eight pixels included as 143

presences is an option in the code on the github repository, but does not substantially 144

affect the overall fit of the model [23]. 145

Although serosurveys of rodents cannot specifically show that LASV is present, they 146

can indicate the absence of LASV (along with all other arenaviruses). Pixels were 147

classified as Lassa negative if five or more *M. natalenis* rodents in total were tested for 148

infection with LASV by RT-PCR, or tested for any previous arenavirus exposure using 149

a serological assay, and all rodents tested were negative. We chose a threshold of five to 150

help reduce the chance of including false negatives (i.e. sites that have LASV but in 151

which only non-exposed rodents were captured). This procedure allowed us to classify 152

62 unique pixels in total: 27 were classified as Lassa negative, and 35 were classified as 153

Lassa positive (Table 2 and Fig 1). 154

Table 2. Summary of LASV positive and LASV negative pixels used in the pathogen layer.

Country	Year	# Pixels	# Neg. Pixels	# Pos. Pixels
Ghana	2010-2011	7	7	0
Guinea	2003-2014	19	6	13
Ivory Coast	2003-2013	4	3	1
Mali	2004-2012	11	7	4
Nigeria	1972-2012	6	3	3
Sierra Leone	1972-2009	15	1	14
Aggregate	1972-2014	62	27	35

Each row aggregates literature and GenBank data sources over a country. # Pos. Pixels indicates the number of unique pixels that had one or more LASV-infected rodents. # Neg. Pixels is the number of pixels in which five or more rodents were tested and found negative for LASV infection or antibody.

Human seroprevalence data

From our full LASV dataset described in the previous section, we collected literature
sources that describe the prevalence of arenavirus antibodies in human populations of
West Africa. As with the rodent LASV infection data, arenavirus antibodies are not
specific to LASV. However, because human serosurveys were often conducted in LASV
endemic areas or near documented locations of LASV-infected rodents, these
serosurveys likely measured the fraction of humans with previous LASV infection, rather
than exposure to another arenavirus. We required that each literature source include
information on the diagnostic method that was used to test individuals (e.g., ELISA,
IFA) and broad details of the survey design. We only included survey studies that were
designed to estimate the seroprevalence in the local community population. This
criterion excluded surveys of hospitals, for example, as well as surveys of missionaries.

Each datum contains latitude and longitude of the serosurvey, the number of
individuals tested, and the number of individuals determined to have arenavirus
antibodies. In total, we collected 94 serosurveys from seven studies (Fig 1) [17, 55–60].
These serosurveys were conducted between 1970 and 2015 and are located in five
countries in West Africa (Table 3 and Fig 1).

Table 3. Summary of human arenavirus serosurveys used in the model.

Country	Year	# Sites	Method	# Tested	% Seropositive	Reference
Ghana	2010-2011	10	ELISA	657	5	[57]
Guinea	2000	30	IFA	977	11	[55]
Guinea	1990-1993	28	ELISA	3276	23	[56]
Liberia	1980-1982	7	IFA	1848	5	[59]
Mali	2015	3	ELISA	600	33	[58]
Sierra Leone	1977-1983	14	IFA	5098	23	[17]
Sierra Leone	1970-1972	2	CF	255	6	[60]
Aggregate	1970-2015	94		12,711	19	

Each row is an individual literature source. For each study, # Sites shows the number
of locations at which arenavirus surveys were performed, # Tested indicates the total
number of individuals tested across sites, and % Seropositive shows the percentage of
individuals that tested positive across all sites.

Predictors

We include predictors that are broadly hypothesized to influence the distributions of *M. natalensis* and LASV. *M. natalensis* is widely distributed across sub-Saharan Africa in savanna and shrubland environments. Within such environments, *M. natalensis* is commonly associated with small rural communities and is considered a serious agricultural pest [19,54]. To allow the model the possibility to learn these relationships, we include predictors that describe MODIS land cover features as predictors, and also include human population density within each pixel. We also include elevation in meters. Because climate seasonality and crop maturation affect the breeding season of *M. natalensis*, we include various measures of the seasonality of the vegetative index (NDVI), precipitation, and temperature [61]. See S1 Appendix for a complete list of environmental variables. LASV is often associated with *M. natalensis*, so we use the same set of predictors for the pathogen layer.

Methods

We developed a model that predicts the rate of LASV infection in humans within individual $0.05°\text{x}0.05°$ pixels across West Africa. An overview of the model framework is depicted in Fig 2. Outputs from the model are generated in two stages. The first stage uses environmental features to estimate different layers of LASV spillover risk. The layers of risk, in turn, are described by: 1) D_M, a classification score indicating the likelihood that a pixel contains the primary rodent reservoir, *M. natalensis*, and 2) D_L, a score indicating the likelihood that LASV circulates within the *M. natalensis* population, conditioned on the rodent being present. Depending on the layer, the response variable for this stage is generated from documented occurrences of *M. natalensis* (D_M layer), or evidence of past LASV infection in *M. natalensis* (D_L layer). These layers are used to define a composite layer of spillover risk D_X, the product of D_M and D_L, that describes the likelihood that a pixel simultaneously contains *M. natalensis* and LASV. The second stage of our framework uses a generalized linear model to regress the estimates of human arenavirus seroprevalence onto the D_X layer. Lastly, we used an epidemiological model to estimate human incidence from the predictions of seroprevalence.

Fig 2. Overview of the model. Ellipses represent datasets, circles represent models, and rectangles represent model predictions.

Each risk layer of the first stage is generated by a separate boosted classification tree ²⁰³ (BCT). The BCT, in turn, uses environmental features within a pixel to infer a ²⁰⁴ classification score, between zero and one, that indicates how likely it is that the pixel is ²⁰⁵ positive for *M. natalensis* (D_M layer) or LASV in *M. natalensis* (D_L layer). BCTs use ²⁰⁶ a stage-wise learning algorithm that, at each stage, trains a new tree model to the ²⁰⁷ residuals of the current model iteration. Each newly fitted tree is added to the ensemble ²⁰⁸ model, thereby reducing the residual deviance between the model predictions and a ²⁰⁹ training set [62]. Boosted trees are commonly used in species and disease distribution ²¹⁰ models because they are simultaneously resistant to over-fitting in scenarios where many ²¹¹ feature variables are implemented and are also able to model complex interactions ²¹² among features [63]. ²¹³

Prior to inclusion in the model-fitting procedure, each feature variable was vetted for ²¹⁴ its ability to distinguish between presences and absences in each of the layers. ²¹⁵ Specifically, for each risk layer's binary response variable, we performed a ²¹⁶ Mann-Whitney U-test on each candidate feature. In doing so, we test the null ²¹⁷ hypothesis that the distribution of a feature is the same between pixels that are ²¹⁸ classified as a presence or (pseudo) absence. We only include predictors for which the ²¹⁹ null hypothesis is rejected at the $\alpha = 0.05$ level. ²²⁰

For a given training set, we fit the BCT model using the gbm.step function of the ²²¹ "dismo" package in the statistical language R [64]. This specific function uses 10-fold ²²² cross-validation to determine the number of successive trees that best model the ²²³ relationship between response and features without over-fitting the data [64]. The ²²⁴ learning rate parameter, which determines the weight given to each successive tree, was ²²⁵ set to small values (D_M: 10^{-2}, D_L: 10^{-3}) that encourage a final model that is ²²⁶ composed of many small incremental improvements. A smaller learning rate was used in ²²⁷ the D_L layer because the corresponding dataset was smaller. The parameter that ²²⁸ describes the maximum number of allowable trees was set to a large value (10^7) to ²²⁹ ensure that the cross-validation fitting process was able to add trees until no further ²³⁰ improvement occurred [62]. ²³¹

For the D_M layer, we trained 25 boosted classification trees to learn how ²³²

environmental predictors influence the suitability of a habitat for *M. natalensis*. Each model was fit by selecting 155 presence pixels and pairing these with 155 background pixels in which only non-Mastomys murids were found. Both presences and background pixels were chosen with replacement. By choosing equal numbers of presences and background pixels for each training set, we encourage each model to learn patterns in features that allow presences to be discriminated from background pixels, rather than having the model learn the (likely biased) distribution of presences and background pixels that are available in the overall dataset [44].

For each model fit for the D_M layer, presence and pseudo-absence pixels that were not used to train the model (i.e., out-of-bag data) were used to test the model using the area-under-the-receiver-curve (AUC). The AUC measures a classifier's ability to assign a high classification score to presences, and a low score to background pixels. A score of one indicates a perfect classifier, and a score of 0.5 indicates a classifier that is no better than chance. A pairwise-distance sampling scheme was used to pair an equal number of test-background pixels to the out-of-bag presences that together comprise the test set. Specifically, for each test presence point, the pairwise distance sampling method chooses a test background point so that the minimum spatial distance between the training presences and test presence is similar to the minimum distance between the test background point and training presences [65]. Compared to random selection of test background points, pairwise distance sampling oftentimes results in a lower AUC score that more accurately measures the model's ability to generalize to new regions [65].

The D_L layer is generated by the averaged predictions of 25 boosted classification tree models, each of which is trained to discriminate between pixels that are Lassa positive or Lassa negative. We trained each model on a dataset comprised of 27 absence locations and 27 presence locations, sampled from the full dataset with replacement. The estimation of error in the D_L layer is similar to that described in the D_M layer. Specifically, we calculate the AUC of the fitted model on an equal number of out-of-bag presences and absences.

Next, we combined the D_M and D_L layers into a composite feature, denoted by D_X, that is indicative of whether a pixel simultaneously has environmental features that are suitable for *M. natalensis*, as well as LASV in *M. natalensis*. The combined feature is defined as $D_X = D_M \times D_L$ and summarizes the realized risk of LASV spillover to

humans within the local environment.

Connection to human seroprevalence and incidence

To connect the new risk parameter D_X to human arenavirus seroprevalence, and to evaluate the ability of the D_X layer to explain historical LASV spillover in humans, we regressed seroprevalence from human arenavirus serosurveys on the D_X layer and an intercept. In doing so, we test whether human seroprevalence is significantly associated with the probability D_X that a pixel contains LASV-infected *M. natalensis*. We used quasi-binomial regression to account for overdispersion in seroprevalence measurements that could otherwise contaminate hypothesis tests on model coefficients [66]. More details on the motivation behind the quasi-binomial regression can be found in the S1 Appendix. In the regression, each seroprevalence estimate is weighted by the number of individuals tested in the serosurvey.

Next, we used an epidemiological model, based on the classic susceptible-infected-recovered framework, to derive an equation that relates a given LASV spillover rate into humans and the resulting seroprevalence in a human population. Throughout, we assume that the seroprevalence measures that were obtained from historical serosurveys describe LASV infection at steady state (i.e., are unchanging in time). This derivation, in turn, is used to translate the regression model's predictions of LASV seroprevalence into spillover infections per year in humans. For the model, we employ several assumptions: 1) humans within each 0.05x0.05° pixel constitute a closed population with constant per-capita death rate d. To facilitate steady-state analysis, we assume that new individuals are born in the pixel at a density-independent birth rate b. Within each pixel, humans are compartmentalized into three non-overlapping classes: susceptible (S), infected with LASV (I), and recovered from LASV infection (R). The size of the human population is assumed to be large enough so that stochastic events (LASV extinction) do not occur. 2) All LASV infections in humans are caused by contact with infectious rodents. Though human-to-human transmission of LASV is common in nosocomial outbreaks, rodent-to-human transmission is believed to be the primary pathway by which the virus is spread outside of hospital environments [15]. 3) Susceptible humans become infected

with LASV at a constant rate FS, where F denotes the rate of infectious contact
between a human and infected *M. natalensis* (i.e., the force of infection). Any seasonal
fluctuation in the contact rate between humans and rodents, as well as fluctuation in
the prevalence of LASV infection in rodents, is assumed to average out over the
decades-long timescales we consider. LASV-infected humans transition out of the
infected class at per-capita rate γ; a fraction μ die from illness associated with Lassa
Fever. 4) The remaining fraction $1 - \mu$ of individuals recover from infection and gain
immunity from LASV.

The duration of LASV immunity in humans is not fully understood. Studies suggest
that LASV immunity is the result of a combination of antibodies and a cell-mediated
immune response [17, 67]. Anecdotal cases have shown that LASV IgG antibodies can
remain in the blood of individuals for decades [10]. However, other studies have
indicated that the level of LASV antibodies, as well as the extent to which an individual
is protected against subsequent LASV infection, can wane with time [17, 67].
Preliminary analyses indicated that the possibility of waning immunity substantially
influenced our model's estimates of LASV infections per year. Because of this
uncertainty, we model the general scenario in which recovered individuals lose immunity
to LASV at per-capita rate λ and transition back into the susceptible class. This more
general model structure includes the scenario of lifelong immunity in the case that $\lambda = 0$.

Within each pixel across West Africa, the assumptions above lead to a system of
equations that describes the number of humans in each of the classes:

$$\begin{aligned}
\frac{dS}{dt} &= b - dS - FS + \lambda R, \\
\frac{dI}{dt} &= FS - dI - \gamma I, \\
\frac{dR}{dt} &= \gamma(1 - \mu)I - dR - \lambda R.
\end{aligned} \tag{1}$$

We assume that, within each pixel, the dynamical system given by System (1) is at
steady state. Consequently, the net rate of mortality is equal to the constant birth rate
b, and each of the classes S, I, and R are not changing with time. The corresponding
steady-state values are found by setting the left-hand-side of equations (1) to zero, and
solving the resulting algebraic equations for each state variable. This yields the

steady-state values

$$S^* = \frac{b(\gamma + d)(d + \lambda)}{d\lambda(\gamma + d + F) + d(\gamma + d)(d + F) + \gamma F \lambda \mu},$$

$$I^* = \frac{bF(d + \lambda)}{d\lambda(\gamma + d + F) + d(\gamma + d)(d + F) + \gamma F \lambda \mu}, \tag{2}$$

$$R^* = \frac{b\gamma F(1 - \mu)}{d\lambda(\gamma + d + F) + d(\gamma + d)(d + F) + \gamma F \lambda \mu}.$$

At steady state, the total population size in a pixel is $P^* = S^* + I^* + R^*$. We can write P^* in terms of the model parameters by plugging in the steady-state values given by equations (2):

$$P^* = b \frac{\gamma \lambda + d^2 + d(\gamma + F + \lambda) + F(\gamma + \lambda - \gamma \mu)}{d\lambda(\gamma + d + F) + d(\gamma + d)(d + F) + \gamma F \lambda \mu}. \tag{3}$$

By dividing R^* by the total population size at steady state, P^*, we derive an equation for the steady-state seroprevalence, denoted Ω^*:

$$\Omega^* = \frac{\gamma F(1 - \mu)}{\gamma \lambda + d^2 + d(\gamma + F + \lambda) + F(\gamma + \lambda - \gamma \mu)}. \tag{4}$$

Now we solve for the total LASV spillover rate FS, given that the steady-state LASV seroprevalence is Ω^*. Solving Eq (4) for F in terms of Ω^* yields:

$$F = -\frac{\Omega^*(\gamma + d)(d + \lambda)}{\Omega^* d + \gamma(-\Omega^* \mu + \Omega^* + \mu - 1) + \Omega^* \lambda}. \tag{5}$$

The rate of new infections is given by

$$\eta := FS^* = \frac{P^* \Omega^*(d + \gamma)(d + \lambda)}{\gamma(1 - \mu)}. \tag{6}$$

These analyses were derived using Mathematica. The notebook file is available in the github repository [23].

By substituting our prediction of human LASV seroprevalence for Ω^*, we can estimate the total human infection rate using Eq (6). Calculating these estimates requires values for d, γ, μ, λ, and P^*. We chose parameters that are broadly in line with the epidemiology of LASV and the demography of humans in West Africa.

We use values of death rate d derived from country-specific lifespan estimates

obtained from WorldBank [68]. For a pixel within a given country, d is set to be the reciprocal mean lifespan of that country's 2018 life expectancy at birth. Studies indicate that the duration of LASV infection is typically about one month, so that $\gamma = 12 \ yr^{-1}$ across all pixels [11]. LASV infection causes mortality in a fraction $\mu = 0.02$ of non-nosocomial infections [17].

The rate of seroreversion is difficult to estimate empirically. McCormick et al. (1987) estimated that $\lambda = 0.064 \ \mathrm{yr}^{-1}$ using a longitudinal study of IgG immune markers in individuals. However, it is unclear whether their results indicated true seroreversion, or whether the reduction of LASV immune markers below detectable levels made it appear as though seroreversion occurred. To better understand the potential consequences of seroreversion in our infection-rate estimates, we focus on two scenarios. In the first, any individual that has recovered from LASV infection remains seropositive for the remainder of their life ($\lambda = 0 \ \mathrm{yr}^{-1}$). In the second scenario, seroreversion occurs at the rate estimated by McCormick et al. (1987) ($\lambda = 0.064 \ \mathrm{yr}^{-1}$). In this case, an individual recovered from LASV is assumed to produce antibodies and maintain LASV immunity for an average duration of 15.6 years. We use the unprocessed WorldPop 2020 population data (described in S1 Appendix) as an estimate of the steady-state population size, P^*, within each pixel of the original $0.0083°$ resolution.

Results

LASV risk layers

The D_M layer is constructed by averaging the predictions of 25 boosted classification tree models. Across all 25 bootstrap fits, the average out-of-bag AUC was 0.68, with a standard deviation of 0.05. This AUC indicates that the model has a modest ability to correctly discriminate pixels in which *M. natalensis* has been captured from background pixels, and is similar to out-of-bag AUC scores obtained in another study with a similar assessment criterion [5]. The algorithm assigned a high likelihood of occurrence to regions with a strong seasonal pattern of vegetation as well as specific levels of rainfall (S1 Appendix). Across 25 fitted models that made up the D_L layer, the average AUC was 0.85, with a standard deviation of 0.08. This indicates a model that is good at

discriminating between Lassa presences and absences. The algorithm primarily used 366
precipitation contingency to determine whether or not a pixel is suitable for endemic 367
LASV in *M. natalensis* (S1 Appendix). 368

Figs 3A-3C show maps of each of the fitted risk layers, as well as the combined layer 369
of realized risk, D_X. As indicated by the IUCN range map for *M. natalensis* [19], most 370
countries of West Africa are predicted to harbor this primary rodent reservoir of LASV 371
(Fig 3A). However, the rodent is predicted to be less prevalent along coastal areas of 372
West Africa and southern Nigeria. Similar to other Lassa risk maps [4,5], our D_L layer 373
predictions indicate that the risk of LASV in rodents is primarily concentrated in the 374
eastern and western extremes of West Africa (Fig 3B). The combined risk, shown in Fig 375
3C, indicates that environmental features suitable for rodent-to-human LASV 376
transmission are primarily located in Sierra Leone, Guinea, and Nigeria. 377

Fig 3. Calculating the combined risk layer. (A) Map shows the likelihood that
each 0.05° pixel in West Africa contains the primary reservoir of Lassa virus, *M.
natalensis*. Pink dots indicate locations of captures that were used to train the model.
Black line indicates the IUCN *M. natalensis* range map. (B) Predicted distribution of
Lassa virus in *M. natalensis*. Dots indicate locations in which *M. natalensis* were
surveyed for the virus. (C) Combined risk, defined as the product of the above two
layers.

Connection to human seroprevalence and spillover

A quasi-binomial regression indicated a significant, positive association between the combined LASV risk predictor D_X, and the human arenavirus seroprevalence measured in serosurveys (coefficient: 1.50, p = 0.000123, Fig 4). The model also indicated the presence of substantial overdispersion in the human seroprevalence dataset ($\phi = 15.1$). More information on the GLM output can be found in the S1 Appendix. By applying the general linear model to the combined LASV risk layer, we extrapolate the human LASV seroprevalence across West Africa (Fig 5). Our results indicate that human LASV seroprevalence is greatest in the eastern and western regions of West Africa, with especially high seroprevalence in Central Guinea, Sierra Leone, and Nigeria.

Fig 4. Human arenavirus seroprevalence vs the combined risk layer. Each circle represents a different serosurvey. The size of the circle indicates the number of humans that were tested. Solid black line shows the quasi-binomial prediction of seroprevalence, and the red dashed lines show the 95% confidence intervals. Confidence intervals were obtained by fitting the model 1000 times on random samples taken from the dataset with replacement.

Fig 5. Predicted human seroprevalence of Lassa virus in West Africa. Dots show locations of human serosurveys, and dot color indicates the residual of the predicted seroprevalence. White dots indicate locations for which measured seroprevalence fell within 0.1 of the prediction. Measured seroprevalence at red dots was 0.1 or more greater than that predicted, and seroprevalence at blue dots was 0.1 or more below the prediction.

Furthermore, by assuming that our predictions are representative of LASV infection at steady state, we can derive the number of LASV infections per year in humans. If the D_X layer accurately describes the spatial heterogeneity of LASV seroprevalence in humans, and if LASV antibody production upon recovery is lifelong, our framework estimates that 897,700 new human infections occur each year. Between 664,300 – 843,800 (i.e., 74 – 94%) of these infections are expected to be sub-clinical or asymptomatic, leaving 53,900 – 233,400 infections that might require hospitalization [17]. Given that 2% of all infections result in fatality, our estimates imply that 18,000 individuals die of Lassa Fever in West Africa each year. Though our model does not account for differences of LASV risk by sex or age, research suggests that hospitalizations may be skewed towards females, and fatalities will be biased towards individuals under 29 years of age but not skewed by gender [69].

Table 4 shows the number of LASV infections per year by country, ordered by number of infections. Our predictions indicate that more than half of new human LASV infections (531,700) in West Africa will occur in Nigeria (Fig 6). This distribution of LASV infection is largely due to the greater population size within Nigeria, as the per person spillover rates do not differ dramatically between countries (Table 4). After

Nigeria, Ghana (60,200 infections per year) and the Ivory Coast (57,700 infections per ⁠405
year), respectively, are predicted to have the highest incidence of human LASV ⁠406
infections. Sierra Leone, Nigeria, and Guinea are predicted to have the highest ⁠407
per-capita rates of LASV infection (Table 4). ⁠408

Fig 6. Predicted spatial density of Lassa virus infections in humans. Map shows the predicted infections per km^2. Yellow colors, representing a high number of infections, tend to occur in areas with high human population density and a high predicted seroprevalence.

Table 4. Predicted annual number of Lassa virus infections and infection rate

Country	1000's of infections	Rate
Nigeria	531.7	2.6
Ghana	60.2	2.0
Ivory Coast	57.7	2.3
Niger	46.9	2.0
Burkina Faso	44.4	2.1
Mali	44.3	2.2
Guinea	35.0	2.5
Benin	27.0	2.2
Sierra Leone	20.7	2.9
Togo	17.9	2.2
Liberia	9.9	2.0
Mauritania	1.0	1.9
Senegal	0.8	2.0

Infection rate is in units of number of infections per year per 1000 people. Estimates in the table are derived assuming seroreversion and reinfection do not occur.

The above estimates are based on the premise that, upon recovery from LASV infection, an individual produces antibodies for the remainder of their life. If, instead, LASV antibody production ceases after an average of 15.6 years as suggested by some longitudinal serosurveys [17], then a given level of seroprevalence implies almost five

times as many infections compared to the scenario with lifelong antibody production. ⁴¹³
Specifically, allowing for seroreversion and subsequent LASV reinfection in the model ⁴¹⁴
implies 4,383,600 infections occur each year. Inclusion of reinfection does not change the ⁴¹⁵
ranking of countries in Table 4. ⁴¹⁶

Discussion ⁴¹⁷

Machine learning approaches that forecast the spatial risk of emerging infectious ⁴¹⁸
diseases such as Lassa virus are often not designed to explain how aspects of the ⁴¹⁹
environment translate into realized pathogen spillover into human populations [4, 5]. ⁴²⁰
Models that specifically predict attributes of the reservoir from the environment, and ⁴²¹
use these predictions to quantify spillover into humans, offer a more mechanistic ⁴²²
understanding of the current and future spatial variation in human disease [70]. Our ⁴²³
forecasting framework advances these approaches by generating predictions of spillover ⁴²⁴
risk based only on data from the primary rodent reservoir of LASV, and rigorously ⁴²⁵
assessing our risk predictions on realized human spillover as measured by human ⁴²⁶
arenavirus serosurveys. As indicated by a generalized linear regression, our ⁴²⁷
reservoir-based model of spillover risk explains a modest and statistically significant ⁴²⁸
amount of the spatial variation in human arenavirus seroprevalence. ⁴²⁹

Using this framework, we are able to generate predictions of the number of new ⁴³⁰
LASV infections within different regions of West Africa. Our results indicate that ⁴³¹
Nigeria contributes the greatest number of new human infections each year, and that ⁴³²
the magnitude of new infections in Nigeria is driven primarily by its greater human ⁴³³
population density, rather than an increased per-capita risk. An assumption that drives ⁴³⁴
this result is the density-dependent form of spillover in the model (i.e., FS), in which ⁴³⁵
rodent-human interactions increase with human population density. This form is ⁴³⁶
appropriate if rodent interactions are well-mixed in the human population. For example, ⁴³⁷
if increases in human density were reflected in an increased number of humans per ⁴³⁸
dwelling, then the LASV risk posed by single rodent in a household would increase with ⁴³⁹
human population size. If these assumptions are correct, Nigeria is likely to represent ⁴⁴⁰
the greatest risk of LASV emergence because the large number of annual spillover ⁴⁴¹
events allows for extensive sampling of viral strain diversity and repeated opportunities ⁴⁴²

for viral adaptation to the human population [71]. 443

Our approach allows us to highlight the regions that contribute most to pathogen 444
spillover, and suggest locations for further surveillance. Our model indicates that the 445
highest per-capita risk to humans occurs in Sierra Leone, Guinea, and Nigeria. Given 446
the data that are currently available, our model suggests that these countries should be 447
prioritized for surveillance of LASV emergence in rodents and at-risk human 448
populations. Human serosurveys of the general population are notably lacking in 449
Nigeria, but have the potential to clarify the true magnitude of LASV spillover in West 450
Africa. Although it is known that certain broad regions of West Africa have a long 451
history of LASV spillover (e.g., Sierra Leone, Guinea, Nigeria), relatively little is known 452
about the prevalence of LASV in rodents or humans in other regions (e.g., Togo, Benin, 453
Mali, Burkina Faso). Our model suggests that Lassa virus infections occur regularly in 454
these under-sampled areas. Human serosurveys and rodent LASV testing from these 455
regions could help modeling approaches clarify the spatial distribution of Lassa fever 456
across West Africa. 457

In addition to identifying the regions most at risk for viral emergence, our model 458
framework provides updated estimates for the rate of LASV spillover across West Africa. 459
Previous estimates of 100,000 – 300,000 infections per year were based on longitudinal 460
studies from communities in Sierra Leone conducted in the 1980s [17]. Using 461
seroprevalence data from studies across West Africa, our model predicts between 462
897,700 – 4,383,600 LASV infections in humans occur each year. As demonstrated by 463
past research focused on estimating LASV infection in humans, where the true value lies 464
within this range depends on whether or not seroreversion and subsequent LASV 465
reinfection are regular features of human LASV epidemiology, and therefore reinforces 466
the need to better understand the scope for LASV reinfection [72]. It is important to 467
realize that our predictions include both symptomatic and asymptomatic infections. 468
Thus, because many human LASV infections result in mild flu-like symptoms or are 469
asymptomatic, it is unsurprising that our predicted values exceed the reported number 470
of confirmed LF cases in Nigeria [73,74]. Several factors may contribute to the 471
discrepancy between previous estimates of LASV spillover, and our revised estimates. 472
McCormick et al. (1987) used seroconversion data from a 15 month period to infer a 473
rate of LASV infection across West Africa. However, the population of West Africa has 474

increased by a factor of 2.4 since that time, making these estimates outdated [75]. ₄₇₅
Furthermore, our estimates are based on human seroprevalence data that comes from ₄₇₆
five countries in West Africa and spans a 45 year time period. Because our dataset was ₄₇₇
obtained from a broader spatial and temporal range, our estimates are less likely to be ₄₇₈
biased by sporadic extremes in LASV spillover. ₄₇₉

Accurate risk predictions could help guide risk-reduction and behavior-change ₄₈₀
communication campaigns, the distribution of future human LASV vaccines, and ₄₈₁
countermeasures directed at the rodent reservoir. In addition to vaccines that prevent ₄₈₂
infection in humans, new vaccine designs are currently being investigated for various ₄₈₃
wildlife pathogens as well, including pathogens in rodents [76, 77]. Wildlife vaccination ₄₈₄
campaigns that use vaccine baits have proven to be effective in the control of rabies in ₄₈₅
red fox (*Vulpes vulpes*) over large land areas, but require substantial planning and ₄₈₆
surveillance of the reservoir population [78]. Rodent population management could be ₄₈₇
another method of attenuating the risk of LASV in an area. Pinpointing areas that are ₄₈₈
most in need of spillover intervention will help overcome the logistical challenges that ₄₈₉
are associated with vaccine distribution to humans or wildlife on large scales. In ₄₉₀
addition to guiding intervention to specific regions, mechanistic forecasts similar to ours ₄₉₁
could help plan the logistics of such operations. ₄₉₂

Our framework sheds light on the connection between LASV spillover in humans and ₄₉₃
the environmental conditions favorable to pathogen and reservoir. The reservoir layer of ₄₉₄
our model identified strong seasonal trends in vegetation (NDVI) as the primary ₄₉₅
explanatory variable that determines where the rodent *M. natalensis* occurs. This ₄₉₆
builds on other work that identified properties of vegetation as important predictors of ₄₉₇
the range of *M. natalensis* [5]. In conjunction with a strong seasonality of vegetation, ₄₉₈
our model identified a range of mean and maximum rainfall values that limit the ₄₉₉
distribution of the LASV reservoir. This is in line with previous ecological studies ₅₀₀
showing that seasonal patterns of precipitation and vegetation are important drivers of ₅₀₁
seasonal breeding in *M. natalensis* [79]. Our model indicates that *M. natalensis* do not ₅₀₂
occur in areas associated with too much rainfall or areas without a clear wet/dry ₅₀₃
seasonality, resulting in a lower risk of LASV spillover in coastal areas of West Africa ₅₀₄
and southern Nigeria. The pathogen layer of our model also indicates that strong ₅₀₅
seasonal precipitation patterns are the leading environmental feature that is associated ₅₀₆

with LASV in *M. natalensis* and the main driver of the LASV's occurrence in only western and eastern West Africa. Though the mechanism by which rainfall affects viral prevalence is unclear, it has been hypothesized, for example, that wetter conditions might facilitate the virus' ability to survive outside the host [4].

Our model of spillover risk predicts a significant, but small amount of the spatial variation in arenavirus seroprevalence studies in humans. The modest relationship between human LASV spillover and predicted risk might be due to the binary classifiers' coarse description of the magnitude of LASV risk. As more longitudinal data become available, these binary models can be upgraded with more nuanced models that predict the time-varying density of *M. natalensis* and the prevalence of LASV among the rodent population. Alternatively, the low correlation could indicate that other predictors like human factors have a large influence on LASV spillover. Geographic differences in housing, cultural practices, and diet likely influence the extent of LASV spillover but are not included in our model. For instance, the use of rodent-proof housing materials (e.g., concrete vs mud) and abstaining from rodent hunting and consumption are known to affect the extent to which LASV is able to transmit between rodents and humans [16, 80]. The residuals of seroprevalence predictions from our model could help guide understanding of which human factors mitigate or facilitate LASV spillover. If human factors like housing type can be readily identified from serosurvey locations within West Africa, they could be incorporated in the human stage of the model that connects spillover risk to human seroprevalence.

Geographic variation in LASV and its primary reservoir may also be responsible for the modest fit of our model. For instance, across West Africa LASV consists of several clades [22]. If certain clades are better at infecting humans, then our model will tend to underestimate the rate of human infections in regions where such highly-infectious clades occur. Similarly, the *M. natalensis* reservoir is also divided into multiple clades [20]. Different *M. natalensis* clades may differ in their contact rates with humans or in their suitability as reservoir, further reducing our model's ability to predict spillover into humans. Some evidence for this latter possibility comes from arenavirsues that preferentially infect certain clades of *M. natalensis* [21]. Because our study region only includes West Africa, it is likely that the *M. natalensis* occurrences that our model is trained on are only from the A-I clade [20]. However, our forecast should be

interpreted with caution in eastern Nigeria, where the transition zone into the A-II ₅₃₉ clade occurs. Future work integrating these factors may help improve our understanding ₅₄₀ of the spatial variation in human seroprevalence that is due to the spatial patterning of ₅₄₁ LASV and reservoir clades. ₅₄₂

Another factor that could influence our model fit is the possibility that rodent ₅₄₃ species other than *M. natalensis* serve as reservoirs or interact with the primary ₅₄₄ reservoir in ways that decrease or increase risk. Though *M. natalensis* is believed to be ₅₄₅ the primary reservoir that contributes to human infection, several species of rodents are ₅₄₆ known to be capable of harboring the virus [48]. Understanding the relationship ₅₄₇ between the habitat suitability of different rodent reservoirs and human LF burden may ₅₄₈ help determine whether *M. natalensis* is the host at which intervention strategies should ₅₄₉ always be directed. Furthermore, other species of rodent may displace *M. natalensis* ₅₅₀ and therefore lower the overall spillover risk of LASV into humans. The layered ₅₅₁ framework we have developed can be easily adapted to include additional reservoir ₅₅₂ species and systematically investigate these possibilities. ₅₅₃

Our model is constructed to learn and explain spatial variation in the average ₅₅₄ historical spillover of LASV, and does not include temporal trends of spillover risk. Due ₅₅₅ to the sparsity of available longitudinal data, our model assumes that the human ₅₅₆ population in West Africa, human LASV seroprevalence, and the rate of LASV spillover, ₅₅₇ are all constant in time. Over decades-long timescales, the rate of LASV spillover is ₅₅₈ likely increasing due to increasing rates of human-rodent interaction that come with ₅₅₉ urban growth, deforestation, or climate change [11, 70]. Estimating the combined ₅₆₀ temporal and spatial variation of infection will require long-term longitudinal studies in ₅₆₁ both rodents and humans across West Africa. With this data, for example, more ₅₆₂ advanced models could mechanistically associate an increasing rate of spillover with ₅₆₃ changes to land cover. ₅₆₄

Another important temporal simplification of our current modeling work is the ₅₆₅ absence of seasonality in LASV spillover. In Sierra Leone, Guinea, and Nigeria, hospital ₅₆₆ admissions attributable to LASV infection generally peak late in the dry ₅₆₇ season [54, 69, 81]. In these regions, the mechanism of seasonal spillover likely involves a ₅₆₈ combination of seasonal rainfall and land use practices, such as crop-harvesting and ₅₆₉ subsequent burning of agricultural fields, that drive rodents into domestic dwellings in ₅₇₀

search of food-stuffs [54, 82]. It is not understood whether these factors operate   571
uniformly across all of West Africa. Temporal fluctuations in the density of the reservoir   572
population, due to seasonal cycles of reproduction, are another potentially important   573
factor that could drive a seasonal spike of human LF cases. However, it is unclear   574
whether the density fluctuations that have been observed outside of the LASV   575
geographic range (e.g., Tanzania [79]) also occur within West Africa. At least in Guinea   576
and Sierra Leone, research on the population dynamics of *M. natalensis* indicates that   577
density fluctuations are much weaker than those in East Africa [83]. In the case of   578
rodent vaccination, understanding population dynamics is particularly important   579
because distributing vaccines at seasonal population lows in wildlife demographic cycles   580
can, in theory, substantially increase the probability of pathogen elimination [83, 84].   581

Although the methods we have used here make efficient use of available data, the   582
accuracy of our risk forecasts remains difficult to rigorously evaluate due to the limited   583
availability of current data from human populations across West Africa. The sparseness   584
of modern human data arises for two reasons: 1) the lack of robust surveillance and   585
testing across much of the region where LASV is endemic and 2) the absence of publicly   586
available databases reporting human cases in those countries that do have relatively   587
robust surveillance in place (i.e., Nigeria). Improving surveillance for LASV across West   588
Africa and developing publicly available resources for sharing the resulting data would   589
allow more robust risk predictions to be developed and facilitate risk reducing   590
interventions. Despite these limitations of existing data, the structured   591
machine-learning models we develop here provide insight into what aspects of   592
environment, reservoir, and virus, contribute to spillover, and the potential risk of   593
subsequent emergence into the human population. By understanding these connections,   594
we can design and deploy more effective intervention and surveillance strategies that   595
work in tandem to reduce disease burden and enhance global health security.   596

Acknowledgements

References

1. Jones KE, Patel NG, Levy MA, Storeygard A, Balk D, Gittleman JL, et al. Global trends in emerging infectious diseases. Nature. 2008;451(7181):990–993. doi:10.1038/nature06536.

2. Plowright RK, Parrish CR, McCallum H, Hudson PJ, Ko AI, Graham AL, et al. Pathways to zoonotic spillover. Nat Rev Microbiol. 2017;15(8):502—-510. doi:10.1038/nrmicro.2017.45.

3. Hughes JM, Wilson ME, Pike BL, Saylors KE, Fair JN, LeBreton M, et al. The origin and prevention of pandemics. Clin Infect Dis. 2010;50(12):1636–1640. doi:10.1086/652860.

4. Fichet-Calvet E, Rogers DJ. Risk maps of Lassa fever in West Africa. PLoS Negl Trop Dis. 2009;3(3):e388. doi:10.1371/journal.pntd.0000388.

5. Mylne AQ, Pigott DM, Longbottom J, Shearer F, Duda KA, Messina JP, et al. Mapping the zoonotic niche of Lassa fever in Africa. Trans R Soc Trop Med Hyg. 2015;109(8):483–492. doi:10.1093/trstmh/trv047.

6. Pigott DM, Deshpande A, Letourneau I, Morozoff C, Reiner Jr RC, Kraemer MU, et al. Local, national, and regional viral haemorrhagic fever pandemic potential in Africa: a multistage analysis. Lancet. 2017;390(10113):2662–2672. doi:10.1016/S0140-6736(17)32092-5.

7. Pigott DM, Golding N, Mylne A, Huang Z, Henry AJ, Weiss DJ, et al. Mapping the zoonotic niche of Ebola virus disease in Africa. Elife. 2014;3:e04395. doi:10.7554/eLife.04395.001.

8. Pigott DM, Millear AI, Earl L, Morozoff C, Han BA, Shearer FM, et al. Updates to the zoonotic niche map of Ebola virus disease in Africa. Elife. 2016;5:e16412. doi:10.7554/eLife.16412.

9. Pigott DM, Bhatt S, Golding N, Duda KA, Battle KE, Brady OJ, et al. Global distribution maps of the leishmaniases. Elife. 2014;3:e02851. doi:10.7554/eLife.02851.001.

10. Bond N, Schieffelin JS, Moses LM, Bennett AJ, Bausch DG. A historical look at the first reported cases of Lassa fever: IgG antibodies 40 years after acute infection. Am J Trop Med Hyg. 2013;88(2):241–244. doi:10.4269/ajtmh.12-0466.

11. Gibb R, Moses LM, Redding DW, Jones KE. Understanding the cryptic nature of Lassa fever in West Africa. Pathog Glob Health. 2017;111(6):276–288. doi:10.1080/20477724.2017.1369643.

12. Peterson TA, Moses LM, Bausch DG. Mapping transmission risk of Lassa fever in West Africa: the importance of quality control, sampling bias, and error weighting. PLoS One. 2014;9(8):e100711. doi:10.1371/journal.pntd.0000388.

13. Maes P, Alkhovsky SV, Bào Y, Beer M, Birkhead M, Briese T, et al. Taxonomy of the family Arenaviridae and the order Bunyavirales: update 2018. Arch Virol. 2018;163(8):2295–2310. doi:10.1007/s00705-018-3843-5.

14. Fisher-Hoch S, Tomori O, Nasidi A, Perez-Oronoz G, Fakile Y, Hutwagner L, et al. Review of cases of nosocomial Lassa fever in Nigeria: the high price of poor medical practice. BMJ. 1995;311(7009):857–859. doi:10.1136/bmj.311.7009.857.

15. Iacono GL, Cunningham AA, Fichet-Calvet E, Garry RF, Grant DS, Khan SH, et al. Using modelling to disentangle the relative contributions of zoonotic and anthroponotic transmission: the case of Lassa fever. PLoS Negl Trop Dis. 2015;9(1):e3398. doi:10.1371/journal.pntd.0003398.

16. Ter Meulen J, Lukashevich I, Sidibe K, Inapogui A, Marx M, Dorlemann A, et al. Hunting of peridomestic rodents and consumption of their meat as possible risk factors for rodent-to-human transmission of Lassa virus in the Republic of Guinea. Am J Trop Med Hyg. 1996;55(6):661–666. doi:10.4269/ajtmh.1996.55.661.

17. McCormick JB, Webb PA, Krebs JW, Johnson KM, Smith ES. A prospective study of the epidemiology and ecology of Lassa fever. J Infect Dis. 1987;155(3):437–444. doi:10.1093/infdis/155.3.437.

18. United Nations. Geographical Regions; 2020.
    https://unstats.un.org/unsd/methodology/m49/#geo-regions.

19. Granjon L. The IUCN red list of threatened species 2016: e.T12868A115107375.
    2016;doi:10.2305/IUCN.UK.2016-3.RLTS.T12868A22425266.en.

20. Colangelo P, Verheyen E, Leirs H, Tatard C, Denys C, Dobigny G, et al. A
    mitochondrial phylogeographic scenario for the most widespread African rodent,
    Mastomys natalensis. Biol J Linn Soc. 2013;108(4):901–916.
    doi:10.1111/bij.12013.

21. Gryseels S, Baird SJ, Borremans B, Makundi R, Leirs H, Goüy de Bellocq J.
    When viruses don't go viral: the importance of host phylogeographic structure in
    the spatial spread of arenaviruses. PLoS Pathog. 2017;13(1):e1006073.
    doi:10.1371/journal.ppat.1006073.

22. Olayemi A, Fichet-Calvet E. Systematics, Ecology, and Host Switching:
    Attributes Affecting Emergence of the Lassa Virus in Rodents across Western
    Africa. Viruses. 2020;12(3):312. doi:10.3390/v12030312.

23. Basinski AJ. Pathogen Spillover Forecast; 2020. Github repository
    https://github.com/54481andrew/pathogen-spillover-forecast.git.

24. Wilson DE, Reeder DM. Mammal species of the world: a taxonomic and
    geographic reference. vol. 1. JHU Press; 2005.

25. Happold D, Happold D. Mammals of Africa. Volume III: Rodents, hares and
    rabbits. 2013;.

26. Granjon L, Duplantier JM, Catalan J, Britton-Davidian J. Systematics of the
    genus Mastomys (Thomas, 1915)(Rodentia: Muridae). A review. Belgian Journal
    of Zoology (Belgium). 1997;.

27. Van de Perre F, Adriaensen F, Terryn L, Pauwels O, Leirs H, Gilissen E, et al..
    African mammalia; 2019.
    http://projects.biodiversity.be/africanmammalia.

28. Centre de Biologie pour la Gestion des Populations. Database on Sahelo-Sudanian rodents; 2017. `http://vminfotron-dev.mpl.ird.fr/bdrss/bdrsspub_form.php`.

29. Brouat C, Loiseau A, Kane M, Bâ K, Duplantier JM. Population genetic structure of two ecologically distinct multimammate rats: the commensal *Mastomys natalensis* and the wild *Mastomys erythroleucus* in southeastern Senegal. Mol Ecol. 2007;16(14):2985–2997. doi:10.1111/j.1365-294X.2007.03353.x.

30. Coulibaly-N'Golo D, Allali B, Kouassi SK, Fichet-Calvet E, Becker-Ziaja B, Rieger T, et al. Novel arenavirus sequences in *Hylomyscus* sp. and *Mus (Nannomys) setulosus* from Côte d'Ivoire: implications for evolution of arenaviruses in Africa. PloS One. 2011;6(6):e20893. doi:10.1371/journal.pone.0020893.

31. Dobigny G, Nomao A, Gautun J. A cytotaxonomic survey of rodents from Niger: implications for systematics, biodiversity and biogeography. Mammalia. 2002;66(4):495–524. doi:10.1515/mamm.2002.66.4.495.

32. Duplantier JM, Britton-Davidian J, Granjon L. Chromosomal characterization of three species of the genus Mastomys in Senegal. J Zool Syst Evol Res. 1990;28(4):289–298. doi:10.1111/j.1439-0469.1990.tb00383.x.

33. Granjon L, Duplantier JM. Les rongeurs de l'Afrique sahélo-soudanienne; 2009.

34. Kronmann KC, Nimo-Paintsil S, Guirguis F, Kronmann LC, Bonney K, Obiri-Danso K, et al. Two novel arenaviruses detected in pygmy mice, Ghana. Emerg Infect Dis. 2013;19(11):1832. doi:10.3201/eid1911.121491.

35. Lecompte E, Brouat C, Duplantier JM, Galan M, Granjon L, Loiseau A, et al. Molecular identification of four cryptic species of Mastomys (Rodentia, Murinae). Biochemical Systematics and Ecology. 2005;33(7):681–689. doi:10.1016/j.bse.2004.12.015.

36. Lecompte E, Fichet-Calvet E, Daffis S, Koulémou K, Sylla O, Kourouma F, et al. Mastomys natalensis and lassa fever, West Africa. Emerg Infect Dis. 2006;12(12):1971. doi:10.3201/eid1212.060812.

37. Olayemi A, Obadare A, Oyeyiola A, Fasogbon S, Igbokwe J, Igbahenah F, et al. Small mammal diversity and dynamics within Nigeria, with emphasis on reservoirs of the Lassa virus. System Biodivers. 2018;16(2):118–127. doi:10.1080/14772000.2017.1358220.

38. Robbins CB, Krebs Jr JW, Johnson KM. Mastomys (Rodentia: Muridae) species distinguished by hemoglobin pattern differences. Am J Trop Med. 1983;32(3):624–630. doi:10.4269/ajtmh.1983.32.624.

39. Yadouleton A, Agolinou A, Kourouma F, Saizonou R, Pahlmann M, Bedié SK, et al. Lassa Virus in Pygmy Mice, Benin, 2016–2017. Emerg Infect Dis. 2019;25(10):1977. doi:10.3201/eid2510.180523.

40. Olayemi A, Akinpelu A. Diversity and distribution of murid rodent populations between forest and derived savanna sites within south western Nigeria. Biodivers Conserv. 2008;17(10):2411. doi:10.1007/s10531-008-9389-1.

41. Safronetz D, Lopez JE, Sogoba N, Traore SF, Raffel SJ, Fischer ER, et al. Detection of Lassa virus, Mali. Emerg Infect Dis. 2010;16(7):1123. doi:10.3201/eid1607.100146.

42. Safronetz D, Sogoba N, Lopez JE, Maiga O, Dahlstrom E, Zivcec M, et al. Geographic distribution and genetic characterization of Lassa virus in sub-Saharan Mali. PLoS Negl Trop Dis. 2013;7(12). doi:10.1371/journal.pntd.0002582.

43. Elith J, Graham CH, Anderson RP, Dudík M, Ferrier S, Guisan A, et al. Novel methods improve prediction of species' distributions from occurrence data. Ecography. 2006;29(2):129–151. doi:10.1111/j.2006.0906-7590.04596.x.

44. Barbet-Massin M, Jiguet F, Albert CH, Thuiller W. Selecting pseudo-absences for species distribution models: how, where and how many? Methods Ecol Evol. 2012;3(2):327–338. doi:10.1111/j.2041-210X.2011.00172.x.

45. Phillips SJ, Dudík M, Elith J, Graham CH, Lehmann A, Leathwick J, et al. Sample selection bias and presence-only distribution models: implications for

background and pseudo-absence data. Ecol Appl. 2009;19(1):181–197. doi:10.1890/07-2153.1.

46. GBIF occurrence download; 2020-10-30. `https://doi.org/10.15468/dl.tbe47y`.

47. Benson DA, Cavanaugh M, Clark K, Karsch-Mizrachi I, Lipman DJ, Ostell J, et al. GenBank. Nucleic Acids Res. 2017;45(D1):D37–D42. doi:10.1093/nar/gkw1070.

48. Fichet-Calvet E, Becker-Ziaja B, Koivogui L, Günther S. Lassa serology in natural populations of rodents and horizontal transmission. Vector Borne Zoonotic Dis. 2014;14(9):665–674. doi:10.1089/vbz.2013.1484.

49. Fichet-Calvet E, Ölschläger S, Strecker T, Koivogui L, Becker-Ziaja B, Camara AB, et al. Spatial and temporal evolution of Lassa virus in the natural host population in Upper Guinea. Scientific reports. 2016;6(1):1–6. doi:10.1038/srep21977.

50. Leski TA, Stockelman MG, Moses LM, Park M, Stenger DA, Ansumana R, et al. Sequence variability and geographic distribution of Lassa virus, Sierra Leone. Emerg Infect Dis. 2015;21(4):609. doi:10.3201/eid2104.141469.

51. Monath TP, Newhouse VF, Kemp GE, Setzer HW, Cacciapuoti A. Lassa virus isolation from *Mastomys natalensis* rodents during an epidemic in Sierra Leone. Science. 1974;185(4147):263–265. doi:10.1126/science.185.4147.263.

52. Olayemi A, Oyeyiola A, Obadare A, Igbokwe J, Adesina AS, Onwe F, et al. Widespread arenavirus occurrence and seroprevalence in small mammals, Nigeria. Parasites Vectors. 2018;11(1):416. doi:10.1186/s13071-018-2991-5.

53. Wulff H, Fabiyi A, Monath T. Recent isolations of Lassa virus from Nigerian rodents. Bull World Health Organ. 1975;52(4-6):609.

54. Fichet-Calvet E, Lecompte E, Koivogui L, Soropogui B, Doré A, Kourouma F, et al. Fluctuation of abundance and Lassa virus prevalence in *Mastomys natalensis* in Guinea, West Africa. Vector Borne Zoonotic Dis. 2007;7(2):119–128. doi:10.1089/vbz.2006.0520.

55. Kernéis S, Koivogui L, Magassouba N, Koulemou K, Lewis R, Aplogan A, et al. Prevalence and risk factors of Lassa seropositivity in inhabitants of the forest region of Guinea: a cross-sectional study. PLoS Negl Trop Dis. 2009;3(11). doi:10.1371/journal.pntd.0000548.

56. Lukashevich I, Clegg J, Sidibe K. Lassa virus activity in Guinea: Distribution of human antiviral antibody defined using enzyme-linked immunosorbent assay with recombinant antigen. J Med Virol. 1993;40(3):210–217. doi:10.1002/jmv.1890400308.

57. Nimo-Paintsil SC, Fichet-Calvet E, Borremans B, Letizia AG, Mohareb E, Bonney JH, et al. Rodent-borne infections in rural Ghanaian farming communities. PloS One. 2019;14(4). doi:10.1371/journal. pone.0215224.

58. Sogoba N, Rosenke K, Adjemian J, Diawara SI, Maiga O, Keita M, et al. Lassa virus seroprevalence in sibirilia commune, Bougouni District, Southern Mali. Emerg Infect Dis. 2016;22(4):657. doi:10.3201/eid2204.151814.

59. Yalley-Ogunro J, Frame J, Hanson A. Endemic Lassa fever in Liberia. VI. Village serological surveys for evidence of Lassa virus activity in Lofa County, Liberia. Trans R Soc Trop Med Hyg. 1984;78(6):764–770. doi:10.1016/0035-9203(84)90013-0.

60. Fraser DW, Campbell CC, Monath TP, Goff PA, Gregg MB. Lassa fever in the eastern province of Sierra Leone, 1970–1972. Am J Trop Med Hyg. 1974;23(6):1131–1139. doi:10.4269/ajtmh.1974.23.1131.

61. Leirs H, Verhagen R, Verheyen W. Productivity of different generations in a population of *Mastomys natalensis* rats in Tanzania. Oikos. 1993; p. 53–60. doi:10.2307/3545308.

62. Hastie T, Tibshirani R, Friedman J. The elements of statistical learning: data mining, inference, and prediction. Springer Science & Business Media; 2009.

63. Elith J, Leathwick JR, Hastie T. A working guide to boosted regression trees. J Anim Ecol. 2008;77(4):802–813. doi:10.1111/j.1365-2656.2008.01390.x.

64. Hijmans RJ, Phillips S, Leathwick J, Elith J. dismo: Species Distribution Modeling; 2017. Available from: `https://CRAN.R-project.org/package=dismo`.

65. Hijmans RJ. Cross-validation of species distribution models: removing spatial sorting bias and calibration with a null model. Ecology. 2012;93(3):679–688. doi:10.1890/11-0826.1.

66. McCullagh P, Nelder JA. Generalized linear models. 2nd ed. CRC Press; 1989.

67. Fisher-Hoch S, Hutwagner L, Brown B, McCormick J. Effective vaccine for Lassa fever. Journal of virology. 2000;74(15):6777–6783. doi:10.1128/JVI.74.15.6777-6783.2000.

68. WorldBank. Life expectancy at birth, total (years); 2020. `https://data.worldbank.org/indicator/SP.DYN.LE00.IN`.

69. Shaffer JG, Grant DS, Schieffelin JS, Boisen ML, Goba A, Hartnett JN, et al. Lassa fever in post-conflict Sierra Leone. PLoS Negl Trop Dis. 2014;8(3):e2748. doi:10.1371/journal.pntd.0002748.

70. Redding DW, Moses LM, Cunningham AA, Wood J, Jones KE. Environmental-mechanistic modelling of the impact of global change on human zoonotic disease emergence: a case study of Lassa fever. Methods Ecol Evol. 2016;7(6):646–655. doi:10.1111/2041-210X.12549.

71. Antia R, Regoes RR, Koella JC, Bergstrom CT. The role of evolution in the emergence of infectious diseases. Nature. 2003;426(6967):658–661. doi:10.1038/nature02104.

72. Richmond JK, Baglole DJ. Lassa fever: epidemiology, clinical features, and social consequences. BMJ. 2003;327(7426):1271–1275. doi:10.1136/bmj.327.7426.1271.

73. CDC. NCDC Lassa cases; 2020. `https://ncdc.gov.ng/data`.

74. Yun NE, Walker DH. Pathogenesis of Lassa fever. Viruses. 2012;4(10):2031–2048. doi:10.3390/v4102031.

75. Worldometers. Western Africa Population;. https://www.worldometers.info/world-population/western-africa-population/.

76. Cross M, Buddle B, Aldwell F. The potential of oral vaccines for disease control in wildlife species. Vet J. 2007;174(3):472–480. doi:10.1016/j.tvjl.2006.10.005.

77. Murphy AA, Redwood AJ, Jarvis MA. Self-disseminating vaccines for emerging infectious diseases. Expert Rev Vaccines. 2016;15(1):31–39. doi:10.1586/14760584.2016.1106942.

78. Freuling CM, Hampson K, Selhorst T, Schröder R, Meslin FX, Mettenleiter TC, et al. The elimination of fox rabies from Europe: determinants of success and lessons for the future. Philos Trans R Soc Lond, B, Biol Sci. 2013;368(1623):20120142. doi:10.1098/rstb.2012.0142.

79. Leirs H, Verheyen W, Michiels M, Verhagen R, Stuyck J. The relation between rainfall and the breeding season of *Mastomys natalensis* (Smith, 1834) in Morogoro, Tanzania. In: Annales de la société Royale Zoologique de Belgique. vol. 119; 1989. p. 59–64.

80. Bonner PC, Schmidt WP, Belmain SR, Oshin B, Baglole D, Borchert M. Poor housing quality increases risk of rodent infestation and Lassa fever in refugee camps of Sierra Leone. Am J Trop Med Hyg. 2007;77(1):169–175. doi:10.4269/ajtmh.2007.77.169.

81. Bausch DG, Demby AH, Coulibaly M, Kanu J, Goba A, Bah A, et al. Lassa fever in Guinea: I. Epidemiology of human disease and clinical observations. Vector Borne Zoonotic Dis. 2001;1(4):269–281. doi:10.1089/15303660160025903.

82. Akhmetzhanov AR, Asai Y, Nishiura H. Quantifying the seasonal drivers of transmission for Lassa fever in Nigeria. Philos Trans R Soc Lond, B, Biol Sci. 2019;374(1775):20180268. doi:10.1098/rstb.2018.0268.

83. Nuismer SL, Remien CH, Basinski AJ, Varrelman T, Layman N, Rosenke K, et al. Bayesian estimation of Lassa virus epidemiological parameters: Implications for spillover prevention using wildlife vaccination. PLoS Negl Trop Dis. 2020;14(9):e0007920. doi:10.1371/journal.pntd.0007920.

84. Schreiner CL, Nuismer SL, Basinski AJ. When to vaccinate a fluctuating wildlife population: is timing everything? J Appl Ecol. 2020;doi:10.1111/1365-2664.13539.

Supporting information

S1 Appendix   Details on the predictors used in the model and model fits.