A method for achieving complete microbial genomes and improving

bins from metagenomics data

Authors:

Lauren M. Lui[1], Torben N. Nielsen[1], Adam P. Arkin[1,2,3]*

Affiliations:

[1]Environmental Genomics and Systems Biology Division, Lawrence Berkeley National

Laboratory, Berkeley, CA, USA.

[2]Department of Bioengineering, University of California, Berkeley, CA, USA

[3]Innovative Genomics Institute, Berkeley, CA, USA

*Correspondence: aparkin@lbl.gov

20
21

Abstract

Metagenomics facilitates the study of the genetic information from uncultured microbes and complex microbial communities. Assembling complete microbial genomes (*i.e.*, circular with no misassemblies) from metagenomics data is difficult because most samples have high organismal complexity and strain diversity. Less than 100 circularized bacterial and archaeal genomes have been assembled from metagenomics data despite the thousands of datasets that are available. Circularized genomes are important for (1) building a reference collection as scaffolds for future assemblies, (2) providing complete gene content of a genome, (3) confirming little or no contamination of a genome, (4) studying the genomic context and synteny of genes, and (5) linking protein coding genes to ribosomal RNA genes to aid metabolic inference in 16S rRNA gene sequencing studies. We developed a method to achieve circularized genomes using iterative assembly, binning, and read mapping. In addition, this method exposes potential misassemblies from k-mer based assemblies. We chose species of the Candidate Phyla Radiation (CPR) to focus our initial efforts because they have small genomes and are only known to have one ribosomal RNA operon. We present 34 circular CPR genomes, one circular Margulisbacteria genome, and two circular megaphage genomes from 19 public and published datasets. We demonstrate findings that would likely be difficult without circularizing genomes, including that ribosomal genes are likely not operonic in the majority of CPR, and that some CPR harbor diverged forms of RNase P RNA. Code and a tutorial for this method is available at https://github.com/lmlui/Jorg.

Introduction

Shotgun metagenomics and marker gene sequencing are powerful tools to survey and study organisms that we cannot yet isolate and culture in the laboratory. This is especially true for environmental samples where culturability estimates for bacterial and archaeal communities range from ~22-53% for soil, ~10-70% for ocean and lakes, and ~8-32% for ocean sediment [1]. Scientists have turned to shotgun metagenomics to provide genome-resolved analysis of complex samples, but assembling genomes from shotgun metagenomics data is inherently more difficult than assembling those from cultured isolates. Challenges in metagenomics assembly arise from the heterogeneity of samples, available sequencing technology, and the limitations of bioinformatics algorithms we use for assembly and genome binning. Metagenomes contain uneven amounts of an unknown number of genomes, which creates a compounded computational problem in terms of simplifying assumptions, time, and computer memory.

In the 1990s when the first genomes were sequenced and assembled, scientists used long reads from Sanger sequencing and overlap layout consensus (OLC) methods for assembly [2]. With the development of next-generation sequencing technologies, we gained the ability to sequence millions of reads at a massively reduced cost, but using traditional OLC algorithms became too computationally intensive. The computational complexity of OLC algorithms scale as the square of the number of input reads (because each read is compared to every other read), so they are impractical for datasets of millions of reads, compared to the thousands of reads generated from Sanger sequencing. To handle the deluge of sequencing data (in terms of the volume of reads and projects) de Bruijn graph assembly methods were developed. The time and memory complexity of de Bruijn based assembly algorithms typically scale with the size of the metagenome instead of the number of reads. However, de Bruijn graph methods can introduce misassemblies; due to the decomposition of reads into k-mers, context is lost and it is

possible for the graphs to contain paths that do not correspond to real genomic sequence [3,4].

Traditional OLC assemblers such as the Celera Assembler [5], SGA [6] and MIRA [7] ensure

that only contigs consistent with actual genome sequence are produced (this is sometimes

referred to as maintaining read coherence). Using long-read sequencing can overcome some of

the issues with k-mer based assembly and newer assemblers for this type of data have started

using OLC assembly methods again [3]. However, long-read sequencing requires much larger

amounts of DNA (micrograms) that is high quality and high molecular weight (10-50kb) as

compared to short read technologies (as little as 1 ng). Extracting enough high molecular weight

DNA can be limited by sampling costs and available biomass, especially for certain types of

environmental samples, so short read sequencing may sometimes be the only option.

Beyond assembly, a key challenge with metagenomics is grouping contigs into genome

bins. We use "contig" in the way it was originally defined by Rodger Staden, where a contig is a

set of overlapping segments of DNA from shotgun sequencing [8]. It is rare for a complete

genome to be assembled into a single piece *de novo* from short reads, so contigs are grouped

into "bins," often based on coverage and tetranucleotide frequencies. If two contigs belong to

the same genome, they are expected to have similar coverage and tetranucleotide profiles [9].

However, coverage has problems for multiple reasons. If a particular microbe is growing rapidly,

some regions may have higher coverage than the rest of the genome [10]. In addition, for

organisms where the copy number of ribosomal RNA (rRNA) operons exceeds unity, the

contig(s) with the rRNA genes will not have the same coverage as the rest of the contigs in the

genome. This is also true of other multi-copy genes and other repetitive elements.

Tetranucleotide frequencies are problematic because horizontally transferred regions may have

different frequencies than the rest of the genome [11] and this can result in such pieces being

put into different bins by the binning algorithm. Despite these issues, binning is helpful in

identifying potential genomes in metagenomics data, especially when using short read

sequencing technologies.

To evaluate the quality of a bin, the metrics of "contamination" and "completeness" are

often used. Completeness and contamination are detected generally by looking for violations of

conserved features of complete isolate genomes. Such features include having complete sets of

universally (or at least phylogenetically) conserved single-copy protein genes without any

duplication or excessive variation in tetranucleotide frequency. Other measures of completeness

have been suggested, such as establishment of a core conserved set of ubiquitous genes.

Tools such as RefineM [12] and CheckM [13] apply these rules to assemblies to determine

completeness and contamination. However, these tools are not always accurate for species

that are not well studied. Candidate Phyla Radiation (CPR) species are often classified as

having 60-80% completeness by these tools, even for circular genomes. To overcome

challenges of binning, scientists have started to assemble circular, complete genomes from

metagenomes [14–23], which are also called CMAGs (complete metagenomic-assembled

genomes) [24]. In comparison to genome bins, a high quality reference collection that controls

for misassemblies and is composed of circular genomes (1) provides more accurate inference

of identity and estimation of capabilities of uncultured microbes within complex microbiomes, (2)

allows more accurate taxonomic assessment of the composition of these microbiomes through

better linkage of marker genes in single organisms, (3) provides high-quality scaffolds on which

reads can be assembled, both to allow measures of strain variation within a microbiome study

and to aid in better assembly of reads across many microbiome samples, and (4) affords the

ability to study synteny and genomic context of genes in these organisms. In addition, while

there are existing methods for generating high-quality MAGs, there is evidence that these MAGs

still contain significant contamination by exogenous sequence and have misassemblies

triggered by lack of read coherence. Circularization of genomes helps assure that there is likely

no contamination in the assemblies. Despite the advantages of circularizing genomes, very few

metagenomics studies to date (<30) have published circular genomes [14,15,24,16–23].

We describe a semi-automated method that facilitates recovery of circular archaeal and

bacterial genomes from metagenomics data and that also provides checks for misassemblies.

In this study we do not focus on how many circularized genomes we can get from a dataset, but

rather the method itself to help circularize bins of interest and to ensure that they are high

quality. To assist with the travails of circularizing genomes, our method overcomes issues from

using k-mer based assembly and automates iterative extension of contigs. Our general

approach is to produce a "standard" metagenomic assembly, bin using a "standard" binning tool,

extract reads based on k-mer similarity and reassemble these using a "standard" isolate-

focused assembler. To demonstrate this method, we have obtained 34 circular CPR genomes,

one circular Margulisbacteria genome, and two circular megaphage genomes from 19 public

and published metagenomics datasets. To our knowledge, only 41 other CPR circularized

genomes have been published from 9 studies [14–22], so we believe this to be the largest

presentation of circularized CPR genomes in a single study. With this set, we demonstrate

findings that would likely be difficult without a large number of unique circularized genomes,

including that ribosomal genes are likely not operonic in the majority of CPR and finding

diverged forms of RNase P RNA in CPR species.

Results

Circularization Method

To have confidence that the genomes we generate match real organisms, we looked for

criteria that would indicate that a genome is circular and complete. The literature is replete with

techniques and proposals for measuring the completeness of genomes and to what level they

are complete [13,16], but these often have difficulty when encountering novel genomes because

their criteria are based on known isolate genomes. We focus on evidence for incompleteness in

terms of missing essential genes that are found across the tree of life. That is, we are more

concerned with ensuring that anything we label a circularized genome meets basic criteria that

indicates that it is not incomplete. We posit that a complete, circularized genome must satisfy at

least the following conditions:

1. The genome is either circular or there is solid evidence that it is linear. While rare, linear

bacterial genomes exist [25].

2. The genome has a full complement of rRNAs (16S, 23S, 5S), transfer RNAs (all amino

acids represented), and RNase P RNA (since this is nearly universally necessary to

process tRNA transcripts). Absence of any of these genes must be explained. We

advocate using these as a check for a complete genome instead of single copy marker

protein genes because checks for single copy marker protein genes can vary by clade;

in only rare instances would these noncoding RNA genes be missing [26,27].

3. There is significant read coverage across the entire genome. Assemblies that rely on

single reads for continuity are prone to error. With some exceptions for very high

coverage organisms, we generally require minimum coverage no lower than 30% of the

average coverage.

To develop and test this method, we mined the Sequence Read Archive (SRA) hosted at the

National Center for Biotechnology Information (NCBI) [28] for metagenomic sequencing data

generated from groundwater samples, where CPR are prevalent (see Table 1 for a list of the

datasets used in this study. We focused on assembling CPR genomes because they are (1)

small and thus easier to assemble, (2) to the best of our knowledge only have one set of rRNA

genes. These two criteria gave us the easiest targets for circularization.

Table 1: Description of Metagenomes in this Study.

We focused on groundwater datasets because they have a higher fraction of CPR. Many of the datasets

are from studies of anthropologically contaminated sites. All identifiers are SRA except for SURF_D.

Identifiers	Study Description	Reference
ERX2165959	Groundwater from monitoring wells from naphthalene contaminated surface sediments, where effluent from the coal-tar contaminated groundwater surfaces	[29]
SRX1085364	Terrestrial subsurface C, N, S and H cycles cross-linked by metabolic handoffs	[16]
SRX1775573 SRX1775577 SRX1775579	Potential for microbial H2 and metal transformations associated with novel bacteria and archaea in deep terrestrial subsurface sediments	[21]
SRX1990955	Groundwater microbial communities from Rifle, Colorado - Rifle Oxygen_injection A2 metagenome	[16]
SRX2838984	Coupling Microbial Communities to Carbon and Contaminant Biogeochemistry in the Groundwater-Surface Water Interaction Zone	[30–32]
SRX3024504 SRX3024507 SRX3024508	DNA from groundwater after nitrate injection, filter size 0.2µm and 0.1µm	[33]
SRX3307784	Subsurface groundwater microbial communities from S. Glens Falls, New York, USA - GMW37 contaminated, 5.8 m metagenome	[29]
SRX3348993	Development of a pipeline for high-throughput recovery of near-complete and complete microbial genomes from complex metagenomic datasets: Groundwater sample from aquifer - Crystal Geyser CG19_WC_8/21/14_NA	[34]
SRX3574179	Investigating microbial roles in methane emission, contaminant degradation, and biogeochemical cycles in an aquifer near a municipal landfill	Laura Hug Lab; https://uwaterloo.ca/hug-research-group/
SRX3602289 SRX3602720	Groundwater microbial communities from the Aspo Hard Rock Laboratory (HRL) deep subsurface site, Sweden	Mark Dopson Lab; https://lnu.se/en/staff/mark.dopson/
SURF_D	Groundwater samples from the Sanford Underground Research Facility (SURF)	[35]

The first steps of the method are standard to regular metagenomics assembly pipelines.

For each metagenome, we trimmed the reads to remove any remaining Illumina adapter

fragments and low quality ends, as well as whole reads that weren't of sufficient quality, using

BBtools (Fig 1A). Next, we assembled the processed reads using SPAdes [36] (Fig 1B). We

proceeded with successful assemblies and used MetaBAT 2 [37] (Fig 1C) to produce a

collection of bins for each. We went through 188 assembled metagenomes and picked bins with

5 or fewer contigs and coverage above 40X, although we made exceptions for bins that looked

promising, such as a bin with many contigs, but with one or two large contigs that comprise

most of the bin's sequence length (Table 2). We used GTDB-Tk [38] to classify the bins and

picked a set of CPR bins. We used these bins as "bait" to select read pairs for use with the

isolate-focused assembler MIRA (Fig 1D).

Fig 1. General method for circularizing genomes from metagenomes.

(A) Reads have adapters trimmed and low quality reads are filtered using BBtools. (B)

Processed reads are assembled into contigs using SPAdes. (C) Contigs are grouped into bins

using MetaBAT 2. (D) After choosing a bin for circularization, reads mapping to the bin are

extracted from the original processed reads and used as input into (E) where they are

assembled into contigs using MIRA. (F) Steps D and E are repeated as necessary until the bin

is deemed to be in a Circularization, Idempotence, or Chaos state. (G) If a bin is deemed

circular, we do final checks for misassemblies using Pilon and for the presence of rRNA, tRNAs,

and RNase P RNA before officially calling the bin a circularized genome.

Read quality filtering and trimming (BBtools) **(A)**

Assembly (SPAdes) **(B)**

Binning (MetaBAT2) **(C)**

Extract reads mapping to a bin (mirabait) **(D)**

Reassemble reads (MIRA) **(E)**

Repeat until **(F)**

1) Circularization

2) Idempotence

3) Chaos

Final Checks
Check for misassemblies using Pilon
Check for presence of tRNAs, RNase P RNA, rRNAs **(G)**

Table 2: List of 35 circularized bacterial genomes in this study.

Thirty-four of the genomes are classified as CPR and one is classified as a Margulisbacteria by GTDB-Tk.

The coverage, original number of contigs, and length of the original bin is also included.

Genome ID	GTDB-TK Taxonomy (Class)	Genome Size (bp)	Original Bin Stats		
			Num of Contigs	Coverage	Size of Bin (bp)
ERX2165959_bin_184	Paceibacteria	523910	2	126X	531828
ERX2165959_bin_23	Microgenomatia	1147419	1	134X	1146985
ERX2165959_bin_53	Microgenomatia	646630	3	116X	780569
ERX2165959_bin_80	Microgenomatia	1014979	1	216X	1024366
SRX1085364_bin_95	Microgenomatia	819458	4	109X	826172
SRX1775573_bin_5	Gracilibacteria	998919	1	104X	999239
SRX1775577_bin_36	Gracilibacteria	999108	1	74X	998727
SRX1775579_bin_0	Dojkabacteria	732899	1	51X	733907
SRX1990955_bin_0	Margulisbacteria (phylum); WOR-1	1676518	1	56X	1673447
SRX1990959_bin_38	Paceibacteria	585024	2	35X	582583
SRX2838984_bin_5	Paceibacteria	1030062	7	274X	1030337
SRX3024504_bin_47	Paceibacteria	672946	1	41X	673298
SRX3024507_bin_14	ABY1	1064268	4	48X	1106392
SRX3024507_bin_96	Paceibacteria	672946	1	29X	673073
SRX3024508_bin_27	Paceibacteria	672946	1	91X	673184
SRX3307784_bin_186	Paceibacteria	581622	4	174X	578873
SRX3307784_bin_197	Paceibacteria	822324	8	203X	962091
SRX3307784_bin_224	Microgenomatia	646579	3	118X	780569
SRX3307784_bin_45	UBA1384	872881	1	93X	872947
SRX3307784_bin_80	Microgenomatia	1013439	1	220X	1024366
SRX3307784_bin_91	Paceibacteria;	523446	3	129X	531688
SRX3348993_bin_93	Saccharimonadia	1005778	1	64X	1005352
SRX3574179_bin_116	Saccharimonadia	949592	4	37X	1000785
SRX3574179_bin_12	ABY1	1027227	1	135X	1028393
SRX3574179_bin_242	Microgenomatia	1127729	3	48X	1130277
SRX3574179_bin_244	Paceibacteria	707009	1	106X	708095
SRX3574179_bin_38	Paceibacteria	682291	5	124X	680516
SRX3574179_bin_63	ABY1	1209998	2	88X	1209921
SRX3574179_bin_75	ABY1	954552	3	152X	979051
SRX3602289_bin_51	Microgenomatia	852061	2	94X	853126
SRX3602720_bin_127	Paceibacteria	750123	18	160X	826132
SRX3602720_bin_74	ABY1	1035066	3	32X	1041342
SRX5650846_bin_20	Microgenomatia	947771	9	53X	949620
SURF_D_bin_21	Microgenomatia	978848	3	193X	978553
SURF_D_bin_31	Microgenomatia	1496927	5	159X	1501138

The purpose for assembling contigs first with SPAdes and then switching to MIRA with a

subset of reads is that the computational requirements of MIRA make it impractical as a

metagenome assembler. This is in part because MIRA does full alignment of the reads during

assembly. We would like to note that OLC metagenomics assemblers exist [3], but their memory

and time requirements are high compared to SPAdes or are not appropriate for assembling

bacterial genomes with paired-end sequencing data. MIRA has been used to extract

mitochondrial genomes [39] from eukaryotic sequencing projects. Our approach is very similar;

instead of providing seed sequences to separate the mitochondrial genomes from the

eukaryotic DNA, we use bins as seeds to separate genomes from the entire metagenomics

dataset. In our experience, MIRA produces superior results for isolates, and it also provides

additional features that benefit our method. MIRA comes with the tool mirabait, which provides

support for extracting read pairs based on k-mer content. MIRA also has a variety of features

that help expose problematic parts of assemblies. For example, MIRA sets tags to indicate parts

of the assembly that may require manual intervention, based on changes in coverage, GC, and

other anomalies. These tags are extremely useful in conjunction with traditional assembly

finishing tools such as Gap4/Gap5 [40].

Perhaps even more critical to this method than MIRA's utilities is the fact that MIRA also

ensures read coherence as an overlap-based assembler, unlike k-mer based assemblers like

SPAdes. SPAdes is commonly used for metagenomics assembly, and in our experience,

produces results that are as good as any other metagenomics assembler that is typically

recommended [4]. However, there are often misassemblies caused by running SPAdes on a

large and heterogeneous collection of metagenomes with the same set of k-mers. Ideally, the

user would conduct tests to find the optimal collection of k-mers for each individual

metagenome, but this step is time consuming. Thus, many users - us included - pick a canonical

set like 21, 33, 55, 77, 99 and 127 that in most cases give the greatest contiguity in the

assembly. Unfortunately, this practice can produce illusionary contiguity if the read coverage

cannot support all of the k-mer sizes [3]. Larger k-mers increase contiguity, but the read

coverage may not support them. By using MIRA, contigs that do not have read coherence may

be exposed.

After we used mirabait to extract read pairs that mapped to selected bins (Fig 1D) and

reassembled them using MIRA (Fig 1E), we iterated these two steps (Fig 1F). This iterative

process results in "digital primer walking" to extend the contigs of the bin, similar to primer or

genome walking that was initially used to sequence genomes in the late 1980s to early 1990s

[41]. At each iteration, reads with a portion mapping to any part of a contig will be included and

can lead to extension or fusion of contigs. We specifically chose to reassemble all of the reads

during each iteration to provide a more robust handling of repeats. On occasion, the extension

of the contigs resulted in overlap with contigs from other bins and unbinned contigs. Manually

including these contigs as part of the bait can speed up the process significantly. However, we

also routinely examined intermediate results and, in some cases, we saw anomalous coverage

values for different contigs indicating possible chimerism. If we saw the bin containing contigs

with significantly different coverage values (>10% difference), we removed the offending contigs

and restarted. We iterated this process until one of these outcomes occurred:

1.   **Circularization.** For us to decide that this had occurred, we looked for a single contig

244   with a significant - and exact - repeat at the ends. In addition, we required that the repeat

245   be at least 100 nt in length, was longer than any other repeat in the contig, and did not

246   match any of the other repeats.

2.   **Idempotence.** In some cases, we observed no change in the assembled contigs after a

round of read pair extraction and reassembly with MIRA. We examined some of these

instances in detail and we believe the change in coverage causes MIRA to refuse to

continue extending contigs. It is possible to adjust MIRA's thresholds of what constitutes

low and high coverage to allow contig extension to continue. However, this modification

increases the risk of collapsing repeats or creating chimeric assemblies.

3. **Chaos.** There are cases where a bin is shattered into a multitude of pieces. We are not

certain as to the exact cause, but this result is likely due to misassemblies from the initial

SPAdes assembly (discussed in more depth in a later section). Chaos appears strongly

correlated with GC and tends to occur more often when the GC content is high. We have

investigated a few in more detail and for some found that the contigs that shatter have

low 127-mer coverage as reported by SPAdes. We believe that Chaos bins are caused

by lack of read coherence in the contigs and if that is indeed the case, there is little we

can do. Once we see Chaos set in, it appears to be permanent.

After circularizing a contig, we did final checks for misassemblies with Pilon (Fig 1G). We

used Pilon [42] on the contig and then we rotated it by half the length to ensure that the ends

were in the middle and applied Pilon again (Fig 1G). We rotate the genomes because Pilon is

not capable of covering the ends of a contig. While Pilon found minor insertions/deletions due to

the circularization, it did not find any other issues in the genomes.

We next searched the genomes for a full complement of ribosomal RNAs (16S, 23S,

5S), tRNAs (all amino acids represented) and RNase P RNA to check that the genome was

correctly circularized and was not missing regions. For RNase P RNA, we needed to manually

reduce score thresholds to find all RNase P RNAs (discussed in more detail in a later section).

We were able to find tRNAs for all amino acids, although some tRNAs had Group I introns,

making them difficult to detect. Structural RNAs are sometimes invaded by Group I introns,

which is particularly true for tRNAs [Patricia Chan, private communication]. When a genome

passed the final check with the detection of the set of non-coding RNAs, we considered the

genome to be accurately circularized.

SRX3307784_bin_197 is an example of a bin that appeared to be circular, but did not

pass the check of having RNase P RNA. The assembled contig had a solid 414 base pairs of

overlap at the ends, ribosomal proteins present and tRNAs for all amino acids. However, we did

not find a copy of RNase P RNA even when we lowered the detection threshold. This caused us

to look closer and we discovered that there was another contig in the assembly which we had

thought was contamination after the initial circularization. This contig has a copy of RNase P

RNA and we were able to incorporate it into the assembly after we discovered a repeat that was

too long for the reads to span and that Pilon did not detect. We came to the conclusion that this

was a case of false circularization. To address the misassembly, we put the bin through more

iterations with mirabait and MIRA, which resulted in a larger genome which passed all of the

final checks.

In approximately 10% of the cases we attempted, we succeeded in creating a

circularized genome out of a bin. However, our selection of bins was not random as we were

heavily biased in favor of bins that we judged easiest to circularize, such as selecting bins

classified as CPR, had relatively few contigs, and had solid coverage. We are confident that we

can assemble more genomes from these datasets because we have also been able to

circularize genomes from archaea and other bacteria; these genomes will be published in future

papers. We intend to make these genomes generally accessible as we finish them. The code for

iterating to pull reads mapping to a bin and reassembly with MIRA are available on Github

(https://github.com/lmlui/Jorg) and we are currently making an app in the Department of Energy

KnowledgeBase [43].

Description of circularized CPR genomes

Using our method, we circularized 34 CPR genomes and one Margulisbacteria genome

(Table 2 and Fig 2). To create a phylogenetic tree, we used a structural alignment 16S rRNA

genes. During this process we found that many of the 16S included large introns with

LAGLIDADG homing endonucleases, an observation that has been noted in other CPR studies

[17].

Fig 2. Phylogenetic Tree based on SSU structural alignment.

At the base of the branches, in the fraction the top value is SH-aLRT which is a branch test [44]

and the bottom value is the bootstrap value. Class is listed for the CPR based on GTDB-TK

taxonomy of that genome. The class UBA1384 is also known as Berkelbacteria. We specified

the Margulisbacteria as the outgroup when creating the tree using IQ-TREE [45].

In general, these genomes are novel, but in one case, SRX1085364_bin_95, we found

that the genome is 100% identical to a previously circularized genome (INSDC Accession

CP011214.1) from that dataset [17]. Assembling the same genome as another group helps

validate our findings and that with careful manual curation two different groups can come to the

same assembly result despite differences with assemblers. Four of the other genomes had 16S

genes that had 100% hits in NCBI. Some of the 16S genes only had percent similarity in the low

80s to other sequences in Genbank.

We compared the genome sizes before and after circularization, and in most cases the

size of the genome decreased after circularization compared to the original bin. Typically the

genome shrank from a few hundred bases to a few thousand, but in some cases the genome

shrank by more than 130kbp (Table 2). This shrinkage may be a result of SPAdes artifacts that

MIRA determines to be lacking in read coherence. We are also aware of cases where SPAdes

generates contigs that are effectively duplicates of each other apart from short stretches at the

ends, and MIRA is able to resolve these into one contig.

Misassembled contigs can be found with MIRA, *i.e.* Chaos

In some cases, when we attempt the reassembly step of a bin with MIRA, we end up

with many more short contigs than what was in the original bin. SRX3024505_bin_48 started

with just 7 contigs with coverage 21X and a GC content of 59%. Superficially, it looks like a

reasonable bin. GTDB-Tk classifies it as a CPR in the Gracilibacteria class. However, after

going through 5 rounds of our method, we ended up with 136 contigs, *i.e.*, this a Chaos bin.

We do not know exactly what happened in the case of SRX3024505_bin_48, but we see

Chaos routinely during reassembly with MIRA. In some cases, we have been able to conclude

that Chaos results from insufficient read support for the largest k-mer used in the original

SPAdes assembly. Put differently, the assembly graph wasn't sufficiently well connected at the

highest k-mer used. To determine if the Chaos of SRX3024505_bin_48 was solely a result of

using MIRA, we used the same reads that we gave to MIRA as input into a SPAdes assembly.

We ended up with 47 contigs, which was still significantly worse than the original bin. It is worth

noting that the size of SRX3024505_bin_48 remained relatively constant during the testing and

reassembly process. Although Chaos is a disappointing result in assembly, knowing that a bin

likely has misassembled contigs is valuable.

Chaos predominantly occurs when the coverage is less than ~30X. Most of the genomes

we successfully circularized have much higher coverage. Based on our experience, we believe

that coverage requirements for successful circularization of genomes from metagenomes are

significantly higher than for isolates.

Likely nearly all CPR have unlinked ribosomal operons

Typically in bacteria and archaea, the 16S, 23S, and 5S ribosomal RNA genes are found

in an operon in the order 16S-23S-5S [46] (Fig 3A). In contrast, we noted that in the CPR

genomes that we circularized, nearly 80% of them had unlinked 16S and 23S genes and

sometimes unlinked 23S and 5S genes (27/34 genomes). We observed the following types of

ribosomal operons in our circularized genomes: (1) operonic, but the 16S and 23S are

separated by tRNAs on the same strand (Fig 3B), (2) operonic, but the 16S and 23S (or 23S

and 5S) are separated by tRNAs and/or protein coding genes on the same strand (Fig 3C), (3)

unlinked by distance, all three ribosomal rRNA genes are on the same strand but the 16S is

separated from the 23S-5S or all three are separated by more than 2000bp and there are no

possible intervening genes in the spacer regions that could connect the ribosomal genes in an

operon (Fig 3D), and (4) unlinked because the 16S is on the opposite strand from the 23S and

5S (Fig 3E). In three cases, tRNA genes and/or protein coding genes on the same strand were

located between the 16S and 23S or between the 23S and 5S, but there are 300-500bp regions

between the genes, so in these cases the ribosomal genes may be uncoupled, but

conservatively we counted them as operonic. In-depth analysis of gene spacing in operons of

these genomes would be required to determine if these cases are operonic or not. In the

SRX1085364_bin_95 genome, we noted that there is a homing endonuclease between both the

16S and 23S, and the 23S and 5S. Given that our genomes span a large part of the CPR

phylogeny, we infer that most of the CPR likely have unlinked ribosomal operons.

Fig 3. Diagram of placement types of ribosomal RNA genes.

Number of genomes in this study for each category are indicated in the rightmost column. (A)

Operonic. The 16S (yellow), 23S (green), and 5S (purple) ribosomal RNA genes are in an

operon. (B) Operonic with tRNAs (blue). The three ribosomal RNA genes are still in an operon,

but one or more tRNAs are located in the spacer between the 16S and 23S genes. (C) Operonic

with tRNAs and protein coding genes (beige). The three ribosomal RNA genes are still in an

operon, but one or more tRNAs or protein coding genes are located in the spacer between the

16S and 23S genes or 23S and 5S genes. It is not unusual to find that the protein coding gene

is a homing endonuclease. (D) Unlinked ribosomal RNA genes by distance. The 16S gene is

unlinked from the 23S and 5S genes, or the 23S is also unlinked from the 5S gene, by enough

distance (>2000bp) and intervening genes on the opposite strand where it is not possible for

them to be transcribed from the same promoter. (E) Unlinked ribosomal RNA genes that are on

opposite strands. The 16S is on the opposite strand from the 23S and 5S genes.

The most common type of bacterial rRNA operons are those where 16S-23S-5S are transcribed together (Fig 3A), sometimes with tRNAs between the 16S and 23S (Fig 3B), so it is notable that most of the genomes in this study have unlinked rRNA operons. Instances where 16S and 23S are decoupled are unusual, although not unknown [46,47]. Separation of 23S and 5S is very unusual in bacteria but typical in archaea [47]. Decoupling between the 16S and 23S is known to occur especially in bacteria and archaea with reduced genomes (<2Mb) [46] such as *Mycoplasma gallisepticum* [48], *Borrelia burgdorferi* [49], *Ferroplasma acidarmanus* [50], as well as obligate symbionts with small genomes such as *Buchnera aphidicola* [51], *Wolbachia pipientis* [52], and *Nanoarchaeaum equitans* [47]. In a recent study of isolate genomes and pairing long reads with metagenomics data, others have also noted that a large percentage of the CPR likely have unlinked ribosomal operons based on analyzing the distance between the 16S and 23S genes [53]. However, to our knowledge, no one else has checked for tRNAs and protein coding genes comprising the operon in this type of analysis. We also do not know of

392 other studies of CPR that have documented possible separation of 23S and 5S genes, proteins

393 in the spacer regions between ribosomal RNA genes, and 16S and 23S on opposite strands.

Diverged forms of RNase P RNA in CPR

RNase P is an RNA-protein endonuclease involved in the maturation of tRNAs by

trimming the 5' leader of pre-tRNAs. The RNA component of this complex is considered

essential for all organisms except for species of the Aquificaceae family, which contain a protein

that does not require the RNA component for tRNA trimming [27], and *Nanoarchaeum equitans*,

an obligate symbiont that does not have any detectable RNase P RNA in its reduced genome,

nor any detectable RNase P activity [26].

Given the otherwise ubiquitous nature of RNase P RNA, we require detection of this

gene as a final quality check of assembled isolate genomes and circularized genomes.

However, in the set of circularized CPR genomes in this study, we found that a significant

number that lacked RNase P RNA (10/35). Absent a high degree of confidence that these are

indeed circular genomes, we would not have noticed this anomaly. We suspected that the

RNase P RNA gene was not being detected by the models because the genomes that lacked

the gene did not fall into a specific clade and the genes that were detected still had many

conserved features of RNase P RNA. To find the missing genes, we reduced the bitscore

threshold below the model noise cutoffs when running cmsearch from the Infernal software

package [54]. The noise cutoff is the score generally considered to be the score of the highest

scoring false positive for that model (Infernal User's Guide, https://infernal.janelia.org). After

reducing the thresholds, we were able to detect the missing RNase P RNAs.

Most of the RNase P RNA genes that we found, even the ones we found initially,

required major manual refolding because of the diverged structures with either large extensions

of helices (S1 Fig) or missing helices. Many are missing P13, P14, P16, and P17, which is not

unusual. However, the RNase P RNA from SRX3307784_bin_224 appears to be missing P12

(Fig 4B), which is highly unusual because this helix is one of the most conserved across the tree

of life [40], and it is only known to be missing in *Mycoplasma fermentans* [55] and members of

the archaeal family Thermoproteaceae [56]. The closely related genome in this study,

ERX2165959_bin_53, is also missing P12 (S2 Fig). Another unusual feature is that

approximately two-thirds of the RNase P RNA (23/35) are missing the UGG motif that binds to

the CCA in pre-tRNAs. This motif tends to be missing from cyanobacteria and chloroplasts,

which may not have the CCA in their pre-tRNAs [57]. Given that cyanobacteria are one of the

closer lineages to the CPR in the bacterial tree, the loss of the UGG motif may be related to

lineage. A final example of a diverged feature is that the RNase P RNA from

SRX1775579_bin_0 appears to be missing nearly the entire P15 helix (S3 Fig). This helix is

responsible for establishing binding to pre-tRNAs in bacteria and typically contains the UGG

motif, although it is missing from all known RNase P RNA in eukaryotes and some archaea.

Fig 4. RNase P RNA can have diverged forms in CPR genomes.

Structures were drawn using VARNA (Visualization Applet for RNA, http://varna.lri.fr/) [58]. (A)

Structure of RNase P RNA from *Escherichia coli K-12* substr. MG1655. Helices P1-P18 are

labeled. The "UGG" sequence in the P15 loop that binds to the 3' end of the pre-tRNAs is

highlighted by a box. (B) Putative Structure of RNase P RNA from SRX3307784_bin_224

genome. Note that the P12, P13, P14, and P18 helices are missing, as well as the UGG motif.

Although it is not uncommon for P13, P14, and P18 to be missing in various bacteria, it is

unusual that P12 is missing. To compare the two structures, large regions of the RNA had to be

refolded manually from the original cmsearch prediction.

A   E. coli RNase P RNA   B   SRX3307784_bin_224

Finding these diverged forms of RNase P RNA would not have been possible without

having confidence that we had a complete genome. Finding these diverged structures also

illustrates that we may find diversity of genes in metagenomics data when we are no longer

restricted by what we can culture in the laboratory.

Detection and Assembly of Megaphage Genomes

In the process of circularizing genomes, we circularized what we first thought were two

novel isolates with small genomes (~0.5 Mb). However, since one of our standard checks is to

run all circular sequences against a full set of Rfam models, these immediately stood out

because the only RNAs detected were tRNAs and a tmRNA. Also, GTDB-tk was unable to

assign a taxonomy. Cursory BLASTX searches of large regions of the genome yielded only

distant hits. Based on this, we decided that they were likely megaplasmids, but have now

concluded that they are megaphage based on a recent publication [59].

SRX3024509_bin_4 is an example of one of these putative megaphages. It is 536,059 nt

long and codes for 74 tRNA sequences along with one tmRNA. We have seen more than 10

similar - in terms of size and RNA content - megaphages in a variety of environments. They

appear to be quite common and if we expanded our size limits, we believe we would find many

more. Our method should in theory perform even better on plasmids and viruses than on normal

genomes since the former are less likely to have repeats. Extraction of plasmids and viruses

from the metagenomes is a matter for future work.

Discussion

We believe it is crucial to have a substantial collection - on the order of hundreds per

phylum - of genomes that approach traditional finished genome standards as closely as

possible, such as having a single circular, contiguous sequence with an error rate less than 1

per 100,000 bp [60]. Given that we have not yet succeeded in isolating many of the species

found in metagenomics datasets, our focus is on extracting their genomes from environmental

metagenomes and enrichments. By checking that assembled genomes are circular and possess

all standard known components of genomes - such as RNAs without which life as we know it

cannot exist - are present, we gain high confidence that we had nothing but the genome and

that the genome was not falsely circularized. We believe that circularity is a top criterion for a

high quality assembly, along with checks for misassemblies. We see a clear need for an

ongoing curation of collections of genomes. As more circularized genomes are generated from

metagenomics data, comparisons will help expose misassemblies and false circularization. For

metagenomics data, checking for misassemblies is crucial because they can produce chimeric

genomes and lead to erroneous conclusions and information in public genomic databases [46].

During the development of our circularization method, we learned some lessons about

when it is the most successful and instances where it will likely fail:

1.   Our method works well for small genomes without repeats of significant length. Exact

repeats longer than the fragment length remain an issue. If the fragment length is less

than the length of a repeat, then it cannot resolve the repeat in the assembly. Once

repeats get above the fragment length, the process will - and should - fail.

2.   We noted that genomes with rRNA copy numbers greater than one will almost always

fail to circularize. We are aware of a single case where MIRA assembled two copies of

an SSU on a single contig, so it is possible to do, but it is rare. Binners almost always fail

to correctly bin multiple copies of rRNA operons as they end up on shorter contigs with

coverage that is a multiple of the single copy stretches. Because we do "digital primer

walking", it is possible to extend a contig to cover a portion of an unbinned contig

containing the ribosomal RNA genes While our method will not result in automatic

circularization in this case, it can set the stage for further manual curation and possible

eventual circularization.

3.   Circularization of genomes from metagenomes depends heavily on coverage. All of the

genomes we circularized had coverage greater than 29X (Table 2), but it may be

possible to circularize a genome with lower coverage. However, in these cases,

circularization will generally require manual intervention and we do not know how it

would be automated.

4.   Circularizing genomes with high GC content is more difficult. This is not particularly

surprising given that all of this data was Illumina sequenced and there are known biases

against high GC content [61]. It is also possible that genomes with high GC content are

larger and hence more difficult to succeed with circularization.

Even for bins that did not achieve circularity, if the bin didn't become Chaos, we suggest

that it likely still improved and, in some cases, significantly in terms of the length of contigs and

number of misassemblies. MIRA is conservative in extending contigs, so in some cases we

believe that with some manual intervention these bins can be circularized.

We are well aware that circularity is not sufficient in and of itself to account for all of the

genetic material of a microbe, *i.e.*, multiple chromosomes, plasmids, etc. may comprise a

genome and not just one chromosome. As long read technology such as PacBio and Nanopore

become more feasible for metagenomics, assembling microbial genomes will also become

easier [18,23]. In addition, long read technologies are starting to yield methylation patterns that

can be used to associate multiple replicons with each other, so it may be possible to resolve if

there are multiple chromosomes and plasmids in a genome [62].

We have at least another 100 CPR genomes that are close to being circularized and we

believe will only require only minor curation to achieve full circularization. We will release these

genomes in future publications. By generating a set of high quality reference genomes, we

intend to use these as scaffolds for future assemblies and comparative genomics. Since these

genomes are complete, we can also begin to study the genomic context of genes, such as

synteny and operons. As we demonstrated with our investigation into RNase P RNA, the study

of non-coding RNAs in the intergenic regions of these genomes is also now more feasible than

ever. Finally, we also plan to link the 16S to the functional potential that we can see in the

genome, allowing us to glean more information from 16S studies in terms of metabolic

inference. As we continue to generate high quality complete genomes from metagenomics

data, we will be able to more accurately analyze the functional potential of microorganisms that

we cannot yet culture.

Materials and Methods

Metagenomics Datasets

525 We used datasets from the NCBI Sequence Read Archive

526 (https://www.ncbi.nlm.nih.gov/sra). Accession numbers are listed in Table 1. In addition, we

527 obtained the SURF datasets from Lily Momper [35].

Read Processing Assembly

Metagenomic reads were preprocessed using BBtools version 38.60 to remove Illumina

adapters, perform quality filtering and trimming, and remove PhiX174 spike-ins. We are not

aware of any published papers documenting these tools. However, it is a standard tool suite

developed at the Department of Energy Joint Genome Institute (JGI) and it is documented at

https://jgi.doe.gov/data-and-tools/bbtools/. Processing was done in two passes. First bbduk.sh

ran with parameters *ktrim=r k=23 mink=11 hdist=1 ref=adapters.fa tbo tpe 2*. This was to

remove any remaining Illumina adapters given in adapters.fa (standard Illumina adapters). Then

bbduk.sh was run again with parameters *bf1 k=27 hdist=1 qtrim=rl trimq=17 cardinality=t*

ref=phix174_Illumina.fa. This was to perform quality filtering and trimming as well as remove

Illumina PhiX174 spike ins given in the file phix174_Illumina.fa.

Genome Assembly and Classification

Assembly was performed using SPAdes version 3.13.0 [36] with parameters *--meta -k*

21,33,55,77,99,127. Following assembly, from BWA version 0.7.17-r1188 [63], we used the

BWA-MEM algorithm with default parameters to map the reads to the set of contigs produced by

the assembly. We did this to obtain the BAM file required by MetaBAT 2 version 2.0 [37]. We

used MetaBAT 2 with parameters *--unbinned --minContig 1500 --maxEdges 500* to bin the

contigs. The iterative assemblies were performed using MIRA 5.0rc1 [7]. The parameters set

were *-NW:cac=warn, -CO:fnic=yes -AS:nop=6:sdlpo=no -KS:fenn=0.3*. We used Pilon version

1.23 [42] with default parameters to run final read coherence checks and clean up issues

created by the circularization. Taxonomic classification was generated using GTDB-Tk version

0.3.3 [38].

We carried out all of the work using standard Haswell architectures with 20 cores and

256 GB of main memory. SPAdes is generally memory limited and that is where the high point

of memory use occurred. Most of the iterative binning work is possible on a standard desktop or

even a laptop with 32 GB of memory as long as the coverage of the candidate genomes doesn't

exceed ~100X.

Gene Annotation

All of the RNA annotations were generated by Infernal 1.1.2 [54] using *cmsearch* with

parameters *--notextw --cut_tc*. We also used in-house scripts to handle RNA clan processing

[64]. We used RFAM version 14.1 for the models except when we used SSU-ALIGN (see

Phylogenetic Tree section below) which uses built-in custom models. For RNase P RNA, we

reduced the required bit score threshold 5 using the bacterial Class A model (RF00010) to find

the diverged forms. Gene calling was done using Prodigal version 2.6.3 [65]. We used *prodigal*

with parameters *-n -p single*.

Phylogenetic Tree

564   A tree was constructed from a structural alignment of the 16S genes generated by SSU-

565 ALIGN version 0.1.1 with default parameters [66,67]. Some 16S genes required manual folding

566 and adjustments to correct the structural alignment. We used IQ-TREE version 2.0-rc1 [45] to

567 generate the tree via the web server at Los Alamos National Laboratory.

Circularization code

569   The code and a tutorial of this method is available on Github

570 (https://github.com/lmlui/Jorg).

Data Availability

Assembled genomes from this study were deposited at NCBI under BioProject accession numbers PRJNA416593, PRJEB22302, PRJNA640834, PRJNA640835, PRJNA640837, PRJNA640839, PRJNA640841, PRJNA640844-PRJNA640846, PRJNA640848, PRJNA640849, PRJNA640851-PRJNA640855, PRJNA640857, PRJNA640860-PRJNA640863, PRJNA640871-PRJNA640876, PRJNA640878, PRJNA640925, PRJNA640926, PRJNA640929, PRJNA641932, PRJNA641935, and PRJNA641936, and PRJNA641940.

.

Acknowledgements

This work was funded by ENIGMA, a Science Focus Area Program supported by the U. S. Department of Energy, Office of Science, Office of Biological and Environmental Research, Genomics Sciences Program: Foundational Science; this research also used resources of the Joint Genome Institute (JGI) and the National Energy Research Scientific Computing Center (NERSC), U.S. Department of Energy Office of Science User Facilities; all managed by Lawrence Berkeley National Laboratory under Contract No. DE-AC02-05CH11231. The authors would also like to thank members of the ENIGMA Science Focus Area as well as Heidi Smith, Patricia Chan, and Norman Pace for helpful feedback and suggestions. We would also like to thank Sean Jungbluth and Lily Momper for giving us access to their datasets.

References

1. Martiny AC. The '1% culturability paradigm' needs to be carefully defined. ISME J. 2020;14: 10–11. doi:10.1038/s41396-019-0507-8

2. Myers Jr EW. A history of DNA sequence assembly. it - Inf Technol. 2016;58. doi:10.1515/itit-2015-0047

3.   Ayling M, Clark MD, Leggett RM. New approaches for metagenome assembly with short

reads. Brief Bioinform. 2020;21: 584–594. doi:10.1093/bib/bbz020

4.   Vollmers J, Wiegand S, Kaster A-K. Comparing and Evaluating Metagenome Assembly

Tools from a Microbiologist's Perspective - Not Only Size Matters! Rodriguez-Valera F,

editor. PLoS One. 2017;12: e0169662. doi:10.1371/journal.pone.0169662

5.   Myers EW, Sutton GG, Delcher AL, Dew IM, Fasulo DP, Flanigan MJ, et al. A whole-

genome assembly of Drosophila. Science (80-). 2000;287: 2196–2204.

doi:10.1126/science.287.5461.2196

6.   Simpson JT, Durbin R. Efficient de novo assembly of large genomes using compressed

data structures. Genome Res. 2012;22: 549–556. doi:10.1101/gr.126953.111

7.   Chevreux B, Pfisterer T, Drescher B, Driesel AJ, Müller WEG, Wetter T, et al. Using the

miraEST assembler for reliable and automated mRNA transcript assembly and SNP

detection in sequenced ESTs. Genome Res. 2004;14: 1147–1159.

doi:10.1101/gr.1917404

8.   Staden R. A new computer method for the storage and manipulation of DNA gel reading

data. Nucleic Acids Res. 1980;8: 3673–3694. Available:

https://www.ncbi.nlm.nih.gov/pmc/articles/PMC324183/

9.   Teeling H, Meyerdierks A, Bauer M, Amann R, Glöckner FO. Application of

tetranucleotide frequencies for the assignment of genomic fragments. Environ Microbiol.

2004;6: 938–947. doi:10.1111/j.1462-2920.2004.00624.x

10.   Korem T, Zeevi D, Suez J, Weinberger A, Avnit-Sagi T, Pompan-Lotan M, et al. Growth

dynamics of gut microbiota in health and disease inferred from single metagenomic

samples. Science (80-). 2015;349: 1101–1106. doi:10.1126/science.aac4812

11.   Tang K, Lu YY, Sun F. Background adjusted alignment-free dissimilarity measures

619   improve the detection of horizontal gene transfer. Front Microbiol. 2018;9.

620   doi:10.3389/fmicb.2018.00711

621   12.   Parks DH, Rinke C, Chuvochina M, Chaumeil PA, Woodcroft BJ, Evans PN, et al.

622   Recovery of nearly 8,000 metagenome-assembled genomes substantially expands the

623   tree of life. Nat Microbiol. 2017;2: 1533–1542. doi:10.1038/s41564-017-0012-7

624   13.   Parks DH, Imelfort M, Skennerton CT, Hugenholtz P, Tyson GW. CheckM: Assessing the

625   quality of microbial genomes recovered from isolates, single cells, and metagenomes.

626   Genome Res. 2015;25: 1043–1055. doi:10.1101/gr.186072.114

627   14.   Chen LX, Anantharaman K, Shaiber A, Murat Eren A, Banfield JF. Accurate and

628   complete genomes from metagenomes. Genome Res. 2020;30: 315–333.

629   doi:10.1101/gr.258640.119

630   15.   Kantor RS, Wrighton KC, Handley KM, Sharon I, Hug LA, Castelle CJ, et al. Small

631   genomes and sparse metabolisms of sediment-associated bacteria from four candidate

632   phyla. MBio. 2013;4. doi:10.1128/mBio.00708-13

633   16.   Anantharaman K, Brown CT, Hug LA, Sharon I, Castelle CJ, Probst AJ, et al. Thousands

634   of microbial genomes shed light on interconnected biogeochemical processes in an

635   aquifer system. Nat Commun. 2016;7: 1–11. doi:10.1038/ncomms13219

636   17.   Brown CT, Hug LA, Thomas BC, Sharon I, Castelle CJ, Singh A, et al. Unusual biology

637   across a group comprising more than 15% of domain Bacteria. Nature. 2015;523: 208–

638   211. doi:10.1038/nature14486

639   18.   Kadnikov V V., Mardanov A V., Beletsky A V., Karnachuk O V., Ravin N V. Complete

640   genome of a member of a new bacterial lineage in the microgenomates group reveals an

unusual nucleotide composition disparity between two strands of dna and limited

metabolic potential. Microorganisms. 2020;8. doi:10.3390/microorganisms8030320

19. Albertsen M, Hugenholtz P, Skarshewski A, Nielsen KL, Tyson GW, Nielsen PH. Genome

sequences of rare, uncultured bacteria obtained by differential coverage binning of

multiple metagenomes. Nat Biotechnol. 2013;31: 533–538. doi:10.1038/nbt.2579

20. Starr EP, Shi S, Blazewicz SJ, Probst AJ, Herman DJ, Firestone MK, et al. Stable isotope

informed genome-resolved metagenomics reveals that Saccharibacteria utilize

microbially-processed plant-derived carbon. Microbiome. 2018;6: 122.

doi:10.1186/s40168-018-0499-z

21. Hernsdorf AW, Amano Y, Miyakawa K, Ise K, Suzuki Y, Anantharaman K, et al. Potential

for microbial H 2 and metal transformations associated with novel bacteria and archaea in

deep terrestrial subsurface sediments. ISME J. 2017;11: 1915–1929.

doi:10.1038/ismej.2017.39

22. Sieber CMK, Probst AJ, Sharrar A, Thomas BC, Hess M, Tringe SG, et al. Recovery of

genomes from metagenomes via a dereplication, aggregation and scoring strategy. Nat

Microbiol. 2018;3: 836–843. doi:10.1038/s41564-018-0171-1

23. Moss EL, Maghini DG, Bhatt AS. Complete, closed bacterial genomes from microbiomes

using nanopore sequencing. Nat Biotechnol. 2020;38: 701–707. doi:10.1038/s41587-

020-0422-6

24. Chivian D, Brodie EL, Alm EJ, Culley DE, Dehal PS, DeSantis TZ, et al. Environmental

genomics reveals a single-species ecosystem deep within earth. Science (80-).

2008;322: 275–278. doi:10.1126/science.1155495

25. Galperin MY. Linear chromosomes in bacteria: No straight edge advantage?

Environmental Microbiology. Wiley-Blackwell; 2007. pp. 1357–1362. doi:10.1111/j.1462-

2920.2007.01328.x

26.   Randau L, Schröder I, Söll D. Life without RNase P. Nature. 2008;453: 120–123.

doi:10.1038/nature06833

27.   Nickel AI, Wäber NB, Gößringer M, Lechner M, Linne U, Toth U, et al. Minimal and RNA-

free RNase P in Aquifex aeolicus. Proc Natl Acad Sci U S A. 2017;114: 11121–11126.

doi:10.1073/pnas.1707862114

28.   Kodama Y, Shumway M, Leinonen R. The sequence read archive: explosive growth of

sequencing data on behalf of the International Nucleotide Sequence Database

Collaboration. Nucleic Acids Res. 2012;40. doi:10.1093/nar/gkr854

29.   Wilhelm RC, Hanson BT, Chandra S, Madsen E. Community dynamics and functional

characteristics of naphthalene-degrading populations in contaminated surface sediments

and hypoxic/anoxic groundwater. Environ Microbiol. 2018;20: 3543–3559.

doi:10.1111/1462-2920.14309

30.   Nelson WC, Graham EB, Crump AR, Fansler SJ, Arntzen E V., Kennedy DW, et al.

Distinct temporal diversity profiles for nitrogen cycling genes in a hyporheic microbiome.

Liu W, editor. PLoS One. 2020;15: e0228165. doi:10.1371/journal.pone.0228165

31.   Graham EB, Crump AR, Kennedy DW, Arntzen E, Fansler S, Purvine SO, et al. Multi

'omics comparison reveals metabolome biochemistry, not microbiome composition or

gene expression, corresponds to elevated biogeochemical function in the hyporheic zone.

Sci Total Environ. 2018;642: 742–753. doi:10.1016/j.scitotenv.2018.05.256

32.   Graham EB, Tfaily MM, Crump AR, Goldman AE, Bramer LM, Arntzen E, et al. Carbon

Inputs From Riparian Vegetation Limit Oxidation of Physically Bound Organic Carbon Via

Biochemical and Thermodynamic Processes. J Geophys Res Biogeosciences. 2017;122:

3188–3205. doi:10.1002/2017JG003967

33. Jewell TNM, Karaoz U, Brodie EL, Williams KH, Beller HR. Metatranscriptomic evidence

of pervasive and diverse chemolithoautotrophy relevant to C, S, N and Fe cycling in a

shallow alluvial aquifer. ISME J. 2016;10: 2106–2117. doi:10.1038/ismej.2016.25

34. Probst AJ, Ladd B, Jarett JK, Geller-Mcgrath DE, Sieber CMK, Emerson JB, et al.

Differential depth distribution of microbial function and putative symbionts through

sediment-hosted aquifers in the deep terrestrial subsurface. Nat Microbiol. 2018;3: 328–

336. doi:10.1038/s41564-017-0098-y

35. Momper L, Jungbluth SP, Lee MD, Amend JP. Energy and carbon metabolisms in a deep

terrestrial subsurface fluid microbial community. ISME J. 2017;11: 2319–2333.

doi:10.1038/ismej.2017.94

36. Bankevich A, Nurk S, Antipov D, Gurevich AA, Dvorkin M, Kulikov AS, et al. SPAdes: A

new genome assembly algorithm and its applications to single-cell sequencing. J Comput

Biol. 2012;19: 455–477. doi:10.1089/cmb.2012.0021

37. Kang DD, Li F, Kirton E, Thomas A, Egan R, An H, et al. MetaBAT 2: An adaptive binning

algorithm for robust and efficient genome reconstruction from metagenome assemblies.

PeerJ. 2019;2019. doi:10.7717/peerj.7359

38. Chaumeil P-A, Mussig AJ, Hugenholtz P, Parks DH. GTDB-Tk: a toolkit to classify

genomes with the Genome Taxonomy Database. Bioinformatics. 2020;36: 1925–1927.

doi:10.1093/bioinformatics/btz848

39. Hahn C, Bachmann L, Chevreux B. Reconstructing mitochondrial genomes directly from

genomic next-generation sequencing reads-a baiting and iterative mapping approach.

Nucleic Acids Res. 2013;41. doi:10.1093/nar/gkt371

40.   Bonfield JK, Whitwham A. Gap5—editing the billion fragment sequence assembly.

Bioinformatics. 2010;26: 1699. doi:10.1093/bioinformatics/btq268

41.   Shapter FM, Waters DLE. Genome walking. In: Henry R, Furtado A, editors. Cereal

Genomics Methods in Molecular Biology (Methods and Protocols). Totowa, NJ: Humana

Press Inc.; 2014. pp. 133–146. doi:10.1007/978-1-62703-715-0_12

42.   Walker BJ, Abeel T, Shea T, Priest M, Abouelliel A, Sakthikumar S, et al. Pilon: An

Integrated Tool for Comprehensive Microbial Variant Detection and Genome Assembly

Improvement. Wang J, editor. PLoS One. 2014;9: e112963.

doi:10.1371/journal.pone.0112963

43.   Arkin AP, Cottingham RW, Henry CS, Harris NL, Stevens RL, Maslov S, et al. KBase:

The United States Department of Energy Systems Biology Knowledgebase. Nat

Biotechnol. 2018;36: 566–569. doi:10.1038/nbt.4163

44.   Guindon S, Dufayard J-FF, Lefort V, Anisimova M, Hordijk W, Gascuel O. New algorithms

and methods to estimate maximum-likelihood phylogenies: Assessing the performance of

PhyML 3.0. Syst Biol. 2010;59: 307–321. doi:10.1093/sysbio/syq010

45.   Trifinopoulos J, Nguyen L, von Haeseler A, Minh B. W-IQ-TREE: A Fast Online

Phylogenetic Tool for Maximum Likelihood Analysis - PubMed. Nucleic Acids Res.

2016;44: W232–W235. Available: https://pubmed.ncbi.nlm.nih.gov/27084950/

46.   Andersson SG, Kurland CG. Genomic evolution drives the evolution of the translation

system. Biochem cell Biol. 1995;73: 775–787. doi:10.1139/o95-086

47.   Londei P. Translational Mechanisms and Protein Synthesis. In: Garrett RA, Klenk H-P,

editors. Archaea. Malden, MA, USA: Blackwell Publishing Ltd; 2007. pp. 217–228.

doi:10.1002/9780470750865.ch19

48.   Chen X, Finch LR. Novel arrangement of rRNA genes in Mycoplasma gallisepticum:

Separation of the 16S gene of one set from the 23S and 5S genes. J Bacteriol. 1989;171:

2876–2878. doi:10.1128/jb.171.5.2876-2878.1989

49.   Davidson MM, Evans R, Ling CL, Wiseman AD, Joss AWL, Ho-Yen DO. Isolation of

Borrelia burgdorferi from ticks in the Highlands of Scotland. J Med Microbiol. 1999;48:

59–65. doi:10.1099/00222615-48-1-59

50.   Allen EE, Tyson GW, Whitaker RJ, Detter JC, Richardson PM, Banfield JF. Genome

dynamics in a natural archaeal population. Proc Natl Acad Sci U S A. 2007;104: 1883–

1888. doi:10.1073/pnas.0604851104

51.   Rouhbakhsh D, Baumann P. Characterization of a putative 23S-5S rRNA operon of

buchnera aphidicola (endosymbiont of aphids) unlinked to the 16S rRNA-encoding gene.

Gene. 1995;155: 107–112. doi:10.1016/0378-1119(94)00910-K

52.   Bensaadi-Merchermek N, Salvado JC, Cagnon C, Karama S, Mouchès C.

Characterization of the unlinked 16S rDNA and 23S-5S rRNA operon of Wolbachia

pipientis, a prokaryotic parasite of insect gonads. Gene. 1995;165: 81–86.

doi:10.1016/0378-1119(95)00539-I

53.   Brewer TE, Albertsen M, Edwards A, Kirkegaard RH, Rocha EPC, Fierer N. Unlinked

rRNA genes are widespread among bacteria and archaea. ISME J. 2020;14: 597–608.

doi:10.1038/s41396-019-0552-3

54.   Nawrocki EP, Eddy SR. Infernal 1.1: 100-fold faster RNA homology searches.

Bioinformatics. 2013;29: 2933–2935. doi:10.1093/bioinformatics/btt509

55.   Siegel RW, Banta AB, Haas ES, Brown JW, Pace NR. Mycoplasma fermentans simplifies

our view of the catalytic core of ribonuclease P RNA. RNA. 1996;2: 452.

56.   Lai LB, Chan PP, Cozen AE, Bernick DL, Brown JW, Gopalan V, et al. Discovery of a

minimal form of RNase P in Pyrobaculum. Proc Natl Acad Sci U S A. 2010;107: 22493–8.

doi:10.1073/pnas.1013969107

57.   Schön A. Ribonuclease P: the diversity of a ubiquitous RNA processing enzyme. FEMS

Microbiol Rev. 1999;23: 391–406. doi:10.1111/j.1574-6976.1999.tb00406.x

58.   Darty K, Denise A, Ponty Y. VARNA: Interactive drawing and editing of the RNA

secondary structure. Bioinformatics. 2009;25: 1974–1975.

doi:10.1093/bioinformatics/btp250

59.   Al-Shayeb B, Sachdeva R, Chen LX, Ward F, Munk P, Devoto A, et al. Clades of huge

phages from across Earth's ecosystems. Nature. 2020;578: 425–431.

doi:10.1038/s41586-020-2007-4

60.   Chain PSG, Grafham D V., Fulton RS, Fitzgerald MG, Hostetler J, Muzny D, et al.

Genome project standards in a new era of sequencing. Science. NIH Public Access;

2009. pp. 236–237. doi:10.1126/science.1180614

61.   Chen Y-C, Liu T, Yu C-H, Chiang T-Y, Hwang C-C. Effects of GC Bias in Next-

Generation-Sequencing Data on De Novo Genome Assembly. Xu Y, editor. PLoS One.

2013;8: e62856. doi:10.1371/journal.pone.0062856

62.   Beaulaurier J, Zhu S, Deikus G, Mogno I, Zhang XS, Davis-Richardson A, et al.

Metagenomic binning and association of plasmids with bacterial host genomes using

DNA methylation. Nat Biotechnol. 2018;36: 61–69. doi:10.1038/nbt.4037

63.   Li H, Durbin R. Fast and accurate short read alignment with Burrows–Wheeler transform.

778   Bioinformatics. 2009;25: 1754–1760. doi:10.1093/bioinformatics/btp324

779   64.   Gardner PP, Daub J, Tate J, Moore BL, Osuch IH, Griffiths-Jones S, et al. Rfam:

780   Wikipedia, clans and the ``decimal'' release. Nucleic Acids Res. 2011;39: D141-5.

781   doi:10.1093/nar/gkq1129

782   65.   Hyatt D, Chen GL, LoCascio PF, Land ML, Larimer FW, Hauser LJ. Prodigal: Prokaryotic

783   gene recognition and translation initiation site identification. BMC Bioinformatics. 2010;11:

784   119. doi:10.1186/1471-2105-11-119

785   66.   Cannone JJ, Gutell RR. RNA Structure Prediction (Comparative Sequence Analysis).

786   Dictionary of Bioinformatics and Computational Biology. Chichester, UK: John Wiley &

787   Sons, Ltd; 2004. doi:10.1002/9780471650126.dob0634.pub2

788   67.   Nawrocki EP. Structural RNA Homology Search and Alignment using Covariance Models.

789   Washington University in Saint Louis. 2009. doi:http://openscholarship.wustl.edu/etd/256

Supplementary figures

S1 Fig. Putative RNase P RNA structure of SRX2838984_bin_5.

This RNase P RNA appears to have an extended P15 helix compared to typical RNase P RNA

(see the *E. coli* RNase P RNA structure in Fig 4A of the main text). Yellow highlights indicate the

portions of the RNA that had to be refolded manually. This amount of refolding was not unusual

for the RNase P RNAs found in this study.

SRX2838984_bin_5 Putative RNase P Structure

S2 Fig. Putative RNase P RNA structure of ERX2165959_bin_53.

This structure is missing P12, P13, P14, and P18. It is not unusual to be missing these helices,

except for P12 which is found in nearly all RNase P RNA structures.

S3 Fig. Putative RNase P RNA structure of SRX1775579_bin_0.

804   This structure appears to be missing most of the P15 helix.

SRX1775579_bin_0 RNase P RNA