Model-based ordination of pin-point cover data: effect of management on

dry heathland

Christian Damgaard[1], Rikke Reisner Hansen[1], and Francis K. C. Hui[2]

1 Department of Bioscience, Aarhus University, Vejlsøvej 25, 8600 Silkeborg, Denmark

2 Research School of Finance, Actuarial Studies and Statistics, Australian National University,

Acton ACT 2601, Australia

Corresponding author: Christian Damgaard, cfd@bios.au.dk, +45-30183153

Abstract

Recently, there has been an increasing interest in model-based approaches for the statistical

modelling of the joint distribution of multi-species abundances. The Dirichlet-multinomial

distribution has been proposed as a suitable candidate distribution for the joint species distribution

of pin-point plant cover data and is here applied in a model-based ordination framework. Unlike

most model-based ordination methods, both fixed and random effects are in our proposed model

structured as p-dimensional vectors and added to the latent variables before the inner product with

the species-specific coefficients. This changes the interpretation of the parameters, so that the fixed

and random effects now measure the relative displacement of the vegetation by the fixed and

random factors in the p-dimensional latent variable space. This parameterization allows statistical

inference of the effect of fixed and random factors in vector space, and makes it easier for

practitioners to perform inferences on species composition in a multivariate setting. The method

was applied on plant pin-point cover data from dry heathlands that had received different

management treatments (burned, grazed, harvested, unmanaged), and it was found that treatment

have a significant effect on heathland vegetation both when considering plant functional groups or

when the taxonomic resolution was at the species level.

Keywords: Model-based ordination; plant pin-point cover data; statistical inferences on fixed and

random effects; management of dry heathlands.

Introduction

The statistical treatment of whole communities with multiple species has traditionally relied on ordination methods that use distance measures to reduce the plot by species abundance matrix into a low dimensional vector of distances among plots. Traditional methods for ordination, such as Non Metric Multidimensional scaling (NMDS) or Principal Component Analysis (PCoA), rely on strictly algorithmic approaches with no underlying statistical model of species abundance (McCune and Grace 2002). To accommodate potential properties inherent in the data (e.g. a strong mean-variance relationship with count data), these methods may apply a variety of distance measures or data transformations. However, commonly applied distance metrics, such as Euclidean, Manhattan, or Bray–Curtis distance, often make implicit assumptions on the mean-variance relationship that may not be entirely compatible with the data at hand. This can result in potentially incorrect conclusions, e.g. location effects may be confounded within the dispersion effects (Warton et al. 2012, Warton and Hui 2017).

Recently, a number of ordination techniques, where the distribution of species abundance is explicitly taken into account, have been developed. Collectively, these new techniques are referred to as model-based ordination. In univariate statistics, for instance, these issues of mean-variance relations in the species response have long been addressed with generalized linear models (GLM's) and their mixed model counterparts, where the mean-variance relationship is modelled for each response variable (Bolker et al. 2009). Model-based ordination can thus be regarded as multivariate extensions of GLM's. This method offers the possibility of adjusting the distribution family to e.g. negative binomial distribution for overdispersed count data and the Bernoulli distribution for presence-absence data to better account for the inherent mean-variance relationship in the species observations (Hui et al. 2015, Warton et al. 2015). Like GLM's, model- based ordination methods further allow for a check of the validity of such model assumptions. Through row/column standardizations, as incorporated in model-based approaches, each taxon or species contribution to the sample location in ordination space is accounted for (Hawinkel et al. 2019) as well as potential clustering of data across different levels of study structure. For instance, hierarchical structures caused by host-parasite dynamics and reflecting nested design experiments (Björk et al. 2018). Hence, additions to the model-based ordination literature are continually emerging (e.g. Sohn and Li 2018, Hawinkel et al. 2019, Niku et al. 2019).

Arguably, the most popular method of model-based ordination that has emerged is latent variable

modelling, which involves modelling community composition through a small set of underlying

latent variables that reduce the dimension of the species abundance data from the number of species

to the dimension of the latent variables (Hui et al. 2015, Warton et al. 2015, Warton et al. 2016,

Niku et al. 2017). In doing so, the latent variables also express the residual variation owing to

factors beyond those of the measured predictors included in the model, e.g. possible biotic

interactions or phylogenetic relatedness between species. More specifically, in latent variable

modelling the species composition of each plot is described by a vector with dimension p that is

much less than the number of species. These latent variables are then fitted to the species abundance

data using the relevant distribution of species abundance. Since the latent variables are not

observed, they are treated as random effects, meaning that they need to be predicted at the same

time that coefficients associated with them are estimated (often called loadings) along with other

potential site, treatment, and environmental effects.

Ecological multivariate data analyzed using these methods often presents as count data or presence-

absence data. Because plants are sessile, they facilitate registration of multiple characteristics

related to vegetation structure and microclimate. The most common way to quantify plant species

abundance in light-open plant communities is to measure the cover, which is the relative area of the

species when projected onto the surface. Unlike plant counts or density, plant cover takes the size of

individuals into account. A relatively objective method for measuring plant cover is the pin-point

(or point-intercept) method in a frame or along a line, which has been widely employed in the field.

This involves vertically inserting a thin pin into the vegetation a number of times in a fixed design,

and the cover of a species is measured by the proportion of the inserted pins that touches the species

(Lindquist 1931, Levy and Madden 1933, Damgaard and Irvine 2019). The pin-point method is not

relevant for measuring the abundance of rare species and has been shown to underestimate species

richness (Bråkenhielm and Qinghong 1995). Importantly, the pin-point method allows unbiased

aggregation of single species cover measures at the pin level into cover data for higher plant taxa,

e.g. the cover of grasses or herbal plants.

Many plant species are spatially aggregated, and it is important to take this spatial aggregation into

account when modelling multi-species pin-point plant cover data. Recently, a reparametrized

Dirichlet-multinomial distribution has been proposed for modelling such cover data (Damgaard

2015, 2018). The Dirichlet distribution is the conjugate prior of the multinomial distribution and a

multivariate generalization of the beta-binomial distribution, which has previously been used with

success to model pin-point cover data of spatially aggregated plant species (Damgaard 2009, 2013,

Damgaard and Irvine 2019). Moreover, the Dirichlet-multinomial distributed model has been

applied for modelling multi-species vegetation dynamics in e.g. heathland ecosystems (Damgaard

2015, Damgaard et al. 2017, Damgaard 2019).

The aim of this study is two-fold; i) to modify the model-based ordination method previously

suggested by among others Hui et al. (2015) so that it is applicable to multi-species pin-point cover

data, and ii) to reparametrize the underlying latent variable model so that the effect of fixed and

random factors on the species composition may be investigated and tested in the same underlying

multidimensional space as the latent variables characterizing the ordination.

The proposed method will be illustrated on multi-species pin-point cover data from six Danish dry

heathlands, where different nature management practices have been applied and, more specifically,

it will be used to assess whether the form of management has had an effect on heathland vegetation.

Model

The objective of the proposed model is to reduce a plant species abundance matrix with k plots and

n species to a $k * p$ matrix of latent variables, where $p < n$ is the number of latent variables, and the

plant abundance of each species is measured by its cover using the pin-point method. The effects of

observed covariates, including e.g. the experimental unit and design effects on species composition,

may be entered as fixed and/or random effects into the mean structure as well. The inclusion of such

effects means that the latent variables then model the so-called residual correlation between species,

i.e. any covariation that cannot be explained by the observed predictors and experimental design

effects (Warton et al. 2015, Niku et al. 2017).

Partly following Hui et al. (2015) and adopting a Bayesian framework for estimation and inference,

we propose that the mean cover of species j in plot i, denoted here as q_{ij}, be modelled as follows

$$logit(q_{ij}) = \left(\boldsymbol{\alpha}_{fixed[i]} + \boldsymbol{\gamma}_{random[i]} + \boldsymbol{z}_i\right)'.\boldsymbol{\theta}_j + \beta_j \qquad (1),$$

where all parameters in bold denote vectors of dimension p.

In this model, the vector $\boldsymbol{\alpha}_{fixed[i]}$ denotes a fixed effect applied to plot i, for instance a treatment

effect, and are assigned weak prior distributions, e.g. $N_p(\mathbf{0}, 100\,\mathbf{I})$, which is a p-dimensional

multivariate normal distribution with zero mean vector and a covariance matrix with a diagonal

matrix with all diagonal elements set to 100. Note that for reasons of parameter identifiability, it is

assumed that these fixed effects are not unique to each plot (which is almost always the case).

Next, the vector $\boldsymbol{\gamma}_{random[i]}$ model denotes a random effect applied to plot i, e.g. the location in a

hierarchical experimental design, and are assumed to be drawn from a common multivariate normal

distribution with a zero mean vector and an unstructured $p*p$ covariance matrix, $N_p(\mathbf{0}, \boldsymbol{\Sigma})$. For

example, when $p = 2$, as is commonly considered for the purposes of ordination, $\boldsymbol{\Sigma} =$

$\begin{pmatrix} \sigma_1^2 & \rho\,\sigma_1\,\sigma_2 \\ \rho\,\sigma_1\,\sigma_2 & \sigma_2^2 \end{pmatrix}$. In this case, the standard deviations σ_i are assigned a uniform positive prior,

while the correlation coefficient ρ is assigned a uniform prior between -1 and 1. As with the fixed

effects, the random effects are assumed not to be at plot level for reasons of parameter

identifiability, e.g. in our case study the random effect is at the level of the site, in which multiple

plots are nested. In the above notation (eq. 1), the model is fitted with only one fixed factor and one

random factor, however, the model may be extended to include either more fixed or random effects,

as well as including a spatial correlated random component. Conversely, the model may be

simplified by either omitting the fixed and random effect.

Finally, the vector $\mathbf{z}_i$ denotes a vector of p latent variables for plot i and, as is standard, is assumed

to come from a standard normal distribution $N_p(\mathbf{0}, \mathbf{I})$, where the zero mean vector and identity

covariance matrix are used to avoid location and scale invariance and ensure that the parameters in

the model are identifiable (Hui et al. 2015, Niku et al. 2017).

The vector $\boldsymbol{\theta}_j$ denotes a p-dimensional vector of coefficients or loading for species j (e.g. Hui et al.

2015) and is, similarly to the fixed effects above, assigned a weak prior $N_p(\mathbf{0}, 100\,\mathbf{I})$. Note that in

order to ensure that the parameters are identifiable, we further apply a standard constraint of

assuming that the upper triangular portion of the $n*p$ matrix of species-specific coefficients is

constrained to be zero, while the diagonal elements are constrained to be positive (see Hui et al.

2015, Niku et al. 2017). The loadings can be interpreted as quantifying the species responses to the

unmeasured latent variables and can also be plotted in conjunction with the ordinations as a model-

based biplot (e.g. Warton et al. 2015). Finally, the quantities β_j denote a vector of constants chosen

a priori, which are required to ensure that $logit(q_{ij})$ is a real number.

The main difference between the present method and the original latent variable ordination

approach proposed by, among others, Hui et al. (2015) and Warton et al. (2015) is that both the

fixed and random effects, $\boldsymbol{\alpha}_{fixed[i]}$ and $\boldsymbol{\gamma}_{random[i]}$ are vectors of dimension p and are added to the

latent variables prior to their inner product with the species-specific coefficients. This changes the

interpretation of the parameters so that the fixed and random effects now measure the relative

displacement of the vegetation by the fixed and random factors in the p-dimensional latent space.

Consequently, the latent variable $\boldsymbol{z}_i$ may now more properly be referred to as modelling the residual

correlation between species, i.e. the covariation after the fixed and random effects are taken into

account. The scale of the displacement by the fixed and random effects is measured relatively to the

residual latent variables, which are fixed to a unit standard variation for ease of interpretation (and

parameter identifiability). This parameterization of the underlying latent variable model uses the

concepts of fixed and random effects in an analogous way as to how the terms are used

in generalized linear mixed effects models (Bolker et al. 2009). It is anticipated that this analogy

will make it easier for practitioners to test e.g. treatment effects on species composition in a

multivariate setting.

The underlying motivation for using model-based ordination instead of traditional ordination is to

take the distributional properties of the species abundance sampling into account such that e.g. the

correct mean-variance relationship is used in the modelling (e.g. Warton et al. 2012, Warton and

Hui 2017). In this study, we consider the multi-species pin-point cover data, and it has previously

been demonstrated that a reparametrized Dirichlet-multinomial distribution is a suitable candidate

distribution to model multi-species pin-point cover data (Damgaard 2015, Damgaard et al. 2017,

Damgaard 2019). The advantage of using the reparametrized Dirichlet-multinomial distribution is

that the degree of spatial aggregation in plant communities is taken into account and explicitly

modelled by a parameter δ (Damgaard 2018). More specifically, the observed hierarchical multi-

species pin-point cover data, Y, is modelled by a mean cover vector of the n-1 first species, q_j, and a

parameter δ, which measures the degree of intra-specific spatial aggregation, by,

$Y \sim Mn\left(\sum_n y_i, (p_1, \ldots, p_{n-1}, 1 - p_1 - \cdots - p_{n-1})\right)$ (2),

$\Lambda \left(p_1, \ldots, p_{n-1}\right) \sim Dir\left(\frac{q_1 - q_1\delta}{\delta}, \ldots, \frac{q_{n-1} - q_{n-1}\delta}{\delta}, \frac{1-\delta}{\delta} - \frac{q_1 - q_1\delta}{\delta} - \cdots - \frac{q_{n-1} - q_{n-1}\delta}{\delta}\right)$

At the limit when $\delta \to 0$, the reparametrized Dirichlet-multinomial distribution degenerates into the

multinomial distribution. Here, the prior distributions of q_j and δ are assumed to be uniformly

distributed between (0, 1) and (0.01, 0.95), respectively. For more details on the properties of the

reparametrized Dirichlet-multinomial distribution, see Damgaard (2018).

Estimation

The proposed latent variable model was estimated using Bayesian Markov Chain Monte Carlo

(MCMC) methods using the Metropolis-Hastings algorithm with normally distributed candidate

distributions. Specifically, we considered a MCMC chain with a burn-in period of 70,000 iterations

followed by 30,000 additional iterations.

Trace plots of the sampling chains of all parameters and latent variables were examined to assess

their mixing properties and convergence of the MCMC chain. Additionally, the overall fitting

properties of the model were checked by examining the regularity and shape of the marginal

distribution of parameters as well as the distribution of the deviance $(= -2 \log L(Y|\theta))$. The

efficiency of the MCMC procedure was assessed by inspecting the evolution in the deviance.

For ordination and when $p = 2$, we constructed plots of the posterior mean values of

$\left(\boldsymbol{\alpha}_{fixed[i]} + \boldsymbol{\gamma}_{random[i]} + \boldsymbol{z}_i\right)$ for each plot along with 95% credibility regions of $\boldsymbol{\alpha}_{fixed[i]}$, noting

that these are p-dimensional regions rather than the unusual one-dimensional credibility intervals

seen in mixed models and the standard latent variable ordination approach. These credibility regions

enabled us to make statistical inferences on the effect of the fixed factors on species composition,

e.g. if the 95% credibility regions of two fixed factors did not overlap. In such cases, we concluded

that there is clear statistical evidence that the effects of the two factors on the species composition

differ substantially. Consequently, the suggested parameterization in equation (1) will allow us to

utilize the model-based ordination framework for testing purposes on the latent space. Furthermore,

posterior means of the species coefficients, $\boldsymbol{\theta}_j$ for each cover class were also extracted and

superimposed on top of the ordination diagram to visualize the contribution of each cover class

(species) to the location of plots in the diagram, although note that the mean of the last species (q_n)

is not estimated in eqn. 2.

All calculations were done using *Mathematica* (Wolfram 2016).

Case study

To assess the effect of heathland management on plant communities, plant species were registered

using the pin-point method. During the fall of 2018, the cover of heathland plant species was

registered at 6 heathland sites encompassing four different heathland management regimes:

harvesting, grazing, burning and unmanaged, i.e. abandonment of management. At each site several

of the four management regimes were applied. Within each site and management combination, the

pin-point measurements were repeated in each of ten randomly positioned plots (Electronic

Supplement, Appendix A). The pin-point frames had 25 pins within an area of 0.5m x 0.5m.

Pinpoint data were then aggregated at the pin-level into five cover classes: Dwarf shrubs,

graminoids, forbs, mosses and lichens. That is, if e.g. two graminoid species where hit by the same

pin, they were aggregated to a single hit (Fig. 1).

We modelled both the aggregated pin-point data and the underlying single species pin-point data

(number of species: 34) using the proposed latent variable ordination approach formulated above,

where the four different heathland management regimes were assumed to be *p*-dimensional fixed

effects, i.e. $\boldsymbol{\alpha}_{treatment[i]}$ in equation (1), and the site was assumed to be a *p*-dimensional random

effect, i.e. $\boldsymbol{\gamma}_{site[i]}$ in equation (1). For the purpose of visualization, the dimension of the latent

variable model (*p*) was set to 2.

Results

The MCMC iterations demonstrated fair mixing properties (Electronic supplement, Appendix B),

and the marginal posterior distribution of selected parameters and compound parameters are

summarized in Table 1 and Fig. 2.

The main results are shown in Fig. 2, where the mean of the posterior distribution of $(\boldsymbol{\alpha}_i + \boldsymbol{\gamma}_i + \boldsymbol{z}_i)$

vectors are displayed for each plot for the aggregated species classes and all species, respectively.

The distance between plots indicates the difference in the cover of the species, where the scale is

relative to the unit standard deviation of the residual latent variables, i.e. the variation in species

cover among plots that could not be explained by the fixed or random factors.

The 95% credibility areas of the effect of fixed factors, i.e. the different management regimes, are

shown by colored ovals in Fig. 2. Since no ovals overlap, we concluded that the four different

management regimes lead to characteristic heathland plant communities, which on average were

distinct from each other. In particular, grazing has a large effect on the composition of the heathland

vegetation, whereas the effects of the other three management regimes were comparably more

similar (Fig. 2). This is further demonstrated in Fig. 1, where grazing leads to a shift in vegetation

cover from a high dwarf-shrub dominance towards forb and graminoid covered heathland. Inclusion

of all species in the model still yielded a distinct plant species composition with non-overlapping

credibility intervals. However, the relative positions of burned and harvested plots were shifted,

indicating that inference of management affects species composition depending on the taxonomic

resolution (functional groups vs. all species). Furthermore, increasing the taxonomic resolution

leads to higher within treatment variation in the unmanaged plots compared to the aggregated

dataset (Fig 2b, larger credibility area).

The scale of the random effects, i.e. the effect of site, is shown in Table 1. Given that the scale of

the random effect is relative to the unit standard deviation of the residual latent variables, then we

conclude that the displacement due to sites tended to be significantly larger than the variation

among the residual latent variables (Table 1, σ_2 is significantly larger than one for both cases). This

indicated a relatively large influence of site-specific species pools. There was also a significant

positive correlation between the two dimensions of the random effects when all species were

analyzed (Table 1).

In both cases, the estimated posterior distribution of δ was dominated by the chosen lower limit of

the prior distribution of the parameter (0.01), and the degree of spatial aggregation could not be

distinguished from zero in a statistical sense, i.e. random expectations.

Discussion

Over the past five years, there has been an increasing interest in model-based approaches for the

statistical modelling of the joint distribution of multi-species abundances (e.g. Clark et al. 2014,

Warton et al. 2015, Warton et al. 2016, Ovaskainen et al. 2017). The current study is an example

reflecting this important trend in community ecology, as the importance of how species abundance

data are distributed, sampled, and driven by a complex interplay of different ecological processes is

given increasing recognition. Plant species are sedentary, and in many terrestrial ecosystems plants

dominate the ground cover, leaving few bare patches. These characteristic features of plant growth

have important consequences for the joint distribution of plant species cover, such as large spatial

aggregation among single species and relatively strong negative correlation among them. To model

these features, the Dirichlet-multinomial distribution has previously been suggested as a suitable

candidate distribution for the joint species distribution of pin-point plant cover data (Damgaard

2015, Damgaard et al. 2017, Damgaard 2019). The Dirichlet-multinomial distribution has markedly

different distributional properties than both the Bernoulli distribution and the negative binomial

distribution, which till now have typically dominated the development of model-based ordination

methods and software (e.g. Niku et al. 2019).

In the suggested reparametrized Dirichlet-multinomial distribution, the spatial aggregation among

plant species is modelled by the parameter δ, which increases with the within-site spatial variation

in cover, i.e. the degree of spatial aggregation. In the present case, the degree of spatial aggregation

could not be distinguished from random expectations, and it was concluded that the observed

covariation pattern among the species was adequately modelled by the residual latent variables.

Consequently, we suggest to test whether the more simple multinomial distribution is suitable in

future model-based ordination of pin-point cover data.

Unlike in Hui et al. (2015) and Warton et al. (2015), among others, both the fixed and random

factors are here modelled as p-dimensional vectors and added to the latent variables before the inner

product with the species-specific coefficients. This means that the estimated displacement effects of

management and sites are estimated and statistical inference performed in the same p-dimensional

latent ordination space. When $p = 2$ in particular, the estimated average displacements of e.g. the

different management regimes may be shown in the plane as ovals of 95% credibility areas. Such

plots, we find, are an illustrative way to show the effects of different treatments on the species

composition. This is in contrast to most latent variable models, which consider the sample scores as

random effects and make prior distributional assumptions on them without incorporating both

random and fixed effects in the same model framework. Naturally, the suggested parameterization

with fixed and random effects may also be applicable for count data or absence-presence data.

Not surprisingly, in our case study the interpretation of the fixed effects is sensitive to taxonomic

resolution of the dataset. This was evident when comparing Fig 2a and b. If data was aggregated

into functional groups (Fig. 2a), it could be concluded that management abandonment leads to relatively homogenous plant communities. However, the model based on species level resolution (Fig. 2b) showed that the variation within the unmanaged plots is high. Regardless of taxonomic resolution, both models exhibited clear evidence that a varied management regime with different management methods produces an overall heterogeneous plant species composition. Hence, interpretation of fixed effects in this model-based framework may provide important information for heathland conservation at both the level of plant functional groups and at the species level resolution.

If $p > 2$, then the 95% credibility regions, which illustrate the mean effect of the fixed and random factors, must be imagined as a p-dimensional ellipsoidal shape. We expect that the statistical power to separate the effects of the different fixed factors on species composition will increase with the dimension of the latent variable used in the ordination. However, this preliminary notion needs to be explored in a more systematic way. Finally, the chosen estimation method is relatively slow, and in cases with many plots and many species faster estimation procedures may be needed and should be developed in the future.

Acknowledgements

We thank K. E. Nielsen for field help providing pin-point data. CFD and RRH both acknowledge the funding provided by Aage V. Jensen foundation. FKCH acknowledges the funding provided by two grants from the Australian Research Council

306 **Tables**

307 Table 1. Marginal credible intervals for the posterior distribution of the parameters in $\Sigma =$

308 $\begin{pmatrix} \sigma_1^2 & \rho\,\sigma_1\,\sigma_2 \\ \rho\,\sigma_1\,\sigma_2 & \sigma_2^2 \end{pmatrix}$

Parameter	2.5%	50%	97.5%
Aggregated cover classes:			
σ_1	0.828	1.355	1.820
σ_2	1.044	1.811	2.651
ρ	-0.710	-0.046	0.563
All species:			
σ_1	0.356	0.753	1.346
σ_2	1.270	1.965	3.005
ρ	0.377	0.870	0.993

Figures

Fig. 1. The cover of the aggregated species at the different management treatments B: burned, G: grazed, H: harvested, U: unmanaged.

Fig. 2. Map of latent variables, posterior means of $(\boldsymbol{\alpha}_i + \boldsymbol{\gamma}_i + \boldsymbol{z}_i)$ and 95% credibility areas of the effect of fixed factors (management), brown: burned, light green: harvested, dark brown: unmanaged, dark green: grazed. Mean species coefficients, $\boldsymbol{\theta}_j$ for each cover class (species) are shown. A: four aggregated species classes. B: All species, 1: *Eriophorum angustifolium*, 2: *Calamagrostis epigejos*, 3: *Trichophorum cespitosum*, 4: *Danthonia decumbens*, 5: *Agrostis capillaris*, 6: *Nardus stricta*, 7: *Festuca ovina*, 8: *Deschampsia flexuosa*, 9: *Festuca rubra*, 10: *Molinium caerulea*, 11: *Poa compressa*, 12: *Carex pilulifera*, 13: *Carex arenaria*, 14: *Carex panacea*, 15: *Carex nigra*, 16: *Luzula campestris*, 17: *Galium saxatile*, 18: *Hieracium pilosella*, 19: *Potentilla erecta*, 20: *Rumex acetosella*, 21: *Hypochoeris radicata*, 22: *Vaccinium vitis-idaea*, 23: *Calluna vulgaris*, 24: *Empetrum nigrum*, 25: *Genista anglica*, 26: *Erica tetralix*, 27: *Dicranum scoparium*, 28: *Pseudoscleropodium purum*, 29: *Pleurozium schreberi*, 30: *Bryum* subsp., 31: *Sphagnum compactum*, 32: *Campylopus introflexus*, 33: *Hypnum cupressiforme*, 34: *Cladonia rangiferina*.

References

Björk, J. R., F. K. C. Hui, R. B. O'Hara, and J. M. Montoya. 2018. Uncovering the drivers of host-associated microbiota with joint species distribution modelling. Molecular Ecology **27**:2714-2724.

Bolker, B. M., M. E. Brooks, C. J. Clark, S. W. Geange, J. R. Poulsen, M. H. H. Stevens, and J.-S. S. White. 2009. Generalized linear mixed models: a practical guide for ecology and evolution. Trends in Ecology & Evolution **24**:127-135.

Bråkenhielm, S. and L. Qinghong. 1995. Comparison of field methods in vegetation monitoring. Water, Air, and Soil Pollution **79**:75-87.

Clark, J. S., A. E. Gelfand, C. W. Woodall, and K. Zhu. 2014. More than the sum of the parts: forest climate response from joint species distribution models. Ecological Applications **24**:990-999.

Damgaard, C. 2009. On the distribution of plant abundance data. Ecological Informatics **4**:76-82.

Damgaard, C. 2013. Hierarchical and spatially aggregated plant cover data. Ecological Informatics **18**:35-39.

Damgaard, C. 2015. Modelling pin-point cover data of complementary vegetation classes. Ecological Informatics **30**:179-184.

Damgaard, C. 2018. The joint distribution of pin-point plant cover data: a reparametrized Dirichlet - multinomial distribution. arXiv e-prints.

Damgaard, C. 2019. Spatio-Temporal Structural Equation Modeling in a Hierarchical Bayesian Framework: What Controls Wet Heathland Vegetation? Ecosystems **22**:152-164.

Damgaard, C. and K. M. Irvine. 2019. Using the beta distribution to analyze plant cover data. Journal of Ecology.

Damgaard, C., K. E. Nielsen, and M. Strandberg. 2017. The effect of nitrogen deposition on the vegetation of wet heathlands. Plant Ecology **218**:373-383.

Hawinkel, S., F.-M. Kerckhof, L. Bijnens, and O. Thas. 2019. A unified framework for unconstrained and constrained ordination of microbiome read count data. PLOS ONE **14**:e0205474.

Hui, F. K. C., S. Taskinen, S. Pledger, S. D. Foster, and D. I. Warton. 2015. Model-based approaches to unconstrained ordination. Methods in Ecology and Evolution **6**:399-411.

Levy, E. B. and E. A. Madden. 1933. The point method of pasture analyses. New Zealand Journal of Agriculture **46**:267-279.

Lindquist, B. 1931. Den skandinaviska bokskogens biologi. Svenska Skogsvårdsföeningens Tidskrift **3**.

McCune, B. and J. Grace. 2002. Analysis of ecological communities. MJM Software Design, Gleneden Beach, OR.

Niku, J., F. K. C. Hui, S. Taskinen, and D. I. Warton. 2019. gllvm: Fast analysis of multivariate abundance data with generalized linear latent variable models in r. Methods in Ecology and Evolution **10**:2173-2182.

Niku, J., D. I. Warton, F. K. C. Hui, and S. Taskinen. 2017. Generalized Linear Latent Variable Models for Multivariate Count and Biomass Data in Ecology. Journal of Agricultural, Biological and Environmental Statistics **22**:498-522.

Ovaskainen, O., G. Tikhonov, A. Norberg, B. F. Guillaume, L. Duan, D. Dunson, T. Roslin, and N. Abrego. 2017. How to make more out of community data? A conceptual framework and its implementation as models and software. Ecology Letters **20**:561-576.

Sohn, B. M. and H. Li. 2018. A GLM-based latent variable ordination method for microbiome samples. Biometrics **74**:448-457.

Warton, D. I., F. G. Blanchet, R. O'Hara, O. Ovaskainen, S. Taskinen, S. C. Walker, and F. K. C. Hui. 2016. Extending Joint Models in Community Ecology: A Response to Beissinger et al. Trends in Ecology & Evolution **31**:737-738.

376   Warton, D. I., F. G. Blanchet, R. B. O'Hara, O. Ovaskainen, S. Taskinen, S. C. Walker, and F. K. C. Hui. 2015. So
377   Many Variables: Joint Modeling in Community Ecology. Trends in Ecology & Evolution **30**:766-779.
378   Warton, D. I. and F. K. C. Hui. 2017. The central role of mean-variance relationships in the analysis of
379   multivariate abundance data: a response to Roberts (2017). Methods in Ecology and Evolution
380   **8**:1408-1414.
381   Warton, D. I., S. T. Wright, and Y. Wang. 2012. Distance-based multivariate analyses confound location and
382   dispersion effects. Methods in Ecology and Evolution **3**:89-101.
383   Wolfram, S. 2016. Mathematica. Wolfram Research, Inc., Champaign, USA.

Electronic supplements

Appendix A: Details of the case study

A map of Jutland showing the location and size of the heathlands investigated in the study. Inset figure shows the location of the selected plot within the management parcel. Inset figure shows the experimental design within each plot. The squares show an example of the randomised locations for pin-point analyses.

An overview of the selected sites and the management carried out in the selected plots. The number under burn, harvest, and unmanaged is the number of years since the management was last applied. For grazing, the number is the number of years it has been grazed. Frequency denotes how often the management is performed.

Site	Harvest	Grazing	Unmanaged	Frequency

	Burn (Years)	(Years)	(Years)	(Years)	
Harrild	3	2		> 50	10 years
Kongenshus	1	1		> 50	5 years
Borris	1	1		> 30	1 year
Ovstrup		2	> 50	> 50	10 years
Randboel		2	> 30	> 100	Once
Noerholm			> 50	> 100	-

398 **Appendix B: Mathematica notebook**