Crystal structure of SARS-CoV-2 nucleocapsid protein RNA binding domain

reveals potential unique drug targeting sites

Sisi KANG[1], Mei YANG[1], Zhongsi HONG[2], Liping ZHANG[3], Zhaoxia HUANG[1], Xiaoxue CHEN[1],

Suhua HE[1], Ziliang ZHOU[1], Zhechong ZHOU[1], Qiuyue CHEN[1], Yan YAN[1], Changsheng ZHANG[3],

Hong SHAN[1,4,5]*, Shoudeng CHEN[1,4,6]*

1. Molecular Imaging Center, Guangdong Provincial Key Laboratory of Biomedical Imaging, The Fifth
Affiliated Hospital, Sun Yat-sen University, 519000, Zhuhai, China.

2. Department of Infectious Diseases, The Fifth Affiliated Hospital, Sun Yat-sen University, 519000, Zhuhai,
China.

3. Key Laboratory of Tropical Marine Bio-resources and Ecology, Guangdong Key Laboratory of Marine
Materia Medica, Institution of South China Sea Ecology and Environmental Engineering, South China Sea
Institute of Oceanology, Chinese Academy of Sciences, 164 West Xingang Road, 510301 , Guangzhou,
China.

4. Guangdong Provincial Engineering Research Center of Molecular Imaging, the Fifth Affiliated Hospital,
Sun Yat-sen University, 519000, Zhuhai, China

5. Department of Interventional Medicine, the Fifth Affiliated Hospital, Sun Yat-sen University, 519000,
Zhuhai, China

6. Department of Experimental Medicine, The Fifth Affiliated Hospital, Sun Yat-sen University, 519000,
Zhuhai, China.

Correspondence:

* Shoudeng CHEN: (chenshd5@mail.sysu.edu.cn ; Tel: +86-13612221254) or

* Hong SHAN (shanhong@mail.sysu.edu.cn ; Tel: +86-18207306672)

Abstract

The outbreak of coronavirus disease (COVID-19) in China caused by SARS-CoV-2 virus

continually lead to worldwide human infections and deaths. It is currently no specific viral protein

targeted therapeutics yet. Viral nucleocapsid protein is a potential antiviral drug target, serving

multiple critical functions during the viral life cycle. However, the structural information of SARS-

CoV-2 nucleocapsid protein is yet to be clear. Herein, we have determined the 2.7 Å crystal

structure of the N-terminal RNA binding domain of SARS-CoV-2 nucleocapsid protein.

Although overall structure is similar with other reported coronavirus nucleocapsid protein N-

terminal domain, the surface electrostatic potential characteristics between them are distinct.

Further comparison with mild virus type HCoV-OC43 equivalent domain demonstrates a unique

potential RNA binding pocket alongside the β-sheet core. Complemented by *in vitro* binding

studies, our data provide several atomic resolution features of SARS-CoV-2 nucleocapsid

protein N-terminal domain, guiding the design of novel antiviral agents specific targeting to

SARS-CoV-2.

Key words: COVID-19, Coronavirus, SARS-CoV-2, Nucleocapsid protein, RNA binding domain,

Crystal structure, Antiviral targeting site

Main Text

Introduction

The ongoing outbreak of coronavirus disease 2019 (COVID-19) is a new emerging human infectious disease caused by a novel coronavirus (severe acute respiratory syndrome coronavirus 2, SARS-CoV-2, previously known as 2019-nCoV). As of 28 February 2020, a cumulative total of 78,961 COVID-19 cases were confirmed with 2791 deaths in China. The emerged global epidemic spread rapidly with 4691 confirmed cases and 67 deaths across 51 countries outside of China (COVID-19 situation Report WHO, 28 Feb 2020). Despite remarkable efforts on containing spread of the virus, there is no specific targeted therapeutic currently.

SARS-CoV-2 is a betacoronavirus with single-stranded RNA genomes, like MERS-CoV and SARS-CoV. The first two-thirds of viral 30kb RNA genome, mainly named as ORF1a/b region, translates into two polyproteins (pp1a and pp1ab) and encodes most of the non-structural proteins (nsp). The rest parts of virus genome encode accessory proteins and four essential structural proteins, including spike (S) glycoprotein, small envelope (E) protein, matrix (M) protein, and nucleocapsid (N) protein[1,2]. Current antiviral drugs developed to treat coronavirus (CoV) infections primarily target S protein, the 3C-like (3CL) and papain-like (PLP) proteases[3,4]. Because mutant viruses in the S protein is prone to escape the targeted therapeutic with different host-cell receptor binding patterns[4], as well as antibody-dependent enhancement (ADE) effects of S protein antibodies are found in MERS coronavirus[5], there are several limitations on targeting S protein for antiviral approaches. Antiviral protease inhibitors may nonspecifically

act on the cellular homologous protease, resulting in host cell toxicity and severe side effects.

Therefore, novel antiviral strategies are needed to combat acute respiratory infections caused

by this novel coronavirus SARS-CoV-2.

The CoV N protein is a multifunctional RNA-binding protein necessary for viral RNA

transcription and replication. It plays many pivotal roles in forming helical ribonucleoproteins

during packaging the RNA genome, regulating viral RNA synthesis during replication,

transcription and modulating infected cell metabolism[6-8]. The primary functions of N protein are

binding to the viral RNA genome, and packing them into a long helical nucleocapsid structure

or ribonucleoprotein (RNP) complex[9,10]. *In vitro* and *in vivo* experiments revealed that N protein

bound to leader RNA, and was critical for maintaining highly ordered RNA conformation suitable

for replicating, and transcribing the viral genome[7,11]. More studies implicated that N protein

regulated host-pathogen interactions, such as actin reorganization, host cell cycle progression,

and apoptosis[12-14]. The N protein is also a highly immunogenic and abundantly expressed protein

during infection, capable of inducing protective immune responses against SARS-CoV and

SARS-CoV-2[15-18].

The common domain architectures of coronavirus N protein are consisting of three distinct but

highly conserved parts: An N-terminal RNA-binding domain (NTD), a C-terminal dimerization

domain (CTD), and intrinsically disordered central Ser/Arg (SR)-rich linker. Previous studies have

revealed that the NTD are responsible for RNA binding, CTD for oligomerization, and (SR)-rich

linker for primary phosphorylation, respectively[19-23]. The crystal structures of SARS-CoV N-NTD[24],

infectious bronchitis virus (IBV) N-NTD[25,26], HCoV-OC43 N-NTD[20] and mouse hepatitis virus

(MHV) N-NTD[27] have been solved. The CoVs N-NTD have been found to associate with the 3'

end of the viral RNA genome, possibly through electrostatic interactions. Additionally, several

critical residues have been identified for RNA binding and virus infectivity in the N-terminal

domain of coronavirus N proteins[24,27-29]. However, the structural and mechanistic basis for newly

emerged novel SARS-CoV-2 N protein remain largely unknown. Understanding these aspects

should facilitate the discovery of agents that specifically block the coronavirus replication,

transcription and viral assembly[30].

At present work, we report the crystal structure of SARS-CoV-2 nucleocapsid N-terminal

domain (termed as SARS-CoV-2 N-NTD), as a model for understanding the molecular

interactions that govern SARS-CoV-2 N-NTD binding to ribonucleotides. Compared with other

solved CoVs N-NTD, we characterized the specificity surface electrostatic potential features of

SARS-CoV-2 N-NTD. Additionally, we further demonstrated the unique RNA binding site

characteristics. Our findings will aid in the development of new drugs that interfere with viral N

protein and viral replication in SARS-CoV-2, and highly related virus SARS-CoV.

Materials and methods

Cloning, expression and purification

The SARS-CoV-2 N-FL plasmid is a gift from Guangdong Medical Laboratory Animal Center.

We designed several constructs including: SARS-CoV-2 N-FL (residues from 1 to 419), SARS-

CoV-2 N-NTD domain (residues from 41 to 174) and SARS-CoV-2 N-NTD domain (residues

from 33 to 180) depending on secondary structure predictions and sequence conservation

characteristics. The constructs were cloned into the pRSF-Duet-1 vector with N-terminal 6xHis-

SUMO tag and expressed in E. coli strain *Rosetta*. SARS-CoV-2 N-NTD was induced with 0.1mM

IPTG and incubated overnight at 16 °C in TB media. After Ni column chromatography followed

by Ulp1 protease digestion for tag removal, SARS-CoV-2 N-NTD (41-174) proteins were further

purified via size-exclusion chromatography (with buffer consisting of 20 mM Tris-HCl (pH 8.0),

150 mM sodium chloride, 1 mM dithiothreitol), and then concentrated by ultrafiltration to a final

concentration of 22 mg/mL.

Crystallization and data collection Crystals were grown by the sitting drop method with 0.3ul

protein（22 mg/mL） mixed with 0.6 μL and 0.3 μL well solution using Mosquito crystallization

robot and after 3 days initial crystallization was performed under 16 °C under 20 mM sodium

acetate, 100 mM sodium cacodylate (pH 6.5), 26 % PEG8000 conditions. Crystals were frozen in

liquid nitrogen in reservoir solution supplemented with 15 % glycerol (v/v) as a cryoprotectant.

X-ray diffraction data were collected at the South China Sea Institute of Oceanology, Chinese

Academy of Sciences by Rigaku X-ray diffraction (XRD) instruments. The structure was solved

by molecular replacement using PHENIX software suite[31]. The X-ray diffraction and structure

refinement statistics are summarized in Table 1.

SPR Analysis. Surface plasmon resonance (SPR) analysis was performed using a Biacore T200

with the CM5 sensor chip (GE Healthcare) at room temperature (25 °C). SARS-CoV-2 N-NTD

(41–174) were exchanged into PBS buffer via gel-filtration. The CM5 chip surface was activated

for 10 min using 0.2 M EDC/ 0.05 M NHS. After the injection of 30 µg/mL protein in 10 mM

sodium acetate (pH 5.5) for three times to immobilize on one of channels of the CM5 chip up

to ~5,800 response units, 10 µL of 1 M ethanolamine (pH 8.0) was used for blocking the

remaining activated groups. Each of the analytes (AMP, GMP, UMP, CMP) were dissolve in PBS

(pH 7.4, 0.05 % NP-20) and flowed through the chip surface at flow rate of 30 µL/min at 25 °C.

30 µL analytes were injected for affinity analysis with 60 s dissociation time. To understanding

dose dependent affinity of analytes and SARS-CoV-2 N-NTD, we tested nine dilutions of

analytes from 0.15625 mM to 10 mM. A blank flow channel was used as a reference channel to

correct the bulk refractive index by subtracting the response signal of the reference channel

from the signals of protein immobilized cells. The equilibrium constant (K_D) for analytes binding

to SARS-CoV-2 N-NTD was determined from the association and dissociation curves of the

sensorgrams, using the BIAevaluation program (Biacore).

Results

Sequence features of SARS-CoV-2 nucleocapsid protein

It is reported that the complete genome of SARS-CoV-2 (MN908947, Wuhan-Hu-1 coronavirus)

is 29.9 kb in length, similarly to 27.9 kb SARS-CoV and 30.1 kb MERS-CoV genome[32,33] (Fig. 1A).

Nucleocapsid (N) protein is translated from the 3' end structural ORF[34-36]. According to Virus

Variation Resource in National Center for Biotechnology Information databank[37], SARS-CoV-2

N protein encoding region are conserved among the known NCBI 103 genome datasets. Only

a few variations (S194L in virus strain Foshan/20SF207/2020, K249I in virus strain Wuhan/IVDC-

HB-envF13-21/2020, P344S in virus strain Guangzhou/20SF206/2020) in N protein are found in

public genomic epidemiology. An overall domain architecture of N protein among four

coronaviruses (SARS-CoV-2, SARS-CoV, MERS-CoV and HCoV-OC43) are shown in Figure 1B,

which indicates that SARS-CoV-2 shares typical characteristics with other coronaviruses. Zoom

into the completed genomic sequence of SARS-CoV-2 N protein encoding region, we find that

the sequence identities between SARS-CoV-2 with SARS-CoV, MERS-CoV, and HCoV-OC43

are 89.74%, 48.59%, 35.62%, respectively (Fig. S1)[38,39]. Since full-length SARS-CoV-2 N protein

aggregated status were found in our expression and purification studies (Fig. S2), as well as

previous reported data on other coronavirus nucleocapsid protein, we next investigate the

structural studies on N-terminal region of SARS-CoV-2 N protein (termed as SARS-CoV-2 N-

NTD).

Crystal structure of SARS-CoV-2 N-NTD

In order to obtain the atomic information of SARS-CoV-2 N-NTD, we solve the structure at 2.7

Å resolution using X-ray crystallography technology. Briefly, 47-173 residues of SARS-CoV-2 N

protein were constructed, expressed and purified as described protocol (Materials and Methods).

The structure of SARS-CoV-2 N-NTD was determined by molecular replacement using the

SARS-CoV N-NTD structure (PDB:2OG3) as the search model[24]. The final structure was refined

to R-factor and R-free values of 0.26 and 0.29, respectively. The complete statistics for data

collection, phasing and refinement are presented in Table 1. Unlike to SARS-CoV N-NTD crystals

packing modes (monoclinic form at 2OFZ, cubic form at 2OG3)[24], SARS-CoV-2 N-NTD crystal

shows orthorhombic crystal packing form with four N-NTD monomers in one asymmetry unit

(Fig. 2A). Although lacks of evidence for real RNP organization in the mature virions, the

differences in the crystal packing patterns may implicate other potential contacts in SARS-CoV-

2 RNP formation process. All four monomers in one asymmetric unit of the SARS-CoV-2 N-

NTD crystal structure shared similar right-handed (loops)-(β-sheet core)-(loops) sandwiched

structure, as conserved among the CoVs N-NTD (Fig. 2B). The β-sheet core is consisted of five

antiparallel β-strands with a single short 3_{10} helix just before strand β2, and a protruding β-

hairpin between strands β2 and β5. The β-hairpin is functionally important for CoV N-NTD,

implicated in mutational analysis of amino acid residues for RNA binding[29] (Fig. 2C). The SARS-

CoV-2 N-NTD is enriched in aromatic and basic residues, folding into a hand shape resembles

with basic fingers that extend far beyond the β-sheet core, a basic palm, and an acidic wrist (Fig.

2D).

Comparison of SARS-CoV-2 N-NTD with related viral N-NTD structures

To obtain more specific information, we first mapped the conserved residues between SARS-

CoV-2 N-NTD with SARS-CoV N-NTD, MERS-CoV N-NTD, HCoV-OC443 N-NTD, respectively

(Fig 3A). The most conserved residues distribute on the basic palm region (Fig. 3A, blue and

green region), while the less conserved residues locate in basic fingers and acidic wrist region

(Fig. 3A, pink and red region). The available CoVs N-NTD crystal structures allowed us to

compare the electrostatic potential on the surface. As shown in Fig. 3B, although CoV N-NTDs

all adapted similar overall organizations, the surface charge distribution patterns are different.

Consisting with our observations, previous modeling of related coronaviral N-NTDs also shown

markedly differ in surface charge distributions[24]. Superimposition of SARS-CoV-2 N-NTD with

three kinds of CoVs N-NTD are shown in Fig. 3C. Compared with SARS-CoV N-NTD, SARS-

CoV-2 N-NTD show a 2.1 Å movement in the β-hairpin region forward to nucleotide binding

site (Fig. 3C, left panel). While compared with MERS-CoV N-NTD, SARS-CoV-2 N-NTD show a

less extended β-hairpin region, and a distinct relax N-treminal tail (Fig. 3C, middle panel). In

consistently, SARS-CoV-2 N-NTD show a distinct relax N-treminal tail, and a 2 Å movement in

the β-hairpin region backward to the opposite side of nucleotide binding site when the structure

is compared with HCoV-OC43 N-NTD (Fig. 3C, right panel). These differences dramatically

change the surface characterizations of the protein, may result in the RNA-binding cleft being

adaptive to its own RNA genome.

A potential unique drug target pocket in SARS-CoV-2 N-NTD

Although there are several CoV N-NTDs structures have been solved, the structural basis for

ribonucleoside 5'-monophosphate binding of N protein had only been described in HCoV-

OC43, a relative type typically causing mild cold symptoms[40]. Since the surface characterizations

of N-NTD between SARS-CoV-2 with HCoV-OC43 are distinct, we next explored the differences

of RNA binding mechanistic basis with superimposition analysis. Previous studies had shown

that, HCoV-OC43 N-NTD contained Adenosine monophosphate (AMP)/ uridine

monophosphate (UMP)/ cytosine monophosphate (CMP)/ guanosine monophosphate (GMP)

binding site alongside the middle two β strands of its β-sheets core[40]. In the complex structure

of HCoV-OC43 N-NTD with ribonucleotides, the phosphate group was bound by Arg 122 and

Gly 68 via ionic interactions, the pentose sugar ribose 2'-hydroxyl group was recognized by

Ser 64 and Arg164, the nitrogenous base inserted into a hydrophobic pocket consisting of Phe

57, Pro61, Tyr 63, Tyr102, Tyr 124, and Tyr 126, mainly interacted with Tyr 124 via π–π stacking

forces (Fig. S3). It is proposed that this ribonucleotide binding mechanism are essential for all

coronavirus N proteins, applying to develop CoV N-NTD-target agents.

To obtain the structure information of SARS-CoV-2 N-NTD ribonucleotide binding site, we

make a superimposition of SARS-CoV-2 N-NTD with HCoV-OC43 N-NTD-AMP complex. As

expectedly, the root mean square deviation (RMSD) between these two structure coordinates

are 1.4 Å over 136 superimposed Cα atoms. However, a number of difference around the

ribonucleotide binding site were shown as superimposition of SARS-CoV-2 N-NTD with HCoV-

OC43 N-NTD. The major difference is N-terminal tail of N-NTD with sequence variation (SARS-

CoV-2: 48 **NNTA** 51 versus HCoV-OC43: 60 **VPYY** 63). In HCoV-OC43 N-NTD, the tail folded

up to compose a nitrogenous base binding channel, whereas this region extended outward in

SARS-CoV-2 one (Fig. 4A). The N-terminal tail movement contributed to the change of N-NTD

surface charge distribution, at which nucleotide binding cavity became easier to accessible in

SARS-CoV-2 N-NTD (Fig. 4B and C). The second difference is on phosphate group binding site,

which SARS-CoV-2 N-NTD have larger sidechain residues (55 **TA** 56) compared with HCoV-

OC43 N-NTD equivalents (67 **SG** 68) (Fig. 4D). Structural superimposition suggested additional

polar properties of Thr 55 and Ala 56 in SARS-CoV-2 N-NTD may increase the steric clash with

ribonucleotide phosphate moiety (Fig. 4E and F). The third difference is on the edge of

nitrogenous base recognized hydrophobic pocket, where SARS-CoV-2 N-NTD had Arg 89

residues compared with HCoV-OC43 N-NTD Tyr 102 equivalents (Fig. 4G). The change of these

residues sidechain may lead to dramatic decreasing of non-polar properties and increasing of

polar properties in the nitrogenous base binding site (Fig. 4H and I). To evaluate these different

observations in our structure, Surface plasmon resonance (SPR) analysis experiments were next

performed to assess the binding affinity between SARS-CoV-2 N-NTD with all four kinds of

ribonucleotide AMP/UMP/CMP/GMP. Intriguingly, all ribonucleoside 5' -monophosphate,

excepted for GMP (K_D value is 8 mM), shown little binding signals in assays (Fig. S4). Taken

together, the above results suggested a potential distinct RNA binding patterns between SARS-

CoV-2 N protein with HCoV-OC43 N protein.

Discussion

Structure-based drug discovery has been shown to be an advance approach for the

development of new therapeutics. Many ongoing studies are developed to treat COVID-19

primarily targeting the spike protein, viral proteases (3C-like protease and papain-like protease).

However, there is little effective targeted therapeutic currently. Recent studies demonstrated

that N proteins will be a good drug-targeting candidate in other CoVs since they process several

critical functions, such as RNA genomic packing, viral transcription and assembly, in the

infectious cell[10]. However, the molecular basis of SARS-CoV-2 N protein is yet to be elucidated.

Here, we present the 2.7 Å crystal structure of SARS-CoV-2 N protein N-terminal domain,

revealing the specific surface charge distributions which may facilitate drug discovery specifically

to SARS-CoV-2 N protein ribonucleotide binding domain.

On the structural basis of SARS-CoV-2 N-NTD, several residues in the ribonucleotide binding

domain were found to distinctly recognize the CoV RNA substrates. The N-terminal tail residues

(Asn 48, Asn 49, Thr 50, and Ala 51) is more flexible and extended outward compared with

equivalent residues in HCoV-OC43 N-NTD, possibly opening up the binding pocket into fitting

with viral RNA genomic high order structure. Residues Arg 89, instead of HCoV-OC43 N-NTD

Tyr 102, may contribute to guanosine base recognition despite the overall ribonucleotide

binding may be excluded by residues Thr 55 and Ala 56 in the phosphate moiety recognition

site.

Up to date, seven coronaviruses have been identified as human-susceptible virus, among which

HCoV-229E, HCoV-NL63, HCoV-HKU1 and HCoV-OC43 with low pathogenicity cause mild

respiratory symptoms similar to common cold, whereas the other three betacoronaviruses,

SARS-CoV-2, SARS-CoV and MERS-CoV lead to severe and potential fatal respiratory tract

infections[32,41,42]. Previous study reported the structural basis of HCoV-OC43 N-NTD with AMP,

GMP, UMP, CMP and a virtual screening-base compound PJ34. However, our data suggested

that SARS-CoV-2 employed a unique pattern for binding RNA with atomic resolution

information. The structure not only help us to understand the RNA-binding mechanisms

between severe infectious coronavirus with mild infectious one, but also guide the design of

novel antiviral agents specific targeting to SARS-CoV-2.

Acknowledgments

We thank for Guangdong Medical Laboratory Animal Center for providing the N-protein

encoding gene plasmids, Dr. Yongzhi Lu from Guangzhou Institutes of Biomedicine and Health

(Chinese Academy of Sciences) for the initial crystals X-ray diffraction screening, supports from

Dr. Xuan Ma of South China Sea Institute of Oceanology (Chinese Academy of Sciences) for

home source X-ray diffraction facility. This work was supported by National Natural Science

Foundation of China (31770801) and Special Fund for Scientific and Technological Innovation

Strategy of Guangdong Province of China (2018B030306029, 2017A030313145) to S.C.

Conflict of interest

The authors declare no conflict of interest

Data Availability Statement

The structures in this paper are deposited to the Protein Data Bank with 6M3M access code.

Figures

Figure 1: Sequence features of SARS-CoV-2 nucleocapsid protein

Figure legend: A. The complete whole genomic features of SARS-CoV-2 isolate Wuhan-Hu-1

(Genebank: MN908947). UTR: untranslated region; orf/ORF: open reading frame; TRS:

transcriptional regulatory sequences; S: spike glycoprotein encoding region; E: envelope protein

encoding region; M: membrane protein encoding region; N: nucleocapsid protein encoding

region. The figure is illustrated by SnapGene Viewer; **B.** Domain architectures of coronavirus

nucleocapsid protein. NTD: N-terminal RNA-binding domain; CTD: C-terminal dimerization

domain; **C.** Multiple sequence alignment of SARS-CoV-2 N-NTD with SARS-CoV N-NTD

(UniProtKB: P59595), MERS-CoV N-NTD (UniProtKB: R9UM87), HCoV-OC43 N-NTD (UniProtKB:

P33469). Red arrows indicate conserved residues for ribonucleotide binding site, dash boxes

indicate variably residues in the structural comparisons.

Figure 2: Structural overview of SARS-CoV-2 N-NTD

A

Monomer A

Monomer B

Monomer C

Monomer D

Molecules in one asymmetry unit

B

β4

β3

β-hairpin

β-sheet core

Loop region 1

β2   β5 β1   β7

β6

Loop region 2

η1

C

β-hairpin   β3   β4   Loop 1

β6   β2   β5   β1   β7   β-sheet core

N   C

Loop 2

η1

D

Basic fingers

Basic palm

Acidic wrist

Right-hand shape structure of N-NTD

Figure legend: A. Ribbon representation of SARS-CoV-2 N-NTD molecules in one asymmetric unit. The four molecules are highlighted with different color respectively; **B.** Superimpositions of four molecules in one asymmetric unit. The dash circles indicate the sandwiched structure composed of Loop region 1, β-sheet core, and Loop region 2. The β-strand is labeled with β1

288   to β7, the 3_{10} helix is labeled with η1; **C.** Topological style illustration of SARS-CoV-2 N-NTD

289   structure; **D.** Electrostatic surface of the SARS-CoV-2 N-NTD. Blue denotes positive charge

290   potential, while red indicates negative charge potential. The potential distribution was calculated

291   by Pymol. The values range from -5 kT (red) to 0 (white) and to +5 kT, where k is the Boltzmann

292   constant and T is the temperature.

 Figure 3: Comparison of SARS-CoV-2 N-NTD with related viral N-NTD structures

 Figure legend: A. Mapping the conserved surfaces of four CoV N-NTDs in SARS-CoV-2 N-

NTD structure. The multiple sequence alignment used for mapping is shown in Fig. 1C. Blue

denotes conserved residues among 4 CoVs N-NTD; green denotes one variation among 4 CoVs

N-NTD; pink denotes two variations among 4 CoVs N-NTD; red denotes three variations among

4 CoVs N-NTD; **B.** Electrostatic surface of the SARS-CoV-2 N-NTD, SARS-CoV N-NTD, MERS-

CoV N-NTD, HCoV-OC43 N-NTD. Blue denotes positive charge potential, while red indicates

negative charge potential; **C.** Overall structural comparison of SARS-CoV-2 N-NTD with related

viral N-NTD structures. Left: superimposition of SARS-CoV-2 N-NTD (blue) to SARS-CoV N-

NTD (green); middle: superimposition of SARS-CoV-2 N-NTD (blue) to MERS-CoV N-NTD

(brown); right: superimposition of SARS-CoV-2 N-NTD (blue) to HCoV-OC43 N-NTD (orange).

Figure 4: A potential unique drug target pocket in SARS-CoV-2 N-NTD

A.

Tyr (110/124)

60-VPYY-63

AMP

48-NNTA-51

Tyr (112/126)

Blue: SARS-CoV-2 N-NTD

Orange: HCoV-OC43 N-NTD

B.

SARS-CoV-2 N-NTD

Surface electrostatic potential

C.

HCoV-OC43 N-NTD with AMP

Surface electrostatic potential

D.

AMP   Tyr (110/124)

Thr 55
Ser 67

Ala 56
Gly 68

Blue: SARS-CoV-2 N-NTD

Orange: HCoV-OC43 N-NTD

E.

AMP   Tyr (110/124)

Thr 55
Ser 67

Ala 56
Gly 68

Blue: SARS-CoV-2 N-NTD

Orange: HCoV-OC43 N-NTD

F.

AMP   Tyr (110/124)

Thr 55
Ser 67

Ala 56
Gly 68

Blue: SARS-CoV-2 N-NTD

Orange: HCoV-OC43 N-NTD

G.

Arg 89
Tyr 102

Tyr (110/124)

AMP

Tyr (112/126)

Blue: SARS-CoV-2 N-NTD

Orange: HCoV-OC43 N-NTD

H.

SARS-CoV-2 N-NTD

Surface electrostatic potential

I.

HCoV-OC43 N-NTD with AMP

Surface electrostatic potential

(Removed N-terminal tail)

Figure legend: A. Detailed view of ribonucleotide binding pocket in superimposition structures

between SARS-CoV-2 N-NTD with HCoV-OC43 N-NTD AMP complex. AMP, interacting

residues and equivalents are highlighted with stick representation; **B.** Electrostatic surface of the

potential ribonucleotide binding pocket on SARS-CoV-2 N-NTD; **C.** Electrostatic surface of the

ribonucleotide binding pocket on HCoV-OC43 N-NTD; **D.** Detailed view of phosphate group

binding site in superimposition structures between SARS-CoV-2 N-NTD with HCoV-OC43 N-

NTD AMP complex; **E.** Dot representation of SARS-CoV-2 residues Thr 55 and Ala 56, which

indicates potential steric clashes with the ribonucleotide phosphate group; **F.** Dot representation

of HCoV-OC43 N-NTD residues Ser 67 and Gly 68; **G.** Detailed view of nitrogenous base binding

site in superimposition structures between SARS-CoV-2 N-NTD with HCoV-OC43 N-NTD AMP

complex; **H.** Electrostatic surface of the potential ribonucleotide nitrogenous base binding

pocket on SARS-CoV-2 N-NTD; **I.** Electrostatic surface of the ribonucleotide nitrogenous base

binding pocket on HCoV-OC43 N-NTD. In electrostatic surface potential panels, blue denotes

positive charge potential, while red indicates negative charge potential. The potential

distribution was calculated by Pymol. The values range from -5 kT (red) to 0 (white) and to +5

kT, where k is the Boltzmann constant and T is the temperature.

Table 1 Data collection and refinement statistics.

	SARS-CoV-2 N-NTD
Protein Data Bank code	6M3M
Wavelength (Å)	1.5418
Resolution range	20.92 - 2.7 (2.796 - 2.7)
Space group	$P\,2_1\,2_1\,2_1$
Unit cell (a, b, c, a, b, g)	58.88, 92.68, 97.32, 90, 90, 90
Total reflections	98913 (10077)
Unique reflections	15133 (1481)
Multiplicity	6.5 (6.8)
Completeness (%)	99.55 (99.80)
Mean I/sigma(I)	14.97 (2.9)
Wilson B-factor	33.94
R-merge	0.1043 (0.3172)
R-meas	0.1135 (0.3436)
R-pim	0.04388 (0.1303)
CC1/2	0.991 (0.96)
CC*	0.998 (0.99)
Reflections used in refinement	15126 (1481)
Reflections used for R-free	1514 (138)

R-work	0.2578 (0.3551)
R-free	0.2934 (0.4058)
CC(work)	0.908 (0.692)
CC(free)	0.851 (0.635)
Number of non-hydrogen atoms	3952
macromolecules	3822
solvent	130
Protein residues	499
RMS(bonds)	0.004
RMS(angles)	0.72
Ramachandran favored (%)	96.48
Ramachandran allowed (%)	3.52
Ramachandran outliers (%)	0.00
Rotamer outliers (%)	0.00
Clashscore	11.15
Average B-factor	31.60
macromolecules	31.86
solvent	24.03

322 Statistics for the highest-resolution shell are shown in parentheses.

Reference

1. Brian DA & Baric RS. Coronavirus genome structure and replication. *Current Topics in Microbiology and Immunology* 2005; **287**:1-30.

2. Cui J, Li F & Shi ZL. Origin and evolution of pathogenic coronaviruses. *Nature Reviews Microbiology* 2019; **17**: 181-192.

3. Ramajayam R, Tan KP & Liang PH. Recent development of 3C and 3CL protease inhibitors for anti-coronavirus and anti-picornavirus drug discovery. *Biochemical Society Transactions* 2011; **39**: 1371-1375.

4. Wrapp D, Wang N, Corbett KS, Goldsmith JA, Hsieh CL, Abiona O et al. Cryo-EM structure of the 2019-nCoV spike in the prefusion conformation. *Science* 2020.

5. Wan Y, Shang J, Sun S, Tai W, Chen J, Geng Q et al. Molecular Mechanism for Antibody-Dependent Enhancement of Coronavirus Entry. *J Virol* 2020; **94**.

6. Nelson GW, Stohlman SA & Tahara SM. High affinity interaction between nucleocapsid protein and leader/intergenic sequence of mouse hepatitis virus RNA. *Journal of General Virology* 2000; **81**: 181-188.

7. Stohlman SA, Baric RS, Nelson GN, Soe LH, Welter LM & Deans RJ. Specific interaction between coronavirus leader RNA and nucleocapsid protein. *Journal of Virology* 1988; **62**: 4288-4295.

8. Cong YY, Ulasli M, Schepers H, Mauthe M, V'kovski P, Kriegenburg F et al. Nucleocapsid Protein Recruitment to Replication-Transcription Complexes Plays a Crucial Role in Coronaviral Life Cycle. *Journal of Virology* 2020; **94**.

9. Masters PS & Sturman LS. Background paper: Functions of the coronavirus nucleocapsid protein. *Advances in Experimental Medicine and Biology* 1990; **276**: 235-238.

10. McBride R, van Zyl M & Fielding BC. The Coronavirus Nucleocapsid Is a Multifunctional Protein. *Viruses-Basel* 2014; **6**: 2991-3018.

11. Tang TK, Wu MPJ, Chen ST, Hou MH, Hong MH, Pan FM et al. Biochemical and immunological studies of nucleocapsid proteins of severe acute respiratory syndrome and 229E human coronaviruses. *Proteomics* 2005; **5**: 925-937.

12. Du L, Zhao G, Lin Y, Chan C, He Y, Jiang S et al. Priming with rAAV encoding RBD of SARS-CoV S protein and boosting with RBD-specific peptides for T cell epitopes elevated humoral and cellular immune responses against SARS-CoV infection. *Vaccine* 2008; **26**: 1644-1651.

13. Surjit M, Liu B, Chow VTK & Lal SK. The nucleocapsid protein of severe acute respiratory

syndrome-coronavirus inhibits the activity of cyclin-cyclin-dependent kinase complex and blocks S phase progression in mammalian cells. *Journal of Biological Chemistry* 2006; **281**: 10669-10681.

14. Hsieh PK, Chang SC, Huang CC, Lee TT, Hsiao CW, Kou YH et al. Assembly of severe acute respiratory syndrome coronavirus RNA packaging signal into virus-like particles is nucleocapsid dependent. *Journal of Virology* 2005; **79**: 13848-13855.

15. Ahmed SF, Quadeer AA & McKay MR. Preliminary Identification of Potential Vaccine Targets for the COVID-19 Coronavirus (SARS-CoV-2) Based on SARS-CoV Immunological Studies. *Viruses* 2020; **12**.

16. Liu SJ, Leng CH, Lien SP, Chi HY, Huang CY, Lin CL et al. Immunological characterizations of the nucleocapsid protein based SARS vaccine candidates. *Vaccine* 2006; **24**: 3100-3108.

17. Shang B, Wang XY, Yuan JW, Vabret A, Wu XD, Yang RF et al. Characterization and application of monoclonal antibodies against N protein of SARS-coronavirus. *Biochem Biophys Res Commun* 2005; **336**: 110-117.

18. Lin Y, Shen X, Yang RF, Li YX, Ji YY, He YY et al. Identification of an epitope of SARS-coronavirus nucleocapsid protein. *Cell Research* 2003; **13**: 141-145.

19. Lo YS, Lin SY, Wang SM, Wang CT, Chiu YL, Huang TH et al. Oligomerization of the carboxyl terminal domain of the human coronavirus 229E nucleocapsid protein. *FEBS Letters* 2013; **587**: 120-127.

20. Chen IJ, Yuann JMP, Chang YM, Lin SY, Zhao J, Perlman S et al. Crystal structure-based exploration of the important role of Arg106 in the RNA-binding domain of human coronavirus OC43 nucleocapsid protein. *Biochimica et Biophysica Acta - Proteins and Proteomics* 2013; **1834**: 1054-1062.

21. Chang CK, Chen CMM, Chiang MH, Hsu YL & Huang TH. Transient Oligomerization of the SARS-CoV N Protein - Implication for Virus Ribonucleoprotein Packaging. *PLoS ONE* 2013; **8**.

22. Chang CK, Sue SC, Yu TH, Hsieh CM, Tsai CK, Chiang YC et al. Modular organization of SARS coronavirus nucleocapsid protein. *Journal of Biomedical Science* 2006; **13**: 59-72.

23. Wootton SK, Rowland RRR & Yoo D. Phosphorylation of the porcine reproductive and respiratory syndrome virus nucleocapsid protein. *Journal of Virology* 2002; **76**: 10569-10576.

24. Saikatendu KS, Joseph JS, Subramanian V, Neuman BW, Buchmeier MJ, Stevens RC et al. Ribonucleocapsid formation of severe acute respiratory syndrome coronavirus through molecular action of the N-terminal domain of N protein. *Journal of Virology* 2007; **81**: 3913-

3921.

25.   Jayaram H, Fan H, Bowman BR, Ooi A, Jayaram J, Collisson EW et al. X-ray structures of the
N- and C-terminal domains of a coronavirus nucleocapsid protein: Implications for
nucleocapsid formation. *Journal of Virology* 2006; **80**: 6612-6620.

26.   Fan H, Ooi A, Tan YW, Wang S, Fang S, Liu DX et al. The nucleocapsid protein of coronavirus
infectious bronchitis virus: Crystal structure of its N-terminal domain and multimerization
properties. *Structure* 2005; **13**: 1859-1868.

27.   Grossoehme NE, Li L, Keane SC, Liu P, Dann Iii CE, Leibowitz JL et al. Coronavirus N Protein
N-Terminal Domain (NTD) Specifically Binds the Transcriptional Regulatory Sequence (TRS)
and Melts TRS-cTRS RNA Duplexes. *Journal of Molecular Biology* 2009; **394**: 544-557.

28.   Keane SC, Lius P, Leibowitzs JL & Giedroc DP. Functional Transcriptional Regulatory
Sequence (TRS) RNA binding and helix destabilizing determinants of Murine Hepatitis Virus
(MHV) Nucleocapsid (N) protein. *Journal of Biological Chemistry* 2012; **287**: 7063-7073.

29.   Tan YW, Fang S, Fan H, Lescar J & Liu DX. Amino acid residues critical for RNA-binding in
the N-terminal domain of the nucleocapsid protein are essential determinants for the infectivity
of coronavirus in cultured cells. *Nucleic Acids Research* 2006; **34**: 4816-4825.

30.   Lin SM, Lin SC, Hsu JN, Chang CK, Chien CM, Wang YS et al. Structure-based stabilization
of non-native protein-protein interactions of coronavirus nucleocapsid proteins in antiviral
drug design. *J Med Chem* 2020.

31.   Liebschner D, Afonine PV, Baker ML, Bunkoczi G, Chen VB, Croll TI et al. Macromolecular
structure determination using X-rays, neutrons and electrons: recent developments in Phenix.
Acta Crystallographica Section D-Structural Biology 2019; **75**: 861-877.

32.   Wu F, Zhao S, Yu B, Chen YM, Wang W, Song ZG et al. A new coronavirus associated with
human respiratory disease in China. *Nature* 2020.

33.   de Wit E, van Doremalen N, Falzarano D & Munster VJ. SARS and MERS: recent insights
into emerging coronaviruses. *Nat Rev Microbiol* 2016; **14**: 523-534.

34.   Gorbalenya AE, Enjuanes L, Ziebuhr J & Snijder EJ. Nidovirales: Evolving the largest RNA
virus genome. *Virus Research* 2006; **117**: 17-37.

35.   Yu LL & Malik Peiris JS. Pathogenesis of severe acute respiratory syndrome. *Current Opinion*
in Immunology 2005; **17**: 404-410.

36.   Tan YJ, Lim SG & Hong W. Characterization of viral proteins encoded by the SARS-
coronavirus genome. *Antiviral Research* 2005; **65**: 69-78.

37.   Hatcher EL, Zhdanov SA, Bao Y, Blinkova O, Nawrocki EP, Ostapchuck Y et al. Virus

421  Variation Resource - improved response to emergent viral outbreaks. *Nucleic Acids Res* 2017;

422  **45**: D482-D490.

423 38. Robert X & Gouet P. Deciphering key features in protein structures with the new ENDscript

424  server. *Nucleic Acids Res* 2014; **42**: W320-324.

425 39. Sievers F, Wilm A, Dineen D, Gibson TJ, Karplus K, Li WZ et al. Fast, scalable generation of

426  high-quality protein multiple sequence alignments using Clustal Omega. *Molecular Systems*

427  *Biology* 2011; **7**.

428 40. Lin SY, Liu CL, Chang YM, Zhao JC, Perlman S & Hou MH. Structural Basis for the

429  Identification of the N-Terminal Domain of Coronavirus Nucleocapsid Protein as an Antiviral

430  Target. *Journal of Medicinal Chemistry* 2014; **57**: 2247-2257.

431 41. Zhu N, Zhang D, Wang W, Li X, Yang B, Song J et al. A Novel Coronavirus from Patients with

432  Pneumonia in China, 2019. *N Engl J Med* 2020.

433 42. Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W et al. A pneumonia outbreak associated

434  with a new coronavirus of probable bat origin. *Nature* 2020.

  Supplementary Figures

Figure S1: Multiple sequence alignment of coronavirus nucleocapsid protein.

Figure legend: The alignment is accomplished at online server (Clustal Omega), and illustrated

with ESPript 3.0 server. The top line with SARS-CoV-2_N label indicates secondary structure

elements extracted from structural coordinate.

Figure S2: Full-length nucleocapsid protein purification.

Figure legend: On the Left panel, a gel filtration chromatography result (HiLoad® Superdex®

200 pg GE Healthcare) of recombined full-length nucleocapsid protein. On the right panel, a

12% SDS-PAGE electrophoresis is used to analyze the purification samples.

Figure S3: Ribonucleotide binding pocket in HCoV-OC43 N-NTD AMP complex.

Figure legend: A. Detailed view of ribonucleotide binding pocket in HCoV-OC43 N-NTD AMP

complex. AMP and its interacting residues are highlighted with stick representation; **B.** Simple

illustration of AMP binding pocket. Nitrogenous base and its binding residues are color with

blue. Pentose sugar and its binding residues are color with red. Phosphate group and its binding

residues are color with green.

Figure S4: *In vitro* **SPR assays for SARS-CoV-2 N-NTD with ribonucleotides.**

Figure legend: A. SPR sensorgram of the binding of varying concentrations of GMP (0, 0.15625,

0.3125, 0.625, 1.25, 2.5, 3.75, 5, 7.5 mM) to SARS-CoV-2 N-NTD captured CM5 chip. The curve

in up-right box represents the binding affinity by fitting the sensorgram to a Langumir binding

rate equations. The similar SPR sensorgrams of the binding of AMP, UMP, and CMP are shown

in panel **B**, **C**, and **D**, respectively.