Repeatable differences in exploratory behaviour predict tick infestation probability in wild great

tits

Robert E. Rollins[a,d], Alexia Mouchet[b], Gabriele Margos[c], Volker Fingerle[c], Noémie S. Becker[a], Niels J.

Dingemanse[b]

[a]Division of Evolutionary Biology, Faculty of Biology, LMU Munich, Grosshaderner Str. 2, 82152

Planegg-Martinsried, Germany

Emails: nbecker@bio.lmu.de (Noémie S. Becker), rollins@bio.lmu.de (Robert E. Rollins)

[b]Behavioural Ecology Group, Ludwig-Maximilians University Munich/Department Biology II,

Grosshaderner Str. 2, 82152 Planegg-Martinsried, Germany

Emails: dingemanse@biologie.uni-muenchen.de (Niels Dingemanse), mouchet@biologie.uni-

muenchen.de (Alexia Mouchet)

[c]National Reference Center for Borrelia, Bayerisches Landesamt fuer Gesundheit und

Lebensmittelsicherheit (LGL), Oberschleissheim Office, Veterinaerstr. 2, 85764 Oberschleissheim,

Germany

Emails: Gabriele.Margos@lgl.bayern.de (Gabrielle Margos), Volker.Fingerle@lgl.bayern.de (Volker

Fingerle)

[d]Corresponding Author:

Robert E. Rollins (rollins@bio.lmu.de)

Division of Evolutionary Biology, Faculty of Biology, LMU Munich, Grosshaderner Str. 2, 82152

Planegg-Martinsried, Germany

Office C01.035, Tel: +49 (0)89 / 2180-74152, Fax: +49 (0)89 / 2180-74104

1. Funding

This work was supported by the Max Planck Society (2012-2017); the German Science Foundation (Grant No. DI 1694/1-1); and the BioNa – Nachwuchsforscherpreis der Fakultät Biologie, Ludwig Maximilian Universität München.

2. Acknowledgments

This is a pre-print of a paper for which the official version is published in Behavioral Ecology and Sociobiology and can be accessed under doi:10.1007/s00265-021-02972-y. We would like to thank all past members of the "Evolutionary Ecology of Variation" group of the Max Plank Institute for Ornithology, the "Behavioural Ecology" and "Evolutionary Biology" groups at the LMU, Alexander Hutfluss, Sabrina Hepner, Janna Wuelbern, all lab technicians and colleagues at the "National Reference Centre for *Borrelia*," field assistants, and students for help in data collection.

3. Data Accessibility Statement

The datasets analysed and R-scripts used during the current study are available through the Dryad Digital repository at https://doi.org/10.5061/dryad.v41ns1rtz.

4. Ethics Statement

All procedures complied with guidelines from the District Government of Upper Bavaria (Regierung von Oberbayern) for Animal Care (Permit Number: ROB-55.2-2532.Vet 02-17-215)

5. Conflict of Interests

The authors have no conflicts of interest to report.

6. Authors Contributions

R.E.R and A.M. collected data, and R.E.R performed all data analysis with the guidance of N.D. and N.S.B. All co-authors helped in developing the study design. V.F. and G.M. provided expert input and guidance for proper model parameters, including development of the screening protocol. R.E.R wrote the manuscript with input and final approval from all co-authors.

Running Title

Increase of tick infestation probability in great tits

Lay Summary

Ticks use many birds as hosts, but why do some individuals have more or fewer ticks? Using a data collected over three years on great tit adults inhabiting 12 different nest-box plots, we showed that more explorative birds and those in highly recreated habitats were more likely to be infested with ticks. Exploratory behaviour and human disturbance could modify great tit habitat choice and, therefore, impact how often a birds and ticks encounter each other. (Word Limit: 75/75)

Abstract

Ticks are parasites that feed on the blood of various vertebrate hosts, including many species of bird. Birds can disperse ticks over short and long distances, therefore impacting tick population dynamics. The likelihood that birds attract ticks should depend on their behaviour and the environment. We studied various key ecological variables (breeding density, human disturbance) and phenotypic traits (exploratory behaviour; body condition) proposed to predict tick burden in great tits (*Parus major*). Our study spanned over three years and 12 human-recreated plots, equipped with nest-boxes in southern Germany. Adult breeders were assessed for exploratory behaviour, tick burden, and body condition. For each plot, human disturbance was quantified as a human recreational pressure index during biweekly nest box inspections by scoring the number of recreants using the plots. Infestation probability but not tick burden increased with exploratory behaviour. We also found moderate support for a positive effect of recreational pressure on infestation probability. Further, body condition negatively predicted tick burden. Individuals were repeatable in tick burden across years. Our study implies that infestation probability and tick burden are governed by distinct ecological and phenotypic drivers. Our findings also highlight the importance of incorporating ecological and individual variation in host phenotypes to predict spatiotemporal distributions of ticks in nature. (207/250-word limit)

Keywords: Exploratory behaviour, great tits, host-tick relationships, infestation probability, tick burden, recreational pressure (6/6 words supplied)

(Full Text Word Count: 5914)

1. Introduction

Ticks are obligate ectoparasites and require a blood-meal between each distinct life stage (Hillyard 1996; Kurtenbach et al. 2006).Thus, tick population dynamics depend on host movement and behaviour (Hillyard 1996; Hasle 2013). In turn, ticks can influence host population dynamics by affecting host fitness (Heylen et al. 2009), condition (Heylen and Matthysen 2008; Müller et al. 2013; Norte et al. 2013), and behaviour (Hoodless et al. 2002; Barber and Dingemanse 2010; Sih et al. 2018) through blood feeding and potential pathogen transfer (Stanek 2009; Norte et al. 2013). Bird species are important hosts of various tick species, disperse ticks over short and large distances (Hasle 2013), and thus affect tick-borne pathogen dynamics (Stanek 2009; Hasle 2013). Insight into the main sources of variation in avian tick uptake and burden is thus required to predict the spatiotemporal distribution of ticks in nature.

The likelihood and number of ticks carried by an individual bird vary as a function of multiple interacting ecological factors affecting tick survival in the breeding habitat, such as temperature or humidity (Lindgren et al. 2000; Oorebeek and Kleindorfer 2008; Kiffner et al. 2011; Titcomb et al. 2017), host breeding density (Oorebeek and Kleindorfer 2008; Takumi et al. 2019), and vegetation type/coverage (Kiffner et al. 2011; K. Tack et al. 2012; W. Tack et al. 2012; Tack et al. 2013), and are thus predicted to covary with habitat modification and climatic change (Lindgren et al. 2000; Randolph 2004; Akimov and Nebogatkin 2016). For example, city-dwelling birds may carry lower tick burdens (Evans et al. 2009; Hamer et al. 2012) as ticks are less abundant in urban environments (Akimov and Nebogatkin 2016). However, birds in worse physiological condition may also attract more ticks because they are more susceptible to take up parasites (Barber and Dingemanse 2010; Abbey-Lee et al. 2016; Hutfluss and Dingemanse 2019). Further studies documenting ecological predictors of tick

burden are therefore required to firmly establish factors with general versus study-specific predictive

effects (Kelly 2006; Nakagawa and Parker 2015). In this study, we quantified tick burdens on an avian

host, the great tit (*Parus major*), within 12 nest box plots surveyed for breeding density and human

disturbance levels across three breeding seasons. This enabled us to investigate the role of these two

key environmental factors in shaping avian tick burden.

Tick burden can also vary among individual birds as a function of individual-specific traits, such as

behaviour (Barber and Dingemanse 2010; Newman et al. 2015; Sih et al. 2018) or body condition

(Norte et al. 2013; Newman et al. 2015). For instance, bird species with larger average body sizes have

increased tick burdens compared to smaller bird species (Newman et al. 2015) as do bird species with a

higher proportion of ground foraging behaviour (Newman et al. 2015; Loss et al. 2016; Kocianová et

al. 2017). Similarly, within populations of the same species, individual-specific differences in foraging

or social behaviour can also predict tick burdens (Sih et al. 2018). For example, individuals that are

more active and explorative in their (foraging) behaviour may encounter more parasites (i.e. ticks), and

consequently recruit higher tick burdens (Barber and Dingemanse 2010; Sih et al. 2018; Taggart et al.

2018). Body condition and exploratory behaviour are both repeatable traits in our great tit populations

(Stuber et al. 2013; Moiron et al. 2018; Moiron et al. 2019). To quantify effects of body condition and

exploratory behaviour on tick burden in this study, we assayed both traits on all birds in each year,

such that effects on tick burden could be estimated among-individuals (Niemelä and Dingemanse).

Among-individual relationships imply that long-term repeatable differences in phenotypic traits predict

tick burden. Sex and age may also play a role in avian tick burden as males (versus females) tend to

have higher tick burdens and older (versus younger) birds have lower tick burdens (Heylen and

Matthysen 2008; Heylen et al. 2013).

The overall aim of this study was to test effects of environmental factors (breeding density and

human disturbance) and phenotypic traits (body condition, exploratory behaviour) on tick burden in the

wild. Our repeated measures design additionally enabled the quantification of the amount of variance

in tick burden attributable to among-year (temporal variation), among-plot (spatial variation), and

among-bird (individual variation) variation, thereby acquiring quantitative estimates of the magnitude

of variation attributable to multiple biological levels of variation.

2. Material and Methods

2.1. Study sites and data collection

Our study was performed on 12 nest box plots within a 10×15 km^2 area in southern Munich (47°

97'N, 11° 21'E) which were established in autumn 2009. Each plot was fitted with 50 nest boxes in a

regular grid covering approximately 9 hectares. Breeding parameters (detailed below) were monitored

for three years (2017-2019) within all plots except one plot that was monitored only in 2017. Each

nest-box was inspected biweekly during the breeding seasons (April-July) for nesting activity. During

such plot inspection, the number of recreants was counted as a measure of human disturbance (detailed

by Hutfluss & Dingemanse, (2019)).

When hatchlings were 10-12 days old, both parents were captured in the nest box using a spring

trap. Un-banded birds were given a unique, numbered band. Birds were then tested for their

exploratory behaviour (detailed below), weighed, morphologically measured (Moiron et al. 2019),

aged based on plumage characteristic as first year breeder or older (Dingemanse et al. 2020), and

assessed for ticks (protocol detailed below). All birds were captured in May (with a few captures in

June in 2018).

2.2. Exploratory behaviour

Exploratory behaviour was measured using a novel environment test adapted to the field from established protocols (Stuber et al. 2013). Birds were transferred to a holding box attached to a cage (61L $\times$ 39W $\times$ 40 H cm^3) fitted with a mesh front and three perches, representing the novel environment. The focal bird was released into the novel environment without handling and recorded for two minutes with the field observer located out of sight. Videos were subsequently scored by dividing the cage into six equal sections and three floor sections as described in (Stuber et al. 2013). Exploration score was calculated as the sum of movements the individual did between these sections (Abbey-Lee and Dingemanse 2019; Dingemanse et al. 2020).

2.3. Tick burden and collection

Ticks tend to concentrate around the eyes and beak of great tits (Heylen and Matthysen 2008). A regional patch examination protocol (reviewed in (Lydecker et al. 2019)) was developed to standardize the screening: 1) screen around the eyes/ears and along the margin of the beak on both sides of the bird, 2) screen underneath the beak, 3) screen along the top margin of the beak, 4) screen the top of the head. All captured birds were screened in 2018-2019, while in 2017 only a sample of birds were screened instead. The total number of ticks carried by each captured bird was recorded. In 2018 and 2019, ticks were collected using fine tweezers and stored in 99% ethanol as part of another study.

2.4. Recreational pressure index

During each plot inspection, all observers recorded each recreant seen and its specific location (see (Hutfluss and Dingemanse 2019)). Each recreant was connected to the first observation to avoid double counts. The probability to observe a recreant during a plot inspection is biased by various factors (Hutfluss and Dingemanse 2019). To obtain an unbiased index of recreational pressure, the binary probability to observe recreants was calculated using all inspections conducted 2010-2019 (n =

3724 inspections). This probability was calculated using a binomial generalized linear mixed effects

model (GLMM) fitting fixed effects for plot inspection duration, the number of observers, and starting

time (in hours from sunrise) (Hutfluss and Dingemanse 2019) We fitted random intercepts for each

unique combination of plot and year (termed plot-year, see (Abbey-Lee et al. 2016; Araya-Ajoy et al.

2016; Araya-Ajoy and Dingemanse 2017)) to acquire an average value of recreational pressure for

each plot in each year, and for date of observation to control for date-specific environmental effects.

We extracted best linear unbiased predictors (BLUPs) for each plot-year between 2017-2019 (n = 34)

and used them in subsequent models as a recreational pressure index. The usage of BLUPs has been

criticised when associated uncertainty is not taken forward; some have proposed to estimate a posterior

distribution of possible BLUP values and thereby take forward uncertainty in subsequent analyses

(Hadfield et al. 2010; Houslay and Wilson 2017). However, recent work has shown that taking forward

uncertainty in BLUP-values resulted in biased estimates, whereas utilizing average BLUPs as fixed

effects was shown to be less precise but unbiased (Dingemanse et al. 2020). Therefore, we here present

the estimated effect of the average BLUP values.

2.5. Data Preparation

The scaled mass index (Peig and Green 2009) was used as our measure of body condition. Briefly,

each body mass measurement was multiplied by the mean tarsus length of the population (separately

for each sex) which was first divided by the individual tarsus length (measured at the same time as

body mass) raised to a scaling exponent calculated from a log transformed regression of tarsus on body

mass measurements as described in Peig & Green, (2009). This regression was calculated using all bird

records from 2010-2019 (n = 4273 records) which included individuals repeatedly captured and

measured.

Certain factors, such as experience (Dingemanse et al. 2012; Dingemanse et al. 2020), influence

exploration scores and need to be corrected for to avoid biased measures. Individual mean exploratory

scores were calculated using a linear mixed effects model (LMM) on exploratory behaviour data from

all years (2010-2019) (n = 4251) (outlined in (Dingemanse et al. 2020)) assuming a Gaussian error

distribution. Sequence number (i.e. the number assays per bird) was fitted as a fixed effect covariate, as

well as individual and plot-year as random intercepts. An average BLUP per individual, representing

an exploratory score corrected for experience, was extracted and used in subsequent analysis as a

measure of an individual's average exploratory score.

Density was calculated as the number of breeding pairs producing first clutches per hectare.

2.6. Statistical Analysis

Tick infestation of great tits was analysed in two parts using GLMMs. First, infestation probability

was estimated using all recorded captures between 2017-2019 (n = 784 records). Second, tick burden

(i.e. absolute number of ticks observed) was modelled using only infested individuals (n = 520

records). For both response variables, the model set up was the same. Fixed effects were fitted for

recreational pressure, breeding density, body condition, sex, age, and mean individual exploration

score. Random intercepts were fitted for year, plot, plot-year, field observer, nest-box, and individual.

Prior to analysis, all fixed effects were variance standardized, such that the statistical intercepts of our

models reflected the value for the average individual in the average environmental condition.

All statistical analyses were performed in R (version 3.5.3) (R Core Team 2019). Models were run

using the functions *lmer* and *glmer* from the package *lme4* (Bates et al. 2015) assuming either binomial

(infestation probability) or Poisson (tick burden) error distributions. Mean estimates and their 95%

credible intervals (CI) were estimated based on 5,000 simulations using the *sim* function from the *arm*

package (Gelman and Su 2016). Residual errors were calculated according to Nakagawa & Schielzeth,

(2010). Adjusted repeatability values were calculated as the proportion of variance unexplained by the

fixed effects that was explained by the focal random effect. Effects of fixed effects were considered

"strongly supported" when their 95% CI did not overlap zero or "moderately supported" if the point

estimate was skewed away from but overlapped zero. Estimates that were centred on zero were viewed

as having strong support for the absence of an effect (Dingemanse et al. 2020).

3. Results

Adult great tits carrying ticks were found in all study plots and all years (Figure 1) with an average

tick burden of (mean ± s.e.) 5.63 ± 0.33 ticks per infested bird (Figure 2). Infestation probability varied

between plots (Table 1) but not tick burden (Table 1). Both tick burden and infestation probability,

varied among nest-boxes (Table 1). Individuals were not repeatable in infestation probability (Table 1)

but were moderately repeatable in tick burden (mean parameter estimate: 0.44; 95% CI: (0.39; 0.50)).

Exploratory behaviour positively predicted infestation probability (0.18 (0.01; 0.35), Figure 3) but

not tick burden (Table 1). Additionally, great tit body condition negatively predicted tick burden (-0.11

(-0.19; -0.02), Figure 4) but not infestation probability (Table 1). Recreational pressure positively

predicted infestation probability with moderate support. (0.21 (-0.03; 0.43), Figure 5). Great tit

breeding density, age, and sex did not predict infestation probability or tick burden (Table 1).

4. Discussion

Birds influence the spread of tick species, thus to understand tick population dynamics, the

underlying drivers of avian tick infestation need to be determined (Hasle 2013; Heylen et al. 2013).

Here we quantified effects of proposed environmental and phenotypic predictors on tick infestation

(Newman et al. 2015) in the great tit. More explorative birds and those living in highly recreated plots

had increased infestation probabilities, while birds with lower body condition had higher tick burdens.

Both tick burden and infestation probability varied across time (years) and space (plots). Individual

tick burden was moderately repeatable across years, implying that individual-specific phenotypic traits

not studied here influence tick infestation in great tits.

Some of the variation in tick infestation could be due to habitat choice of the great tits. Several

ecological factors might influence in which habitats those birds nest, such as resource availability,

conspecific density (Spiegel et al. 2015; Spiegel et al. 2017; Sih et al. 2018), or human disturbance

(Hutfluss and Dingemanse 2019). Great tit population density and human disturbance might push some

individuals to occupy habitats more suitable for ticks, e.g. more humid habitats with denser vegetation

(K. Tack et al. 2012; W. Tack et al. 2012), therefore influencing tick infestation. Breeding density did

not influence tick infestation; however, human disturbance might have as tick infestation was higher in

plots with higher recreational pressure and varied among nest-boxes within plots. A previous study

showed that, great tits preferentially nested further away from heavily recreated paths (Hutfluss and

Dingemanse 2019) supporting the fact that recreational pressure can modify habitat choice.

Recreational pressure however, as breeding density, did not predict tick burden. Tick burden did vary

among plots and nest-boxes implying that other environmental factors influence great tit tick burden.

For example, several ecological factors, such as relative humidity (Lindgren et al. 2000; Oorebeek and

Kleindorfer 2008; Kiffner et al. 2011), vegetation (Kiffner et al. 2011; W. Tack et al. 2012; K. Tack et

al. 2012; Tack et al. 2013), or host availability (i.e. rodents, roe deer, etc) (Oorebeek and Kleindorfer

2008; Kiffner et al. 2011; Takumi et al. 2019) might impact tick survival and distribution, leading to

spatial heterogeneity in tick abundance and, consequently, tick burden on hosts.

Various phenotypic traits varying among individual birds might also influence tick infestation.

Tick infestation probability, but not tick burden, increased with individual exploratory behaviour. This

pattern has previously been observed in Eastern grey squirrels (*Sciurus carolinensis*) (Santicchia et al.

2019). Exploratory behaviour is known to predict foraging style (Verbeek et al. 1994; Aplin et al.

2014) but to which extent is not fully understood (Moyers et al. 2018). *Ixodes ricinus* (the most

common tick infesting vertebrates in Europe (Hillyard 1996)) quests for hosts vertically up to 1.5m but

with decreasing abundance with increasing distance from the ground (Mejlon and Jaenson 1997). More

explorative birds might forage more within this questing range and closer to the ground and, therefore,

increase their encounter rate with ticks. Other research has argued that other parasite burdens than tick

burden should increase with body mass instead of exploration (Santicchia et al. 2019). Some studies

have indeed found that parasite burdens can increase with body mass (Kiffner et al. 2011; Kiffner et al.

2013; Mysterud et al. 2015). However, this relationship has not been observed in great tits (Heylen and

Matthysen 2008; Heylen et al. 2013) and other commonly studied species (Kiffner et al. 2013).

Moreover, effects of both exploratory behaviour and body mass have not been simultaneously

investigated and thus not disentangled. By jointly fitting exploratory behaviour and body condition in

our analysis we could properly differentiate between these two effects. Previously, the absence of a

relationship between great tit body mass and tick burden has been supported (Heylen and Matthysen

2008). Our study partially confirms this result as body condition did not predict infestation probability,

though it negatively predicted tick burden. This negative relationship between body condition and tick

burden has been observed in other species and has been attributed to health reductions due to blood

loss (Norte et al. 2013). Both age and sex did not influence tick infestation in our study. Previous

literature showed that males tend to have increased tick burdens (Hoodless et al. 2003) and that tick

burden decreases with age (Heylen et al. 2013). Other work however showed no effect of age on tick

burden as in our study (Heylen and Matthysen 2008). The null effect of sex though could be due to the

time of data collection where adults might be in a lower condition due to chick rearing (Ots and Horak

1996), and thus any sex differences might be masked. Infested individuals were, however, repeatable

for their tick burdens meaning that other phenotypic traits than those measured here (i.e. exploratory

behaviour and body condition) might explain tick burden of infested birds. This could be other

behaviours such as aggression (Zohdy et al. 2017) or foraging behaviour (Newman et al. 2015; Loss et al. 2016; Kocianová et al. 2017) as both play roles in tick burden.

It is important to mention that birds were captured approximately over a one-month period (range: 25-29 days) mostly during May each year which corresponds to the peak of tick abundance (Kurtenbach et al. 2006). However, some variation in tick infestation could be due to weather related conditions when the birds encountered the ticks. Weather and climatic variables affect questing activity of ticks (Mejlon and Jaenson 1997) and consequently the probability for a bird to encounter a tick and ultimately overall tick infestation. Additionally, both breeding density and bird age where shown to have null effects on tick infestation. However, this null result could be due to effects at different levels of biological variation being confounded (van de Pol and Wright 2009). To test if this was the case, we split age into among- and within-individual variation in age and split breeding density into temporal- and spatial-variation in density. These level-specific analyses also failed to show effects of these factors (Supplemental Text 1) supporting the absence of these effects.

5. Conclusions

We showed that exploratory behaviour predicted tick infestation probability in great tits, with more explorative birds having an increased infestation probability. However, tick burden of infested birds was not influenced by exploratory behaviour, sex, or age but was negatively predicted by body condition. We additionally showed that individuals were repeatable for tick burden, meaning that other phenotypic traits not analysed here might explain consistent differences in tick burden of infested individuals. We also found moderate support for a positive effect of recreational pressure on infestation probability. Our results highlight the complex nature of tick infestation and how various factors influence infestation probability and tick burden.

6. Ethics Statement

All procedures complied with guidelines from the District Government of Upper Bavaria (Regierung von Oberbayern) for Animal Care (Permit Number: ROB-55.2-2532.Vet 02-17-215)

7. Conflict of Interests

The authors have no conflicts of interest to report.

8. References

Abbey-Lee RN, Dingemanse NJ. 2019. Adaptive individual variation in phenological responses to perceived predation levels. Nat Commun. 10(1601):1–10. doi:10.1038/s41467-019-09138-5.

Abbey-Lee RN, Mathot KJ, Dingemanse NJ. 2016. Behavioral and morphological responses to perceived predation risk: A field experiment in passerines. Behav Ecol. 27(3):857–864. doi:10.1093/beheco/arv228.

Akimov IA, Nebogatkin I V. 2016. Ixodid Ticks (Acari, Ixodidae) in Urban Landscapes. A review. Vestn Zool. 50(2):155–162.

Aplin LM, Farine DR, Mann RP, Sheldon BC. 2014. Individual-level personality influences social foraging and collective behaviour in wild birds. Proc R Soc B. 281:1–9.

Araya-Ajoy YG, Dingemanse NJ. 2017. Repeatability, heritability, and age-dependence of seasonal plasticity in aggressiveness in a wild passerine bird. J Anim Ecol. 86(2):227–238. doi:10.1111/1365-2656.12621.

Araya-Ajoy YG, Kuhn S, Mathot KJ, Mouchet A, Mutzel A, Nicolaus M, Wijmenga JJ, Kempenaers B, Dingemanse NJ. 2016. Sources of (co)variation in alternative siring routes available to male great tits (*Parus major*). Evolution (N Y). 70(10):2308–2321.

Barber I, Dingemanse NJ. 2010. Parasitism and the evolutionary ecology of animal personality. Philos Trans R Soc B. 365:4077–4088.

Bates D, Maechler M, Bolker B, Walker S. 2015. Fitting Linear Mixed-Effects Models Using lme4. J

Stat Softw. 67(1–48).

Dingemanse NJ, Bouwman KM, Pol M Van De, Overveld T Van, Patrick SC, Matthysen E, Quinn JL.

2012. Variation in personality and behavioural plasticity across four populations of the great tit *Parus*

major. J Anim Ecol. 81:116–126.

Dingemanse NJ, Moiron M, Araya-Ajoy YG. 2020. Individual variation in age-dependent

reproduction: fast explorers live fast but senesce young? J Anim Ecol. 89:601–613.

Evans KL, Gaston KJ, Sharp SP, McGowan A, Simeoni M, Hatchwell BJ. 2009. Effects of

urbanisation on disease prevalence and age structure in blackbird *Turdus merula* populations. Oikos.

118(5):774–782. doi:10.1111/j.1600-0706.2008.17226.x.

Gelman A, Su Y-S. 2016. arm: Data Analysis Using Regression and Multilevel/Hierarchical Models.

https://cran.r-project.org/package=arm.

Hadfield JD, Wilson AJ, Garant D, Sheldon BC, Kruuk LEB. 2010. The misuse of BLUP in ecology

and evolution. Am Nat. 175(1):116–125. doi:10.1086/648604.

Hamer SA, Lehrer E, Magle SB. 2012. Wild Birds as Sentinels for Multiple Zoonotic Pathogens Along

an Urban to Rural Gradient in Greater Chicago, Illinois. Zoonoses Public Health. 59(5):355–364.

doi:10.1111/j.1863-2378.2012.01462.x.

Hasle G. 2013. Transport of ixodid ticks and tick-borne pathogens by migratory birds. Front Cell Infect

Microbiol. 3(September):1–6.

Heylen D, Adriaensen F, Dauwe T, Eens M, Matthysen E. 2009. Offspring quality and tick infestation

load in brood rearing great tits *Parus major*. Oikos. 118(10):1499–1506. doi:10.1111/j.1600-

0706.2009.17606.x.

Heylen D, Adriaensen F, Van Dongen S, Sprong H, Matthysen E. 2013. Ecological factors that

determine *Ixodes ricinus* tick burdens in the great tit (*Parus major*), an avian reservoir of *Borrelia*

burgdorferi s.l. Int J Parasitol. 43:603–611.

Heylen D, Matthysen E. 2008. Effect of tick parasitism on the health status of a passerine bird. Funct

Ecol. 22(6):1099–1107. doi:10.1111/j.1365-2435.2008.01463.x.

Hillyard PD. 1996. Ticks of North-West Europe. Barnes RSK, Crothers JH, editors. London: Feild

Studies Council, Schrewsbury.

Hoodless AN, Kurtenbach K, Nuttall PA, Randolph SE. 2003. Effects of tick *Ixodes ricinus* infestation

on pheasant Phasianus colchicus breeding success and survival. Wildlife Biol. 9(4):171–178.

doi:10.2981/wlb.2003.046.

Hoodless Andrew N, Kurtenbach Klaus, Nuttall PA, Randolph Hoodless SE, Hoodless A N,

Kurtenbach K, Randolph SE. 2002. The impact of ticks on pheasant territoriality. OIKOS. 96:245–250.

Houslay TM, Wilson AJ. 2017. Avoiding the misuse of BLUP in behavioural ecology. Behav Ecol.

28(4):948–952. doi:10.1093/beheco/arx023.

Hutfluss A, Dingemanse NJ. 2019. Human recreation reduces clutch size in great tits Parus major

regardless of risk-taking personality. Behav Ecol. 30(6):1751–1760. doi:10.1093/beheco/arz145.

Kelly CD. 2006. Replicating empirical research in behavioral ecology: How and why it should be done

but rarely ever is. Q Rev Biol. 81(3):221–236. doi:10.1086/506236.

Kiffner C, Stanko M, Morand S, Khokhlova IS, Shenbrot GI, Laudisoit A, Leirs H, Hawlena H,

Krasnov BR. 2013. Sex-biased parasitism is not universal: Evidence from rodent-flea associations

from three biomes. Oecologia. 173(3):1009–1022.

Kiffner C, Vor T, Hagedorn P, Niedrig M, Rühe F. 2011. Factors affecting patterns of tick parasitism

on forest rodents in tick-borne encephalitis risk areas, Germany. Parasitol Res. 108(2):323–335.

doi:10.1007/s00436-010-2065-x.

Kocianová E, Rus V, Haruˇ D, Spitalská E. 2017. Seasonal infestation of birds with immature stages of

Ixodes ricinus and *Ixodes arboricola*. Ticks Tick Borne Dis. 8:423–431.

doi:10.1016/j.ttbdis.2017.01.006.

Kurtenbach K, Hanincová K, Tsao JI, Margos G, Fish D, Ogden NH. 2006. Fundamental processes in

the evolutionary ecology of Lyme borreliosis. Nat Rev Microbiol. 4(9):660–669.

Lindgren E, Tälleklint L, Polfeldt T. 2000. Impact of climatic change on the northern latitude limit and

population density of the disease-transmitting European tick *Ixodes ricinus*. Environ Health Perspect.

108(2):119–123. doi:10.2307/3454509.

Loss SR, Noden BH, Hamer GL, Hamer SA. 2016. A quantitative synthesis of the role of birds in

carrying ticks and tick-borne pathogens in North America. Oecologia. 182(4):947–959.

doi:10.1007/s00442-016-3731-1.

Lydecker HW, Banks PB, Hochuli DF. 2019. Counting Ticks (Acari: Ixodida) on Hosts Is Complex: A

Review and Comparison of Methods. J Med Entomol. 56(6):1–7.

Mejlon HA, Jaenson TGT. 1997. Questing behaviour of *Ixodes ricinus* ticks (Acari: Ixodidae). Exp

Appl Acarol. 21:747–754.

Moiron M, Araya-Ajoy YG, Mathot KJ, Mouchet A, Dingemanse NJ. 2019. Functional relations

between body mass and risk-taking behavior in wild great tits. Behav Ecol. 30(3):617–623.

Moiron M, Mathot KJ, Dingemanse NJ. 2018. To eat and not be eaten: diurnal mass gain and foraging

strategies in wintering great tits. Proc R Soc B. 285.

Moyers SC, Adelman JS, Farine DR, Moore IT, Hawley DM. 2018. Exploratory behavior is linked to

stress physiology and social network centrality in free-living house finches (*Haemorhous mexicanus*).

Horm Behav. 102:105–113.

Müller W, Heylen D, Eens M, Fabio Rivera-Gutierrez H, G Groothuis TG, Rivera-Gutierrez HF,

Groothuis TGG. 2013. An experimental study on the causal relationships between (ecto-)parasites,

testosterone and sexual signalling. Behav Ecol Sociobiol. 67(11):1791–1798. doi:10.1007/s00265-013-

1586-6.

Mysterud A, Byrkjeland R, Qviller L, Viljugrein H. 2015. The generalist tick *Ixodes ricinus* and the

specialist tick *Ixodes trianguliceps* on shrews and rodents in a northern forest ecosystem- a role of

body size even among small hosts. Parasites and Vectors. 8(1):1–10.

Nakagawa S, Parker TH. 2015. Replicating research in ecology and evolution: Feasibility, incentives,

and the cost-benefit conundrum. BMC Biol. 13(1):1–6. doi:10.1186/s12915-015-0196-3.

http://dx.doi.org/10.1186/s12915-015-0196-3.

Nakagawa S, Schielzeth H. 2010. Repeatability for Gaussian and non-Gaussian data: A practical guide

for biologists. Biol Rev. 85(4):935–956. doi:10.1111/j.1469-185X.2010.00141.x.

Newman EA, Eisen L, Eisen RJ, Fedorova N, Hasty JM, Vaughn C, Lane RS. 2015. *Borrelia*

burgdorferi Sensu Lato Spirochetes in Wild Birds in Northwestern California: Associations with

Ecological Factors, Bird Behavior and Tick Infestation. PLoS One. 10(2):1–26.

doi:10.1371/journal.pone.0118146.

Niemelä PT, Dingemanse NJ. Meta-analysis reveals weak associations between intrinsic state and

personality. doi:10.1098/rspb.2017.2823.

Norte AC, Lobato DNC, Braga EM, Antonini Y, Lacorte G, Gonçalves M, Lopes De Carvalho I, Gern

L, Núncio MS, Ramos JA. 2013. Do ticks and *Borrelia burgdorferi* s.l. constitute a burden to birds?

Parasitol Res. 112(5):1903–1912. doi:10.1007/s00436-013-3343-1.

Oorebeek M, Kleindorfer S. 2008. Climate or host availability: What determines the seasonal

abundance of ticks? Parasitol Res. 103(4):871–875. doi:10.1007/s00436-008-1071-8.

Ots I, Horak P. 1996. Great tits *Parus major* trade health for reproduction. Proc R Soc B Biol Sci.

263(1376):1443–1447.

Peig J, Green AJ. 2009. New perspectives for estimating body condition from mass/length data: The

scaled mass index as an alternative method. Oikos. 118(12):1883–1891.

van de Pol M, Wright J. 2009. A simple method for distinguishing within- versus between-subject

effects using mixed models. Anim Behav. 77:753–758.

R Core Team. 2019. R: A Language and Environment for Statistical Computing. Vienna, Austria: R

Foundation for Statistical Computing.

Randolph SE. 2004. Tick ecology: Processes and patterns behind the epidemiological risk posed by

ixodid ticks as vectors. Parasitology. 129:S37–S65. doi:10.1017/S0031182004004925.

Santicchia F, Romeo C, Ferrari N, Matthysen E, Vanlauwe L, Wauters LA, Martinoli A. 2019. The

price of being bold? Relationship between personality and endoparasitic infection in a tree squirrel.

Mamm Biol. 97:1–8.

Sih A, Spiegel O, Godfrey S, Leu S, Bull CM. 2018. Integrating social networks, animal personalities,

movement ecology and parasites: a framework with examples from a lizard. Anim Behav. 136:195–

205.

Spiegel O, Leu ST, Bull CM, Sih A. 2017. What's your move? Movement as a link between

personality and spatial dynamics in animal populations. Ecol Lett. 20(1):3–18. doi:10.1111/ele.12708.

Spiegel O, Leu ST, Sih A, Godfrey SS, Bull CM. 2015. When the going gets tough: Behavioural type-

dependent space use in the sleepy lizard changes as the season dries. Proc R Soc B Biol Sci.

282(1819).

Stanek G. 2009. Büchse der Pandora: Krankheitserreger in Ixodes ricinus-Zecken in Mitteleuropa.

Wien Klin Wochenschr. 121:673–683. doi:10.1007/s00508-009-1281-9.

Stuber EF, Araya-Ajoy YG, Mathot KJ, Mutzel A, Nicolaus M, Wijmenga JJ, Mueller JC,

Dingemanse NJ. 2013. Slow explorers take less risk: a problem of sampling bias in ecological studies.

Behav Ecol. 24(5):1092–1098.

Tack W, Madder M, Baeten L, Vanhellemont M, Gruwez R, Verheyen K. 2012. Local habitat and

landscape affect *Ixodes ricinus* tick abundances in forests on poor, sandy soils. For Ecol Manage.

265:30–36. doi:10.1016/j.foreco.2011.10.028.

Tack W, Madder M, Baeten L, Vanhellemont M, Verheyen K. 2013. Shrub clearing adversely affects

the abundance of Ixodes ricinus ticks. Exp Appl Acarol. 60(3):411–420.

Tack K, Madder M, Baeten L, Frenne P DE, Verheyen K. 2012. The abundance of Ixodes ricinus ticks

depends on tree species composition and shrub cover. Parasitology. 139(10):1273–1281.

Taggart P, Leu S, Spiegel O, Sih A. 2018. Endure your parasites: sleepy lizard movement is not

affected by their parasites. Can J Zool. 96(12):1309–1316.

Takumi K, Sprong H, Hofmeester TR. 2019. Impact of vertebrate communities on Ixodes ricinus-

borne disease risk in forest areas. Parasit Vectors. 12(1):1–12.

Titcomb G, Allan BF, Ainsworth T, Henson L, Hedlund T, Pringle RM, Palmer TM, Njoroge L,

Campana MG, Fleischer RC, et al. 2017. Interacting effects of wildlife loss and climate on ticks and

tick-borne disease. Proc R Soc B. 284:1–10.

Verbeek ME, Drent PJ, Wiepkema PR. 1994. Consistent individual differences in early exploratory

behaviour of male great tits. Anim Behav. 48:1113–1121.

Zohdy S, Bisanzio D, Tecot S, Wright PC, Jernvall J. 2017. Aggression and hormones are associated

with heterogeneity in parasitism and parasite dynamics in the brown mouse lemur. Anim Behav.

132:109–119. doi:10.1016/j.anbehav.2017.08.002.

Figure Legends:

Figure 1. Percentages of all captured great tit adults infested with ticks from the 2017-2019 breeding

seasons. Data for plot 7 were only available in 2017.

Figure 2. Average tick burden per infested great tit adult captured in each study plot during each field

season (years 2017-2019). The error bars represent standard error and the numbers above each bar

reflect the absolute number of infested great tit adults per plot. Data for plot 7 were only available in

2017.

Figure 3. Tick presence in relation to exploratory score. For graphical purposes only, data were split

into 10 quantiles and were averaged to better show the linear relationship. Each point refers to the

mean tick presence and exploratory score for that quantile including data from all collection years

(2017-2019). Error bars report the standard error for the respective mean tick presence. The numbers

report the number of records in the respective bin. The line reports the linear relationship from the

model.

Figure 4. Tick burden of infested birds in relation to birds scaled mass index (i.e. body condition).

Each point refers to an individual infested bird record. Records from all collection years (2017-2019)

are included. The line reports the linear relationship from the model.

Figure 5. Tick presence in relation to recreational pressure index. For graphical purposes only, data

were split into 10 quantiles and were averaged to better show the linear relationship. Each point refers

to the mean tick presence and recreational pressure index for that quantile including data from all

collection years (2017-2019). Error bars report the standard error for the respective mean tick presence.

The numbers report the number of records in the respective bin. The line reports the linear relationship

from the model.

Table 1. Effect size estimates (β) and 95% credible intervals (CI) for great tit tick infestation as the
probability of infestation (n = 784 records) and tick burden of infested individuals (n = 520 records).

	Infestation Probability	Tick Burden
Fixed Effects	β (95% CI)	β (95% CI)
Intercept	0.89 (0.42, 1.37)	1.20 (0.75, 1.64)
Recreational pressure	0.21 (-0.03, 0.43)	-0.07 (-0.22, 0.08)
Breeding density	0.08 (-0.26, 0.43)	-0.04 (-0.25, 0.17)
Exploratory behaviour	0.18 (0.01, 0.35)	-0.02 (-0.12, 0.09)
Body condition	0.06 (-0.11, 0.24)	-0.11 (-0.19, -0.02)
Sex[a]	-0.06 (-0.42, 0.24)	-0.06 (-0.24, 0.12)
Age	-0.07 (-0.24, 0.09)	0.01 (-0.08, 0.10)
Random Effects	σ^2 (95% CI)	σ^2 (95% CI)
Year	0.04 (0.00, 0.17)	0.13 (0.03, 0.36)
Plot	0.47 (0.18, 0.94)	0.05 (0.02, 0.09)
Plot-Year	0.08 (0.04, 0.12)	0.07 (0.04, 0.12)
Observer	0.01 (0.00, 0.01)	0.13 (0.06, 0.23)
Nest-box	0.64 (0.55, 0.74)	0.28 (0.23, 0.32)
Individual	0.00 (0.00, 0.00)	0.45 (0.40, 0.51)
Residual	$0.35\pi^2$ ($0.35\pi^2$, $0.34\pi^2$)	1.27 (1.39, 1.18)
Adjusted Repeatability	r (95% CI)	r (95% CI)
Year	0.01 (0.00, 0.03)	0.05 (0.01, 0.13)
Plot	0.10 (0.04, 0.18)	0.02 (0.01, 0.03)

Plot-Year	0.02 (0.01, 0.02)	0.03 (0.02, 0.04)
Observer	0.00 (0.00, 0.00)	0.05 (0.03, 0.08)
Nest-box	0.14 (0.13, 0.14)	0.12 (0.11, 0.11)
Individual	0.00 (0.00, 0.00)	0.19 (0.18, 0.18)
Residual	0.74 (0.82, 0.63)	0.53 (0.64, 0.42)

[a]Differences between sexes (reference category = female)

535 **1.**

537 **2.**

543   **3.**

4.

548   **5.**