A validated quantitative method for the assessment of neuroprotective barrier

impairment in neurodegenerative disease models

Vinod Kumar[1*], John D. Lee[1*], Elizabeth J. Coulson[1, 2] and Trent M. Woodruff[1, 2§]

[1]School of Biomedical Sciences, the University of Queensland, Brisbane, QLD, 4072,

Australia

[2]Queensland Brain Institute, the University of Queensland, St Lucia, Brisbane, QLD, 4072,

Australia

[§]Correspondence to Trent M. Woodruff, School of Biomedical Sciences, Skerman Building,

University of Queensland, St. Lucia QLD 4072. Australia.

Phone: +61-7-336 52924

Fax: +61-7-336-51766

Email: t.woodruff@uq.edu.au

*Equal contribution

Running title: Na-Fl is a suitable indicator of BBB/BSCB breakdown

Keywords: blood-brain barrier, blood-spinal cord barrier, LPS, neurodegeneration, Na-Fl

Abbreviations used

AD, Alzheimer's disease; ALS, amyotrophic lateral sclerosis; BBB, blood brain barrier; BSCB, blood spinal cord barrier; CNS, central nervous system; FITC, fluorescein isothiocyanate; HD, Huntington's disease; LPS, lipopolysaccharide; Na-Fl, sodium fluorescein; PD, Parkinson's disease; RRID, Research Resource Identifier (see scicrunch.org); WT, wild-type

ORCID ID

Vinod Kumar: 0000-0002-7263-9917

John D. Lee: 0000-0002-9976-7396

Elizabeth J. Coulson: 0000-0003-3783-6197

Trent M. Woodruff: 0000-0003-1382-911X

Abstract

The blood brain barrier (BBB) and blood spinal cord barrier (BSCB) are highly specialised

structures that limit molecule entry from the blood and maintain homeostasis within the central

nervous system (CNS). BBB and BSCB breakdown are associated with multiple

neurodegenerative diseases. Given the key role of neuroprotective barrier impairment in

neurodegeneration, it is important to identify an effective quantitative method to assess barrier

integrity in animal models. In the present study, we developed and validated a quantitative

method for assessing BBB and BSCB integrity using sodium fluorescein, a compound that

outperformed other fluorescent dyes. We demonstrated using this method that multiple CNS

regions progressively increase in permeability in models of Huntington's disease and

amyotrophic lateral sclerosis, while biphasic disruption occurred in a mouse model of

Alzheimer's disease with disease progression. Collectively, we report a quantitative

fluorometric marker with validated reproducible experimental methods, that allows the

effective assessment of BBB and BSCB integrity in animal models. This method could be

useful to further the understanding of the contribution of these neuroprotective barriers to

neurodegeneration processes.

1 Introduction

The blood brain barrier (BBB) and blood spinal cord barrier (BSCB) are highly specialised

epithelial structures linked through intercellular junctional complexes, termed tight junctions.

These structures maintain homeostasis within the central nervous system (CNS) by ensuring

strict regulation of molecule transport between circulating blood and CNS tissues (Zhao *et al.*

2015). The primary physiological function of these barriers is to maintain the chemical

composition of the CNS, and protect against toxic molecules, to support neuronal cell health.

BBB and BSCB breakdown, due to disruption of tight junctions or epithelial cell integrity, can

lead to altered transport of molecules between the CNS and blood, which can contribute to the

pathological processes resulting in progressive structural and functional loss of neurons in

multiple neurodegenerative diseases (Ellrichmann *et al.* 2013; Harry & Kraft 2008; Hirsch &

Hunot 2009; Lobsiger & Cleveland 2007).

There is evidence to support the notion that neuroprotective barrier impairment occurs within

the progression of these neurodegenerative diseases, including in Alzheimer's disease (AD)

(Zlokovic 2011), Parkinson's disease (PD) (Brochard *et al.* 2009; Carvey *et al.* 2005; Chen *et*

al. 2008; Sarkar *et al.* 2014), Huntington's disease (HD) (Drouin-Ouellet *et al.* 2015) and

amyotrophic lateral sclerosis (ALS) (Garbuzova-Davis *et al.* 2007b). Damaged BBB and BSCB

integrity and resulting increased permeability can lead to oedema, neurotoxic molecule

accumulation, as well as enhancing the infiltration of immune cells to the CNS, which can

accelerate the process of neurodegeneration (Arima *et al.* 2013; Bell *et al.* 2010; Stokum *et al.*

2016). This interaction between neuroinflammation, neurodegeneration and neuroprotective

barrier integrity can also influence drug bioavailability to the CNS, which may affect the

activity of neuroprotective therapies. For example, Tarenflurbil used in AD was shown to have

poor BBB penetration, which may have led to its failure in clinical trials (Wan *et al.* 2009).

Given the key involvement of neuroprotective barrier impairment in neurological and

neurodegenerative conditions, it is therefore important to identify an effective quantitative

method to assess barrier impairment and permeability changes during disease progression in

animal models of these conditions.

Previous studies have utilized different dyes to measure the permeability of BBB and BSCB,

with Evans blue dye being the most commonly reported (Saunders *et al.* 2015). However,

numerous studies have indicated that Evans blue is a substandard marker for BBB permeability,

due to significant binding to tissues (Dallal & Chang 1994; Lindner & Heinle 1982), solution

instability, *in vivo* toxicity (demonstrated in rats, monkeys, dogs, cats and rabbits), and lack of

a validated quantitative determination method. Furthermore, Evans blue relies on binding to

the large plasma protein albumin, and subsequent transport across the BBB, which only occurs

during later stages of barrier disruption. In contrast to Evans blue, more recent fluorescent-

based dyes have been used extensively as a marker for permeability studies including sodium

fluorescein (Na-Fl) (Hoffman & Olszewski 1961; Malmgren & Olsson 1980), fluorescein

isothiocyanate (FITC) labelled albumin (Boutoille *et al.* 2009; King-VanVlack *et al.* 2003), and

FITC labelled dextran-70 (Hoffmann *et al.* 2011; Oliver *et al.* 1984). However, no study to date

has directly compared the relative utility of these dyes in assessing BBB and BSCB impairment

and provided a robust validated quantitative method.

In the present study, we therefore initially examined these three different dyes (Na-Fl, FITC

labelled albumin and FITC labelled dextrans-70) as potential markers for neuroprotective

permeability in the brains of lipopolysaccharide (LPS)-injected mice to determine the most

suitable and sensitive marker for BBB and BSCB permeability studies. We developed a

quantitative method that demonstrated Na-Fl as the most sensitive marker for measuring

neuroprotective barrier integrity. We then utilized this method in mouse models of AD, HD, and ALS, where we identified perturbed barrier integrity in different regions of the brain and spinal cord of these animals. Our data therefore reveal a novel quantitative method for monitoring BBB and BSCB integrity in animal models of neurodegeneration.

Material and methods

Animals

All animal procedures were approved by the University of Queensland Animal Ethics Committee (Permit Number: 291/15) and conducted in accordance with the Australian Code of Practice for the Care and Use of Animals for Scientific Purposes (8th edition, 2013). All experiments were conducted in compliance with ARRIVE guidelines. This study was not pre-registered, with no randomisation nor blinding performed. Animals were group housed 3 ~ 4 per cage on arrival in the University of Queensland animal facility. Mice were housed in a 12-h light: dark phase (on at 06:00 h and off at 18:00 h) with room temperature maintained at 24 ± 2 °C. Animals had free access to *ad-libitum* food and water. All experimental procedures were conducted between 10:00 h and 12:00 h. Animal sources and details are outlined below.

Wild-type mice: C57BL/6J male mice aged 10-12 weeks old were obtained from the Animal Resources Centre in Canning Vale, Australia. We used a total of 82 mice for selecting and assessing appropriate fluorescent markers by comparing uptake results of the markers FITC dextran, FITC albumin and Na-Fl in the brain and serum of mice over time after an intraperitoneal LPS injection.

Transgenic R6/1 mice (transgenic mouse model for Huntington's disease):

Transgenic R6/1 mice (B6-CBA-Tg (HDexon1) 61Gpb/1J) expressing a transgene containing

the human HD gene were obtained from Jackson Laboratory (RRID: IMSR_JAX:002809;

(Mangiarini *et al.* 1996)) and bred with female mice on CBA/BL6 background to produce R6/1

transgenic and non-transgenic (wild-type; WT) mice. Male transgenic R6/1 and matched WT

littermate mice were used at various stages of disease progression (6 weeks, 12 weeks, 20 weeks

and 32 weeks) (Naver *et al.* 2003). We used a total of 40 mice (20 WT and 20 R6/1) to assess

BBB/BSCB integrity at four different stages of disease progression.

Transgenic hSOD1^{G93A} mice (transgenic mouse model for amyotrophic lateral sclerosis):

Transgenic hSOD1^{G93A} mice (B6-Cg-Tg (SOD1-G93A) 1Gur/J) carrying a mutant form of the

human *SOD1* gene were obtained from Jackson Laboratory (RRID: IMSR_JAX:004435;

(Rosen *et al.* 1993)) and bred on C57BL/6J background to produce transgenic hSOD1^{G93A} and

WT littermates as control mice. Male hSOD1^{G93A} and WT mice were used at various stages of

disease progression (30 days, 70 days, 130 days and 150 – 175 days) as described previously

(Lee *et al.* 2013). We used a total of 40 mice (20 WT and 20 hSOD1^{G93A}) to assess BBB/BSCB

integrity at four different stages of disease progression.

Transgenic APP/PS1 mice (transgenic mouse model for Alzheimer's disease):

Double transgenic APP/PS1 mice (B6-Cg-Tg (APPswe, PSEN1dE9)85Dbo/Mmjax)

expressing a chimeric mouse/human amyloid precursor protein and a mutant human presenilin

1 were source from Jackson Laboratory (RRID: MMRRC_034832 – JAX; (Jankowsky *et al.*

2004)) and bred on C57BL/6J background to produce transgenic APP/PS1 and WT littermates.

Male APP/PS1 and WT mice were used at 4 and 18 months, ages correlating to early, and late

stage disease phenotypes (Radde *et al.* 2006; Serneels *et al.* 2009; Rupp *et al.* 2011). We used

a total of 20 mice (10 WT and 10 APP/PS1) to assess BBB/BSCB integrity at early and late stages of disease.

Experimental conditions for quantification and selection of fluorescent marker permeability studies

Fluorescence stability of markers during the extraction and analysis process were first tested by studying effect of different solvents as a loading solution, the effect of solvents during the extraction process, extraction efficiency of the process, the influence of matrix/solvents on the fluorescence of markers, and loss of fluorescence during the extraction and analysis steps (refer to Supplementary methods for full methodology and results). From these validation steps, TRIS buffer (pH 7.4) was identified as the most suitable solvent.

The selection of an appropriate fluorescent marker was accomplished by comparing uptake results of the markers in the brain and serum of mice after an intraperitoneal LPS injection (Clone 0111:B4; Sigma, Cat#L2630, 4 mg kg^{-1} of mice body weight). After 24 hours, mice were then anaesthetized with intraperitoneally injected zolazapam (50 mg kg^{-1}) and xylazine (12 mg kg^{-1}) to minimize animal suffering during experiments. This was followed by 100 µl intravenous injection of FITC albumin (Sigma, Cat#A9771), FITC dextran (Sigma, Cat#FD40S) and Na-Fl (Sigma, Cat#46960) at 100 mg mL^{-1}. After one hour, blood samples were collected via cardiac puncture and allowed to clot at room temperature for 15 minutes. Samples were then centrifuged at 2,000 x g for 10 minutes at 4°C for serum collection. After blood collection, mice were immediately perfused transcardially with 50 mL of 1 x PBS to remove any remaining fluorescent marker in the circulation and euthanized via cervical dislocation. Whole brain (which also consists of extracellular fluids such as the cerebrospinal fluid (CSF)) was carefully collected, washed with 1 x PBS, dried with Whatman® gel blotting paper and homogenized in

an equal volume of TRIS buffer (pH 7.4). Serum samples were deproteinized with acetonitrile

(1:3; Merck, Cat# 1000292500), vortexed, sonicated and centrifuged at 13,000 x g for 15

minutes at room temperature. Supernatant samples were evaporated to a dry state using a

centrivap concentrator (Labconco) at room temperature and reconstituted in an equal volume

of TRIS buffer (pH 7.4). Brain tissue samples were homogenized with milli Q water (1:2 ratio

to the tissue weight). Homogenized brain samples were deproteinized with acetonitrile (1:6

ratio to the tissue weight) and processed via a similar method used for serum samples. Dried

brain samples were reconstituted in equal weight-volume of TRIS buffer (pH 7.4). 10 µl of

reconstituted solutions were diluted with 200 µl of TRIS buffer (pH 7.4) in 96-well black plate

(Corning Inc, Cat#3603) along with standard curve samples in same matrix. The fluorescence

levels were measured using a microplate reader (FlexStation 3) with λ_{ex} 460 nm and λ_{em} 515

nm for Na-Fl, λ_{ex} 495 nm and λ_{em} 520 nm for FITC-albumin and λ_{ex} 490 nm and λ_{em} 530 nm

for FITC-dextran respectively. The amount of fluorescent marker was calculated from

fluorescence levels via a standard curve prepared in a similar manner using brain and serum

obtained from mice without any treatment. Raw fluorescent data values obtained for serum and

brain are located at Open Science Framework (DOI 10.17605/OSF.IO/Q6XGB). Fluorescence

uptake was calculated by comparing ratio of fluorescent marker level in the brain with the level

of fluorescent marker in the serum. Fluorescence values were corrected with the fluorescence

of blank brain and serum samples obtained from mice without any fluorescent marker or LPS

treatment and treated with the same method as above.

Application of a validated quantitative fluorometric method in mouse models of neurodegeneration

The applicability of the developed and validated quantitative fluorometric method was assessed

by performing neuroprotective barrier integrity studies in various chronic mouse models of

neurodegeneration. Mice ($n = 5$) at various disease stages (Table 1) were anaesthetized with an

intraperitoneal injection of zolazapam (50 mg kg^{-1}) and xylazine (12 mg kg^{-1}) to minimize

animal suffering during the experiments. Anaesthetized mice were then intravenously

administered with 100 µl of Na-Fl (100 mg mL^{-1}) via tail vein. After 15 minutes from injection,

blood samples were collected by cardiac puncture for serum extraction. Animals were

transcardially perfused immediately with 1 x PBS to remove circulating Na-Fl and euthanized

via cervical dislocation. Following this, whole brain and spinal cord tissue was collected and

different brain regions (cortex, striatum, hippocampus and cerebellum) were isolated on ice

using a mouse coronal brain matrix (Cat#U040-C, ProSciTech) under the microscope as per

the suggested protocol (Spijker 2011). Samples were stored at -80°C for further processing and

analysis using the developed quantitative fluorometric method as described above.

Fluorescence levels in brain and spinal cord samples were measured as describe above.

Fluorescence values were back calculated for the level of Na-Fl in the serum and tissues using

the standard curve samples prepared in the same matrix and analyzed as above. Results are

presented as relative levels of fluorescence uptake in R6/1, hSOD1^{G93A} and APP/PS1 mice

compared with expression in their respective litter-matched non-transgenic WT controls.

Complete data values obtained prior to normalisation to WT controls are also found at Open

Science Framework (DOI 10.17605/OSF.IO/Q6XGB).

Statistical analysis

All results are reported as mean ± standard error of mean and statistical analysis performed

using GraphPad Prism 8.3.1 (RRID:SCR_002798). No sample size calculation was performed

prior to starting the experiments. No tests for outliers (i.e. no exclusion of data points) and no

exclusion criteria were pre-determined for the study. Normality was tested using Shapiro –

Wilk test. The statistical difference between FITC albumin, FITC dextran and Na-Fl for

percentage uptake in the brain were analyzed using a one-way ANOVA and a *post-hoc* Tukey's test. The statistical difference for levels of fluorescence uptake between WT and R6/1, WT and hSOD1^{G93A} and WT and APP/PS1 mice were analyzed using a two-tailed Student's t-test for each stage of disease progression. The differences were considered significant when $P < 0.05$.

Results

Sodium fluorescein has greater sensitivity as an indicator of BBB disruption than FITC-dextran or FITC-albumin

Selection of an appropriate fluorescent marker for permeability studies was based upon the percentage uptake of the fluorescent marker, which was quantitatively determined via a validated fluorometric method. We utilized peripheral administration of LPS to induce BBB disruption after 24 hours as previously documented (Banks *et al.* 2015; Nonaka *et al.* 2005). We then determined the uptake of fluorescent dye into the brain 1 hour after intravenous dye injection. The percentage uptake of Na-Fl (~50%) was found to be the highest compared to either FITC-dextran-70 (~1%) or FITC-albumin (~17%) (**Figure 1A**). We next extended this analysis by comparing BBB permeability changes using FITC-albumin and Na-Fl (both having detectible uptake in the brain after 1 hour), over 24 hours. Fluorescence uptake of FITC-albumin in the brain was significantly increased at 0.25 hour (1.4% vs. 4.8%), 1 hour (6.8% vs. 16.7%) and 24 hours (8.4% vs. 24%) in saline-injected versus LPS-injected mice respectively (**Figure 1B**). Fluorescence uptake of Na-Fl in the brain was also significantly increased at 0.25 hour (8.5% vs. 40%), 1 hour (10.4% vs. 67.7%) and 24 hours (3.8% vs. 17.2%) in saline-injected versus LPS-injected mice respectively (**Figure 1C**). Notably, the fold-increases in dye uptake between saline and LPS groups, were significantly larger for Na-Fl than FITC-albumin, peaking 1 hour after dye injection. We therefore identified 1-hour Na-Fl to be an ideal method to progress for future validation.

Progressive breakdown of neuroprotective barriers in the R6/1 transgenic mouse model of Huntington's disease

We first applied our method to investigate BBB and BSCB integrity at different disease stages in the R6/1 transgenic mouse model of HD. We observed no change in Na-Fl uptake in cortex, striatum, cerebellum and spinal cord between WT and R6/1 transgenic mice at 6 weeks of age, which is considered to be a pre-symptomatic stage of disease. Na-Fl uptake in the cortex and striatum progressively increased in R6/1 mice at 12 (1.7-fold, 1.8-fold), 20 (2.1-fold, 2.4-fold) and 32 (2.3-fold, 2.9-fold) weeks when compared to WT mice (**Figure 2A** and **2B**). The cerebellum was less affected, with Na-Fl uptake only increased by 2.0-fold in R6/1 mice at 32 weeks of age (**Figure 2C**). Interestingly, Na-Fl was also increased in the spinal cord of R6/1 mice at 20 and 32 weeks by 2.7-fold respectively (**Figure 2D**).

Breakdown of neuroprotective barriers in the hSOD1^{G93A} transgenic mouse model of amyotrophic lateral sclerosis

Next, we assessed neuroprotective barrier integrity in hSOD1^{G93A} transgenic mice. Na-Fl uptake was measured in cortex, striatum, cerebellum and spinal cord at progressive ages of disease progression. Similar to R6/1 mice, we observed no changes in Na-Fl uptake in cortex, striatum, cerebellum and spinal cord between WT and hSOD1^{G93A} transgenic mice at the pre-symptomatic age of 30 days. However, Na-Fl uptake progressively increased in the cortex of hSOD1^{G93A} mice at 70 (1.2-fold), 130 (1.5-fold) and 175 (1.6-fold) days of age (**Figure 3A**). Na-Fl also increased in the striatum of hSOD1^{G93A} mice at these same ages (**Figure 3B**). BBB permeability increases were also observed in the cerebellum of hSOD1^{G93A} mice, with significant Na-Fl uptake observed at 70, 130 and 175 days (**Figure 3C**). Finally, BSCB permeability was also detected, with Na-Fl uptake in the spinal cord gradually increased in hSOD1^{G93A} mice by 1.1-, 1.3- and 2.0-fold at 70, 130 and 175 days respectively (**Figure. 3D**).

Neuroprotective barriers in the APP/PS1 transgenic mouse model of Alzheimer's disease

display biphasic alterations at different diseases stages

To measure possible alterations in neuroprotective barriers of APP/PS1 transgenic mice, Na-Fl uptake was measured in cortex, striatum, hippocampus, cerebellum and spinal cord at an early (4 months) and late (18 months) stage of disease. Brain and spinal cord tissues from APP/PS1 transgenic mice at the early stages of disease showed an overall increase in barrier permeability (**Figure 4**), with significant increases in Na-Fl uptake observed in the striatum (**Figure 4C**) and spinal cord (**Figure 4E**). Notably, by 18 months of age, these same tissues had reduced levels of Na-Fl, with decreases of up to 50% observed in brain regions and spinal cord (**Figure 4B**).

Discussion

BBB and BSCB breakdown are associated with a number of neurodegenerative diseases, including AD, PD, HD and ALS (Desai *et al.* 2007; Di Pardo *et al.* 2017; Garbuzova-Davis *et al.* 2007a; Garbuzova-Davis & Sanberg 2014; Garbuzova-Davis *et al.* 2007b; Gray & Woulfe 2015). The understanding of the underlying mechanisms that initiates the opening or closure of the BBB and BSCB is therefore of therapeutic significance in many neurodegenerative diseases and is commonly investigated by researchers. However, methods to investigate BBB and BSCB permeability *in-vivo* are often associated with technical difficulties that depend on subjective quantification of fluorescence images, or semi-quantitative evaluation of endogenous molecules. To address this, prior studies have utilized an abundance of fluorescent markers for *in-vivo* and *in-vitro* permeability studies, each with their own strength and weaknesses (Saunders *et al.* 2015; de Vries *et al.* 1995; Ravenstijn *et al.* 2008; de Lange *et al.* 1998). Methods used for the detection of these fluorescent markers also display great variation in their quantitation methodology between studies. Selection of an ideal validated method for assessing

neuroprotective barrier permeability is therefore difficult due to lack of prior published results

for quantitative determination. The present research, supported by previously published

methods (de Lange *et al.* 1998; Ravenstijn *et al.* 2008; de Vries *et al.* 1995), therefore instigated

the selection and identification of a suitable fluorescent marker for CNS permeability studies

using a validated fluorometric method of quantitative analysis. The method was developed

using an acute LPS-induced BBB breakdown model and then its application verified by

assessing neuroprotective barrier permeability in various mouse model of neurodegeneration.

The BBB and BSCB are highly specialized epithelial structures that are important in

maintaining the CNS homeostasis and preventing the entry of foreign toxic substances. It

undergoes degeneration during disease processes of neurodegenerative diseases and has a re-

emerging interest as a mechanistic and therapeutic target for intervention. In spite of their

limitations, use of dyes to detect barrier breakdown have remained widespread, where

numerous studies have utilized different dyes including, Evans blue, fluorescein, horse radish

peroxidase, dextran, albumin and trypan blue (Saunders *et al.* 2015). To the best of our

knowledge, our study is the first to compare three widely used fluorescent dye markers (FITC-

dextran, FITC-albumin and Na-Fl) to determine the most suitable marker to quantitatively

evaluate BBB integrity. We demonstrate that 1-hour after the injection of these dyes to LPS-

induced BBB-injured mice, Na-Fl showed the greatest percentage uptake in the brain. Although

this has not been directly compared previously, this is consistent with previous studies

suggesting that Na-Fl only binds weakly to proteins, and is therefore considered to be an

effective low molecular weight marker for BBB studies in contrast to dyes that bind to proteins

(Wolman *et al.* 1981). Although Na-Fl cannot be detected by visual examination, a quantitative

fluorometric method utilized in the current study may enable detection of subtle alterations in

BBB/BSCB permeability and be the earliest and the most sensitive indicator of BBB/BSCB

disruption (Kaya & Ahishali 2011). Our studies also demonstrated that Na-Fl uptake in the brain was reduced after 24 hours of injection (when compared to 15 minutes and 1 hour), suggesting rapid clearance of Na-Fl in the brain parenchyma. Collectively, as small molecule, Na-Fl is considered to be less toxic than other dye tracers, is freely diffusible, minimally binds to proteins and can be evaluated quantitatively and reproducibly. We therefore posit this method provides an ideal marker of assessing neuroprotective barrier integrity. In contrast to Na-Fl, FITC-albumin reflects a significant increase in time dependant percentage uptake, supporting its potential as a marker for longer-duration permeability studies.

In pathological conditions, BBB permeability is increased due to various biochemical mediators (Abbott 2000). BBB permeability also increases with ageing (Erdő *et al.* 2017), and our data obtained from WT mice from the transgenic mouse colonies across different ages supports this (Supplementary Figure 1). Increased permeability favours the transport of peripheral immune cells, inflammatory products and other circulating substances to the brain and spinal cord, resulting in breakdown of tight junction molecules and ultimately, disruption of BBB and BSCB (Ballabh *et al.* 2004; Abbott *et al.* 2010; de Vries *et al.* 1997). BBB and BSCB disruption and malfunction are involved in various neurological disorders including AD, PD, HD and ALS (Desai *et al.* 2007; Di Pardo *et al.* 2017; Garbuzova-Davis *et al.* 2007a; Garbuzova-Davis & Sanberg 2014; Garbuzova-Davis *et al.* 2007b; Gray & Woulfe 2015). Using different mouse models of HD, ALS and AD, we investigated barrier integrity in different regions of the brain and spinal cord using our quantitative method of Na-Fl extravasation. We demonstrated that there is an early disruption of BBB and BSCB permeability in R6/1 mouse model of HD as we showed increased Na-Fl uptake in cortex, striatum and spinal cord, in parallel with the cognitive and motor deficits present in this model (Ayton *et al.* 2019; Brooks *et al.* 2012). This early disruption of BBB is further supported by

previous studies showing early vascular impairment and BBB dysfunction in R6/2 mice, with a reduction in expression of tight junction proteins claudin-5 and occludin, leading to increased paracellular permeability (Di Pardo *et al.* 2017; Drouin-Ouellet *et al.* 2015; Luissint *et al.* 2012). This validates the applicability of this fluorometric method to measure subtle changes in BBB/BSCB integrity.

We also demonstrated that there is an early disruption of BBB and BSCB permeability in hSOD1[G93A] mouse model of ALS based on increased Na-Fl uptake in cortex, striatum, cerebellum and spinal cord. Similar to HD, these findings support previous studies demonstrating damage to BSCB integrity in hSOD1[G93A] mice via ultrastructural capillary alterations and increased leakage in the spinal cord at both early and late stages of disease (Garbuzova-Davis *et al.* 2007a; Garbuzova-Davis *et al.* 2007b; Lee *et al.* 2017; Zhong *et al.* 2008). These findings are further supported by numerous studies demonstrating decreased expression of tight junction proteins integral to the integrity of the BSCB, such as zona occludens-1, occludin and claudin-5 in the spinal cord of hSOD1 transgenic mice and ALS patients (Henkel *et al.* 2009; Meister *et al.* 2015; Winkler *et al.* 2014; Zhong *et al.* 2008). Collectively, our study complements these previous findings given our method was able to detect early barrier impairments very early in disease, at ages that precede significant motor neuron death (Lee *et al.* 2013), which may have pathophysiological relevance. As neuroinflammation and microgliosis are commonly correlated with barrier impairment (Erdő *et al.* 2017), it will be of interest to explore these association in future studies, particularly as neuroinflammation is a key driver of pathology in hSOD1[G93A] mice (Chiarotto *et al.* 2019).

We also demonstrated an apparent biphasic BBB/BSCB disruption in APP/PS1 mouse model of AD. Unlike HD and MND, neuroprotective barrier disruption in APP/PS1 mouse model of

AD was only evident in early in the disease, at an age where there is minimal cognitive dysfunction (Zhu *et al.* 2017; Bien-Ly *et al.* 2015). Interestingly, unlike the other disease models studied, there was a notable decrease in Na-Fl uptake in brain and spinal cord regions at the late stage of disease, an age where is significant cognitive decline (Gengler *et al.* 2010; Milne *et al.* 2019). Although it is possible that the result is due to a faster clearance of the dye from a highly permeable BBB, the decrease in dye uptake in APP/PS1 mice late in disease is consistent with previous studies demonstrating thickening of the basement membrane in AD model mice and humans (Hawkes *et al.* 2013; Lepelletier *et al.* 2017). Although the causes of the basement membrane thickening in AD have not been fully elucidated to date, decreased clearance of amyloid beta or increased astrocyte production of basement membrane proteins like perlecan and fibronectin have been suggested to contribute to this thickening of basement membrane (Deane & Zlokovic 2007; Okoye & Watanabe 1982; Thomsen *et al.* 2017). Importantly, the progressive molecular changes in barrier permeability of multiple neurodegenerative disease models tested in previous studies mirrored the changes in Na-Fl levels in the brain and spinal cord of our current study. This strongly validates the sensitivity and accuracy of our validated fluorometric method to measure subtle changes in BBB/BSCB integrity. This may have implications for future preclinical studies that aim to correlate the extent of barrier impairment and permeability changes with disease progression.

In summary, we have identified a validated, effective quantitative method for assessing BBB and BSCB integrity using Na-Fl. We demonstrated using this method that multiple CNS regions progressively increase in permeability in the R6/1 mouse model of HD and the hSOD1[G93A] mouse model of ALS. By contrast, whilst increased BBB/BSCB permeability was observed in young APP/PS1 mouse model of AD, later stage diseased mice showed marked reduction in CNS permeability. Collectively, we report a quantitative fluorometric marker with validated

reproducible experimental methods, that allows the effective assessment of BBB and BSCB integrity in animal models. This method could be useful to further the understanding of the contribution of these neuroprotective barriers to neurodegeneration processes, and to assess the efficacy of neuroprotective agents.

Acknowledgements and conflict of interest disclosure

The research was funded by grants from the MNDRIA to TMW (GIA1865) and JDL (GIA1830), and a National Health and Medical Research Council (NHMRC) Project grant (APP1082271) to TMW. The authors would like to sincerely thank the University of Queensland Biological Resources for the animal care and husbandry. We also thank Maryam Shayegh for her technical support with genotyping mice. The authors declare that they have no competing interests. An earlier version of this manuscript was uploaded to bioRxiv: https://www.biorxiv.org/content/10.1101/2020.03.06.979930v1.

Author's contributions

VK, JDL, and TMW conceived the project. VK, JDL, EJC and TMW designed the study. VK and JDL performed experiments. All authors contributed to the analyses and/or interpreted the data. VK and JDL wrote the paper with TMW, with editing contributions from EJC. All authors read and approved the final manuscript.

References

Abbott, N. J. (2000) Inflammatory mediators and modulation of blood-brain barrier permeability. *Cell Mol Neurobiol* **20,** 131-147.

Abbott, N. J., Patabendige, A. A., Dolman, D. E., Yusof, S. R. and Begley, D. J. (2010) Structure and function of the blood-brain barrier. *Neurobiol Dis* **37,** 13-25.

Arima, Y., Kamimura, D., Sabharwal, L., Yamada, M., Bando, H., Ogura, H., Atsumi, T. and Murakami, M. (2013) Regulation of immune cell infiltration into the CNS by regional neural inputs explained by the gate theory. *Mediators Inflamm* **2013,** 898165.

Ayton, S., Lei, P., Appukuttan, A. T., Renoir, T., Foliaki, S., Chen, F., Adlard, P. A., Hannan, A. J. and Bush, A. I. (2019) Brain Zinc Deficiency Exacerbates Cognitive Decline in the R6/1 Model of Huntington's Disease. *Neurotherapeutics*.

Ballabh, P., Braun, A. and Nedergaard, M. (2004) The blood-brain barrier: an overview: structure, regulation, and clinical implications. *Neurobiol Dis* **16,** 1-13.

Banks, W. A., Gray, A. M., Erickson, M. A. et al. (2015) Lipopolysaccharide-induced blood-brain barrier disruption: roles of cyclooxygenase, oxidative stress, neuroinflammation, and elements of the neurovascular unit. *J Neuroinflammation* **12,** 223.

Bell, R. D., Winkler, E. A., Sagare, A. P., Singh, I., LaRue, B., Deane, R. and Zlokovic, B. V. (2010) Pericytes control key neurovascular functions and neuronal phenotype in the adult brain and during brain aging. *Neuron* **68,** 409-427.

Bien-Ly, N., Boswell, C. A., Jeet, S. et al. (2015) Lack of Widespread BBB Disruption in Alzheimer's Disease Models: Focus on Therapeutic Antibodies. *Neuron* **88,** 289-297.

Boutoille, D., Marechal, X., Pichenot, M., Chemani, C., Guery, B. and Faure, K. (2009) FITC-albumin as a marker for assessment of endothelial permeability in mice: comparison with 125I-albumin. *Experimental lung research* **35,** 263-271.

Brochard, V., Combadiere, B., Prigent, A. et al. (2009) Infiltration of CD4+ lymphocytes into the brain contributes to neurodegeneration in a mouse model of Parkinson disease. *J Clin Invest* **119,** 182-192.

Brooks, S. P., Janghra, N., Workman, V. L., Bayram-Weston, Z., Jones, L. and Dunnett, S. B.

(2012) Longitudinal analysis of the behavioural phenotype in R6/1 (C57BL/6J)

Huntington's disease transgenic mice. *Brain Res Bull* **88,** 94-103.

Carvey, P. M., Zhao, C. H., Hendey, B., Lum, H., Trachtenberg, J., Desai, B. S., Snyder, J.,

Zhu, Y. G. and Ling, Z. D. (2005) 6-Hydroxydopamine-induced alterations in blood-

brain barrier permeability. *Eur J Neurosci* **22,** 1158-1168.

Chen, X., Lan, X., Roche, I., Liu, R. and Geiger, J. D. (2008) Caffeine protects against MPTP-

induced blood-brain barrier dysfunction in mouse striatum. *J Neurochem* **107,** 1147-

1157.

Chiarotto, G. B., Cartarozzi, L. P., Perez, M., Biscola, N. P., Spejo, A. B., Gubert, F., França

Junior, M., Mendez-Otero, R. and de Oliveira, A. L. R. (2019) Tempol improves

neuroinflammation and delays motor dysfunction in a mouse model (SOD1G93A) of

ALS. *Journal of Neuroinflammation* **16,** 218.

Dallal, M. M. and Chang, S. W. (1994) Evans blue dye in the assessment of permeability-

surface area product in perfused rat lungs. *J Appl Physiol (1985)* **77,** 1030-1035.

de Lange, E. C., de Bock, G., Schinkel, A. H., de Boer, A. G. and Breimer, D. D. (1998) BBB

transport and P-glycoprotein functionality using MDR1A (-/-) and wild-type mice.

Total brain versus microdialysis concentration profiles of rhodamine-123. *Pharm Res*

15, 1657-1665.

de Vries, H. E., Eppens, E. F., Prins, M., Kuiper, J., van Berkel, T. J. C., de Boer, A. G. and

Breimer, D. D. (1995) Transport of a Hydrophilic Compound into the Cerebrospinal

Fluid During Experimental Allergic Encephalomyelitis and After Lipopolysaccharide

Administration. *Pharmaceutical Research* **12,** 1932-1936.

de Vries, H. E., Kuiper, J., de Boer, A. G., Van Berkel, T. J. and Breimer, D. D. (1997) The

blood-brain barrier in neuroinflammatory diseases. *Pharmacol Rev* **49,** 143-155.

Deane, R. and Zlokovic, B. V. (2007) Role of the blood-brain barrier in the pathogenesis of Alzheimer's disease. *Curr Alzheimer Res* **4,** 191-197.

Desai, B. S., Monahan, A. J., Carvey, P. M. and Hendey, B. (2007) Blood-brain barrier pathology in Alzheimer's and Parkinson's disease: implications for drug therapy. *Cell Transplant* **16,** 285-299.

Di Pardo, A., Amico, E., Scalabri, F. et al. (2017) Impairment of blood-brain barrier is an early event in R6/2 mouse model of Huntington Disease. *Sci Rep* **7,** 41316.

Drouin-Ouellet, J., Sawiak, S. J., Cisbani, G. et al. (2015) Cerebrovascular and blood-brain barrier impairments in Huntington's disease: Potential implications for its pathophysiology. *Annals of neurology* **78,** 160-177.

Ellrichmann, G., Reick, C., Saft, C. and Linker, R. A. (2013) The role of the immune system in Huntington's disease. *Clin Dev Immunol* **2013,** 541259.

Erdő, F., Denes, L. and de Lange, E. (2017) Age-associated physiological and pathological changes at the blood-brain barrier: A review. *Journal of cerebral blood flow and metabolism : official journal of the International Society of Cerebral Blood Flow and Metabolism* **37,** 4-24.

Garbuzova-Davis, S., Haller, E., Saporta, S., Kolomey, I., Nicosia, S. V. and Sanberg, P. R. (2007a) Ultrastructure of blood-brain barrier and blood-spinal cord barrier in SOD1 mice modeling ALS. *Brain research* **1157,** 126-137.

Garbuzova-Davis, S. and Sanberg, P. R. (2014) Blood-CNS Barrier Impairment in ALS patients versus an animal model. *Front Cell Neurosci* **8,** 21.

Garbuzova-Davis, S., Saporta, S., Haller, E., Kolomey, I., Bennett, S. P., Potter, H. and Sanberg, P. R. (2007b) Evidence of compromised blood-spinal cord barrier in early and late symptomatic SOD1 mice modeling ALS. *PLoS One* **2,** e1205.

1   Gengler, S., Hamilton, A. and Holscher, C. (2010) Synaptic plasticity in the hippocampus of a

2   APP/PS1 mouse model of Alzheimer's disease is impaired in old but not young mice.

3   *PLoS One* **5,** e9764.

4   Gray, M. T. and Woulfe, J. M. (2015) Striatal blood-brain barrier permeability in Parkinson's

5   disease. *J Cereb Blood Flow Metab* **35,** 747-750.

6   Harry, G. J. and Kraft, A. D. (2008) Neuroinflammation and microglia: considerations and

7   approaches for neurotoxicity assessment. *Expert Opin Drug Metab Toxicol* **4,** 1265-

8   1277.

Hawkes, C. A., Gatherer, M., Sharp, M. M., Dorr, A., Yuen, H. M., Kalaria, R., Weller, R. O.

and Carare, R. O. (2013) Regional differences in the morphological and functional

effects of aging on cerebral basement membranes and perivascular drainage of amyloid-

beta from the mouse brain. *Aging Cell* **12,** 224-236.

Henkel, J. S., Beers, D. R., Wen, S., Bowser, R. and Appel, S. H. (2009) Decreased mRNA

expression of tight junction proteins in lumbar spinal cords of patients with ALS.

Neurology **72,** 1614-1616.

Hirsch, E. C. and Hunot, S. (2009) Neuroinflammation in Parkinson's disease: a target for

neuroprotection? *Lancet Neurol* **8,** 382-397.

Hoffman, H. J. and Olszewski, J. (1961) Spread of sodium fluorescein in normal brain tissue.

A study of the mechanism of the blood-brain barrier. *Neurology* **11,** 1081-1085.

Hoffmann, A., Bredno, J., Wendland, M., Derugin, N., Ohara, P. and Wintermark, M. (2011)

High and Low Molecular Weight Fluorescein Isothiocyanate (FITC)-Dextrans to

Assess Blood-Brain Barrier Disruption: Technical Considerations. *Transl Stroke Res* **2,**

106-111.

Jankowsky, J. L., Fadale, D. J., Anderson, J. et al. (2004) Mutant presenilins specifically elevate the levels of the 42 residue beta-amyloid peptide in vivo: evidence for augmentation of a 42-specific gamma secretase. *Human molecular genetics* **13,** 159-170.

Kaya, M. and Ahishali, B. (2011) Assessment of permeability in barrier type of endothelium in brain using tracers: Evans blue, sodium fluorescein, and horseradish peroxidase. *Methods Mol Biol* **763,** 369-382.

King-VanVlack, C. E., Mewburn, J. D., Chapler, C. K. and MacDonald, P. H. (2003) Hemodynamic and proinflammatory actions of endothelin-1 in guinea pig small intestine submucosal microcirculation. *Am J Physiol Gastrointest Liver Physiol* **284,** G940-948.

Lee, J. D., Kamaruzaman, N. A., Fung, J. N., Taylor, S. M., Turner, B. J., Atkin, J. D., Woodruff, T. M. and Noakes, P. G. (2013) Dysregulation of the complement cascade in the hSOD1G93A transgenic mouse model of amyotrophic lateral sclerosis. *J Neuroinflammation* **10,** 119.

Lee, J. D., Kumar, V., Fung, J. N., Ruitenberg, M. J., Noakes, P. G. and Woodruff, T. M. (2017) Pharmacological inhibition of complement C5a-C5a1 receptor signalling ameliorates disease pathology in the hSOD1(G93A) mouse model of amyotrophic lateral sclerosis. *British journal of pharmacology* **174,** 689-699.

Lepelletier, F. X., Mann, D. M., Robinson, A. C., Pinteaux, E. and Boutin, H. (2017) Early changes in extracellular matrix in Alzheimer's disease. *Neuropathol Appl Neurobiol* **43,** 167-182.

Lindner, V. and Heinle, H. (1982) Binding properties of circulating Evans blue in rabbits as determined by disc electrophoresis. *Atherosclerosis* **43,** 417-422.

Lobsiger, C. S. and Cleveland, D. W. (2007) Glial cells as intrinsic components of non-cell-autonomous neurodegenerative disease. *Nat Neurosci* **10,** 1355-1360.

Luissint, A. C., Artus, C., Glacial, F., Ganeshamoorthy, K. and Couraud, P. O. (2012) Tight

junctions at the blood brain barrier: physiological architecture and disease-associated

dysregulation. *Fluids Barriers CNS* **9,** 23.

Malmgren, L. T. and Olsson, Y. (1980) Differences between the peripheral and the central

nervous system in permeability to sodium fluorescein. *J Comp Neurol* **191,** 103-107.

Mangiarini, L., Sathasivam, K., Seller, M. et al. (1996) Exon 1 of the HD gene with an

expanded CAG repeat is sufficient to cause a progressive neurological phenotype in

transgenic mice. *Cell* **87,** 493-506.

Meister, S., Storck, S. E., Hameister, E., Behl, C., Weggen, S., Clement, A. M. and Pietrzik, C.

U. (2015) Expression of the ALS-causing variant hSOD1(G93A) leads to an impaired

integrity and altered regulation of claudin-5 expression in an in vitro blood-spinal cord

barrier model. *J Cereb Blood Flow Metab* **35,** 1112-1121.

Milne, M. R., Qian, L., Turnbull, M. T., Kinna, G., Collins, B. M., Teasdale, R. D. and Coulson,

E. J. (2019) Downregulation of SNX27 expression does not exacerbate

amyloidogenesis in the APP/PS1 Alzheimer's disease mouse model. *Neurobiol Aging*

77, 144-153.

Naver, B., Stub, C., Møller, M., Fenger, K., Hansen, A. K., Hasholt, L. and Sørensen, S. A.

(2003) Molecular and behavioral analysis of the r6/1 huntington′s disease transgenic

mouse. *Neuroscience* **122,** 1049-1057.

Nonaka, N., Shioda, S. and Banks, W. A. (2005) Effect of lipopolysaccharide on the transport

of pituitary adenylate cyclase activating polypeptide across the blood-brain barrier. *Exp*

Neurol **191,** 137-144.

Okoye, M. I. and Watanabe, I. (1982) Ultrastructural features of cerebral amyloid angiopathy.

Hum Pathol **13,** 1127-1132.

Olesen, L. Ø., Bouzinova, E. V., Severino, M., Sivasaravanaparan, M., Hasselstrøm, J. B.,

Finsen, B. and Wiborg, O. (2016) Behavioural Phenotyping of APPswe/PS1δE9 Mice:

Age-Rrelated Changes and Effect of Long-Term Paroxetine Treatment. *PLOS ONE* **11,**

e0165144.

Oliver, J. M., Berlin, R. D. and Davis, B. H. (1984) Use of horseradish peroxidase and

fluorescent dextrans to study fluid pinocytosis in leukocytes. *Methods in enzymology*

108, 336-347.

Radde, R., Bolmont, T., Kaeser, S. A. et al. (2006) Abeta42-driven cerebral amyloidosis in

transgenic mice reveals early and robust pathology. *EMBO Rep* **7,** 940-946.

Ravenstijn, P. G. M., Merlini, M., Hameetman, M. et al. (2008) The exploration of rotenone as

a toxin for inducing Parkinson's disease in rats, for application in BBB transport and

PK–PD experiments. *Journal of Pharmacological and Toxicological Methods* **57,** 114-

130.

Rosen, D. R., Siddique, T., Patterson, D. et al. (1993) Mutations in Cu/Zn superoxide dismutase

gene are associated with familial amyotrophic lateral sclerosis. *Nature* **362,** 59-62.

Rupp, N. J., Wegenast-Braun, B. M., Radde, R., Calhoun, M. E. and Jucker, M. (2011) Early

onset amyloid lesions lead to severe neuritic abnormalities and local, but not global

neuron loss in APPPS1 transgenic mice. *Neurobiol Aging* **32,** 2324.e2321-2326.

Sarkar, S., Chigurupati, S., Raymick, J. et al. (2014) Neuroprotective effect of the chemical

chaperone, trehalose in a chronic MPTP-induced Parkinson's disease mouse model.

Neurotoxicology **44,** 250-262.

Saunders, N. R., Dziegielewska, K. M., Mollgard, K. and Habgood, M. D. (2015) Markers for

blood-brain barrier integrity: how appropriate is Evans blue in the twenty-first century

and what are the alternatives? *Front Neurosci* **9,** 385.

Serneels, L., Van Biervliet, J., Craessaerts, K. et al. (2009) gamma-Secretase heterogeneity in the Aph1 subunit: relevance for Alzheimer's disease. *Science* **324,** 639-642.

Spijker, S. (2011) Dissection of Rodent Brain Regions. In: *Neuroproteomics*, (K. W. Li ed.)*,* pp. 13-26. Humana Press, Totowa, NJ.

Stokum, J. A., Gerzanich, V. and Simard, J. M. (2016) Molecular pathophysiology of cerebral edema. *J Cereb Blood Flow Metab* **36,** 513-538.

Thomsen, M. S., Routhe, L. J. and Moos, T. (2017) The vascular basement membrane in the healthy and pathological brain. *J Cereb Blood Flow Metab* **37,** 3300-3317.

Wan, H. I., Jacobsen, J. S., Rutkowski, J. L. and Feuerstein, G. Z. (2009) Translational medicine lessons from flurizan's failure in Alzheimer's disease (AD) trial: Implication for future drug discovery and development for AD. *Clin Transl Sci* **2,** 242-247.

Winkler, E. A., Sengillo, J. D., Sagare, A. P. et al. (2014) Blood-spinal cord barrier disruption contributes to early motor-neuron degeneration in ALS-model mice. *Proc Natl Acad Sci U S A* **111,** E1035-1042.

Wolman, M., Klatzo, I., Chui, E., Wilmes, F., Nishimoto, K., Fujiwara, K. and Spatz, M. (1981) Evaluation of the dye-protein tracers in pathophysiology of the blood-brain barrier. *Acta Neuropathol* **54,** 55-61.

Zhao, Z., Nelson, A. R., Betsholtz, C. and Zlokovic, B. V. (2015) Establishment and Dysfunction of the Blood-Brain Barrier. *Cell* **163,** 1064-1078.

Zhong, Z., Deane, R., Ali, Z. et al. (2008) ALS-causing SOD1 mutants generate vascular changes prior to motor neuron degeneration. *Nat Neurosci* **11,** 420-422.

Zhu, S., Wang, J., Zhang, Y., He, J., Kong, J., Wang, J. F. and Li, X. M. (2017) The role of neuroinflammation and amyloid in cognitive impairment in an APP/PS1 transgenic mouse model of Alzheimer's disease. *CNS Neurosci Ther* **23,** 310-320.

1   Zlokovic, B. V. (2011) Neurovascular pathways to neurodegeneration in Alzheimer's disease

2   and other disorders. *Nature reviews. Neuroscience* **12,** 723-738.

Figures

Figure 1. Brain fluorescence uptake profile of three different fluorescent markers. (A)

Fluorescence uptake of sodium fluorescein (Na-Fl), FITC-albumin and FITC-dextran in LPS

treated mice after 1 hour of *i.v.* injection of fluorescent marker through tail vein (*n* = 5 animals).

Fluorescence uptake of Na-Fl **(B)** and FITC-albumin **(C)** in control (blue) and LPS-

administered mice (red) after 0.25, 1 and 24 hours of *i.v.* injection of fluorescent marker through

tail vein (*n* = 5 – 7 animals). Uptake values are expressed as % ratio of fluorescent marker level

in brain to serum. Data are expressed as mean ± SEM. * *P* < 0.05, ** *P* < 0.01, **** *P* <

0.0001, one-way ANOVA.

Figure 2: Progressive increase of Na-Fl in the brain and spinal cord of R6/1 Huntington's

mice throughout disease progression. **A** and **B** show increased Na-Fl uptake in cortex and

striatum of R6/1 mice at 12, 20 and 32 weeks when compared to wild-type (WT) mice ($n = 5$

animals). **C** shows an increased Na-Fl extravasation in cerebellum of R6/1 mice at 32 weeks

when compared to WT mice ($n = 5$ animals). **D** shows increased Na-Fl uptake in the spinal cord

of R6/1 mice at 20 and 32 weeks of age when compared to WT mice ($n = 5$ animals). Data are

presented as mean ± SEM. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$, Student's

t-test for each time point.

Figure 3: Increased uptake of Na-Fl in the brain and spinal cord of hSOD1^{G93A} amyotrophic lateral sclerosis mice throughout disease progression. **A**, **B** and **C** shows increased fluorescence uptake of Na-Fl in cortex, striatum and cerebellum of hSOD1^{G93A} (SOD1) mice at 70, 130 and 175 days when compared to wild-type (WT) mice (*n* = 5 animals). **D** shows increased Na-Fl uptake in the spinal cord of SOD1 mice at 130 and 175 days of age when compared to WT mice (*n* = 5 animals). Data are presented as mean ± SEM. * *P* < 0.05, ** *P* < 0.01, **** *P* < 0.0001, Student's t-test for each time point.

Figure 4: Biphasic brain and spinal cord Na-Fl uptake in the APP/PS1 mouse model of

Alzheimer's disease. Fluorescence was examined in the cortex of APP/PS1 and wild-type (WT)

mice at an early (4 months) and late (18 months) age (n = 5 animals). Na-Fl uptake was

examined in the (**A**) cortex, (**B**) hippocampus, (**C**) striatum, (**D**) cerebellum, and (**E**) spinal cord

(n = 5 animals). Data are presented as mean ± SEM. * P < 0.05, ** P < 0.01, Student's t-test

for each time point. P values < 0.1 are also identified on the graph.

Table 1: Different stages of neurodegeneration in transgenic mouse models used in this study

Disease	Mouse model	Disease Stage				References
		Pre-symptomatic (PS)	Onset (OS)	Mid-symptomatic (MS)	End stage (ES)	
Huntington's disease	R6/1 mice	6 weeks (no signs of cognitive and motor deficits)	12 weeks (first signs of cognitive deficits)	20 weeks (cognitive and motor deficits)	32 weeks (significant reductions in brain and body weights associated with marked neuronal loss in the striatum)	(Naver *et al.* 2003; Mangiarini *et al.* 1996)
Amyotrophic lateral sclerosis	hSOD1^{G93A} mice	30 days (no signs of motor deficit)	70 days (Initial signs of motor deficit grip strength)	125-135 days (Weakness in hind limbs & tremor when suspended by tail)	150-175 days (Full paralysis of lower limbs and loss of righting reflex.)	(Lee *et al.* 2013)
Alzheimer's disease	APP/PS1 mice		4 months (early signs of cognitive deficits; initiation of Aβ deposition)		18 months (reduced locomotion and anxiety-like behaviour, reduced social activity, reduced explorative behaviour and impaired hippocampal tests)	(Radde *et al.* 2006; Rupp *et al.* 2011; Serneels *et al.* 2009; Olesen *et al.* 2016)