A validated quantitative method for the assessment of neuroprotective barrier impairment in neurodegenerative disease models

Vinod Kumar[1*], John D. Lee[1*], Elizabeth J. Coulson[1, 2] and Trent M. Woodruff[1, 2§]

[1]School of Biomedical Sciences, the University of Queensland, Brisbane, QLD, 4072, Australia

[2]Queensland Brain Institute, the University of Queensland, St Lucia, Brisbane, QLD, 4072, Australia

[§]Correspondence to Trent M. Woodruff, School of Biomedical Sciences, Skerman Building, University of Queensland, St. Lucia QLD 4072. AUSTRALIA

*Equal contribution

Email addresses: Vinod Kumar - v.kumar1@uq.edu.au

John D Lee - j.lee9@uq.edu.au

Elizabeth J Coulson - e.coulson@uq.edu.au

Trent M Woodruff[§] - t.woodruff@uq.edu.au

Highlights

-   Developed and validated a quantitative method for assessing BBB and BSCB integrity

using sodium fluorescein

-   Sodium fluorescein has greater sensitivity as an indicator of BBB/BSCB disruption

than FITC-dextran or FITC-albumin

-   Progressive increase in permeability of different CNS regions in Huntington's disease

and amyotrophic lateral sclerosis mice

-   Biphasic disruption in BBB/BSCB permeability in Alzheimer's disease mice

Abstract

The blood-brain barrier (BBB) and blood-spinal cord barrier (BSCB) are highly specialised

structures that limit molecule entry from the blood, and maintain homeostasis within the

central nervous system (CNS). BBB and BSCB breakdown are associated with multiple

neurodegenerative diseases. Given the key role of neuroprotective barrier impairment in

neurodegeneration, it is important to identify an effective quantitative method to assess

barrier integrity in animal models. In the present study, we developed and validated a

quantitative method for assessing BBB and BSCB integrity using sodium fluorescein, that

outperformed other fluorescent dyes. We demonstrated using this method that multiple CNS

regions progressively increase in permeability in models of Huntington's disease and

amyotrophic lateral sclerosis, while biphasic disruption occurred in a mouse model of

Alzheimer's disease with disease progression. Collectively, we report a quantitative

fluorometric marker with validated reproducible experimental methods, that allows the

effective assessment of BBB and BSCB integrity in animal models. This method could be

useful to further the understanding of the contribution of these neuroprotective barriers to

neurodegeneration processes.

Keywords: blood-brain barrier, blood-spinal cord barrier, LPS, neurodegeneration, Na-Fl

Introduction

The blood-brain barrier (BBB) and blood-spinal cord barrier (BSCB) are highly specialised

epithelial structures linked through intercellular junctional complexes, termed tight junctions.

These structures maintain homeostasis within the central nervous system (CNS) by ensuring

strict regulation of molecule transport between circulating blood and CNS tissues (Zhao et al.,

2015). The primary physiological function of these barriers is to maintain the chemical

composition of the CNS, and protect against toxic molecules, to support neuronal cell health.

BBB and BSCB breakdown, due to disruption of tight junctions or epithelial cell integrity,

can lead to altered transport of molecules between the CNS and blood, which can contribute

to the pathological processes resulting in progressive structural and functional loss of neurons

in multiple neurodegenerative diseases (Ellrichmann et al., 2013; Harry and Kraft, 2008;

Hirsch and Hunot, 2009; Lobsiger and Cleveland, 2007).

There is evidence to support the notion that neuroprotective barrier impairment occurs within

the progression of these neurodegenerative diseases, including in Alzheimer's disease (AD)

(Zlokovic, 2011), Parkinson's disease (PD) (Brochard et al., 2009; Carvey et al., 2005; Chen

et al., 2008; Sarkar et al., 2014), Huntington's disease (HD) (Drouin-Ouellet et al., 2015) and

amyotrophic lateral sclerosis (ALS) (Garbuzova-Davis et al., 2007b). Damaged BBB and

BSCB integrity and resulting increased permeability can lead to oedema, neurotoxic molecule

accumulation, as well as enhancing the infiltration of immune cells to the CNS, which can

accelerate the process of neurodegeneration (Arima et al., 2013; Bell et al., 2010; Stokum et

al., 2016). This interaction between neuroinflammation, neurodegeneration and

neuroprotective barrier integrity can also influence drug bioavailability to the CNS, which

may affect the activity of neuroprotective therapies. For example, Tarenflurbil used in AD

was shown to have poor BBB penetration, which may have led to its failure in clinical trials

(Wan et al., 2009). Given the key involvement of neuroprotective barrier impairment in

neurological and neurodegenerative conditions, it is therefore important to identify an

effective quantitative method to assess barrier impairment and permeability changes during

disease progression in animal models of these conditions.

Previous studies have utilized different dyes to measure the permeability of BBB and BSCB,

with Evans blue dye being the most commonly reported (Saunders et al., 2015). However,

numerous studies have indicated that Evans blue is a substandard marker for BBB

permeability, due to significant binding to tissues (Dallal and Chang, 1994; Lindner and

Heinle, 1982), solution instability, *in vivo* toxicity (demonstrated in rats, monkeys, dogs, cats

and rabbits), and lack of a validated quantitative determination method. Furthermore, Evans

blue relies on binding to the large plasma protein albumin, and subsequent transport across

the BBB, which only occurs during later stages of barrier disruption. In contrast to Evans blue,

more recent fluorescent-based dyes have been used extensively as a marker for permeability

studies including sodium fluorescein (Na-Fl) (Hoffman and Olszewski, 1961; Malmgren and

Olsson, 1980), fluorescein isothiocyanate (FITC) labelled albumin (Boutoille et al., 2009;

King-VanVlack et al., 2003), and FITC labelled dextran-70 (Hoffmann et al., 2011; Oliver et

al., 1984). However, no study to date has directly compared the relative utility of these dyes

in assessing BBB and BSCB impairment, and provided a robust validated quantitative method.

In the present study, we therefore initially examined these three different dyes (Na-Fl, FITC

labelled albumin and FITC labelled dextrans-70) as potential markers for neuroprotective

permeability in the brains of lipopolysaccharide (LPS)-injected mice to determine the most

suitable and sensitive marker for BBB and BSCB permeability studies. We developed a

quantitative method that demonstrated Na-Fl as the most sensitive marker for measuring

neuroprotective barrier integrity. We then utilized this method in mouse models of AD, HD,

and ALS, where we identified perturbed barrier integrity in different regions of the brain and

spinal cord of these animals. Our data therefore reveal a novel quantitative method for

monitoring BBB and BSCB integrity in animal models of neurodegeneration.

Material and methods

Animals

All animal procedures were approved by the University of Queensland Animal Ethics

Committee and conducted in accordance with the Australian Code of Practice for the Care

and Use of Animals for Scientific Purposes (8th edition, 2013). ARRIVE guidelines have

been followed in the preparation of the manuscript. Animals were group housed 3 ~ 4 per

cage on arrival in the University of Queensland animal facility. Mice were housed in a 12-h

light: dark phase (on at 06:00 h and off at 18:00 h) with room temperature maintained at 24 ±

2 °C. Animals had free access to *ad-libitum* food and water. Animal sources and details are

outlined below.

Wild-type mice: C57BL/6J male mice aged 10-12 weeks old were obtained from the Animal

Resources Centre in Canning Vale, Australia.

Transgenic R6/1 mice (transgenic mouse model for Huntington's disease):

Transgenic R6/1 mice (B6-CBA-Tg (HDexon1) 61Gpb/1J) expressing a transgene containing

the human HD gene were obtained from Jackson Laboratory (Stock No: 002809; Bar Harbour,

ME, USA; (Mangiarini et al., 1996)) and bred with female mice on CBA/BL6 background to

produce R6/1 transgenic and non-transgenic (wild-type; WT) mice. Male transgenic R6/1 and

matched WT littermate mice were used at various stages of disease progression (6 weeks, 12

weeks, 20 weeks and 32 weeks).

Transgenic hSOD1^{G93A} mice (transgenic mouse model for amyotrophic lateral sclerosis):

Transgenic hSOD1^{G93A} mice (B6-Cg-Tg (SOD1-G93A) 1Gur/J) carrying a mutant form of

the human *SOD1* gene were obtained from Jackson Laboratory (Stock No: 004435; Bar

Harbor, ME, USA; (Rosen et al., 1993)) and bred on C57BL/6J background to produce

transgenic hSOD1^{G93A} and WT littermates as control mice. Male hSOD1^{G93A} and WT mice

were used at various stages of disease progression (30 days, 70 days, 130 days and 150 – 175

days) as described previously (Lee et al., 2013).

Transgenic APP/PS1 mice (transgenic mouse model for Alzheimer's disease):

Double transgenic APP/PS1 mice (B6-Cg-Tg (APPswe, PSEN1de9)85Dbo/Mmjax)

expressing a chimeric mouse/human amyloid precursor protein and a mutant human

presenilin 1 were source from Jackson Laboratory (MMRRC Stock No: 34832 – JAX; Bar

Harbor, ME, USA; (Jankowsky et al., 2004)) and bred on C57BL/6J background to produce

transgenic APP/PS1 and WT littermates. Male APP/PS1 and WT mice were used at 4 and 18

months, ages correlating to early, and late stage disease phenotypes.

Experimental conditions for quantification and selection of fluorescent marker

permeability studies

Fluorescence stability of markers during the extraction and analysis process were first tested

by studying effect of different solvents as a loading solution, the effect of solvents during the

extraction process, extraction efficiency of the process, the influence of matrix/solvents on

the fluorescence of markers, and loss of fluorescence during the extraction and analysis steps.

From these preliminary steps, TRIS buffer (pH 7.4) was identified as the most suitable

solvent.

The selection of an appropriate fluorescent marker was accomplished by comparing uptake

results of the markers in the brain and serum of mice after an intraperitoneal LPS injection (4

mg kg^{-1} of mice body weight). After 24 hours, mice were then anaesthetized with

intraperitoneally injected zolazapam (50 mg kg^{-1}) and xylazine (12 mg kg^{-1}) followed by 100

µl intravenous injection of the fluorescent markers (100 mg mL^{-1}). After one hour, blood

samples were collected via cardiac puncture and allowed to clot at room temperature for 15

minutes. Samples were then centrifuged at 2,000 x g for 10 minutes at 4˚C for serum

collection. After blood collection, mice were immediately perfused transcardially with 50 mL

of 1 x PBS to remove any remaining fluorescent marker in the circulation. Whole brain was

dissected and homogenized in an equal volume of TRIS buffer (pH 7.4). Serum samples were

deproteinized with acetonitrile (1:3), vortexed, sonicated and centrifuged at 13,000 x g for 15

minutes at room temperature. Supernatant samples were evaporated to a dry state using a

centrivap concentrator (Labconco, U.S.A.) at room temperature and reconstituted in an equal

volume of TRIS buffer (pH 7.4). Brain tissue samples were homogenized with milli Q water

(1:2 ratio to the tissue weight). Homogenized brain samples were deproteinized with

acetonitrile (1:6 ratio to the tissue weight) and processed via a similar method used for serum

samples. Dried brain samples were reconstituted in equal weight-volume of TRIS buffer (pH

7.4). 10 µl of reconstituted solutions were diluted with 200 µl of TRIS buffer (pH 7.4) in 96-

well black plate (Corning Inc, U.S.A.) along with standard curve samples in same matrix. The

fluorescence levels were measured using a microplate reader (FlexStation 3, U.S.A.) with λ_{ex}

460 nm and λ_{em} 515 nm for Na-Fl, λ_{ex} 495 nm and λ_{em} 520 nm for FITC-albumin and λ_{ex} 490

nm and λ_{em} 530 nm for FITC-dextran respectively. The amount of fluorescent marker was

calculated from fluorescence levels via a standard curve prepared in a similar manner using

brain and serum obtained from mice without any treatment. Fluorescence uptake was

calculated by comparing ratio of fluorescent marker level in the brain with the level of

fluorescent marker in the serum. Fluorescence values were corrected with the fluorescence of

blank brain and serum samples obtained from mice without any fluorescent marker or LPS

treatment and treated with the same method as above.

Application of a validated quantitative fluorometric method in mouse models of

neurodegeneration

The applicability of the developed and validated quantitative fluorometric method was

assessed by performing neuroprotective barrier integrity studies in various chronic mouse

models of neurodegeneration. Mice ($n = 5$) at various disease stages were anaesthetized with

an intraperitoneal injection of zolazapam (50 mg kg^{-1}) and xylazine (12 mg kg^{-1}).

Anaesthetized mice were then intravenously administered with 100 µl of Na-Fl (100 mg mL^{-1})

via tail vein. After 15 minutes from injection, blood samples were collected by cardiac

puncture for serum extraction. Animals were transcardially perfused immediately with 1 x

PBS to remove circulating Na-Fl followed by whole brain and spinal cord tissue collection.

Different brain regions were isolated on ice using a mouse coronal brain matrix (ProSciTech)

under the microscope. Samples were stored at -80°C for further processing and analysis using

the developed quantitative fluorometric method as described above. Fluorescence levels in

brain and spinal cord samples were measured as describe above. Fluorescence values were

back calculated for the level of Na-Fl in the serum and tissues using the standard curve

samples prepared in the same matrix and analyzed as above. Results are presented as relative

levels of fluorescence uptake in R6/1, hSOD1^{G93A} and APP/PS1 mice compared with

expression in their respective litter-matched non-transgenic WT controls.

Data analysis

All results are reported as mean ± standard error of mean and statistical analysis performed using GraphPad Prism 8.3.1 (LaJolla, CA, USA). The statistical difference between FITC labelled albumin, FITC labelled dextran-70 and Na-Fl for percentage uptake in the brain were analyzed using a one-way ANOVA and a *post-hoc* Tukey's test. The statistical difference for levels of fluorescence uptake between WT and R6/1, WT and hSOD1^{G93A} and WT and APP/PS1 mice were analyzed using a two-tailed Student's t-test for each stage of disease progression. The differences were considered significant when $P < 0.05$.

Results

Sodium fluorescein has greater sensitivity as an indicator of BBB disruption than FITC-dextran or FITC-albumin

Selection of an appropriate fluorescent marker for permeability studies was based upon the percentage uptake of the fluorescent marker, which was quantitatively determined via a validated fluorometric method. We utilized peripheral administration of LPS to induce BBB disruption after 24 hours as previously documented (Banks et al., 2015; Nonaka et al., 2005). We then determined the uptake of fluorescent dye into the brain 1 hour after intravenous dye injection. The percentage uptake of Na-Fl (~50%) was found to be the highest compared to either FITC-dextran-70 (~1%) or FITC-albumin (~17%) (**Figure 1A**). We next extended this analysis by comparing BBB permeability changes using FITC-albumin and Na-Fl (both having detectible uptake in the brain after 1 hour), over 24 hours. Fluorescence uptake of FITC-albumin in the brain was significantly increased at 0.25 hour (1.4% vs. 4.8%), 1 hour (6.8% vs. 16.7%) and 24 hours (8.4% vs. 24%) in saline-injected versus LPS-injected mice respectively (**Figure 1B**). Fluorescence uptake of Na-Fl in the brain was also significantly increased at 0.25 hour (8.5% vs. 40%), 1 hour (10.4% vs. 67.7%) and 24 hours (3.8% vs.

17.2%) in saline-injected versus LPS-injected mice respectively (**Figure 1C**). Notably, the

fold-increases in dye uptake between saline and LPS groups, were significantly larger for Na-

Fl than FITC-albumin, peaking 1 hour after dye injection. We therefore identified 1-hour Na-

Fl to be an ideal method to progress for future validation.

Progressive breakdown of neuroprotective barriers in the R6/1 transgenic mouse model

of Huntington's disease

We first applied our method to investigate BBB and BSCB integrity at different disease

stages in the R6/1 transgenic mouse model of HD. We observed no change in Na-Fl uptake in

cortex, striatum, cerebellum and spinal cord between WT and R6/1 transgenic mice at 6

weeks of age, which is considered to be a pre-symptomatic stage of disease. Na-Fl uptake in

the cortex and striatum progressively increased in R6/1 mice at 12 (1.7-fold, 1.8-fold), 20

(2.1-fold, 2.4-fold) and 32 (2.3-fold, 2.9-fold) weeks when compared to WT mice (**Figure 2A**

and **2B**). The cerebellum was less affected, with Na-Fl uptake only increased by 2.0-fold in

R6/1 mice at 32 weeks of age (**Figure 2C**). Interestingly, Na-Fl was also increased in the

spinal cord of R6/1 mice at 20 and 32 weeks by 2.7-fold respectively (**Figure 2D**).

Breakdown of neuroprotective barriers in the hSOD1^{G93A} transgenic mouse model of

amyotrophic lateral sclerosis

Next, we assessed neuroprotective barrier integrity in hSOD1^{G93A} transgenic mice. Na-Fl

uptake was measured in cortex, striatum, cerebellum and spinal cord at progressive ages of

disease progression. Similar to R6/1 mice, we observed no changes in Na-Fl uptake in cortex,

striatum, cerebellum and spinal cord between WT and hSOD1^{G93A} transgenic mice at the pre-

symptomatic age of 30 days. However, Na-Fl uptake progressively increased in the cortex of

hSOD1^{G93A} mice at 70 (1.2-fold), 130 (1.5-fold) and 175 (1.6-fold) days of age (**Figure 3A**).

Na-Fl also increased in the striatum of hSOD1^{G93A} mice at these same ages (**Figure 3B**). BBB permeability increases were also observed in the cerebellum of hSOD1^{G93A} mice, with significant Na-Fl uptake observed at 70, 130 and 175 days (**Figure 3C**). Finally, BSCB permeability was also detected, with Na-Fl uptake in the spinal cord gradually increased in hSOD1^{G93A} mice by 1.1-, 1.3- and 2.0-fold at 70, 130 and 175 days respectively (**Figure. 3D**).

Neuroprotective barriers in the APP/PS1 transgenic mouse model of Alzheimer's disease display biphasic alterations at different diseases stages

To measure possible alterations in neuroprotective barriers of APP/PS1 transgenic mice, Na-Fl uptake was measured in cortex, striatum, hippocampus, cerebellum and spinal cord at an early (4 months) and late (18 months) stage of disease. Brain and spinal cord tissues from APP/PS1 transgenic mice at the early stages of disease showed an overall increase in barrier permeability (**Figure 4**), with significant increases in Na-Fl uptake observed in the striatum (**Figure 4C**) and spinal cord (**Figure 4E**). Notably, by 18 months of age, these same tissues had reduced levels of Na-Fl, with decreases of up to 50% observed in brain regions and spinal cord (**Figure 4B**).

Discussion

BBB and BSCB breakdown are associated with a number of neurodegenerative diseases, including AD, PD, HD and ALS (Desai et al., 2007; Di Pardo et al., 2017; Garbuzova-Davis et al., 2007a; Garbuzova-Davis and Sanberg, 2014; Garbuzova-Davis et al., 2007b; Gray and Woulfe, 2015). The understanding of the underlying mechanisms that initiates the opening or closure of the BBB and BSCB is therefore of therapeutic significance in many neurodegenerative diseases, and is commonly investigated by researchers. However, methods to investigate BBB and BSCB permeability *in-vivo* are often associated with technical

difficulties that depend on subjective quantification of fluorescence images, or semi-

quantitative evaluation of endogenous molecules. To address this, prior studies have utilized

an abundance of fluorescent markers for *in-vivo* and *in-vitro* permeability studies, each with

their own strength and weaknesses (Saunders et al., 2015). Methods used for the detection of

these fluorescent markers also display great variation in their quantitation methodology

between studies. Selection of an ideal validated method for assessing neuroprotective barrier

permeability is therefore difficult due to lack of prior published results for quantitative

determination. The present research therefore instigated the selection and identification of a

suitable fluorescent marker for CNS permeability studies using a validated fluorometric

method of quantitative analysis. The method was developed using an acute LPS-induced BBB

breakdown model and then its application verified by assessing neuroprotective barrier

permeability in various mouse model of neurodegeneration.

The BBB and BSCB are highly specialized epithelial structures that are important in

maintaining the CNS homeostasis and preventing the entry of foreign toxic substances. It

undergoes degeneration during disease processes of neurodegenerative diseases and has a re-

emerging interest as a mechanistic and therapeutic target for intervention. In spite of their

limitations, use of dyes to detect barrier breakdown have remained widespread, where

numerous studies have utilized different dyes including, Evans blue, fluorescein, horse radish

peroxidase, dextran, albumin and trypan blue (Saunders et al., 2015). To the best of our

knowledge, our study is the first to compare three widely used fluorescent dye markers

(FITC-dextran, FITC-albumin and Na-Fl) to determine the most suitable marker to

quantitatively evaluate BBB integrity. We demonstrate that 1-hour after the injection of these

dyes to LPS-induced BBB-injured mice, Na-Fl showed the greatest percentage uptake in the

brain. Although this has not been directly compared previously, this is consistent with

previous studies suggesting that Na-Fl only binds weakly to proteins, and is therefore considered to be an effective low molecular weight marker for BBB studies in contrast to dyes that bind to proteins (Wolman et al., 1981). Although Na-Fl cannot be detected by visual examination, a quantitative fluorometric method utilized in the current study may enable detection of subtle alterations in BBB/BSCB permeability and be the earliest and the most sensitive indicator of BBB/BSCB disruption (Kaya and Ahishali, 2011). Our studies also demonstrated that Na-Fl uptake in the brain was reduced after 24 hours of injection (when compared to 15 minutes and 1 hour), suggesting rapid clearance of Na-Fl in the brain parenchyma. Collectively, as small molecule, Na-Fl is considered to be less toxic than other dye tracers, is freely diffusible, minimally binds to proteins and can be evaluated quantitatively and reproducibly. We therefore posit this method provides an ideal marker of neuroprotective barrier integrity.

In pathological conditions, BBB permeability is increased due to various biochemical mediators (Abbott, 2000). Increased permeability favours the transport of peripheral immune cells, inflammatory products and other circulating substances to the brain and spinal cord, resulting in breakdown of tight junction molecules and ultimately, disruption of BBB and BSCB (Abbott et al., 2010; Ballabh et al., 2004; de Vries et al., 1997). BBB and BSCB disruption and malfunction are involved in various neurological disorders including AD, PD, HD and ALS (Desai et al., 2007; Di Pardo et al., 2017; Garbuzova-Davis et al., 2007a; Garbuzova-Davis and Sanberg, 2014; Garbuzova-Davis et al., 2007b; Gray and Woulfe, 2015). Using different mouse models of HD, ALS and AD, we investigated barrier integrity in different regions of the brain and spinal cord using our quantitative method of Na-Fl extravasation. We demonstrated that there is an early disruption of BBB and BSCB permeability in R6/1 mouse model of HD as we showed increased Na-Fl uptake in cortex,

striatum and spinal cord, in parallel with the cognitive and motor deficits present in this

model (Ayton et al., 2019; Brooks et al., 2012). This early disruption of BBB is further

supported by previous studies showing early vascular impairment and BBB dysfunction in

R6/2 mice, with a reduction in expression of tight junction proteins claudin-5 and occludin,

leading to increased paracellular permeability (Di Pardo et al., 2017; Drouin-Ouellet et al.,

2015; Luissint et al., 2012). This validates the applicability of this fluorometric method to

measure subtle changes in BBB/BSCB integrity.

In the present study, we also demonstrated that there is an early disruption of BBB and BSCB

permeability in hSOD1^{G93A} mouse model of ALS based on increased Na-Fl uptake in cortex,

striatum, cerebellum and spinal cord. Similar to HD, these findings support previous studies

demonstrating damage to BSCB integrity in hSOD1^{G93A} mice via ultrastructural capillary

alterations and increased leakage in the spinal cord at both early and late stages of disease

(Garbuzova-Davis et al., 2007a; Garbuzova-Davis et al., 2007b; Lee et al., 2017; Zhong et al.,

2008). These findings are further supported by numerous studies demonstrating decreased

expression of tight junction proteins integral to the integrity of the BSCB, such as zona

occludens-1, occludin and claudin-5 in the spinal cord of hSOD1 transgenic mice and ALS

patients (Henkel et al., 2009; Meister et al., 2015; Winkler et al., 2014; Zhong et al., 2008).

Collectively, our study complements these previous findings as our validated fluorometric

methods was able to detect early barrier impairments very early in disease, at ages that

precede significant motor neuron death (Lee et al., 2013), which may have pathophysiological

relevance.

We also demonstrated an apparent biphasic BBB/BSCB disruption in APP/PS1 mouse model

of AD. Unlike HD and MND, neuroprotective barrier disruption in APP/PS1 mouse model of

AD was only evident in early in the disease, at an age where there is minimal cognitive

dysfunction (Zhu et al., 2017). Interestingly, unlike the other disease models studied, there

was a notable decrease in Na-Fl uptake in brain and spinal cord regions at the late stage of

disease, an age where is significant cognitive decline (Gengler et al., 2010; Milne et al., 2019).

Although it is possible that the result is due to a faster clearance of the dye from a highly

permeable BBB, the decrease in dye uptake in APP/PS1 mice late in disease is consistent

with previous studies demonstrating thickening of the basement membrane in AD model mice

and humans (Hawkes et al., 2013; Lepelletier et al., 2017). Although the causes of the

basement membrane thickening in AD have not been fully elucidated to date, decreased

clearance of amyloid beta or increased astrocyte production of basement membrane proteins

like perlecan and fibronectin have been suggested to contribute to this thickening of basement

membrane (Deane and Zlokovic, 2007; Okoye and Watanabe, 1982; Thomsen et al., 2017).

Importantly, the progressive molecular changes in barrier permeability of multiple

neurodegenerative disease models tested in previous studies mirrored the changes in Na-Fl

levels in the brain and spinal cord of our current study. This strongly validates the sensitivity

and accuracy of our validated fluorometric method to measure subtle changes in BBB/BSCB

integrity. This may have implications for future preclinical studies that aim to correlate the

extent of barrier impairment and permeability changes with disease progression.

Conclusions

In summary, we have identified a validated, effective quantitative method for assessing BBB

and BSCB integrity using Na-Fl, that outperforms other fluorescent dyes. We demonstrated

using this method that multiple CNS regions progressively increase in permeability in the

R6/1 mouse model of HD and the hSOD1^{G93A} mouse model of ALS. By contrast, whilst

increased BBB/BSCB permeability was observed in young APP/PS1 mouse model of AD,

later stage diseased mice showed marked reduction in CNS permeability. Collectively, we

report a quantitative fluorometric marker with validated reproducible experimental methods,

that allows the effective assessment of BBB and BSCB integrity in animal models. This

method could be useful to further the understanding of the contribution of these

neuroprotective barriers to neurodegeneration processes, and to assess the efficacy of

neuroprotective agents.

Declarations

Ethics approval and consent to participate

All experimental procedures were approved by the University of Queensland Animal Ethics

Committee and complied with the policies and regulations regarding animal experimentation.

They were conducted in accordance with the Queensland Government Animal Research Act

2001, associated Animal Care and Protection Regulations (2002 and 2008), and the

Australian Code of Practice for the Care and Use of Animals for Scientific Purposes, 8th

Edition (National Health and Medical Research Council, 2013). ARRIVE guidelines have

been followed in the preparation of the manuscript.

Consent for publication

Not applicable

Availability of data and material

Not applicable

Competing interests

The authors declare that they have no competing interests

Funding

TMW is supported by a NHMRC Career Development Fellowship. The research was funded by grants from the MNDRIA to TMW (GIA1865) and JDL (GIA1830), and a National Health and Medical Research Council (NHMRC) Project grant (APP1082271) to TMW.

Author's contributions

VK, JDL, and TMW conceived the project. VK, JDL, EJC and TMW designed the study. VK and JDL performed experiments. All authors contributed to the analyses and/or interpreted the data. VK and JDL wrote the paper with TMW, with editing contributions from EJC. All authors read and approved the final manuscript.

Acknowledgements

The authors would like to sincerely thank the University of Queensland Biological Resources for the animal care and husbandry. We also thank Maryam Shayegh for her technical support with genotyping mice.

References

Abbott, N.J., 2000. Inflammatory mediators and modulation of blood-brain barrier permeability. Cell Mol Neurobiol 20(2), 131-147.

Abbott, N.J., Patabendige, A.A., Dolman, D.E., Yusof, S.R., Begley, D.J., 2010. Structure and function of the blood-brain barrier. Neurobiol Dis 37(1), 13-25.

Arima, Y., Kamimura, D., Sabharwal, L., Yamada, M., Bando, H., Ogura, H., Atsumi, T., Murakami, M., 2013. Regulation of immune cell infiltration into the CNS by regional neural inputs explained by the gate theory. Mediators Inflamm 2013, 898165.

Ayton, S., Lei, P., Appukuttan, A.T., Renoir, T., Foliaki, S., Chen, F., Adlard, P.A., Hannan,

A.J., Bush, A.I., 2019. Brain Zinc Deficiency Exacerbates Cognitive Decline in the R6/1

Model of Huntington's Disease. Neurotherapeutics.

Ballabh, P., Braun, A., Nedergaard, M., 2004. The blood-brain barrier: an overview:

structure, regulation, and clinical implications. Neurobiol Dis 16(1), 1-13.

Banks, W.A., Gray, A.M., Erickson, M.A., Salameh, T.S., Damodarasamy, M., Sheibani, N.,

Meabon, J.S., Wing, E.E., Morofuji, Y., Cook, D.G., Reed, M.J., 2015. Lipopolysaccharide-

induced blood-brain barrier disruption: roles of cyclooxygenase, oxidative stress,

neuroinflammation, and elements of the neurovascular unit. J Neuroinflammation 12, 223.

Bell, R.D., Winkler, E.A., Sagare, A.P., Singh, I., LaRue, B., Deane, R., Zlokovic, B.V.,

2010. Pericytes control key neurovascular functions and neuronal phenotype in the adult brain

and during brain aging. Neuron 68(3), 409-427.

Boutoille, D., Marechal, X., Pichenot, M., Chemani, C., Guery, B., Faure, K., 2009. FITC-

albumin as a marker for assessment of endothelial permeability in mice: comparison with

125I-albumin. Experimental lung research 35(4), 263-271.

Brochard, V., Combadiere, B., Prigent, A., Laouar, Y., Perrin, A., Beray-Berthat, V.,

Bonduelle, O., Alvarez-Fischer, D., Callebert, J., Launay, J.M., Duyckaerts, C., Flavell, R.A.,

Hirsch, E.C., Hunot, S., 2009. Infiltration of CD4+ lymphocytes into the brain contributes to

neurodegeneration in a mouse model of Parkinson disease. J Clin Invest 119(1), 182-192.

Brooks, S.P., Janghra, N., Workman, V.L., Bayram-Weston, Z., Jones, L., Dunnett, S.B.,

2012. Longitudinal analysis of the behavioural phenotype in R6/1 (C57BL/6J) Huntington's

disease transgenic mice. Brain Res Bull 88(2-3), 94-103.

Carvey, P.M., Zhao, C.H., Hendey, B., Lum, H., Trachtenberg, J., Desai, B.S., Snyder, J.,

Zhu, Y.G., Ling, Z.D., 2005. 6-Hydroxydopamine-induced alterations in blood-brain barrier

permeability. Eur J Neurosci 22(5), 1158-1168.

Chen, X., Lan, X., Roche, I., Liu, R., Geiger, J.D., 2008. Caffeine protects against MPTP-induced blood-brain barrier dysfunction in mouse striatum. J Neurochem 107(4), 1147-1157.

Dallal, M.M., Chang, S.W., 1994. Evans blue dye in the assessment of permeability-surface area product in perfused rat lungs. J Appl Physiol (1985) 77(2), 1030-1035.

de Vries, H.E., Kuiper, J., de Boer, A.G., Van Berkel, T.J., Breimer, D.D., 1997. The blood-brain barrier in neuroinflammatory diseases. Pharmacol Rev 49(2), 143-155.

Deane, R., Zlokovic, B.V., 2007. Role of the blood-brain barrier in the pathogenesis of Alzheimer's disease. Curr Alzheimer Res 4(2), 191-197.

Desai, B.S., Monahan, A.J., Carvey, P.M., Hendey, B., 2007. Blood-brain barrier pathology in Alzheimer's and Parkinson's disease: implications for drug therapy. Cell Transplant 16(3), 285-299.

Di Pardo, A., Amico, E., Scalabri, F., Pepe, G., Castaldo, S., Elifani, F., Capocci, L., De Sanctis, C., Comerci, L., Pompeo, F., D'Esposito, M., Filosa, S., Crispi, S., Maglione, V., 2017. Impairment of blood-brain barrier is an early event in R6/2 mouse model of Huntington Disease. Sci Rep 7, 41316.

Drouin-Ouellet, J., Sawiak, S.J., Cisbani, G., Lagace, M., Kuan, W.L., Saint-Pierre, M., Dury, R.J., Alata, W., St-Amour, I., Mason, S.L., Calon, F., Lacroix, S., Gowland, P.A., Francis, S.T., Barker, R.A., Cicchetti, F., 2015. Cerebrovascular and blood-brain barrier impairments in Huntington's disease: Potential implications for its pathophysiology. Annals of neurology 78(2), 160-177.

Ellrichmann, G., Reick, C., Saft, C., Linker, R.A., 2013. The role of the immune system in Huntington's disease. Clin Dev Immunol 2013, 541259.

Garbuzova-Davis, S., Haller, E., Saporta, S., Kolomey, I., Nicosia, S.V., Sanberg, P.R., 2007a. Ultrastructure of blood-brain barrier and blood-spinal cord barrier in SOD1 mice modeling ALS. Brain research 1157, 126-137.

Garbuzova-Davis, S., Sanberg, P.R., 2014. Blood-CNS Barrier Impairment in ALS patients

versus an animal model. Front Cell Neurosci 8, 21.

Garbuzova-Davis, S., Saporta, S., Haller, E., Kolomey, I., Bennett, S.P., Potter, H., Sanberg,

P.R., 2007b. Evidence of compromised blood-spinal cord barrier in early and late

symptomatic SOD1 mice modeling ALS. PLoS One 2(11), e1205.

Gengler, S., Hamilton, A., Holscher, C., 2010. Synaptic plasticity in the hippocampus of a

APP/PS1 mouse model of Alzheimer's disease is impaired in old but not young mice. PLoS

One 5(3), e9764.

Gray, M.T., Woulfe, J.M., 2015. Striatal blood-brain barrier permeability in Parkinson's

disease. J Cereb Blood Flow Metab 35(5), 747-750.

Harry, G.J., Kraft, A.D., 2008. Neuroinflammation and microglia: considerations and

approaches for neurotoxicity assessment. Expert Opin Drug Metab Toxicol 4(10), 1265-1277.

Hawkes, C.A., Gatherer, M., Sharp, M.M., Dorr, A., Yuen, H.M., Kalaria, R., Weller, R.O.,

Carare, R.O., 2013. Regional differences in the morphological and functional effects of aging

on cerebral basement membranes and perivascular drainage of amyloid-beta from the mouse

brain. Aging Cell 12(2), 224-236.

Henkel, J.S., Beers, D.R., Wen, S., Bowser, R., Appel, S.H., 2009. Decreased mRNA

expression of tight junction proteins in lumbar spinal cords of patients with ALS. Neurology

72(18), 1614-1616.

Hirsch, E.C., Hunot, S., 2009. Neuroinflammation in Parkinson's disease: a target for

neuroprotection? Lancet Neurol 8(4), 382-397.

Hoffman, H.J., Olszewski, J., 1961. Spread of sodium fluorescein in normal brain tissue. A

study of the mechanism of the blood-brain barrier. Neurology 11, 1081-1085.

Hoffmann, A., Bredno, J., Wendland, M., Derugin, N., Ohara, P., Wintermark, M., 2011. High and Low Molecular Weight Fluorescein Isothiocyanate (FITC)-Dextrans to Assess Blood-Brain Barrier Disruption: Technical Considerations. Transl Stroke Res 2(1), 106-111.

Jankowsky, J.L., Fadale, D.J., Anderson, J., Xu, G.M., Gonzales, V., Jenkins, N.A., Copeland, N.G., Lee, M.K., Younkin, L.H., Wagner, S.L., Younkin, S.G., Borchelt, D.R., 2004. Mutant presenilins specifically elevate the levels of the 42 residue beta-amyloid peptide in vivo: evidence for augmentation of a 42-specific gamma secretase. Human molecular genetics 13(2), 159-170.

Kaya, M., Ahishali, B., 2011. Assessment of permeability in barrier type of endothelium in brain using tracers: Evans blue, sodium fluorescein, and horseradish peroxidase. Methods Mol Biol 763, 369-382.

King-VanVlack, C.E., Mewburn, J.D., Chapler, C.K., MacDonald, P.H., 2003. Hemodynamic and proinflammatory actions of endothelin-1 in guinea pig small intestine submucosal microcirculation. Am J Physiol Gastrointest Liver Physiol 284(6), G940-948.

Lee, J.D., Kamaruzaman, N.A., Fung, J.N., Taylor, S.M., Turner, B.J., Atkin, J.D., Woodruff, T.M., Noakes, P.G., 2013. Dysregulation of the complement cascade in the hSOD1G93A transgenic mouse model of amyotrophic lateral sclerosis. J Neuroinflammation 10(1), 119.

Lee, J.D., Kumar, V., Fung, J.N., Ruitenberg, M.J., Noakes, P.G., Woodruff, T.M., 2017. Pharmacological inhibition of complement C5a-C5a1 receptor signalling ameliorates disease pathology in the hSOD1(G93A) mouse model of amyotrophic lateral sclerosis. British journal of pharmacology 174(8), 689-699.

Lepelletier, F.X., Mann, D.M., Robinson, A.C., Pinteaux, E., Boutin, H., 2017. Early changes in extracellular matrix in Alzheimer's disease. Neuropathol Appl Neurobiol 43(2), 167-182.

Lindner, V., Heinle, H., 1982. Binding properties of circulating Evans blue in rabbits as determined by disc electrophoresis. Atherosclerosis 43(2-3), 417-422.

Lobsiger, C.S., Cleveland, D.W., 2007. Glial cells as intrinsic components of non-cell-autonomous neurodegenerative disease. Nat Neurosci 10(11), 1355-1360.

Luissint, A.C., Artus, C., Glacial, F., Ganeshamoorthy, K., Couraud, P.O., 2012. Tight junctions at the blood brain barrier: physiological architecture and disease-associated dysregulation. Fluids and barriers of the CNS 9(1), 23.

Malmgren, L.T., Olsson, Y., 1980. Differences between the peripheral and the central nervous system in permeability to sodium fluorescein. J Comp Neurol 191(1), 103-107.

Mangiarini, L., Sathasivam, K., Seller, M., Cozens, B., Harper, A., Hetherington, C., Lawton, M., Trottier, Y., Lehrach, H., Davies, S.W., Bates, G.P., 1996. Exon 1 of the HD gene with an expanded CAG repeat is sufficient to cause a progressive neurological phenotype in transgenic mice. Cell 87(3), 493-506.

Meister, S., Storck, S.E., Hameister, E., Behl, C., Weggen, S., Clement, A.M., Pietrzik, C.U., 2015. Expression of the ALS-causing variant hSOD1(G93A) leads to an impaired integrity and altered regulation of claudin-5 expression in an in vitro blood-spinal cord barrier model. J Cereb Blood Flow Metab 35(7), 1112-1121.

Milne, M.R., Qian, L., Turnbull, M.T., Kinna, G., Collins, B.M., Teasdale, R.D., Coulson, E.J., 2019. Downregulation of SNX27 expression does not exacerbate amyloidogenesis in the APP/PS1 Alzheimer's disease mouse model. Neurobiology of aging 77, 144-153.

Nonaka, N., Shioda, S., Banks, W.A., 2005. Effect of lipopolysaccharide on the transport of pituitary adenylate cyclase activating polypeptide across the blood-brain barrier. Exp Neurol 191(1), 137-144.

Okoye, M.I., Watanabe, I., 1982. Ultrastructural features of cerebral amyloid angiopathy. Hum Pathol 13(12), 1127-1132.

Oliver, J.M., Berlin, R.D., Davis, B.H., 1984. Use of horseradish peroxidase and fluorescent dextrans to study fluid pinocytosis in leukocytes. Methods in enzymology 108, 336-347.

Rosen, D.R., Siddique, T., Patterson, D., Figlewicz, D.A., Sapp, P., Hentati, A., Donaldson, D., Goto, J., O'Regan, J.P., Deng, H.X., et al., 1993. Mutations in Cu/Zn superoxide dismutase gene are associated with familial amyotrophic lateral sclerosis. Nature 362(6415), 59-62.

Sarkar, S., Chigurupati, S., Raymick, J., Mann, D., Bowyer, J.F., Schmitt, T., Beger, R.D., Hanig, J.P., Schmued, L.C., Paule, M.G., 2014. Neuroprotective effect of the chemical chaperone, trehalose in a chronic MPTP-induced Parkinson's disease mouse model. Neurotoxicology 44, 250-262.

Saunders, N.R., Dziegielewska, K.M., Mollgard, K., Habgood, M.D., 2015. Markers for blood-brain barrier integrity: how appropriate is Evans blue in the twenty-first century and what are the alternatives? Front Neurosci 9, 385.

Stokum, J.A., Gerzanich, V., Simard, J.M., 2016. Molecular pathophysiology of cerebral edema. J Cereb Blood Flow Metab 36(3), 513-538.

Thomsen, M.S., Routhe, L.J., Moos, T., 2017. The vascular basement membrane in the healthy and pathological brain. J Cereb Blood Flow Metab 37(10), 3300-3317.

Wan, H.I., Jacobsen, J.S., Rutkowski, J.L., Feuerstein, G.Z., 2009. Translational medicine lessons from flurizan's failure in Alzheimer's disease (AD) trial: Implication for future drug discovery and development for AD. Clin Transl Sci 2(3), 242-247.

Winkler, E.A., Sengillo, J.D., Sagare, A.P., Zhao, Z., Ma, Q., Zuniga, E., Wang, Y., Zhong, Z., Sullivan, J.S., Griffin, J.H., Cleveland, D.W., Zlokovic, B.V., 2014. Blood-spinal cord barrier disruption contributes to early motor-neuron degeneration in ALS-model mice. Proc Natl Acad Sci U S A 111(11), E1035-1042.

Wolman, M., Klatzo, I., Chui, E., Wilmes, F., Nishimoto, K., Fujiwara, K., Spatz, M., 1981. Evaluation of the dye-protein tracers in pathophysiology of the blood-brain barrier. Acta neuropathologica 54(1), 55-61.

Zhao, Z., Nelson, A.R., Betsholtz, C., Zlokovic, B.V., 2015. Establishment and Dysfunction

of the Blood-Brain Barrier. Cell 163(5), 1064-1078.

Zhong, Z., Deane, R., Ali, Z., Parisi, M., Shapovalov, Y., O'Banion, M.K., Stojanovic, K.,

Sagare, A., Boillee, S., Cleveland, D.W., Zlokovic, B.V., 2008. ALS-causing SOD1 mutants

generate vascular changes prior to motor neuron degeneration. Nat Neurosci 11(4), 420-422.

Zhu, S., Wang, J., Zhang, Y., He, J., Kong, J., Wang, J.F., Li, X.M., 2017. The role of

neuroinflammation and amyloid in cognitive impairment in an APP/PS1 transgenic mouse

model of Alzheimer's disease. CNS Neurosci Ther 23(4), 310-320.

Zlokovic, B.V., 2011. Neurovascular pathways to neurodegeneration in Alzheimer's disease

and other disorders. Nature reviews. Neuroscience 12(12), 723-738.

Figure Legends

Figure 1. Brain fluorescence uptake profile of three different fluorescent markers. (A)

Fluorescence uptake of sodium fluorescein (Na-Fl), FITC-albumin and FITC-dextran in LPS

treated mice after 1 hour of *i.v.* injection of fluorescent marker through tail vein ($n = 5$).

Fluorescence uptake of (**B**) Na-Fl and (**C**) FITC-albumin in control and LPS-administered

mice after 0.25, 1 and 24 hours of *i.v.* injection of fluorescent marker through tail vein ($n = 5$).

Uptake values are expressed as % ratio of fluorescent marker level in brain to serum. Data

are expressed as mean ± SEM. * $P < 0.05$, ** $P < 0.01$, **** $P < 0.0001$, one-way ANOVA.

Figure 2: Progressive increase of Na-Fl in the brain and spinal cord of R6/1

Huntington's mice throughout disease progression. A and **B** show increased Na-Fl uptake

in cortex and striatum of R6/1 mice at 12, 20 and 32 weeks when compared to wild-type (WT)

mice ($n = 5$). **C** shows an increased Na-Fl extravasation in cerebellum of R6/1 mice at 32

weeks when compared to WT mice ($n = 5$). **D** shows increased Na-Fl uptake in the spinal

cord of R6/1 mice at 20 and 32 weeks of age when compared to WT mice ($n = 5$). Data are

presented as mean ± SEM. * $P < 0.05$, ** $P < 0.01$, *** $P < 0.001$, **** $P < 0.0001$,

Student's t-test for each time point.

Figure 3: Increased uptake of Na-Fl in the brain and spinal cord of hSOD1^{G93A}

amyotrophic lateral sclerosis mice throughout disease progression. A, **B** and **C** shows

increased fluorescence uptake of Na-Fl in cortex, striatum and cerebellum of hSOD1^{G93A}

(SOD1) mice at 70, 130 and 175 days when compared to wild-type (WT) mice ($n = 5$). **D**

shows increased Na-Fl uptake in the spinal cord of SOD1 mice at 130 and 175 days of age

when compared to WT mice ($n = 5$). Data are presented as mean ± SEM. * $P < 0.05$, ** $P <$

0.01, **** $P < 0.0001$, Student's t-test for each time point

Figure 4: Biphasic brain and spinal cord Na-Fl uptake in the APP/PS1 mouse model of Alzheimer's disease. Fluorescence was examined in the cortex of APP/PS1 and wild-type (WT) mice at an early (4 months) and late (18 months) age ($n = 5$). Na-Fl uptake was examined in the (**A**) cortex, (**B**) hippocampus, (**C**) striatum, (**D**) cerebellum, and (**E**) spinal cord ($n = 5$). Data are presented as mean ± SEM. * $P < 0.05$, ** $P < 0.01$, Student's t-test for each time point. *P* values < 0.1 are also identified on the graph.