Genotyping of Russian isolates of fungal pathogen *Trichophyton rubrum*, based on simple sequence repeat and single nucleotide polymorphism

Running title: Population structure in *Trichophyton rubrum*

Ivan M. Pchelin[1], Yuri V. Mochalov[1], Daniil V. Azarov[2], Sofya A. Romanyuk[3], Galina A. Chilina[1], Irina V. Vybornova[1], Tatiyana V. Bogdanova[4], Vasily V. Zlatogursky[5], Svetlana V. Apalko[3], Natalya V. Vasilyeva[1,4], Anastasia E. Taraskina[1]

[1]Kashkin Research Institute of Medical Mycology, North-Western State Medical University named after I.I. Mechnikov, Saint Petersburg, Russia

[2]Department of Epidemiology, Parasitology and Disinfectology, North-Western State Medical University named after I.I. Mechnikov, Saint Petersburg, Russia

[3]City Hospital No. 40, Saint Petersburg, Russia

[4]Department of Medical Microbiology, North-Western State Medical University named after I.I. Mechnikov, Saint Petersburg, Russia

[5]Department of Invertebrate Zoology, Faculty of Biology, St. Petersburg State University, Saint Petersburg, Russia

Correspondence

Ivan M. Pchelin, Kashkin Research Institute of Medical Mycology, North-Western State Medical University named after I.I. Mechnikov, Saint Petersburg, Russia

Email: arcella.oraia@gmail.com

Funding information

Russian Foundation for Basic Research, Grant Number: 18-34-00153

Highlights

- Russian population of fungal pathogen *Trichophyton rubrum* is formed by two global genetic lineages
- Population structure in *T. rubrum* can be studied by improved microsatellite typing assay along with SNP typing and whole genome sequencing
- In *T. rubrum*, there is seemingly no association between genetic lineages and types of infection, places of geographic origin, aldolase gene expression and urease activity
- In *T. rubrum*, microevolution is driven by strong directional selection of yet unknown nature

Abstract

The *Trichophyton rubrum* species group consists of prevalent causative agents of human skin, nail and hair infections, including *T. rubrum* sensu stricto and *T. violaceum*, as well as other less well established or debatable taxa like *T. soudanense*, *T. kuryangei* and *T. megninii*. We aimed to describe genetic lineages in Russian *T. rubrum* sensu stricto, to find whether these lineages possess specific characteristics and to identify factors, shaping the population structure. We assessed polymorphism of 12 simple sequence repeat (SSR, or microsatellite) loci and TERG_02941 protein-coding gene in 70 *T. rubrum* isolates and also performed phylogenomic analysis. In both cases, two genetic lineages were found. The analysis of molecular variance (AMOVA) on the basis of SSR typing data indicated that 22-48% of the variability was between groups within *T. rubrum*. The pattern of phylogenetic changes in polymorphic protein-coding loci and SSR typing results revealed the pressure of strong directional selection, but its sources remained elusive. Our results suggest that Russian

50   population of *T. rubrum* consists of two cosmopolitan genetic lineages, with no clear

51   connection of population structure with types of infection, places of geographic origin,

52   aldolase gene expression and urease activity.

Keywords

55   microsatellites, population structure, dermatophyte, *Trichophyton rubrum*

1. Introduction

Trichophyton rubrum is an ascomycete from the order Onygenales. It is considered to be prevalent fungus of medical importance in most countries due to its major contribution to etiology of toenail onychomycosis and foot skin mycosis (Sigurgeirsson, Baran 2014). *Trichophyton rubrum* sensu stricto is a part of species group, including also *T. violaceum* and *T. soudanense* along with several forms, potentially deserving species status. All of these can be differentiated by ribosomal DNA internal transcribed spacer (ITS) region sequences (Summerbell et al. 1999; Li et al. 2008; Zhan et al. 2018; Packeu et al. 2019; Su et al. 2019). Statistical analysis of genetic polymorphism in simple sequence repeats (SSR, or microsatellites) revealed clonal mode of reproduction in the species (Gräser et al. 2007; Gong et al. 2016). It is expectedly accompanied by low genetic polymorphism (Gräser et al. 1999). A number of studies of distribution of mating type locus idiomorphs in *T. rubrum* population found only one mating type locus variant, from two needed for mating (Li et al. 2010; Kano et al. 2013; Persinoti et al. 2018). A certain proportion of *T. rubrum* isolates, belonging to the so-called "raubitschekii" morphotype, possess urease activity. This trait is believed to be ancestral, but has unclear value for taxonomy (Arabatzis et al. 2005).

To the best of our knowledge, by the moment there are two papers, reporting bipartite population structure in *T. rubrum* sensu stricto. In previous study, we reported a concordance between ribosomal non-transcribed spacer region (NTS) genotypes and nonsynonymous single nucleotide polymorphisms (SNPs) in two protein-coding loci, TERG_02941 and TERG_03298 in *T. rubrum* isolates from a city in North-Western Russia (Pchelin et al. 2018a). The two genetic lines were named ST1 and ST2. Also, Zheng et al. (2019) studied a sample of *T. rubrum* isolates, covering major part of China, by whole genome sequencing

(WGS). A number of other studies obtained *T. rubrum* WGS data from wide variety of

geographic locations, covering Europe, Asia and North America (Martinez et al. 2012; Latka

et al. 2015; Persinoti et al. 2018; Zhan et al. 2018; Zheng et al. 2019). However, there have

been no studies, bringing together data on SNP, SSR, WGS and NTS polymorphism in

T. rubrum.

Our study aimed to describe genetic lineages of *T. rubrum* sensu stricto in Russia, to

find whether these lineages possess specific characteristics and, finally, to identify factors,

shaping the population structure. We argue that Russian *T. rubrum* population is represented

by ST1 and ST2 lineages sensu Pchelin et al. (2018a). This structure could not explain

observed differences in types of infection, places of geographic origin, aldolase gene

expression and urease activity. The population structure in *T. rubrum* was seemingly shaped

by strong directional selection, but its sources remained elusive.

2. Materials and Methods

2.1. Fungal isolates and species identification

Clinical isolates of *T. rubrum* have been obtained in the clinic of Kashkin Research

Institute of Medical Mycology, Saint Petersburg (n = 44) and in Sverdlovsk Regional

Dermatovenereologic Dispensary, Yekaterinburg (n = 23) from October 2015 to January 2018

(Supplement S1) and subcultured on Sabouraud agar medium, with the addition of 2%

dextrose. We also used two reference strains RCPF 1280 and RCPF 1408 and the type strain

CBS 392.58 up to a total of 70 strains in the sample. This sample shared 19 isolates with our

earlier study (Pchelin et al. 2018a). The following clinical *T. rubrum* isolates were deposited

in the Russian Collection of Pathogenic Fungi (RCPF): D15P33, RCPF 1864; D15P44, RCPF

1865; D15P50, RCPF 1866; D15P62, RCPF 1867; D15P71, RCPF 1868; Y1013, RCPF 1869;

D15P139, RCPF 1870. The urease activity was tested twice in Christensen's urea broth,

supplemented with chloramphenicol to the final concentration of 40 µg/ml.

Species identification was performed by phenotypic methods and confirmed by DNA

sequencing of ITS region. To extract genomic DNA, we used GeneJET Plant Genomic DNA

Purification Mini Kit (Thermo Fisher Scientific, Lithuania). Aerial mycelium and fungal

spores were scraped by pipette tips, placed into microtubes containing Lysis Buffer A, beaten

with two glass beads in Minilys tissue homogenizer (Bertin Technologies, France),

supplemented with Lysis Buffer B and RNase A and further treated according to the kit

manufacturer's protocol. Amplification of ITS region with ITS5 and ITS4 primers (White et

al. 1990) was performed in 50-µl volumes, using standard PCR diagnostics reaction kit

(Syntol, Russia). Reaction mixture consisted of 100 mM Tris/HCl (pH 9.0), 250 mM KCl, 10

mM Tween-20, 2.5 mM $MgCl_2$, 250 µM each dNTP and contained 5 pmol of primers (Table

1), 1 U Taq polymerase, and 2.5 µl template DNA. The thermal cycler C1000 Touch (Bio-

Rad, USA) was programmed as follows: initial denaturation step for 5 min at 95 °C, 34 cycles

of amplification: denaturation for 30 s at 95 °C, annealing 30 s at 56 °C, extension for 1 min

at 72 °C, and a final extension step for 10 min at 72 °C. The products of PCRs were purified

by isopropanol precipitation in the presence of glycogen. PCR products were sequenced on

both strands using BigDye Terminator v3.1 Cycle Sequencing Kit (Thermo Fisher Scientific,

USA) on a 3500 Genetic Analyzer (Thermo Fisher Scientific, USA).

Table 1. Primers, used in the study.

Locus	Description	Appli-cation	Sequences, 5'-3' direction	Amplicon length, bp	Reference
ITS	rDNA ITS region	DNA seq	F: GGAAGTAAAAGTCGTAACAAGG	716	White et al.
			R: TCCTCCGCTTATTGATATGC		(1990)
TERG_02941 §	Hypothetical protein	— " —	F: CGGCCTCGTTCTGCGTTAAC	449	Pchelin et
			R: CTTGCCGGCTTCCAAGTCTTT		al. (2018a)
— " —	— " —	TMA	F: TACCACGCCCGTCAAAGG	71	This paper
			R: CCACTTGTTCTTGCCGGCTT		
			Probe A:		
			ROX-TCTTCCGTTTCACTTTGCGAGTG-**BHQ1**		
			Probe G:		
			FAM-TCTTCCGTTTCACTTTGCGAGCG-**BHQ2**		
TERG_05742	DNA-dependent RNA polymerase II, *rpb2*	qPCR	F: TGCAGGAGCTGGTGGAAGA	59	Jacob et al.
			R: GCTGGGAGGTACTGTTTGATCAA		(2012)
TERG_07904	tubulin beta chain, β-*tub*	— " —	F: AACATGATGGCTGCCACTGA	253	— " —
			R: AAGATGGCAGAGCAGGTAAGGT		
TERG_06744	fructose 1,6-bisphosphate aldolase	— " —	F: TACTTTGCCGGAAAGGGAGT	139	This paper
			R: TTCTTAGCACAGTGGTCGGT		

123   DNA seq, Sanger DNA sequencing; TMA, TaqMan SNP genotyping assay; qPCR,

124   quantitative real-time PCR. § Pchelin et al. (2018a) referred to this locus as A7C99_6411.

2.2. Antifungal susceptibility testing

Antifungal susceptibility testing was performed according to EUCAST E.Def 9.3.1

document (Arendrup et al. 2017), in RPMI-1640 medium with the addition of glucose to

a final concentration of 2%. We tested the activity of terbinafine, fluconazole and itraconazole

against six *T. rubrum* isolates with newly obtained WGS sequences. Terbinafine was tested at

concentration range 0.007-4 μg/ml, fluconazole and itraconazole were tested at 0.015-8

μg/ml.

2.3. Molecular strain typing

We employed two strain typing approaches. Firstly, we probed the G793A nucleotide

substitution in protein-coding locus TERG_02941 (A7C99_6411). The amplification was

carried out under following conditions: 5 min at 95 °C; 39 cycles of 30 s at 95 °C, 1 min 5 s at

64 °C; and final elongation for 5 min at 72 °C (Pchelin et al. 2018a). All other steps were

performed as described in the section 2.1 for ITS region sequencing. To optimize the

genotyping process, we developed TaqMan SNP genotyping assay with two allele-specific

probes (Table 1). Samples were run using a C1000 Touch thermal cycler equipped with a

CFX96 real-time PCR detection system (Bio-Rad, USA) according to the following program:

initial denaturation step for 3 min at 95 °C; and 39 cycles of 10 s at 95 °C and 30 s at 59 °C.

Secondly, we performed microsatellite typing. We used eight microsatellite loci from

the papers by Gräser et al. (2007) and Gong et al. (2016) and developed four additional loci

(Table 2). To find new polymorphic loci, we searched genome contigs for repeat sequences

and made nucleotide alignments. In total, 170 loci were screened. Primers for amplification of

nine candidate loci were designed with the use of Primer-BLAST tool (Ye et al. 2012). Five

loci had uniform lengths in 21-56 isolates and were excluded from further experiments.

Amplification of microsatellite loci was performed in 25-μl volumes. The primers used are

listed in the Table 2.

Table 2. Primers, used for SSR amplification in *Trichophyton rubrum* strains. Fluorescent
modifications are given in bold.

Locus	Sequences, 5'-3' direction	Amplicon length, b.p.	Repeat unit	Length of flanking regions, b.p.	Reference
Tr001	F: **FAM**-GTGGGCTCGAACCGAGACAAAC R: GGTGACTGGAGGGTGAAGGG	263–286	GA	212	Gong et al. 2016
Tr002	F: **HEX**-CTCGCATTTGGCCTTGAGTGT R: CTGGCTGGTGGAAGGACATAG	248–281	CTT, ACT	138	— " —
Tr003	F: **FAM**-GCCAGTGATCCTGCATCGTCC R: CGGGCTTCTCGTCTTGCTCTTC	233–242	ACT	187	— " —
Tr004	F: **HEX**-CAGGAGGGTAGAAGACGAGAAG R: AAAGAAAAACAAAGTCAAGCACGC	295–300	GCA, GAA	228	— " —
Tr005	F: **FAM**-AAAGTACCTCCTGTTTACCATC R: GGCATTCCGGGCAGTTGAAGC	304–310	CT	276	— " —
Tr006	F: **HEX**-TAAGACAGGGATCTCAGAATG R: CGGAACCGCTACAAGCAGAAT	191–214	GTT	133	— " —
Tr007	F: **TAMRA**-TGGGATACAGCCATGCCGTT R: CTCAGCCCTACCAGACGACC	361–364	AG	325	This paper
Tr008	F: **TAMRA**-AGGCCCCCAAGGTAGAGGAT R: GTGCCTGCGCGAAATACGAA	195–204	GA	151	— " —
Tr009	F: **HEX**-ACGACGATACTGGCTGGACAA R: GTTATGACTTCCACGGTGTCACG	348–368	AAG	318	— " —
Tr(CT)₂₀	F: **FAM**-GCGGTGGCGTCTTCTATC R: CAGCAGACAACATAGCAGTCG	134–139	GA	101	Gräser et al. 2007
Tr(GA)₂₅	F: **HEX**-GCGATGGTTGGAGGAGTTG R: GCCTGTCGCTGCTTACTTG	143–150	GA	92	— " —
STR*Tr9*	F: **TAMRA**-ACACTGCTCTTCGCATCTTGA R: GGCTGTTGTTGTTCGCTGCT	240–342	CAAG	197	This paper

The amplification program included initial denaturation at 95 °C for 10 min, followed

by 30 cycles of 50 s at 95 °C, 1 min at 58 °C, and 1 min at 72 °C, and a terminal extension

step of 72 °C for 10 min. The set of fluorescent dyes used in this work appeared to be

suboptimal because of the need to check data for issues of spectral overlapping between

TAMRA and ROX channels.

2.4. Whole genome sequencing

Genomic DNA of six *T. rubrum* and one *T. benhamiae* isolates was extracted using

GeneJET Plant Genomic DNA Purification Mini Kit (Thermo Fisher Scientific, Lithuania).

Preparation of DNA sample library was done with the use of Kapa HTP Library Preparation

Kit (Roche, USA), according to the manufacturer's instructions. Paired end sequencing (250

× 250 cycles) was performed using the MiSeq sequencer (Illumina, USA). The reads obtained

were prepared for the analyses as described in the paper by Pchelin et al. (2019a). Quality

control of the reads was performed using FastQC 0.11.8 (Andrews 2010). It was followed by

trimming low-quality reads and TruSeq2 adapters with Trimmomatic 0.39 (Bolger et al.

2014). De novo assembly was performed with SPAdes 3.9.0 with the "--careful option" (Nurk

et al. 2013). Raw reads were submitted to the Sequence Read Archive of National Center for

Biotechnology Information (BioProject ID PRJNA552357).

2.5. Phylogenomic reconstruction

The whole genome dataset included six genomes of Russian isolates, as well as the

genomes of isolate CBS 118892, BioProject accession ACPH00000000 (Martinez et al.

2012), isolates CBS 202.88 AOKX00000000, CBS 288.86 AOKU00000000, CBS 289.86

AOKV00000000, CBS 100081 AOKT00000000, D6 AOLB00000000, MR850

AOKR00000000, MR1448 AOKZ00000000, MR1459 AOLC00000000 (Persinoti et al.

2018), isolate CMCC(F)T1i LHPM00000000 (Zhan et al. 2018), IGIB-SBL-CI1

JPGR00000000 (Latka et al. 2015). The raw read data for eight strains from the project of

Zheng et al. (2019) have been downloaded from NCBI SRA database and assembled *de novo*

for the analysis. This number included the strains LZ164 (NCBI SRA accession

SRX5814844), GZ0580 (SRX5814815), NJ785 (SRX5814822) from one cluster, and the

strains LZ155 (SRX5814845), NJ786 (SRX5814823), NJ823 (SRX5814824), DL7506

(SRX5814836) from another cluster, revealed in the original publication. The eighth strain

GZ0663 (SRX5814819) from the study of Zheng et al., analyzed here, had unclear cluster

affiliation. The sequences of *T. rubrum* morphotype "megninii" CBS 735.88

JHQM00000000, *T. soudanense* CBS 452.61 AOKW00000000 (Persinoti et al. 2018) and

T. violaceum CMCC(F)T3l LHPN00000000 (Zhan et al. 2018) were used as an outgroup.

Single-nucleotide polymorphism analysis in core genome was done in Harvest Parsnp 1.2

software (Treangen et al. 2014). Parsnp run with *T. rubrum* CBS 118892 assembly

GCA_000151425.1 as a reference and with default parameters. Whole-genome phylogenetic

tree was reconstructed by Maximum Likelihood method in FastTree2 (Price et al. 2010), as

implemented in Harvest suite. Branch support values were calculated by performing

Shimodaira-Hasegawa test on three alternate topologies around particular splits, with a total

of 1000 resamples.

2.6. Analysis of genetic polymorphism

A matrix of mutation probability-based genetic distances (Bruvo et al. 2004) was

calculated in polysat 1.7-3 package for R (Clark, Jasieniuk 2011) using microsatellite

polymorphism data. Distances in terms of weighted divergence of microsatellite loci were

calculated in Microsoft Excel 2010 according to the formula, modified for haploids from

Rozenfeld et al. (2007). The dissimilarity d_i between haplotypes *A* and *B* at particular locus *i*

was calculated as absolute value of arithmetic difference between the lengths of PCR products

A_i and B_i

$$d_i(A, B) = |A_i - B_i|$$

The genetic distance D between haplotypes A and B was based on the sum of

dissimilarities d between haplotypes at all loci, weighted by dividing this sum by the number

of loci taken into analysis, k

$$D(A,B) = \frac{1}{k}\sum_{i=1}^{k} d_i$$

Both distance matrices were further employed in the construction of phylogenetic

network in SplitsTree 4.14.2 program (Huson, Bryant 2006). Neighbor-Joining tree on the

basis of Bruvo distances was calculated in Mega 5.2 (Tamura et al. 2013).

The allele association index was calculated from the lengths of microsatellite loci in

poppr 2.8.1 package for R (Kamvar et al. 2014), using the Multilocus Style Permutation

method, after deleting one non-informative locus and performing clonal correction. The null

hypothesis was the absence of association between the lengths of microsatellite loci.

Simpson's index of genotype diversity within upper and lower bounds for a 95% confidence

interval was calculated in the package polysat on the basis of Bruvo genetic distances, in

equal groups of 21 isolates for ST1 and ST2 lineages. The analysis of molecular variation

(AMOVA) was done on the basis of Bruvo and Rozenfeld distance matrices in GenAlEx 6.5

package (Peakall, Smouse 2006, 2012) with 999 standard permutations.

TERG_03298 genotypes, MAT idiomorphs and TERG_05790 urease gene variants

were determined through Blast searches in whole genome sequences (Zhang et al. 2000). For

mating type determination, TERG_02406 gene coding for α-domain protein was probed by

AB759084 sequence (Kano et al. 2013). The absence of HMG protein-coding gene was

verified by the searches with *T. simii* HMG gene, GenBank accession number AB605766

(Kano et al. 2012). The boundaries of the minisatellite in TERG_08771 gene were determined

with the use of Tandem Repeats Finder 4.09 (Benson 1999).

To address evolutionary pattern in *T. rubrum* complex, we prepared a phylogenetic

tree on the basis of seven protein-coding genes sequences with predicted nonsynonymous

mutations among *T. rubrum* complex isolates. This number included genes, coding for

hypothetical proteins TERG_02941 (A7C99_6411), TERG_03298 (A7C99_6714),

TERG_03748, copper radical oxidase TERG_02718, LIM domain-containing protein

TERG_02913, mismatch repair protein TERG_03581, serine/threonine protein kinase

TERG_03779. The same sequences were extracted by blast searches from the genomes of *T.*

mentagrophytes, *T. interdigitale*, *T. tonsurans*, *T. equinum*, *T. benhamiae*, *T. verrucosum* and

Nannizzia gypsea (Martinez et al. 2012; Gallo et al. 2017; Alshahni et al. 2018; Persinoti et al.

2018; Pchelin et al. 2019b).

2.7. Quantitative RT-PCR

The sample of six isolates was grown on Sabouraud agar medium for two weeks at 28

°C. RNA was isolated with the use of GeneJet Plant RNA Purification Mini Kit (Thermo

Fisher Scientific, Lithuania). The quantitative RT-PCR experiments were performed in

triplicates on two independently grown colonies for each isolate in the sample with the use of

OneTube RT-PCR SYBR Kit (Evrogen, Russia). The reactions were carried out on CFX96

Touch Real-Time PCR Detection System (Bio-Rad, USA) with the following cycling

conditions: reverse transcription step at 55 °C for 15 min, polymerase activation and reverse

transcriptase inactivation step at 95 °C for 1 min, and main amplification stage for 40 cycles,

including denaturation at 95 °C for 15 s, annealing at 60 °C for 20 s and elongation at 72 °C

for 15 s. Melting curve analysis was performed to check whether one PCR product was

obtained. Relative gene expression levels were assessed using beta-tubulin and DNA-

dependent RNA polymerase II for normalization. The sequences of primers used are given in

the Table 1.

3. Results

The majority of 69 *T. rubrum* isolates from Russia and the type strain CBS 392.58 had

the same ribosomal ITS region sequence, identical to KT285224. There were two exceptions,

the isolate D15P139 had 247insC mutation, and another isolate had additional TA element in

the microsatellite near the 3'-end of the sequence (Fig. S1). Our preliminary study showed

discrepancies between the results of strain typing by TERG_02941 locus polymorphism and

the microsatellite typing assay of Gong et al. (2006), based on eight loci (Pchelin et al.

2018b). We hypothesised that this discrepancy was due to insufficient informativeness of the

assay and developed four new polymorphic loci, up to a total of 12. The stability of

microsatellite loci was assessed in 20 isolates for the time span of 1.5 years. Observed

differences of less than one repeat length were considered insignificant, since those

differences could be particularly associated with the rounding of measurements, fluctuating

near the half-integers. At the endpoint, mutations were observed in five loci. The most

extensive changes were noticed in $Tr(CT)_{20}$ locus, with 30% of isolates with mutations. The

most pronounced indels were in Tr006 and Tr008 loci, covering up to 3 repeat lengths.

Our first approach to calculating microsatellite polymorphism-based genetic distances

employed the probability of mutation events, described with the use of stepwise mutation

model (Bruvo et al. 2004). The split network, inferred using mutation probability-based

distances, was divided into three major partitions. One partition contained TERG_02941

793A isolates and another two partitions contained TERG_02941 793G isolates (Fig. 1). The

second approach was based on the comparison of the lengths of microsatellite loci (Rozenfeld

et al. 2007). The resulting phylogenetic network also had pronounced division into

TERG_02941 793A and 793G parts (Fig. 2). In Rozenfeld distances-based network, further

subdivision of TERG_02941 793G partition was less evident, but still discernable. From 22

TERG_02941 793G isolates, 18-21 located to the same subgroups in both networks.

For the two genetic lines, the average lengths of PCR products of Tr001, Tr002,

Tr003, Tr004, Tr005, Tr007, Tr008, Tr009, Tr(CT)$_{20}$ and Tr(GA)$_{25}$ loci differed by less than

one repeat unit length. Average Tr006 lengths differed by a unit length. For STR*Tr*9 locus,

this difference was 7.5 unit lengths (Fig. S2A). This difference should be obviously explained

by the pressure of natural selection. Since the analysis of intraspecific divergence in most

cases is carried out using neutral loci (Leliaert et al. 2014), we performed calculation of

genetic distances without STR*Tr*9 length data. This broke clear grouping of isolates in both

phylogenetic networks. However, a certain tendency to clustering of TERG_02941 793G

isolates persisted (Figs. 3, 4).

The analysis of molecular variance with 12 microsatellite loci revealed neglectable

contribution of TERG_02941 793G subdivision to intergroup and overall variability (Table

3). With the removal of STR*Tr*9 from the analysis, the percent of variability, explained by the

differences between groups, dropped drastically. But, this was accompanied by significant

increase in explanatory power of TERG_02941 793G group subdivision. A half of the

between-group variability in the dataset with 11 loci depended on it.

Table 3. The analysis of molecular variance (AMOVA) in 70 *T. rubrum* isolates, performed on microsatellite polymorphism-based genetic distances. The numbers reflect percent of variability among TERG_02941 793A and 793G groups of isolates from the total amount of variability between and within groups. To obtain three groups, TERG_02941 793G isolates were further divided according to the topology of Bruvo distances-based network (left-hand numbers in columns with data on 3 lineages) or by Rozenfeld distances-based network (right-hand numbers).

	Bruvo et al. (2004)		Rozenfeld et al. (2007)	
	2 groups	3 groups	2 groups	3 groups
12 loci	22	27-24	48	48-48
11 loci	9	18-14	10	19-20

The CAAG repeats of STR*Tr*9 ended at the position -278 in the 5'-region of fructose 1,6-bisphosphate aldolase gene TERG_06744, wherein the adenine of the ATG translation initiation codon is considered +1. Therefore, we hypothesized that STR*Tr*9 locus promotes microevolution in *T. rubrum* through regulation of aldolase gene expression, which in its turn is one of the key enzymes of glycolysis (Nelson, Cox 2008). We determined relative TERG_06744 gene expression in six *T. rubrum* isolates. The isolate D15P44 with maximal length of STR*Tr*9 locus demonstrated the lowest level of TERG_06744 transcripts (Fig. S2B). The highest levels of TERG_06744 transcripts were observed in D15P62 isolate with the microsatellite length, very close to minimal one. In the same time, three isolates with the same STR*Tr*9 length demonstrated significantly lower level of TERG_06744 gene expression. Also, there were no clear differences between TERG_06744 transcript levels in isolates of different sequence types. Since urease activity has been regarded as phylogenetically important trait in the dermatophytes (Summerbell 2000), we assessed urease activity in our sample and found three urease-positive strains (Supplement S1).

In the sample of 70 isolates, the association index value $\bar{r}_d = 0.0248$ lay beyond the right boundary of the distribution of expected frequencies. The probability $\bar{r}_d$ was estimated as

0.002, which made it possible to reject the null hypothesis of the absence of connections

between the markers. Hence, the studied *T. rubrum* population was considered to be clonal.

Also, six Russian isolates, subjected here to whole genome sequencing, harbored *MAT1-1*

idiomorph of mating type locus. Simpson index was calculated on the basis of data from 11

neutral SSR loci. Its value within 95% confindence interval was estimated as 0.59-1.03 for

ST1 lineage and 0.38-0.91 for ST2 lineage. The Fisher exact test showed no significant

association between TERG_02941 genotypes and the localization of the lesion on patient's

body. Also, Yates-corrected χ^2 test showed no connection between TERG_02941 genotype

and a city of origin.

Russian sample of *T. rubrum* isolates was placed in global context by whole genome

sequencing in six *T. rubrum* isolates. To get a representative sample, we selected isolates

D15P33, D15P44, D15P50, D15P62, D15P71 and Y1013, covering the whole span of

microsatellite distances-based Neighbour-Joining tree (not shown). These isolates were

susceptible to tested antifungals (Table 4).

Table 4. Annotation and antifungal susceptibility data for six *T. rubrum* isolates, subjected to
whole genome sequencing.

Isolate	Year of isolation	Place of isolation	Clinical picture	MIC$_{50}$ of antifungals (μg/ml)		
				TER	FCZ	ICZ
D15P33	2015	St. Petersburg	OM, TP	≤ 0.01	0.125	0.06
D15P44	2015	St. Petersburg	OM	≤ 0.01	0.25	0.125
D15P50	2016	St. Petersburg	OM	≤ 0.01	0.125	0.06
D15P62	2016	St. Petersburg	OM	≤ 0.01	0.25	0.06
D15P71	2016	St. Petersburg	OM	≤ 0.01	0.06	0.125
Y1013	2017	Yekaterinburg	OM	≤ 0.01	0.25	0.125

Clinical picture: OM, toenail onychomycosis; TP, tinea pedis. Antifungals: TER, terbinafine;
ICZ, itraconazole; FCZ, fluconazole.

The phylogenomic tree included *T. rubrum* sensu stricto genomes from Europe, Asia

and North America (Fig. 5) and contained well-supported monophyletic branch of ST1

isolates and basal paraphyletic ST2 branch. Both branches contained European and Asian

isolates. In a representative selection of seven isolates from the study of Zheng et al. (2019),

cluster affiliation, determined in the original study, matched their TERG_02941/03298

genotypes. Depending on the inclusion or the exclusion of *T. violaceum* CMCC(F)T3l and

T. rubrum "megninii" CBS 735.88 from the phylogenomic analysis, the depth of branches

within *T. rubrum* sensu stricto clade related to *T. rubrum* sensu stricto — *T. soudanense* CBS

452.61 branch as 1:63 or 1:337.

The study of Zheng et al. (2019) has associated two SNPs in TERG_08771 with

voriconazole resistance. Unfortunately, these mutations were located in continuous

degenerated minisatellite, spanning 1699 bp in long read-based assembly LHPM01 and

2248 bp in reference assembly ACPH01. We were unable to assemble the whole span of

TERG_08771 gene unambiguously and perform the alignment, needed to determine the

mutations. Also, in the same study the position NW_003456428.1 1020017, residing in

TERG_00393 gene, has been associated with griseofulvin resistance. In our dataset, this

position fell into a microsatellite and wasn't polymorphic.

There is an observation of the 68-bp intron and the 9-bp insertion in a urease-positive

T. rubrum strain (Hiruma et al. 2013). However, similarly to the results of Adamski et al.

(2014), we found identical urease gene sequences in all *T. rubrum* sensu stricto isolates,

shown on the Fig. 5, including urease-positive D15P50 and CBS 202.88. *Trichophyton*

rubrum "megninii" CBS 735.88 (Persinoti et al. 2018) harbored two point mutations in the

urease gene, A685G in the intron and synonymous substitution G803A in the second exon.

All our three isolates, positive for urease activity, came from the cases of toenail and foot skin

infections and belonged to both major genetic lineages of *T. rubrum*.

We studied seven protein-coding loci with predicted nonsynonymous SNPs,

differentiating *T. rubrum* strains: TERG_02718, TERG_02913, TERG_02941, TERG_03298,

TERG_03581, TERG_03748 and TERG_03779 (Fig. 6). The nonsynonymous SNPs in loci

TERG_02718, TERG_02941, TERG_03298, TERG_03581 and TERG_03748 allowed

distinguishing groups of isolates, whereas mutations in TERG_02913 and TERG_03779

occurred in single isolates. Unlike *T. mentagrophytes* / *T. interdigitale* species group isolates,

T. rubrum isolates demonstrated consecutive branching pattern, suggestive of strong

directional selection of discriminative SNPs (Ito, Dieckmann 2007; Edwards 2009).

In our previous work, we associated single-banded TRS-1 pattern with the length of

approximately 450 bp with particular TERG_02941/03298 genotype, substantiating the

presence of ST1 and ST2 genetic lines in Russian population of *T. rubrum* (Pchelin et al.

2018a). Later on, Zheng et al. 2019 observed the subdivision of Chinese *T. rubrum* population

into two groups. Here through phylogenomic reconstruction we showed that those groups not

only match Russian ones, but also have cosmopolitan distribution. Therefore we believed that

mapping global distribution of *T. rubrum* sequence types sensu Pchelin et al. (2018a) is

possible (Fig. 7). The map of global distribution was prepared with the use of strain typing

data from the following publications: Jackson et al. (2000), Iceland, Japan, the UK; Baeza et

al. (2006), Brazil; de Assis Santos et al. (2007), Brazil; Rad et al. (2008), Iran; Hryncewicz-

Gwóźdź et al. (2011), Poland; Takahashi et al. (2014), Japan; Ramaraj et al. (2017), India;

Pchelin et al. (2018a), Russia; Suzuki et al. (2018), Japan; Zheng et al. (2019), China.

4. Discussion

Until recently, low genetic diversity of *T. rubrum* precluded persuasive description of

its population structure. For this species, our earlier work (Pchelin et al. 2018a) seemingly

represented the first successful attempt to obtain congruent results by two independent

molecular strain typing techniques. However, we have not performed DNA sequencing of

ribosomal ITS region to clarify whether the sample contained only *T. rubrum* sensu stricto

isolates or not. Here we filled this gap by demonstrating that Russian population of the fungus

is represented by only «classic» *T. rubrum* genotype. The two genetic lines of Russian

T. rubrum population, found here, demonstrated significant divergence. The AMOVA

analysis showed relatively high proportion of variability, explained by the division of

T. rubrum sample into ST1 and ST2 groups, 22-48% depending on the distance metrics. It

was not less, than in other studies on clonal filamentous ascomycetes with 19-21% of

variability between groups (Ahmadpour et al. 2017; Singh et al. 2019) or filamentous

ascomycetes with mixed reproduction mode, with 2-13% of total variability explained by

differences between intraspecific groups (Kashyap et al. 2020; Sexton & Howlett, 2004).

Intraspecific genetic lines in fungi sometimes demonstrate restricted geographic

ranges (Abdel-Rachman et al. 2010; Mironenko et al. 2016; Gultyaeva et al. 2017; Taghipour

et al. 2019). Here, we compiled the map of global distribution of *T. rubrum* genetic lineages

(Fig. 7). We largely built upon existing literature on NTS typing in *T. rubrum* and departed

from the fact that single 450 bp PCR product of TRS-1 locus is associated with ST1 genetic

lineage (Pchelin et al. 2018a). We didn't observe a specific distribution pattern of the two

genetic lineages, but in most parts of the World, the derived group of *T. rubrum*, called ST1,

presumably had higher prevalence and therefore demonstrated evolutionary success (Fig. 7).

Therefore, we studied the pattern of evolutionary changes in *T. rubrum*. The branching pattern

of phylogenetic tree, based upon seven polymorphic protein-coding loci consisted of

consequential branching events (Fig. 6). There were no inner branches, as in the

T. mentagrophytes / *T. interdigitale* species group. This pattern can be explained by clonal

divergence, driven by directional selection. Intriguingly, this pattern obviously can't be

explained neither by the selective pressure of climatic factors, nor by the adaptation to the

host.

The affiliation of *T. rubrum* isolates to genetic lineages can be probed not only by

NTS, SNP and WGS-based typing techniques, but also by our modified scheme of

microsatellite analysis. But, this result may be of limited interest for healthcare practitioners,

since there is seemingly no association of lineage affiliation and common isolate descriptors,

like geography of origin and clinical picture of respective infection. Also, in the study of

Zheng et al. (2019), there was no association between cluster affiliation and antifungal

susceptibility levels. It is widely accepted that truly independent species of clonal microbes

occupy distinct ecological niches (Tibayrenc, 2006; Achtman & Wagner, 2008; Giraud et al.

2008). In present study, there seemingly have been no clinical differences between the

subpopulations, contrary to what we has claimed earlier (Pchelin et al. 2018a). In that

previous study, the association of STs and clinical forms of dermatophytosis might have

occurred due to some random factors. From this perspective, the two genetic lines described

here belong to the same fungal species.

The STR*Tr*9 locus was paramount for the analysis of the structure of the *T. rubrum*

population by microsatellite analysis. The fact that its length is significantly different between

the isolates of two genetic lines meant that it was under the pressure of natural selection. The

regulatory role of this microsatellite could not be confirmed experimentally. But, there were

significant differences in relative aldolase gene expression levels between different isolates.

Also, in our sample there were three urease-positive isolates, a proportion of 4% from the

total. We did not find peculiarities in urease gene sequences of urease-positive isolates

D15P50 and CBS 202.88. Positive strains belonged to both ST1 and ST2 groups. Therefore,

our results suggest that urease-positive strains present in certain proportion in the population

of the fungus, and represent natural variation of *T. rubrum* sensu stricto, not an independent

entity. Lackner et al. (2019) obtained comparable results, studying virulence potential of

closely related *Aspergillus* species from the section *Terrei*. They concluded that virulence is a

strain-specific rather than species-specific feature. Lineage-specific differences in STR*Tr9*

locus length may still hallmark different adaptation strategies, given that glycolysis pathway

regulation is connected with mechanisms of keratin degradation (Martins et al. 2020).

Urease activity is an ancestral trait, commonly observable in geophilic dermatophytes

(Summerbell 2000) and we expected to find it only in basal ST2 lineage. However, in our

sample one of three urease-positive isolates belonged to ST1 lineage. Also, genetic diversity

in the two lineages proved to be the same, which is contrary to what might be expected if ST2

is an ancestral lineage. It can be probably due to a sort of bottleneck, occurred in the

population on the way from South-East Asia to Russia. In future studies, it is necessary to

sample isolates from South and South-East Asia, in order to find the territory with highest

genetic diversity and thereby conclusively determine the geographic region of the origin of

the species.

Thus, we provided the description of the two intraspecific global lineages of *T. rubrum*

sensu stricto. They can be revealed by SNP, SSR and WGS-based typing techniques (and

NTS, with a certain amount of ambiguity). This population structure is shaped by pressure of

natural selection of yet unknown nature. Also, the species is polymorphic in terms of ecology

and physiology, but this polymorphism is not associated with genetic population structure.

Acknowledgements

The authors would like to thank the CBS-KNAW culture collection of Westerdijk Fungal Biodiversity Institute, Utrecht, the Netherlands, for providing the strain CBS 392.58. We also appreciate valuable comments from Adéla Čmoková and Ali Rezaei-Matehkolaei. This work was supported by Russian Foundation for Basic Research [grant number 18-34-00153].

Author contributions

I.P. designed the study. I.P., Y.M., I.V., G.C., S.R. and T.B. performed the experiments. I.P., D.A. and Y.M. analyzed the data with the participation of A.T. N.V., A.T. and S.A. performed supervision and funding acquisition. I.P. wrote the manuscript with the help of V.Z.

Declarations of interest:

none

Data accessibility

Raw WGS reads were submitted to NCBI SRA database, BioProject PRJNA552357: the isolate D15P33, accession SRX6389058; D15P44, SRX6389060; D15P50, SRX6389057; D15P62, SRX6389059; D15P71, SRX6389062; Y1013, SRX6389061.

ORCID

Svetlana V. Apalko https://orcid.org/0000-0002-3853-4185
Daniil V. Azarov https://orcid.org/0000-0003-2483-5144
Ivan M. Pchelin https://orcid.org/0000-0003-3062-3316
Vasily V. Zlatogursky https://orcid.org/0000-0002-2688-3900
Anastasia E. Taraskina https://orcid.org/0000-0002-1725-8433
Natalya V. Vasilyeva https://orcid.org/0000-0003-3693-5468

References

Abdel-Rahman, S. M., Sugita, T., González, G. M., Ellis, D., Arabatzis, M., Vella-Zahra, L.,
... Preuett, B. (2010). Divergence among an international population of *Trichophyton*
tonsurans isolates. *Mycopathologia, 169*(1), 1–13. https://doi.org/10.1007/s11046-009-9223-7

Achtman, M., & Wagner, M. (2008). Microbial diversity and the genetic nature of microbial
species. *Nature Reviews Microbiology, 6*(6), 431–440. https://doi.org/10.1038/nrmicro1872

Adamski, Z., Kowalczyk, M. J., Adamska, K., Kubisiak-Rzepczyk, H., Bowszyc-
Dmochowska, M., Banaszak, A., … Zaba, R. (2014). The first non-African case of
Trichophyton rubrum var. *raubitschekii* or a urease-positive *Trichophyton rubrum* in Central
Europe? *Mycopathologia, 178*(1-2), 91–96. https://doi.org/10.1007/s11046-014-9751-7

Ahmadpour, A., Castell-Miller, C., Javan-Nikkhah, M., Naghavi, M. R., Dehkaei, F. P., Leng,
Y., ... Zhong, S. (2018). Population structure, genetic diversity, and sexual state of the rice
brown spot pathogen *Bipolaris oryzae* from three Asian countries. *Plant Pathology, 67*(1),
181–192. https://doi.org/10.1111/ppa.12714

Alshahni, M. M., Yamada, T., Yo, A., Murayama, S. Y., Kuroda, M., Hoshino, Y., …
Makimura, K. (2018). Insight into the draft whole-genome sequence of the dermatophyte
Arthroderma vanbreuseghemii. *Scientific Reports, 8*(1), 15127.
https://doi.org/10.1038/s41598-018-33505-9

Andrews, S. (2010). FastQC: a quality control tool for high throughput sequence data.
http://www.bioinformatics.babraham.ac.uk/projects/fastqc. accessed: 22.02.2020.

Arabatzis, M., Velegraki, A., Kantardjiev, T., Stavrakieva, V., Rigopoulos, D., & Katsambas,
A. (2005). First report on autochthonous urease-positive *Trichophyton rubrum* (*T.
raubitschekii*) from South-east Europe. *British Journal of Dermatology, 153*(1), 178–182.
https://doi.org/10.1111/j.1365-2133.2005.06615.x

Arendrup, M. C., Meletiadis, J., Mouton, J. W., Lagrou, K., Hamal, P., Guinea, J., … Rogers,
T. (2017). EUCAST definitive document E.Def 9.3.1 method for the determination of broth
dilution minimum inhibitory concentrations of antifungal agents for conidia forming moulds.
EUCAST.

Baeza, L. C., Matsumoto, M. T., Almeida, A. M., & Mendes-Giannini, M. J. (2006). Strain
differentiation of *Trichophyton rubrum* by randomly amplified polymorphic DNA and
analysis of rDNA nontranscribed spacer. *Journal of Medical Microbiology, 55*(4), 429–436.
https://doi.org/10.1099/jmm.0.46236-0

Benson, G. (1999). Tandem repeats finder: a program to analyze DNA sequences. *Nucleic
Acid Research, 27*(2), 573–580. https://doi.org/10.1093/nar/27.2.573

Bolger, A. M., Lohse, M., & Usadel, B. (2014). Trimmomatic: a flexible trimmer for Illumina
sequence data. *Bioinformatics, 30*(15), 2114–2120.
https://doi.org/10.1093/bioinformatics/btu170

Bruvo, R., Michiels, N. K., D'Souza, T. G., & Schulenburg, H. (2004). A simple method for
the calculation of microsatellite genotype distances irrespective of ploidy level. *Molecular*
Ecology, 13(7), 2101–2106. https://doi.org/10.1111/j.1365-294X.2004.02209.x

Clark, L., & Jasieniuk, M. (2011). Polysat: an R package for polyploid microsatellite analysis.
Molecular Ecology Resources, 11(3), 562–566. https://doi.org/10.1111/j.1755-
0998.2011.02985.x

de Assis Santos, D., de Carvalho Araújo, R. A., Kohler, L. M., Machado-Pinto, J., Hamdan, J.
S., & Cisalpino, P. S. (2007). Molecular typing and antifungal susceptibility of *Trichophyton*
rubrum isolates from patients with onychomycosis pre- and post-treatment. *International*
Journal of Antimicrobial Agents, 29(5), 563–569.
https://doi.org/10.1016/j.ijantimicag.2006.09.028

Edwards, S. V. (2009). Natural selection and phylogenetic analysis. *Proceedings of the*
National Academy of Sciences, 106(22), 8799–8800.
https://doi.org/10.1073/pnas.0904103106

Gallo, J. G., Woods, M., Graham, R. M., & Jennison, A. V. (2017). A severe transmissible
Majocchi's granuloma in an immunocompetent returned traveler. *Medical Mycology Case*
Reports, 18, 5–7. https://doi.org/10.1016/j.mmcr.2017.07.003

Giraud, T., Refrégier, G., Le Gac, M., de Vienne, D. M., & Hood, M. E. (2008). Speciation in
fungi. *Fungal Genetics and Biology, 45*(6), 791–802.
https://doi.org/10.1016/j.fgb.2008.02.001

Gong, J., Wu, W., Ran, M., Wang, X., Liu, W., Wan, Z., … Li, R. (2016). Population
differentiation and genetic diversity of *Trichophyton rubrum* as revealed by highly
discriminatory microsatellites. *Fungal Genetics and Biology, 95*, 24–29.
https://doi.org/10.1016/j.fgb.2016.08.002

Gräser, Y., Fröhlich, J., Presber, W., & de Hoog, S. (2007). Microsatellite markers reveal
geographic population differentiation in *Trichophyton rubrum*. *Journal of Medical*
Microbiology, 56(8), 1058–1065. https://doi.org/10.1099/jmm.0.47138-0

Gräser, Y., Kühnisch, J., & Presber, W. (1999). Molecular markers reveal exclusively clonal
reproduction in *Trichophyton rubrum*. *Journal of Clinical Microbiology, 37*(11), 3713–3717.

Gultyaeva, E. I., Aristova, M. K., Shaidayuk, E. L., Mironenko, N. V., Kazartsev, I. A.,
Akhmetova, A. & Kosman, E. (2017). Genetic differentiation of *Puccinia triticina* Erikss. in
Russia. *Russian Journal of Genetics, 53*(9), 998–1005. https://10.1134/S1022795417070031

Hiruma M, Kano R, Sugita T, Mochizuki T, Hasegawa A, Hiruma M. Urease gene of
Trichophyton rubrum var. *raubitschekii*. J Dermatol. 2013; 40: 111–113.
https://10.1111/1346-8138.12017

Hryncewicz-Gwóźdź, A., Jagielski, T., Sadakierska-Chudy, A., Dyląg, M., Pawlik, K., Baran,
E., & Szepietowski, J. C. (2011). Molecular typing of *Trichophyton rubrum* clinical isolates
from Poland. *Mycoses, 54*(6), e726–e736. https://doi.org/10.1111/j.1439-0507.2010.02007.x

Huson, D. H., & Bryant, D. (2006). Application of phylogenetic networks in evolutionary
studies. *Molecular Biology and Evolution, 23*(2), 254–267.
https://doi.org/10.1093/molbev/msj030

Jackson, C. J., Barton, R. C., Kelly, S. L., & Evans, E. G. V. (2000). Strain identification of
Trichophyton rubrum by specific amplification of subrepeat elements in the ribosomal DNA
nontranscribed spacer. *Journal of Clinical Microbiology, 38*(12), 4527–4534.

Jacob, T. R., Peres, N. T., Persinoti, G. F., Silva, L. G., Mazucato, M., Rossi, A., & Martinez-
Rossi, N. M. (2012). rpb2 is a reliable reference gene for quantitative gene expression
analysis in the dermatophyte *Trichophyton rubrum. Medical Mycology, 50*(4), 368–377.
https://doi.org/10.3109/13693786.2011.616230

Ito, H. C., & Dieckmann, U. (2007). A new mechanism for recurrent adaptive radiations. *The
American Naturalist, 170*(4), E96–E111. https://doi.org/10.1086/521229

Kamvar, Z. N., Tabima, J. F., & Grünwald, N. J. (2014). Poppr: an R package for genetic
analysis of populations with clonal, partially clonal, and/or sexual reproduction. *PeerJ, 2*,
e281. http://dx.doi.org/10.7717/peerj.281

Kano, R., Isizuka, M., Hiruma, M., Mochizuki, T., Kamata, H., & Hasegawa, A. (2013).
Mating type gene (*MAT1-1*) in Japanese isolates of *Trichophyton rubrum. Mycopathologia,
175*(1-2), 171–173. https://doi.org/10.1007/s11046-012-9603-2

Kano, R., Kawasaki, M., Mochizuki, T., Hiruma, M., & Hasegawa, A. (2012). Mating genes
of the *Trichophyton mentagrophytes* complex. *Mycopathologia, 173*(2-3), 103–112.
https://doi.org/10.1007/s11046-011-9487-6

Kashyap, P. L., Kumar, S., Kumar, R. S., Tripathi, R., Sharma, P., Sharma, A., ... Singh, G. P.
(2020). Identification of novel microsatellite markers to assess the population structure and
genetic differentiation of *Ustilago hordei* causing covered smut of barley. *Frontiers in
Microbiology, 10*, 2929. https://doi.org/10.3389/fmicb.2019.02929

Lackner, M., Obermair, J., Naschberger, V., Raschbichler, L. M., Kandelbauer, C., Pallua, J.,
... Binder, U. (2019). Cryptic species of *Aspergillus* section *Terrei* display essential
physiological features to cause infection and are similar in their virulence potential in
Galleria mellonella. Virulence, 10(1), 542–554.
https://doi.org/10.1080/21505594.2019.1614382

Latka, C., Dey, S. S., Mahajan, S., Prabu, R., Jangir, P. K., Gupta, C., … Taneja, B. (2015).
Genome sequence of a clinical isolate of dermatophyte, *Trichophyton rubrum* from India.
FEMS Microbiology Letters, 362(8), fnv039. https://doi.org/10.1093/femsle/fnv039

Leliaert, F., Verbruggen, H., Vanormelingen, P., Steen, F., López-Bautista, J. M., Zuccarello,
G. C. & De Clerck, O. (2014). DNA-based species delimitation in algae. *European Journal of
Phycology, 49*(2), 179–196. http://dx.doi.org/10.1080/09670262.2014.904524

Li, H. C., Bouchara, J. P., Hsu, M. M., Barton, R., Su, S., & Chang, T. C. (2008).
Identification of dermatophytes by sequence analysis of the rRNA gene internal transcribed

spacer regions. *Journal of Medical Microbiology, 57*(5), 592–600.
https://doi.org/10.1099/jmm.0.47607-0

Li, W., Metin, B., White, T. C., & Heitman, J. (2010). Organization and evolutionary
trajectory of the mating type (MAT) locus in dermatophyte and dimorphic fungal pathogens.
Eukaryotic Cell, 9(1), 46–58. https://doi.org/10.1128/EC.00259-09

Martinez, D. A., Oliver, B. G., Gräser, Y., Goldberg, J. M., Li, W., Martinez-Rossi, N. M., …
White, T. C. (2012). Comparative genome analysis of *Trichophyton rubrum* and related
dermatophytes reveals candidate genes involved in infection. *MBio, 3*(5), e00259–12.
https://doi.org/10.1128/mBio.00259–12

Martins, M. P., Rossi, A., Sanches, P. R., Bortolossi, J. C., & Martinez-Rossi, N. M. (2020).
Comprehensive analysis of the dermatophyte *Trichophyton rubrum* transcriptional profile
reveals dynamic metabolic modulation. *Biochemical Journal,* pii: BCJ20190868.
https://doi.org/10.1042/BCJ20190868

Mironenko, N. V., Baranova, O. A., Kovalenko, N. M., Mikhailova, L. A., & Rosseva, L. P.
(2016). Genetic structure of the Russian populations of *Pyrenophora tritici-repentis*,
determined by using microsatellite markers. *Russian Journal of Genetics, 52*(8), 771–779.
https://doi.org/10.1134/S1022795416080093

Nelson, D. L. & Cox, M. M. (2008). Lehninger principles of biochemistry. WH Freeman.
New York.

Nurk, S., Bankevich, A., Antipov, D., Gurevich, A., Korobeynikov, A., Lapidus, A., ...
Stepanauskas, R. (2013). Assembling genomes and mini-metagenomes from highly chimeric
reads. In M. Deng, R. Jiang, F. Sun, & X. Zhang (Eds.), *Annual International Conference on
Research in Computational Molecular Biology* (pp. 158-170). Springer, Berlin, Heidelberg.
https://doi.org/10.1007/978-3-642-37195-0_13

Packeu, A., Stubbe, D., Roesems, S., Goens, K., Van Rooij, P., de Hoog, S., & Hendrickx, M.
(2019). Lineages within the *Trichophyton rubrum* complex. *Mycopathologia.*
https://doi.org/10.1007/s11046-019-00386-z

Pchelin, I. M., Azarov, D. V., Chilina, G. A., Dmitriev, K. A., Vasilyeva, N. V., & Taraskina,
A. E. (2018a). Single-nucleotide polymorphism in a local population of *Trichophyton rubrum*.
Medical Mycology, 56(1), 125–128. https://doi.org/10.1093/mmy/myx009

Pchelin, I. M., Azarov, D. V., Churina, M. A., Ryabinin, I. A., Vibornova, I. V., Apalko, S.
V., … Vasilyeva, N. V. (2019a). Whole genome sequence of first *Candida auris* strain,
isolated in Russia. *Medical Mycology*. https://doi.org/10.1093/mmy/myz078

Pchelin, I. M., Azarov, D. V., Churina, M. A., Scherbak, S. G., Apalko, S. V., Vasilyeva, N.
V., & Taraskina, A. E. (2019b). Species boundaries in the *Trichophyton mentagrophytes / T.
interdigitale* species complex. *Medical Mycology, 57*(6), 781–789.
https://doi.org/10.1093/mmy/myy115

Pchelin, I. M., Kryuchkova, M. A., Bogdanova, T. V., Chilina, G. A., Boronina, L. G., Olina
E. S., … Taraskina A. E. (2018b). We need more powerful microsatellite assay for population
genetic studies of *Trichophyton rubrum*. *Medical Mycology, 56*(S2), S119.

Peakall, R. & Smouse P. E. (2006). GENALEX 6: genetic analysis in Excel. Population
genetic software for teaching and research. *Molecular Ecology Notes, 6*(1), 288–295.
https://doi.org/10.1111/j.1471-8286.2005.01155.x

Peakall, R. & Smouse, P. E. (2012). GenAlEx 6.5: genetic analysis in Excel. Population
genetic software for teaching and research – an update. *Bioinformatics, 28*(19), 2537–2539.
https://doi.org/10.1093/bioinformatics/bts460

Persinoti, G. F., Martinez, D. A., Li, W., Döğen, A., Billmyre, R. B., Averette, A., … Cuomo,
C. A. (2018). Whole-genome analysis illustrates global clonal population structure of the
ubiquitous dermatophyte pathogen *Trichophyton rubrum*. *Genetics, 208*(4), 1657–1669.
https://doi.org/10.1534/genetics.117.300573

Price, M. N., Dehal, P. S., & Arkin, A. P. (2010). FastTree 2–approximately maximum-
likelihood trees for large alignments. *PLoS One, 5*, e9490.
https://doi.org/10.1371/journal.pone.0009490

Rad, M. M., Mohammadi, A. M. A., & Barton, R. C. (2008). PCR typing of *Trichophyton*
rubrum isolates by specific amplification of subrepeat elements in ribosomal DNA
nontranscribed spacer. *Iranian Journal of Dermatology, 11*(1), 17–20.

Ramaraj, V., Vijayaraman, R. S., Elavarashi, E., Rangarajan, S., & Kindo, A. J. (2017).
Molecular strain typing of clinical isolates, *Trichophyton rubrum* using non transcribed spacer
(NTS) region as a molecular marker. *Journal of Clinical and Diagnostic Research, 11*(5),
DC04–DC09. https://doi.org/10.7860/jcdr/2017/21994.9843

Rozenfeld, A. F., Arnaud-Haond, S., Hernández-García, E., Eguíluz, V. M., Matías, M. A.,
Serrão, E., & Duarte, C. M. (2007). Spectrum of genetic diversity and networks of clonal
organisms. *Journal of the Royal Society Interface, 4*(17), 1093–1102.
https://doi.org/10.1098/rsif.2007.0230

Sexton, A. C., & Howlett, B. J. (2004). Microsatellite markers reveal genetic differentiation
among populations of *Sclerotinia sclerotiorum* from Australian canola fields. *Current*
Genetics, 46(6), 357–365. https://doi.org/10.1007/s00294-004-0543-3

Sigurgeirsson, B., & Baran, R. (2014). The prevalence of onychomycosis in the global
population: a literature study. *Journal of the European Academy of Dermatology and*
Venereology, 28(11), 1480–1491. https://doi.org/10.1111/jdv.12323

Singh, N., Anand, G., & Kapoor, R. (2019). Virulence and genetic diversity among *Fusarium*
oxysporum f. sp. *carthami* isolates of India using multilocus RAPD and ISSR markers.
Tropical Plant Pathology, 44(5), 409–422. https://doi.org/10.1007/s40858-019-00303-1

Su, H., Packeu, A., Ahmed, S. A., Al-Hatmi, A. M. S., Blechert, O., İlkit, M., … de Hoog, S.
(2019). Species distinction in the *Trichophyton rubrum* complex. *Journal of Clinical*
Microbiology, 57(9), e00352-19. https://doi.org/10.1128/jcm.00352-19

Summerbell, R. C. (2000). Form and function in the evolution of dermatophytes. In:
Kushwaha RKS, Guarro J, editors. *Biology of dermatophytes and other keratinophilic fungi.*
*Revista Iberoamericana de Micología.*17 Suppl: 30–43.

Summerbell, R. C., Haugland, R. A., Li, A., & Gupta, A. K. (1999). rRNA gene internal
transcribed spacer 1 and 2 sequences of asexual, anthropophilic dermatophytes related to
Trichophyton rubrum. Journal of Clinical Microbiology, 37(12), 4005–4011.

Suzuki, S., Mano, Y., Furuya, N., & Fujitani, K. (2018). Molecular epidemiological analysis
of the spreading conditions of *Trichophyton* in long-term care facilities in Japan. *Japanese*
Journal of Infectious Diseases, 71(6), 462–466. https://doi.org/10.7883/yoken.jjid.2018.090

Taghipour, S., Pchelin, I. M., Zarei Mahmoudabadi, A., Ansari, S., Katiraee, F., Rafiei, A., …
Rezaei-Matehkolaei, A. (2019). *Trichophyton mentagrophytes* and *T. interdigitale* genotypes
are associated with particular geographic areas and clinical manifestations. *Mycoses, 62*(11),
1084–1091. https://doi.org/10.1111/myc.12993

Takahashi, I., Fukushima, K., Miyaji, M., Nishimura, K., Asano, K., & Iizuka, H. (2014).
Species identification and strain typing of dermatophytes by single-strand conformation
polymorphism (SSCP) analysis of the ribosomal DNA and polymerase chain reaction analysis
of subrepeat elements in the intergenic spacer region of *Trichophyton rubrum. Asahikawa*
Medical University Repository, 15, 27–36.

Tamura, K., Stecher, G., Peterson, D., Filipski, A., & Kumar, S. (2013). MEGA6: molecular
evolutionary genetics analysis version 6.0. *Molecular Biology and Evolution, 30*(12), 2725–
2729. https://doi.org/10.1093/molbev/mst197

Tibayrenc, M. (2006). The species concept in parasites and other pathogens: a pragmatic
approach? *Trends in Parasitology, 22*(2), 66–70. https://doi.org/10.1016/j.pt.2005.12.010

Treangen, T. J., Ondov, B. D., Koren, S., & Phillippy, A. M. (2014). The Harvest suite for
rapid core-genome alignment and visualization of thousands of intraspecific microbial
genomes. *Genome Biology, 15*(11), 524. https://doi.org/10.1186/s13059-014-0524-x

White, T. J., Bruns, T., Lee, S., & Taylor, J. W. (1990). Amplification and direct sequencing
of fungal ribosomal RNA genes for phylogenetics. In M. Innis, D. Gelfand, J. Sninsky, & T.
White (Eds.), *PCR protocols: A guide to methods and applications* (pp. 315–322). Orlando,
FL: Academic Press.

Ye, J., Coulouris, G., Zaretskaya, I., Cutcutache, I., Rozen, S., & Madden, T. (2012). Primer-
BLAST: A tool to design target-specific primers for polymerase chain reaction. *BMC*
Bioinformatics, 13, 134. https://doi.org/10.1186/1471-2105-13-134

Zhan, P., Dukik, K., Li, D., Sun, J., Stielow, J. B., Gerrits van den Ende, B., … de Hoog, G.
S. (2018). Phylogeny of dermatophytes with genomic character evaluation of clinically
distinct *Trichophyton rubrum* and *T. violaceum. Studies in Mycology, 89*, 153–175.
https://doi.org/10.1016/j.simyco.2018.02.004

Zhang, Z., Schwartz, S., Wagner, L., & Miller, W. (2000). A greedy algorithm for aligning
DNA sequences. *Journal of Computational Biology, 7*(1-2): 203–214.
https://doi.org/10.1089/10665270050081478

Zheng, H., Blechert, O., Mei, H., Ge, L., Liu, J., Tao, Y., … Liu, W. (2019). Whole-genome
resequencing of *Trichophyton rubrum* provides insights into population differentiation and
drug resistance. *Mycopathologia*. https://doi.org/10.1007/s11046-019-00384-1

Supplementary data

706   Figures S1 and S2.

707   Supplement S1. Molecular strain typing in 70 *T. rubrum* isolates.

Figures

Figure 1. A phylogenetic network, calculated from the matrix of mutation probability-based
genetic distances (Bruvo et al. 2004), 12 microsatellite loci. Urease-positive isolates are
marked by asterisks.

Figure 2. A phylogenetic network, calculated from the matrix of distances, based on the
lengths of microsatellite loci (Rozenfeld et al. 2007), 12 microsatellite loci. Urease-positive
isolates are marked by asterisks.

Figure 3. A phylogenetic network, calculated from the matrix of mutation probability-based
genetic distances (Bruvo et al. 2004), 11 microsatellite loci. Urease-positive isolates are
marked by asterisks.

Figure 4. A phylogenetic network, based on Rozenfeld genetic distances, 11 microsatellite
loci. Urease-positive isolates are marked by asterisks.

Figure 5. Phylogenomic tree of *T. rubrum* isolates. ST1 and ST2 lineages are defined by non-
synonymous substitutions in TERG_02941 and TERG_03298 protein-coding loci. Russian
population shares both clades with West European and Asian countries. Single and double
arrowheads mark Chinese isolates, fallen into two different clusters in the original publication
by Zheng et al. (2019). Asterisks mark isolates with urease activity. Branch support values $\geq$
75% are marked by open circles and values $\geq$ 95% are marked by solid circles. Newly
sequenced isolates are given in bold. The tree is rooted by *T. soudanense* CBS 452.61 whole
genome sequence (the branch isn't shown).

Figure 6. *Trichophyton* spp. A tree, based on concatenated sequences of TERG_02718, TERG_02913, TERG_02941, TERG_03298, TERG_03581, TERG_03748 and TERG_03778 loci. In the *T. mentagrophytes* / *T. interdigitale* species group, ITS region genotypes are indicated (Taghipour et al. 2019). Asterisks mark isolates with urease activity. Branch support values ≥ 75% are marked by open circles and values ≥ 95% are marked by solid circles. Two clades are shown separately at different scales.

Figure 7. Hypothetical distribution of *Trichophyton rubrum* genetic lines. A total of 443 isolates, from 17 to 70 isolates for a country. The abbreviations above pie charts indicate the sources of variability: SNP, single nucleotide polymorphism of TERG_02941 and TERG_03298 genes; TRS-1, the length of TRS-1 locus of ribosomal non-transcribed spacer; WGS, whole genome sequence.

Figure S1. Nucleotide polymorphism of *Trichophyton rubrum* ITS region sequences. *A*, 247insC mutation in D15P139 isolate (asterisk). *B*, an additional TA repeat near the 5'-end of the sequence in RCPF 1280 strain, 595insTA (double asterisk).

Figure S2. STR*Tr*9 microsatellite locus length and TERG_6744 gene expression in *T. rubrum* genetic lineages. *A*, Average length of STR*Tr*9 amplification products differs between ST1 and ST2 isolates. *B*, Relative expression of TERG_6744 gene, coding for fructose 1,6-bisphosphate aldolase, varies between strains irrespectively of their genetic lineage affiliation. From the left to the right: D15P44 and D15P71 (ST1), D15P46 and D15P50 ("ancestral" ST2), D15P33 and D15P62 ("derived" ST2). There is no immediate consequence of STR*Tr*9 locus length for TERG_06744 transcript level.

0.05

D15P33

Y927 *

D15P51

Y1013

RCPF 1408

D15P71

D15P62

D15P139

Y2

RCPF 1280 *

D15P134

D15P44

Y1932

CBS 392.58

D15P46

D15P50 *

TERG_02941 793A 793G

0.5

D15P71

Y1932

D15P44

D15P62

D15P139

Y2

Y927 *

D15P33

D15P46

RCPF 1280 *

Y1013

CBS 392.58

D15P134

D15P51

RCPF 1408

D15P50 *

TERG_02941 793A 793G

Hypothetical lineage affiliation: ST1 ST2