A fast and efficient smoothing approach to Lasso regression and an application in statistical genetics: polygenic risk scores for chronic obstructive pulmonary disease (COPD)

Georg Hahn, Sharon M. Lutz, Nilanjana Laha, Michael H. Cho,
Edwin K. Silverman, and Christoph Lange

Abstract

High dimensional linear regression problems are often fitted using Lasso approaches. Although the Lasso objective function is convex, it is not differentiable everywhere, making the use of gradient descent methods for minimization not straightforward. To avoid this technical issue, we apply Nesterov smoothing to the original (unsmoothed) Lasso objective function. We introduce a closed-form smoothed Lasso which preserves the convexity of the Lasso function, is uniformly close to the unsmoothed Lasso, and allows us to obtain closed-form derivatives everywhere for efficient and fast minimization via gradient descent. Our simulation studies are focused on polygenic risk scores using genetic data from a genome-wide association study (GWAS) for chronic obstructive pulmonary disease (COPD). We compare accuracy and runtime of our approach to the current gold standard in the literature, the FISTA algorithm. Our results suggest that the proposed methodology provides estimates with equal or higher accuracy than the the FISTA algorithm while having the same asymptotic runtime scaling.

Keywords: COPD; FISTA; LASSO; Nesterov; Penalized linear regression; Polygenic risk scores; Smoothing.

1   Introduction

Many substantive research questions in health, economic and social science require solving a classical linear regression problem $X\beta = y$. Here, the data matrix $X \in \mathbb{R}^{n \times p}$, the parameter vector $\beta \in \mathbb{R}^p$, and the response vector $y \in \mathbb{R}^n$ encode $n \in \mathbb{N}$ linear equations in $p \in \mathbb{N}$ variables. The approach remains one of the most widely used statistical analysis tools.

Traditionally, linear regression problems have been solved by finding parameter estimates β that minimize the squared error, leading to the least squares estimate $\arg\min_\beta \|X\beta - y\|_2^2$. However, their lack of robustness, as well as sparsity requirements in high dimensional settings with $p \gg n$, are two problems that have led to the development of alternative estimation approaches, e.g. the *least absolute shrinkage and selection operator* (Lasso) of Tibshirani (1996) or *least-angle regression* (LARS) of Efron et al. (2004).

This article focuses on Lasso regression, which obtains regression estimates $\hat{\beta}$ by solving

$$\hat{\beta} = \arg\min_\beta \frac{1}{n} \|X\beta - y\|_2^2 + \lambda \|\beta\|_1, \tag{1}$$

where $\|\cdot\|_2$ is the Euclidean norm, $\|\cdot\|_1$ is the L_1 norm, and $\lambda > 0$ is a tuning parameter (called the Lasso regularization parameter) controlling the sparseness of the solution $\hat{\beta}$.

As the objective function in eq. (1) is convex, minimization via steepest descent (quasi-Newton) methods is sensible. However, many applications in biostatistics, especially those that are focused on "big data", such as the simultaneous analysis of genome-wide association studies (Wu et al., 2009) or the calculation of polygenic risk scores (Mak et al., 2016), involve data sets with several thousand parameters and are often sparse. In such applications, conventional gradient-free solvers can lose accuracy. This is due to the non-differentiability of the L_1 penalty term in eq. (1), i.e. the term $\|\beta\|_1 = \sum_{i=1}^p |\beta_i|$.

We address this issue by smoothing the Lasso objective function. We apply Nesterov smoothing (Nesterov, 2005) to the non-differentiable $\|\beta\|_1$ penalty term in eq. (1). This will result in an approximation of the Lasso objective function that is differentiable everywhere. The Nesterov formalism depends on a smoothing parameter μ that controls the smoothness level.

Our approach has three major advantages: (1) The smoothing preserves the convexity of the Lasso objective function; (2) it allows us to obtain closed-form derivatives of the smoothed function everywhere which we use in a gradient descent algorithm; and (3) it provides uniform error bounds on the difference between the smoothed and the original objective functions. The error bounds depend only on the smoothing parameter μ, and by letting $\mu \to 0$, the smoothed Lasso objective function can be made arbitrarily close to the original Lasso objective function, thus justifying its use as a surrogate of the original Lasso objective.

We benchmark our proposed smoothing approach against the current gold standard for minimizing the Lasso objective function, the FISTA algorithm of Beck and Teboulle (2009). FISTA is a proximal gradient version of the algorithm of Nesterov (1983) which combines the basic iterations of the Iterative Shrinkage-Thresholding Algorithm (Daubechies et al., 2004) with a Nesterov acceleration step. Among others, the algorithm is implemented in the R-package *fasta* on CRAN (Chi et al., 2018). We use this package as a benchmark. The FISTA algorithm requires the separate specification of the smooth and non-smooth parts of the objective function including their explicit gradients. In contrast to our approach, a variety of tuning parameters need to be selected by the user, e.g. an initial starting value, an initial stepsize, parameters determining the lookback window for non-monotone line search, and a shrinkage parameter for the stepsize.

This article is structured as follows. Section 2 applies Nesterov smoothing to the Lasso objective function. Using polygenic risk scores as an example, we evaluate the proposed methodology in a simulation study in Section 3. The article concludes with a discussion in Section 4. The proof of our theoretical result is given in Appendix 5.

Throughout the article, $X_{\cdot,i}$ denotes the ith column of a matrix X. Similarly, $X_{I,\cdot}$ (and y_I) denote the submatrix (subvector) consisting of all rows of X (entries of y) indexed in the set I. Moreover, $X_{-I,\cdot}$ (and y_{-I}) denote the submatrix (subvector) consisting of all rows of X (entries of y) not indexed in the set I. Finally, $|\cdot|$ denotes the absolute value, and $\|\cdot\|_\infty$ denotes the supremum norm.

2 Smoothing the Lasso objective function

This section lays the theoretical foundation for the modified Lasso approach we propose to address the non-differentiability of the L_1 penality term in eq. (1), while the smooth L_2 term remains unchanged. The substitute of the term $\|\beta\|_1 = \sum_{i=1}^p |\beta_i|$ in eq. (1) will be computed with the help of a technique called Nesterov smoothing, whose details are given in Section 2.1. In Section 2.2, we simplify the general Nesterov smoothing approach for the particular case of the Lasso penalty and show how our approach results in explicit closed-form expressions for both the smoothed Lasso and its gradient. The uniform closeness of the smoothed Lasso we propose and the original Lasso objective function is proven in Section 2.3.

2.1 Summary of Nesterov smoothing

We briefly summarize the idea of Nesterov smoothing. We are given a piecewise affine and convex function $f : \mathbb{R}^q \to \mathbb{R}$ which we aim to smooth, where $q \in \mathbb{N}$. We assume that f is composed of $k \in \mathbb{N}$ linear pieces (components) and can thus be expressed as

$$f(z) = \max_{i=1,\ldots,k} \left(A[z,1]^\top \right)_i,$$

where $A \in \mathbb{R}^{k \times (q+1)}$ is a matrix whose rows contain the linear coefficients for each of the k linear pieces (with the constant coefficients being in column $q + 1$), $z \in \mathbb{R}^q$, and $[z, 1] \in \mathbb{R}^{q+1}$ denotes the vector obtained by concatenating z and the scalar 1.

Let $\mu \geq 0$ be the Nesterov smoothing parameter. Using a so-called proximity (or prox) function ρ, which is assumed to be nonnegative, continuously differentiable, and strongly convex, f is replaced by the approximation

$$f^\mu(z) = \max_{w \in Q_k} \left\{ \langle A[z,1]^\top, w \rangle - \mu \rho(w) \right\},$$

where $Q_k \subseteq \mathbb{R}^k$ is the unit simplex in k dimensions defined as

$$Q_k = \left\{ w \in \mathbb{R}^k : \sum_{i=1}^k w_i = 1, w_i \geq 0 \ \forall i = 1, \ldots, k \right\}.$$

As shown in (Nesterov, 2005, Theorem 1), f^μ is both smooth and uniformly close to f, meaning that for all $z \in \mathbb{R}^q$, the approximation error is uniformly upper bounded by

$$\sup_{z \in \mathbb{R}^q} |f(z) - f^\mu(z)| \leq \mu \sup_{w \in Q_k} \rho(w) = O(\mu). \tag{2}$$

The larger the value of μ the more f is smoothed, while $\mu = 0$ recovers the original function $f = f^0$. Two choices of the prox function are given a closer look in Section 2.2. Smoothing with the so-called entropy prox function results in the smooth approximation f_e^μ having a closed-form

expression given by

$$f_e^\mu(z) = \mu \log \left(\frac{1}{k} \sum_{i=1}^k e^{\frac{\left(A[z,1]^\top\right)_i}{\mu}} \right), \tag{3}$$

which satisfies the uniform bound

$$\sup_{z \in \mathbb{R}^q} |f(z) - f_e^\mu(z)| \leq \mu \log(k). \tag{4}$$

Similarly, the smooth approximation of f with the help of the so-called squared error prox function given by $\rho_s(w) = \frac{1}{2} \sum_{i=1}^k \left(w_i - \frac{1}{k}\right)^2$ for $w \in \mathbb{R}^k$ can be written as

$$f_s^\mu(z) = \langle \hat{c}(z), A[z,1]^\top \rangle - \mu \rho_s(\hat{c}(z)). \tag{5}$$

In eq. (5), $\hat{c}(z) \in \mathbb{R}^k$ is the Michelot projection (Michelot, 1986) of the vector $c(z) = (c_1(z), \ldots, c_k(z))$, given componentwise by $c_i(z) = 1/\mu \cdot \left(A[z,1]^\top\right)_i - 1/k$ for $i \in \{1, \ldots, k\}$, onto the k-dimensional unit simplex Q_k. The approximation f_s^μ via squared error prox function satisfies the uniform bound

$$\sup_{z \in \mathbb{R}^q} |f(z) - f_s^\mu(z)| \leq \mu \left(1 - \frac{1}{k}\right). \tag{6}$$

2.2 Application to the Lasso objective function

For given $X \in \mathbb{R}^{n \times p}$, $y \in \mathbb{R}^n$, and $\lambda > 0$, according to eq. (1), the Lasso objective function $L : \mathbb{R}^p \to \mathbb{R}$ given by

$$L(\beta) = \frac{1}{n} \|X\beta - y\|_2^2 + \lambda \|\beta\|_1 \tag{7}$$

is smooth in its first term but non-differentiable in the penalty term $\|\beta\|_1 = \sum_{i=1}^p |\beta_i|$. We thus smooth the latter, where it suffices to apply Nesterov smoothing to each absolute value independently.

Let $k = 2$. Using one specific choice of the matrix $A \in \mathbb{R}^{2 \times 2}$ given by

$$A = \begin{pmatrix} -1 & 0 \\ 1 & 0 \end{pmatrix},$$

we rewrite the (one dimensional) absolute value as $f(z) = \max\{-z, z\} = \max_{i=1,2} \left(A[z, 1]^\top \right)_i$, where here and in the following subsections $z \in \mathbb{R}$ is a scalar.

2.2.1 Entropy prox function

For the entropy prox function, eq. (3) with A as in Section 2.2 simplifies to

$$f_e^\mu(z) = \mu \log \left(\frac{1}{2} e^{-z/\mu} + \frac{1}{2} e^{z/\mu} \right),$$

which according to eq. (4) satisfies the approximation bound

$$\sup_{z \in \mathbb{R}} |f(z) - f_e^\mu(z)| \leq \mu \log(2). \tag{8}$$

The first and second derivatives of f_e^μ are given by

$$\frac{\partial}{\partial z} f_e^\mu(z) = \frac{-e^{-z/\mu} + e^{z/\mu}}{e^{-z/\mu} + e^{z/\mu}} =: g_e^\mu(z),$$

$$\frac{\partial^2}{\partial z^2} f_e^\mu(z) = \frac{4}{\mu(e^{-z/\mu} + e^{z/\mu})^2} =: h_e^\mu(z).$$

Together, smoothing eq. (7) with the entropy prox function results in

$$L_e^\mu(\beta) = \frac{1}{n} \|X\beta - y\|_2^2 + \lambda \sum_{i=1}^p f_e^\mu(\beta_i),$$

$$\frac{\partial L_e^\mu}{\partial \beta_i}(\beta) = -\frac{2}{n} \langle y - X\beta, X_{\cdot,i} \rangle + \lambda g_e^\mu(\beta_i), \tag{9}$$

$$\frac{\partial^2 L_e^\mu}{\partial \beta_i \partial \beta_j}(\beta) = \frac{2}{n} (X^\top X)_{ij} + \mathbb{I}(i = j) \cdot \lambda h_e^\mu(\beta_i),$$

where the entropy prox smoothed Lasso is L_e^μ, its explicit gradient is $\partial L_e^\mu/\partial \beta_i$, and its Hessian matrix is given by $\partial^2 L_e^\mu/\partial \beta_i \partial \beta_j$. The function $\mathbb{I}(\cdot)$ denotes the indicator function.

In principle, the Lasso objective can be minimized using a (second order) Newton-Raphson or

a (first order) quasi-Newton approach. However, since $X \in \mathbb{R}^{n \times p}$ with $n < p$, the matrix $X^\top X$ is singular, meaning that for the Hessian to be invertible one needs the added diagonal elements $\lambda h_e^\mu(\beta_i)$ to be large. However, this is usually not true, since if β_i is nonzero, then in a neighborhood of the true Lasso estimate the term $(e^{-\beta_i/\mu} + e^{\beta_i/\mu})^{-2}$ will be exponentially small. Thus to make $\lambda h_e^\mu(\beta_i)$ large for a fixed μ, we need λ to be exponentially large. Likewise, given λ and β_i, too small or too large values of μ will make h_e^μ vanish. However, since typically λ and μ are fixed, the Hessian in eq. (9) will be singular except for a few artificial cases and thus the second order Newton-Raphson method will not be applicable. In the simulations we therefore focus on quasi-Newton methods which require only L_e^μ and its gradient $\partial L_e^\mu / \partial \beta_i$.

2.2.2 Squared error prox function

Similarly, eq. (5) with A as in Section 2.2 simplifies to

$$f_s^\mu(z) = \langle \hat{c}(z), [-z, z] \rangle - \mu \rho_s(\hat{c}(z)).$$

where $\hat{c}(z) \in \mathbb{R}^2$ is the Michelot projection of the vector $c(z) = 1/\mu \cdot [-z, z] - 1/k$ onto the two-dimensional unit simplex Q_2. According to eq. (6), we obtain the approximation bound

$$\sup_{z \in \mathbb{R}} |f(z) - f_s^\mu(z)| \leq \frac{1}{2}\mu. \tag{10}$$

The derivative of f_s^μ is given by

$$\frac{\partial}{\partial z} f_s^\mu(z) = \langle \hat{c}(z), [-1, 1] \rangle =: g_s^\mu(z),$$

see (Hahn et al., 2017, Lemma 4) for a proof of this result. The second derivative $\partial^2 f_s^\mu / \partial z^2$ does not have a closed form expression, though it can be approximated numerically. Analogously to the results for the entropy prox function, smoothing eq. (7) with the squared error prox function

results in

$$L_s^\mu(\beta) = \frac{1}{n}\|X\beta - y\|_2^2 + \lambda \sum_{i=1}^p f_s^\mu(\beta_i),$$

$$\frac{\partial L_s^\mu}{\partial \beta_i}(\beta) = -\frac{2}{n}\langle y - X\beta, X_{\cdot,i}\rangle + \lambda g_s^\mu(\beta_i), \tag{11}$$

$$\frac{\partial^2 L_s^\mu}{\partial \beta_i \partial \beta_j}(\beta) = \frac{2}{n}(X^\top X)_{ij} + \mathbb{I}(i=j)\cdot\lambda \left.\frac{\partial^2 f_s^\mu}{\partial z^2}\right|_{z=\beta_i}$$

where as before, the smoothed Lasso via squared error prox function is L_s^μ, its explicit gradient is $\partial L_s^\mu/\partial \beta_i$, and its Hessian matrix is $\partial^2 L_s^\mu/\partial\beta_i\partial\beta_j$.

As in Section 2.2.1 we observe that the Hessian matrix is singular since $X^\top X$ is singular, and since the additional diagonal entries stemming from $\lambda\partial^2 f_s^\mu/\partial z^2$ are usually too small to make the Hessian invertible. We therefore again resort to a quasi-Newton method to minimize L_s^μ with the help of its gradient $\partial L_s^\mu/\partial\beta_i$ only.

2.3 Theoretical error bounds

The approximation bounds on the absolute value, given in eqs. (8) and (10), carry over to bounds on the distance between the unsmoothed and smoothed Lasso objective functions, as shown in the following proposition.

Proposition 1. *The smoothed Lasso objective functions with entropy and squared error prox, given in eqs. (9) and (11), satisfy the following uniform bounds:*

$$\sup_{\beta\in\mathbb{R}^p} |L_e^\mu(\beta) - L(\beta)| \le \lambda p\mu\log(2),$$

$$\sup_{\beta\in\mathbb{R}^p} |L_s^\mu(\beta) - L(\beta)| \le \frac{\lambda p\mu}{2}.$$

Moreover, both L_e^μ and L_s^μ are strictly convex.

In Proposition 1 the Lasso parameter $\lambda > 0$ and the dimension p are fixed for a particular estimation problem, thus allowing to make the approximation error arbitrarily small as the smoothing parameter $\mu \to 0$. This justifies the use of our proposed smoothed Lasso as a surrogate of the original (non smooth) Lasso objective function. Moreover, the convexity of the Lasso is preserved.

3 Simulation studies for polygenic risk scores for COPD

In this section, we evaluate three approaches to compute Lasso estimates on simulated data (Section 3.1) as well as real data coming from a genome-wide association study for COPDGENE (Regan et al., 2010), in which polygenic risk scores are computed and evaluated (Section 3.2). Of the three approaches we consider, the first two utilize existing methodology, while the last approach implements the methodology we developed in this paper:

1. We carry out the minimization of eq. (1) using R's function *optim*. The *optim* function implements the quasi-Newton method *BFGS* for which we supply the explicit (though non-smooth) gradient $\partial L/\partial \beta_i = -2/n \cdot \langle y - X\beta, X_{\cdot,i}\rangle + \lambda \text{sign}(\beta_i)$. This approach will be referred to as the *unsmoothed Lasso*;

2. We use the FISTA algorithm as implemented in the *fasta* R-package (Chi et al., 2018), available on *The Comprehensive R Archive Network* (R Core Team, 2014);

3. We minimize the smoothed Lasso objective function of eq. (9) using its explicit gradient.

The main function of the *fasta* package which implements the FISTA algorithm, also called *fasta*, requires the separate specification of the smooth and non-smooth parts of the objective function including their explicit gradients. We follow Example 2 in the *vignette* of the *fasta* R-package in Chi et al. (2018) and supply both as specified in eq. (7). Additionally, we employ a uniform random starting value as done for the unsmoothed and smoothed Lasso. The initial stepsize is set to $\tau = 10$ as in Example 2 of Chi et al. (2018). The lookback window for non-monotone line search and the shrinkage parameter for the stepsize are left at their default values.

Three potential implementations are unconsidered in this simulation section for the following reasons. The *glmnet* algorithm of Friedman et al. (2010), available in the R-package *glmnet*, is a variant of FISTA which performs a cyclic update of all coordinates, whereas FISTA updates all coordinates per iteration. We thus focus on FISTA. Since the R-package *SIS* accompanying Fan and Li (2001) does itself rely on *glmnet* for computing regression estimates, we omit it in this section. The LARS algorithm of Efron et al. (2004) is implemented in the R-package *lars* on CRAN (Hastie and Efron, 2013). As remarked in Friedman et al. (2010), LARS is slower than glmnet/

Figure 1: L_2 norm of parameter estimate to truth (left) and runtime in seconds (right) as a function of $n \in [1, 10000]$ while $p = 1000$. Logarithmic scale on the y-axes.

FISTA. Additionally, since the implementation of Hastie and Efron (2013) always computes a full Lasso path, it is considerably slower than the other methods.

All results are averages over 100 repetitions. The choice of the Lasso regularization parameter λ varies in each experiment and is given individually.

3.1 Application to simulated data

We designed our simulation study so that it mimics the application to a genome-wide association study, i.e. we simulate $X \in \mathbb{R}^{n \times p}$ from a multidimensional normal distribution, where the entries of the mean vector of the multidimensional normal distribution are sampled independently from a uniform distribution in $[0, 0.5]$. To ensure positive definiteness of the covariance matrix Σ of the multidimensional normal distribution, we set $\Sigma = \frac{1}{2}(A + A^\top) + nD_n$, where D_n is a $n \times n$ matrix with ones on its diagonal, and zeros otherwise. The added term nD_n ensures positive definiteness.

After X is obtained, we generate the entries of the true β independently from a standard normal distribution, and set all but $nz \in \{0, \ldots, p\}$ out of the p entries to zero. The number of non-zero entries $nz \in \mathbb{N}$ is a parameter of the simulations. The response $y \in \mathbb{R}^n$ is then easily obtained as $y = X\beta + \epsilon$, where the entries of the noise vector $\epsilon \in \mathbb{R}^n$ are generated independently from a

Normal distribution with mean zero and some variance σ^2. The smaller the variance, the easier the recovery of β will be. We will employ $\sigma^2 = 0.1$ in our simulations. In this subsection, we fix the number of true non-zero parameters at 20% (that is, $nz = 0.2p$), resulting in sparse parameter vectors.

The regularization parameter of the Lasso was chosen as $\lambda = 1$ for the experiments in this subsection.

Figure 1 (left) shows results on simulated data of dimension $n \in [1, 10000]$ while keeping $p = 1000$ fixed. We measure the accuracy of the obtained Lasso estimates through their L_2 norm to the generated true parameters. We observe that the unsmoothed and smoothed Lasso approaches seem to suffer from numerical instabilities for small n. As n increases, both the unsmoothed and smoothed Lasso approaches stabilize. The FISTA algorithm yields stable estimates for all n. The smoothed Lasso approach and FISTA roughly draw equal in accuracy for large n. As expected, all methods become more accurate as the number of data points n increases.

Figure 1 (right) shows that the unsmoothed and smoothed Lasso approaches have an almost identical runtime. We find that although highly optimized, FISTA is only a low multiple factor faster than our approaches and, more importantly, its asymptotic runtime scaling is comparable to the one of the smoothed Lasso.

Similarly to the previous experiment, in Figure 2 (left) we keep $n = 1000$ fixed and investigate the dependence of all four approaches on $p \in [1, 5000]$. Again, the unsmoothed and smoothed Lasso approaches seem to suffer from numerical instabilities for smaller p, while FISTA is very stable. Importantly, the smoothed Lasso seems to outperform FISTA for very large p. As expected, while keeping the data size n fixed, estimation becomes more challenging for all methods as p increases.

Figure 2 (right) confirms the timing results seen in the assessment of the dependence on n. The unsmoothed and smoothed approaches have virtually equal speeds, while FISTA is again slightly faster. Notably, the scaling of the runtimes seems to be roughly equal for all four methods.

Since the unsmoothed Lasso does not come with the guarantee we established for our smoothing approach (see Section 2.3) and is outperformed by the smoothed Lasso, we will focus in the remainder of the simulations on FISTA and smoothed Lasso only.

Figure 2: L_2 norm of parameter estimate to truth (left) and runtime in seconds (right) as a function of $p \in [1, 5000]$ while $n = 1000$. Logarithmic scale on the y-axes.

3.2 Application to polygenic risk scores

We evaluate the smoothed Lasso approach on polygenic risk scores of the COPDGene study (genetic epidemiology of COPD), a multi-center case-control study designed to identify genetic determinants of COPD and COPD-related phenotypes (Regan et al., 2010). The study has been sequenced as part of the TOPMed Project. The data is available through NHLBI TOPMed (2018). For the study, non-Hispanic Whites and African Americans aged 45 to 80 were recruited as COPD cases and controls. The dataset contains $n = 4010$ individuals (rows), all of which had at least 10 pack-years of smoking history. For each individual, we observe $p = 9153$ datapoints, among them the covariates *age*, *sex*, *packyears*, *height* and five PCA vectors. The remaining entries are SNP data per individual. The input data are summarized in a matrix $X \in \mathbb{R}^{n \times p}$. The response $y \in \mathbb{R}^n$ is the *fev1* ratio, also called the Tiffeneau-Pinelli index, per individual. It describes the proportion of lung volume that a person can exhale within the first second of a forced expiration in a spirometry (pulmonary function) test.

The regularization parameter of the Lasso was chosen as $\lambda = 0.05$ for the following two experiments.

method	L_2 norm	runtime [s]
FISTA	284.6	66.6
smoothed Lasso	26.3	116.6

Table 1: L_2 norm of fitted to true response and runtime in seconds for a single application of any method to the dataset of polygenic risk scores.

Figure 3: L_2 norm of predicted to withheld data in cross-validation (left) and runtime in seconds (right) as a function of the number of withheld entries. Dataset of polygenic risk scores. Logarithmic scale on the y-axes.

3.2.1 Results from a single run

We solve $y = X\beta$ for the given X and y using our smoothed Lasso approach of eq. (9), and compare it to the FISTA algorithm.

Table 1 shows results for a single application of both algorithms. After computing the estimate $\hat{\beta}$ with each method, we consider $\|y - X\hat{\beta}\|_2$, the L_2 norm between the fitted and generated (true) response. We observe that FISTA with standard choices of its tuning parameters seems to have trouble locating the minimum of the Lasso objective function, and is thus worse than smoothed Lasso. However, the FISTA method beats our approach in terms of runtime.

3.2.2 Cross validation

To quantify the accuracy of the methods further, we perform a simple cross-validation experiment in which we withhold a random set of row indices I (of varying size) of the dataset X and the corresponding entries of the response y and fit a linear model to the rest of the data, that is we fit $y_{-I} = X_{-I,.}\beta$. After obtaining an estimate $\hat{\beta}$, we use the withheld rows of X to predict the withheld entries of y, that is we compute $X_{I,.}\hat{\beta}$. We evaluate the quality of the prediction by computing the L_2 norm $\|X_{I,.}\hat{\beta} - y_I\|_2$ between predicted and withheld data.

Figure 3 (left) shows results of this cross-validation experiment. We observe that FISTA with default choices of its tuning parameters seems to have difficulties to converge to the minimum of the Lasso objective function, whereas smoothed Lasso performs better. Not surprisingly, the quality of the prediction becomes worse in general as the number of withheld entries increases, since predictions are based on fewer and fewer datapoints. However, this decrease in accuracy seems to be more pronounced for FISTA than for our smoothed Lasso.

Finally, Figure 3 (right) displays runtime measurements for both approaches. We observe that smoothed Lasso seems to be rather insensitive to the number of withheld entries apart from a very large number of withheld entries. The FISTA algorithm is the faster method, however it exhibits a greater sensitivity to the number of withheld entries, that is the size of the estimation problem.

3.3 Application to synthetic polygenic risk scores

We aim to extend the simulations of Section 3.2 in order to vary the sparsity of the parameter estimate $\hat{\beta}$. To this end, we change the simulation setting as follows. Leaving $X \in \mathbb{R}^{n \times p}$ unchanged ($n = 4010$ individuals and $p = 9153$ datapoints), we simulate a parameter vector $\beta \in \mathbb{R}^p$ in which $nz \in \{0, \ldots, p\}$ entries are drawn from a Beta distribution with shape parameters 1.5 and 10. This will produce nonzero entries in the vector β of magnitude around 0.15, which is realistic in practice. The remaining $p - nz$ entries are set to zero. We then calculate the response as $y = X\beta + \epsilon$, where the entries of the noise vector $\epsilon \in \mathbb{R}^n$ are generated independently from a Normal distribution with mean 0 and standard deviation 0.1.

After generating X and y, we again use FISTA and smoothed Lasso to recover an estimate $\hat{\beta}$. We evaluate the quality of the estimate using $\|\beta - \hat{\beta}\|_2$, that is using the L_2 norm between truth

Figure 4: L_2 norm of parameter estimate to truth (left) and runtime in seconds (right) as a function of the number of non-zero entries in the simulated true parameter vector. Dataset of polygenic risk scores. Logarithmic scale on the y-axis of the time plot.

and estimate.

Figure 4 (left) shows results as a function of the number nz of non-zero entries in the generated true β. We observe that in this experiment, and in contrast to Figures 1 and 2, the smoothed Lasso yields considerably more stable estimates than FISTA (as expressed through a lower deviation in L_2 norm).

Figure 4 (right) shows runtime results for both approaches, showing that in this experiment the runtime scalings of both methods seem to be rather insensitive to the simulation scenario. Interestingly, our smoothed Lasso draws equal in speed with the FISTA algorithm.

Overall, we conclude from the simulations that in the scenarios we investigated, our smoothed Lasso approach yields estimates with equal or improved accuracy in comparison to the gold standard FISTA, exhibits roughly the same runtime scaling as FISTA, and comes with a guarantee on its accuracy.

4   Discussion

This article investigated a smoothing approach for penalized regression using the Lasso. The smoothing approach allows us to obtain smooth gradients everywhere, which facilitate minimization of the convex but non-smooth Lasso objective function.

Importantly, our approach comes with a guarantee of correctness. We prove a uniform bound on the distance between the unsmoothed and smoothed Lasso objective function, thus justifying the use of our smoothed Lasso as a surrogate of the original (non-smooth) Lasso objective function.

Simulations show that in the scenarios we investigated, our proposed smoothing approach yields Lasso estimates with equal or improved accuracy than the gold standard in the literature, the FISTA algorithm of Beck and Teboulle (2009).

5   Proof of Proposition 1

Proof. The bounds on L_e^μ and L_s^μ follow from eq. (8) and eq. (10) after a direct calculation. In particular, for the entropy prox function,

$$
\begin{aligned}
\sup_{\beta \in \mathbb{R}^p} |L_e^\mu(\beta) - L(\beta)| &= \sup_{\beta \in \mathbb{R}^p} \left| \lambda \sum_{i=1}^p f_s^\mu(\beta_i) - \lambda \sum_{i=1}^p |\beta_i| \right| \\
&\leq \sup_{\beta \in \mathbb{R}^p} \lambda \sum_{i=1}^p \left| f_s^\mu(\beta_i) - f(\beta_i) \right| \\
&\leq \sup_{\beta \in \mathbb{R}^p} \lambda \sum_{i=1}^p \mu \log(2) \\
&\leq \lambda p \mu \log(2),
\end{aligned}
$$

where f is as defined in Section 2.2 and where it was used that $\lambda \geq 0$. The result for the squared error prox smoothed L_s^μ is proven analogously.

Since both f_e^μ and f_s^μ are convex according to (Nesterov, 2005, Theorem 1), and since the least squares term $\frac{1}{2}\|X\beta - y\|_2^2$ is convex, it follows that both L_e^μ and L_s^μ remain convex as the sum of two convex functions.

To be precise, strict convexity holds true. Observe that the second derivative of the entropy

smoothed absolute value of Section 2.2.1 is given by

$$\frac{\partial^2}{\partial z^2} f_e^\mu(z) = \frac{4e^{2x/\mu}}{\mu \left(e^{2x/\mu} + 1\right)^2},$$

which is always positive, thus making f_e^μ strictly convex. Therefore, L_e^μ is strictly convex as the sum of a convex function and a strictly convex function. Similar arguments show that L_s^μ is strictly convex. □

References

Beck, A. and Teboulle, M. (2009). A Fast Iterative Shrinkage-Thresholding Algorithm for Linear Inverse Problems. *SIAM J Imaging Sciences*, 2(1):183–202.

Chi, E., Goldstein, T., Studer, C., and Baraniuk, R. (2018). fasta: Fast Adaptive Shrinkage/Thresholding Algorithm. R-package version 0.1.0.

Daubechies, I., Defrise, M., and Mol, C. (2004). An iterative thresholding algorithm for linear inverse problems with a sparsity constraint. *Comm. Pure Appl. Math.*, 57(11):1413–1457.

Efron, B., Hastie, T., Johnstone, I., and Tibshirani, R. (2004). Least angle regression. *Ann Stat*, 32(2):407–499.

Fan, J. and Li, R. (2001). Variable Selection via Nonconcave Penalized Likelihood and its Oracle Properties. *J Am Stat Assoc*, 96(456):1348–1360.

Friedman, J., Hastie, T., and Tibshirani, R. (2010). Regularization Paths for Generalized Linear Models via Coordinate Descent. *Journal of Statistical Software*, 33(1):1–22.

Hahn, G., Banerjee, M., and Sen, B. (2017). Parameter Estimation and Inference in a Continuous Piecewise Linear Regression Model.

Hastie, T. and Efron, B. (2013). lars: Least Angle Regression, Lasso and Forward Stagewise. R-package version 1.2.

Mak, T., Porsch, R., Choi, S., Zhou, X., and Sham, P. (2016). Polygenic scores via penalized regression on summary statistics. *Genet Epidemiol*, 41(6):469–480.

Michelot, C. (1986). A finite algorithm for finding the projection of a point onto the canonical simplex of $\mathbb{R}^n$. *J Optimiz Theory App*, 50(1):195–200.

Nesterov, Y. (1983). A method of solving a convex programming problem with convergence rate $O(1/k^2)$. *Dokl Akad Nauk SSSR*, 269(3):543–547.

Nesterov, Y. (2005). Smooth minimization of non-smooth functions. *Math. Program. Ser. A*, 103:127–152.

NHLBI TOPMed (2018). Boston Early-Onset COPD Study in the National Heart, Lung, and Blood Institute (NHLBI) Trans-Omics for Precision Medicine (TOPMed) Program.

R Core Team (2014). *R: A Language and Environment for Statistical Computing*. R Foundation for Stat Comp, Vienna, Austria.

Regan, E., Hokanson, J., Murphy, J., Make, B., Lynch, D., Beaty, T., Curran-Everett, D., Silverman, E., and Crapo, J. (2010). Genetic epidemiology of copd (copdgene) study design 2. *COPD*, 7:32–43.

Tibshirani, R. (1996). Regression Shrinkage and Selection Via the Lasso. *J Roy Stat Soc B Met*, 58(1):267–288.

Wu, T., Chen, Y., Hastie, T., Sobel, E., and Lange, K. (2009). Genome-wide association analysis by lasso penalized logistic regression. *Bioinformatics*, 25(6):714–721.