1 **Title:** Intergenerational transmission of the patterns of functional and structural brain
2 networks
3
4 **Authors and addresses:**

5 Yu Takagi[1,2*], Naohiro Okada[1,3], Shuntaro Ando[1,4], Noriaki Yahata[1,5,6], Kentaro Morita[1,7],
6 Daisuke Koshiyama[1], Shintaro Kawakami[1], Kingo Sawada[8], Shinsuke Koike[1,9], Kaori Endo[10],
7 Syudo Yamasaki[4], Atsushi Nishida[4], Kiyoto Kasai[1,3], Saori C Tanaka[2]
8
[1] Department of Neuropsychiatry, Graduate School of Medicine, University of Tokyo, Tokyo,
Japan
[2] ATR Brain Information Communication Research Laboratory Group, Kyoto, Japan
[3] International Research Center for Neurointelligence (WPI-IRCN), University of Tokyo
Institutes for Advanced Study (UTIAS), University of Tokyo, Tokyo, Japan
[4] Department of Psychiatry and Behavioral Science, Tokyo Metropolitan Institute of Medical
Science, Tokyo, Japan
[5] Institute for Quantum Life Science, National Institutes for Quantum and Radiological Science
and Technology, Chiba, Japan
[6] Department of Molecular Imaging and Theranostics, National Institute of Radiological
Sciences, National Institutes for Quantum and Radiological Science and Technology, Chiba,
Japan
[7] Department of Rehabilitation, University of Tokyo Hospital, Tokyo, Japan
[8] Office for Mental Health Support, Mental Health Unit, Division for Practice Research, Center
for Research on Counseling and Support Services, University of Tokyo, Tokyo, Japan
[9] University of Tokyo Institute for Diversity and Adaptation of Human Mind (UTIDAHM),
University of Tokyo, Tokyo, Japan
[10] Unit for Mental Health Promotion, Research Center for Social Science & Medicine, Tokyo
Metropolitan Institute of Medical Science, Tokyo, Japan
28
29 ***Corresponding author**: yutakagi322@gmail.com (Yu Takagi, Ph.D.)
30 **Competing Interests**: The authors declare no competing interests.
31

Abstract

There is clear evidence of intergenerational transmission of life values, cognitive traits, psychiatric disorders, and even aspects of daily decision making. To investigate biological substrates of this phenomenon, brain has received increasing attention as a measurable biomarker and potential target for intervention. However, no previous study has quantitatively and comprehensively investigated the effects of intergenerational transmission on functional and structural brain networks from parents to their children. Here, by employing an unusually large cohort dataset, we show that patterns of functional and structural brain networks are preserved over a generation. We also demonstrate that several demographic and behavioural phenotypes have effects on brain similarity. Collectively, our results provide a comprehensive picture of neurobiological substrates of parent-child similarity, and demonstrate the usability of our dataset for investigating the neurobiological substrates of intergenerational transmission.

Introduction

There is clear evidence of intergenerational transmission of socio-economic status[1], intelligence[2], personality[3], parenting style[4], job-selection[5], and psychiatric disorders[6]. This correspondence between parents and their children is not confined to the period in which children are young and live with their parents, but is found over the course of their lives[7]. Although genetic and non-genetic environmental effects are clearly transferred to children from their parents, the mechanisms of parent-child similarity are poorly understood[8].

In recent years, brain has received increasing attention as a target for monitoring and intervention because genetic and epigenetic effects occur at the molecular level and are distal from complex behaviour[9]. Several previous studies have shown that functional connectivity (FC or edge) during wakeful rest obtained by functional magnetic resonance imaging (fMRI) is associated with individual differences in diverse cognitive traits[10–19]. In parallel to functional brain information, individual differences in brain structure have also been characterised and related to diverse cognitive traits[20]. Importantly, previous studies have reported that grey matter volume (GMV) at specific locations in the brain is associated with individual differences in cognitive traits[20–24].

In addition to individual differences in cognitive traits, previous studies also showed that FC[25–43] and GMV[25,44–57] are heritable. These studies have typically used genome wide association study (GWAS) or family/twin study. Most studies have assessed the genetic effects on each edge- or region-level, i.e. univariate analysis, and typically considered demographic or behavioural information as covariates. Importantly, previous studies have not directly focused on the effects of intergenerational transmission from parents to their children.

More recent studies have started to directly examine the effects of intergenerational transmission on the brain using datasets of parent-child dyads[8,59–62]. For example, Lee et al. and Yamagata et al. investigated the similarity of parent-child dyads in FC and GMV, respectively[59,62]. However, these studies involved several limitations. First, no study has quantitatively compared the similarity of different brain networks in detail. Second, because none of these studies examined both functional and structural data together, it remains unclear how functional and structural information are interrelated. Third, no study has comprehensively investigated the effects of demographic and behavioural effects on

similarity. Overall, it is currently unclear whether, to what extent, and how the brains of
parent-child dyads are similar. This situation has arisen, in part, because investigating the
above questions requires a large number of parent-child dyads to provide neuroimaging
datasets with rich behavioural phenotypes. Furthermore, such an approach requires a formal
analytical framework with rigorous statistical analyses and rich computational resources.

In the current study, we sought to understand the neurobiological substrates of
parent-child similarity by combining a statistical framework that allowed us to investigate
network-level similarities and a rich dataset from a subsample of a large population-based
longitudinal cohort (N = 84 parent-child dyads) consisting of resting-state fMRI, structural MRI,
and behavioural phenotypes of parents and their children[63,64]. We sought to answer several
questions: Can a parent-child dyad be identified based on their brains? If so, which brain
networks are more similar compared with other networks? Is the similarity solely driven by
functional or structural information? How do demographic and behavioural factors affect
brain similarity?

Using a dataset consisting of parents and their children, we quantitatively
investigated the brain similarity of parent-child dyads in detail. The present findings
demonstrated that it is possible to reliably identify a parent-child dyad based on the similarity
of their brains. This effect was not solely driven by either functional or structural brain
similarity alone: although functional and structural information had comparable accuracy,
they exhibited important differences, and played complementary roles. Children's basic
demographic factors, including age and sex, testosterone level, and questionnaire-based
developmental scores affected parent-child brain similarity. Collectively, our results provide a
detailed picture of how the brains of children and their parents are similar, and demonstrate
the usability of our unique cohort dataset for investigating the neurobiological substrates of
intergenerational transmission.

Results

We tested 84 parent-child dyads who participated in the "population-neuroscience study of the Tokyo TEEN Cohort (pn-TTC)," a longitudinal study exploring the neurobiological substrates of development during adolescence[63,64]. In the pn-TTC study, neuroimaging and non-imaging behavioural phenotypes were collected from children every 2 years from the age of 11, and from their primary parents (Figure 1a; see *Methods: Overview of the dataset*). Here, we used three brain datasets from the pn-TTC study: children at the ages of 11 and 13 years, and their primary parents. Parents' brains were scanned when their children were 11 years old. The basic demographic data are shown in Table 1.

Table 1: Demographic data. [a] Ages of parents when their children were 11 years old are shown.

	N	Age[a] (mean ± s.t.d)	Sex (Male/Female)
Child age at 11		11.59±0.66	45/39
Child age at 13	84	13.63±0.62	
Parent		43.35±3.95	3/81

We first defined the functional and structural whole-brain patterns for each individual (Figure 1a; see *Methods: Information extraction*). For fMRI, we used a functional atlas defining 268 regions of interest (ROIs) covering the entire brain[12,65]. The FC between these ROIs was estimated using Pearson's correlation coefficient, resulting in a 268 × 268 FC matrix for each subject. As the FC matrix is symmetrical, only the strictly lower triangular part of each matrix was kept, resulting in 35,778 (= 268 × 267 / 2) unique entries. We regressed potential confounds including total GMV and head motion. To further avoid the effects of motion artefacts, we employed a "scrubbing" procedure to identify and exclude any frames exhibiting excessive head motion[66]. We also obtained GMV for each ROI using T1w images, then averaged within each region. We used the same 268 ROIs as in the fMRI, resulting in a vector with a size of 268 for each subject.

To quantitatively evaluate the brain similarity of parent-child dyads, we proposed a framework to compare network-level similarity between parent-child dyads (Figure 1b; See *Methods: Similarity analysis* for details), inspired by a recently proposed approach for individual identification based on the brain[12]. To calculate the similarity of parent-child dyads,

we first calculated the correlations between a child's FC and/or GMV vector to all parents' vectors including the child's own parent. We next assessed whether the similarity of the parent-child dyad (child and his/her own parent) was larger than that of a stranger-child dyad (child and another child's parent). We then calculated the winning rate of the similarity between parent-child dyad, which was referred to as "accuracy", because it can be considered as a "pairwise classification accuracy" when we randomly sampled a parent-child dyad and another parent, then conduct classification (See *Methods: Similarity analysis* for details). We repeated this procedure across all dyads and averaged these accuracies. Compared with conventional individual identification methods, our proposed framework has more statistical power, as described later. We performed 1,000-times bootstrapping to estimate 95% confidence intervals of accuracy by randomly subsampling 90% of the subjects in each iteration. To determine whether accuracy was achieved at above-chance levels, we used 1,000-times permutation testing to generate a null distribution by randomly shuffling the parent-child mapping.

Figure 1: **Analysis procedure of parent-child brain similarity**. (**a**) We employed a dataset obtained from the "population-neuroscience study of the Tokyo TEEN Cohort (pn-TTC)" study, which consists of resting-state fMRI

and T1w images of parents and their children. To obtain a functional connectivity (FC) matrix, signals were

extracted from all subjects using resting-state fMRI data from 268 ROIs. The signals were then turned into an FC

matrix via covariance estimation. To obtain grey matter volume (GMV) vectors, T1w images were first segmented

into grey matter, white matter, and cerebrospinal fluid. Using the grey matter, GMV of each ROI was obtained by

averaging values within the ROI. We used the same 268 ROIs as in the FC. **(b)** For each parent-child dyad, we first

calculated the similarity of FC and/or GMV vectors based on their Pearson's correlation. We next calculated

similarities between the child and another child's parent. We then calculated whether the similarity of the

parent-child dyad is greater than that of stranger-child dyads. Finally, we calculated the winning rate of the

parent-child dyad ("accuracy"), and repeated this procedure across all parent-child dyads.

Whole-brain analysis

We first assessed the similarity of parent-child dyads using whole-brain FC and GMV. When

we used a dataset of children at age 11, accuracies were 64.6% for FC (estimated via 1,000-

times bootstrapping; 95% CI = [62.5, 66.8]; P < 0.001, 1,000-times permutation test) and

70.3% for GMV (95% CI = [68.9, 71.8]; P < 0.001) (Figure 2a). When we used a dataset of

children at age 13, the accuracies were 66.7% for FC (95% CI = [64.5, 68.9]; P < 0.001) and

73.8% for GMV (95% CI = [72.1, 75.6]; P < 0.001) (Figure 2b). Thus, we provided the first strong

evidence that it is possible to identify parent-child dyads based on their functional and

structural brain information.

We next assessed the importance of information for performance of specific edges

for FC and regions for GMV, respectively (Figures 2c-2f). To quantify the extent to which

different edges and regions contribute to similarity, we derived two measures: the differential

power (DP) which calculates how characteristic edges and regions tend to be, and group

consistency (φ) which quantifies edges and regions that are highly consistent across all parent-

child pairs in a dataset[12] (see *Methods: Similarity analysis*). For visualisation purposes, we

show the structural locations of DP and φ in the 99.75th and 90th percentile, for edges (FC)

and regions (GMV) respectively. For both FC and GMV, significant edges or regions tended to

be distributed across the entire brain. This pattern was stable across a range of thresholds

(Supplementary Figure 1). Note that, for visualisation purposes, we excluded the brainstem

from the figure for GMV because all regions in this area had extremely high φ values, possibly

because of the much lower magnitudes of signals compared with the other regions

(Supplementary Figure 2).

Given that head motion confounds analyses of FC[66], we confirmed that qualitatively

similar results were obtained when we excluded parent-child dyads whose children's head

movements were in the top 25%, either at age 11 or 13, resulting in the inclusion of 41.25%

of the total sample (Supplementary Figure 3). We also confirmed that accuracy obtained by

the distribution of their frame-to-frame motion during fMRI scans[12] was not above chance

level (51.0% for age 11, 95% CI = [48.9, 53.0]; 48.7% for age at 13, 95% CI = [47.1, 50.1]). Thus,

it is unlikely that the identification power of FC was based on idiosyncratic patterns related to

motion in the scanner.

Figure 2: **Successful identification of parent-child dyad based on their functional and structural brain**

information. Box plots of parent-child identification accuracy and factors affecting accuracy for **(a)** children at

age 11 and **(b)** at age 13 using whole-brain (268-node) for functional connectivity (FC: red box) and grey matter

volume (GMV: blue box). Directly to the right of these boxes (grey box) are the results of the 1,000-times

permutation testing. The bottom and top edges of the box indicate the 25th and 75th percentiles obtained via

bootstrapping, respectively. The crosses denote outliers, and the whiskers extend to the most extreme data

points not considered outliers. **(c–f)** Factors affecting identification accuracy. For FC, the top 99.75th percentile

of differential power (DP: highly discriminative; yellow) edges and group consistency (φ: highly similar, or least

helpful; green) edges are shown (circle plot; in which nodes are grouped according to anatomic location). For

195   GMV, the top 90th percentile ROIs of DP and φ were calculated, then normalised by dividing the number of ROIs

196   in each anatomical group (bar plot). A legend indicating the approximate anatomical "lobe" is shown. PFC,

197   prefrontal; Mot, motor; Ins, insula; Par, parietal; Tem, temporal; Occ, occipital; Lim, limbic (including cingulate

198   cortex, amygdala, and hippocampus); Cer, cerebellum; Sub, subcortical (including thalamus and striatum); Bsm,

199   brainstem; L, left hemisphere; R, right hemisphere.

Network-based similarity

We further investigated the contributions of specific networks to this similarity. We grouped

the whole regions into 10 sub-networks[67] (Figure 3a), and subsequently performed the same

analyses using only the edges and regions from a given network. Note that we calculated the

null distribution for each network via permutation testing, thus taking differences of the

number of edges/regions among networks into account.

For FC, medial frontal and frontoparietal networks led to high accuracies (Figure 3b).

In contrast, for GMV, default mode, subcortical, cerebellum and visual networks led to high

accuracy. Compared with FC, GMV achieved modestly higher accuracy than FC at age 11

(Figure 3c; paired sample t-test, t(10) = −2.16, P = 0.056, Hedge's g = −0.72) and significantly

higher accuracy at age 13 (Figure 3c; paired sample t-test, t(10) = −3.10, P = 0.011 , Hedge's g

= −0.89). To further assess the importance of each network, we next assessed performance

using between-network pairs of edges. We observed that edges between the medial frontal–

frontoparietal networks and medial frontal–motor networks resulted in higher accuracies

than the other between-network pairs (Supplementary Figure 4).

The results confirmed that our proposed method had greater statistical power than

conventional methods for individual identification (Supplementary Figure 5). Specifically, for

FC, six and five of 11 networks were significant at age 11 and 13, respectively, using the

conventional method, whereas nine and 10 networks, respectively, were significant in our

proposed method. For GMV, six and nine of 11 networks were significant at age 11 and 13,

respectively, using the conventional method, whereas all networks were significant using our

proposed method.

Figure 3: **Network-based analyses demonstrated that almost all brain networks were highly similar between parents and their children**. (**a**) We utilised a 268-node functional atlas. Nodes were further grouped into the 10 functional networks. Network names are shown to the left. (**b**) Box plots of accuracies using within-network edges of FC analysis (top row; networks 1–10 and whole-brain (ALL); indicated below the x-axis of each graph) and within-network nodes of GMV (bottom row). The bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The crosses denote outliers, and the whiskers extend to the most extreme data points not considered outliers. (**c**) Comparison between accuracies of child age at 11 and 13 for FC (top row) and GMV (bottom row). Each scatter shows each network and line connected the same network. Bold lines indicate ALL. * paired sample t-test, P < 0.05, n.s. non-significant, uncorrected.

Function and structure provide complementary information

Although FC and GMV revealed comparable performance in the above analyses, it remained unclear whether they contained similar information. This raises the following question: if parents and their children exhibit similar patterns of structural brain information, do they also exhibit similar patterns of functional brain information? Indeed, although functional and structural brain information is interrelated, they contain exclusive information that putatively

characterises distinct properties of individual differences[68]. To test this question, we

investigated the relationship between parent-child similarity defined by FC and that of GMV.

We found that the Pearson's correlation between the similarities defined by FC and GMV was

low: only two of 22 networks were significant (Supplementary Figure 6; $P = 0.026$ for visual

and $P = 0.021$ for whole-brain when we used data from children at age 13; the other networks

were not significant, $P > 0.05$, 1,000-times permutation test, uncorrected). Thus, although

both FC and GMV were similar between parent-child dyads, their characteristics were

dissimilar.

Given that FC and GMV appeared to contain independent information, we further

investigated whether they contained complementary information. To test this question, we

conducted the same analyses using both FC and GMV simultaneously by concatenating the

two vectors (hereafter referred to as "COMB"). COMB achieved the highest accuracies in more

than half of cases (15/22), compared with function (4/22) and structure (3/22) alone (Table

2). This number is significantly greater than chance (Supplementary Figure 7; $P < 0.001$, 1,000-

times permutation test). Overall, these results suggest that patterns of functional and

structural information contained complementary information in terms of parent-child brain

similarity.

Table 2: **COMB achieved higher accuracies compared with FC and GMV for many networks in children at age 11 and at age 13**. Means and 95% confidence intervals estimated via 1,000-times bootstrapping are shown. **<u>Bold underlined</u>** text indicates the best performance. MF, medial frontal; FP, frontoparietal; DMN, default mode network; Mot, motor; Vis I, visual I; Vis II, visual II; Vis A, visual association; Cing, cingulo-opercular; Sub, subcortical; Cer, cerebellum.

Network	Age	FC	GMV	COMB
ALL	11	64.60 [62.54,66.83]	70.30 [68.86,71.75]	**<u>70.64 [69.21,72.16]</u>**
	13	66.69 [64.50,68.86]	**<u>73.82 [72.14,75.55]</u>**	73.57 [71.89,75.30]
1. MF	11	**<u>61.14 [58.83,63.66]</u>**	57.56 [56.02,58.94]	58.32 [56.77,59.66]
	13	**<u>64.26 [62.20,66.36]</u>**	59.02 [57.06,60.70]	59.14 [57.14,60.92]
2. FP	11	**<u>64.35 [62.40,66.52]</u>**	60.52 [59.01,62.20]	61.90 [60.38,63.55]
	13	60.57 [58.61,62.49]	60.94 [59.35,62.65]	**<u>61.79 [60.25,63.46]</u>**
3. DMN	11	62.77 [61.06,64.65]	63.49 [61.93,64.94]	**<u>63.66 [62.04,65.21]</u>**
	13	59.07 [57.03,61.03]	62.53 [60.77,64.23]	**<u>62.84 [61.05,64.50]</u>**
4. Mot	11	57.95 [55.60,60.29]	61.30 [59.80,62.85]	**<u>62.27 [60.70,63.82]</u>**
	13	59.78 [57.89,61.77]	63.56 [61.95,65.14]	**<u>64.27 [62.63,65.98]</u>**
5. Vis I	11	55.42 [53.19,57.59]	63.98 [62.40,65.59]	**<u>64.91 [63.23,66.58]</u>**
	13	54.60 [52.61,56.77]	65.37 [63.60,67.21]	**<u>65.98 [64.20,67.77]</u>**
6. Vis II	11	56.44 [54.41,58.61]	58.14 [56.20,60.04]	**<u>58.95 [57.14,60.81]</u>**
	13	56.20 [54.14,58.25]	59.07 [56.95,61.03]	**<u>60.33 [58.22,62.32]</u>**
7. Vis A	11	**<u>57.52 [55.35,59.86]</u>**	56.81 [55.51,58.02]	57.13 [55.84,58.31]
	13	52.78 [50.50,55.06]	**<u>59.05 [57.28,60.56]</u>**	58.92 [57.28,60.41]
8. Cing	11	53.28 [51.08,55.66]	**<u>61.54 [59.89,62.99]</u>**	61.53 [59.91,62.94]
	13	62.04 [59.93,64.34]	63.48 [61.51,65.17]	**<u>63.71 [61.68,65.46]</u>**
9. Sub	11	60.43 [58.52,62.59]	64.34 [62.67,65.95]	**<u>64.66 [62.97,66.23]</u>**
	13	55.45 [53.23,57.62]	63.08 [61.41,64.65]	**<u>63.61 [62.00,65.19]</u>**
10. Cer	11	53.32 [51.23,55.37]	62.13 [60.38,63.73]	**<u>62.86 [61.24,64.54]</u>**
	13	56.24 [53.87,58.68]	61.83 [59.91,63.71]	**<u>62.50 [60.67,64.34]</u>**
# best		4	3	15

Effects of demographic factors on the brain similarity

264   The results described above indicated strong intergenerational transmission effects
265 on the brain from parents to their children. However, it was still unclear whether all parent-
266 child dyads were equally similar. Therefore, we then investigated which factors influence brain
267 similarity. Here, we focused on fundamental demographic factors: age and sex.

268   We found that accuracies at age 11 were significantly lower than those at age 13 for
269 GMV (Figure 3c bottom; paired sample t-test, $t(10) = -2.39$, $P = 0.038$, Hedge's $g = -0.25$),
270 whereas they were no different for FC (Figure 3c top; paired sample t-test, $t(10) = -0.03$, $P = $
271 0.97, Hedge's $g = -0.01$).

272   We then divided children into males and females (Figure 4). The results confirmed
273 that both male and female children exhibited significant accuracies for almost all networks, as
274 in the previous analyses (Figure 4a). When we compared males and females (Figure 4b), the
275 accuracies of female children were significantly higher than those of male children for FC, at
276 both age 11 (paired sample t-test, $t(10) = -2.34$, $P = 0.04$, Hedge's $g = -0.88$) and age 13
277 (paired sample t-test, $t(10) = -3.80$, $P = 0.003$, Hedge's $g = -1.21$). Female children also
278 exhibited higher accuracy than male children for GMV at age 13 (paired sample t-test, $t(10) = $
279 -2.47 , $P = 0.03$, Hedge's $g = -0.83$), but not at age 11 (paired sample t-test, $t(10) = -0.25$, $P = $
280 0.81 , Hedge's $g = 0.07$).

281   When we compared accuracies at age 11 and age 13 for male and female children
282 separately (Figure 4c), female children at age 13 had significantly greater accuracy than those
283 at age 11 for GMV (paired sample t-test, $t(10) = -3.75$, $P = 0.004$, Hedge's $g = -0.76$). All other
284 comparisons were not significant: male, FC (paired sample t-test, $t(10) = 0.37$, $P = 0.72$,
285 Hedge's $g = 0.13$); male, GMV (paired sample t-test, $t(10) = 2.05$, $P = 0.068$, Hedge's $g = 0.23$);
286 female, FC (paired sample t-test, $t(10) = -0.73$, $P = 0.48$, Hedge's $g = -0.18$)

Figure 4: Male and female children show different trends of brain similarity across development. (a) Accuracies split by children's sex for FC and GMV. For each network, accuracies for males (left boxes) and females (right boxes) are shown. (b) Comparison between males and females. (c) Comparison between children age at 11 and 13. Each scatter shows each network and line connected the same network. Bold lines indicate ALL. M(11), males at age 11; F(11), females at age 11; M(13), males at age 13; M(13), males at age 13; * Paired sample t-test, P < 0.05; ** P < 0.01. n.s. non-significant, uncorrected.

Effects of behavioural phenotypes on brain similarity

Finally, we examined whether behavioural phenotypes have effects on the brain similarity of parent-child dyads. Here, we used two important behavioural phenotypes for adolescents: hormone level and questionnaire-based developmental score (Figure 5; see

Methods: Testosterone and Child Behavior Checklist). We used COMB for this analysis as brain

information.

We examined testosterone as a hormone level, because it is known that the

pubertal period is a sensitive period for testosterone-dependent organisation of the brain[69].

We found that children with high testosterone exhibited significantly higher accuracy

compared with children with low testosterone at age 11 (paired sample t-test, $t(10) = -2.91$,

$P = 0.016$, Hedge's $g = -1.03$) but not at age 13 (paired sample t-test, $t(10) = -0.60$, $P = 0.56$,

Hedge's $g = -0.22$).

We next investigated the effects of a questionnaire-based development score (Child

Behavior Checklist: CBCL)[70]. The CBCL is a parental-report assessment used to screen for

emotional, behavioural, and social problems, and to predict psychiatric illnesses[71]. We found

that children with high CBCL had significantly higher accuracy compared with children with

low CBCL at age 11 (paired sample t-test, $t(10) = -2.81$, $P = 0.018$, Hedge's $g = -1.05$) but not

at age 13 (paired sample t-test, $t(10) = -0.99$, $P = 0.34$, Hedge's $g = -0.33$).

Figure 5: Behavioural phenotypes have effects on brain similarity. Accuracies split by testosterone and

CBCL. Left panels: Boxplots of accuracies split by scores of behavioural phenotypes (left: upper-half

children; right: lower-half children). Right panels: Comparisons between upper- (left) and lower-half (right)

children. Each scatter shows each network and line connected the same network. Bold lines indicate ALL. *

Paired sample t-test, P < 0.05. n.s. non-significant, uncorrected.

Discussion

The present study revealed that patterns of functional and structural brain information are preserved over a generation. Although the effects of the intergenerational transmission have been investigated in various research fields, including developmental psychology, educational psychology, and economics, no study has comprehensively and quantitatively investigated the neurobiological substrates of intergenerational transmission. The current results revealed that, despite substantial differences between parents and their children, their brains are sufficiently similar that we can identify parent-child dyads based on information about their brains. We employed a rigorous statistical framework and an unusually large functional and structural neuroimaging dataset of parent-child dyads with rich behavioural phenotypes (N = 84 parent-child dyads)[63,64]. Although both functional and structural brain information has comparable levels of accuracy, their characteristics were different but complementary. Demographic factors and behavioural phenotypes also have large effects on brain similarity. Taken together, our results provide a detailed picture of whether, to what extent, and how brains of parent-child dyads are similar.

Previous studies have reported that brain information is heritable. Several genome-wide association studies (GWAS) have been conducted to identify genetic risk variants for GMV[25,44,50–56] and FC[25,26]. Although heritable regions or edges have been successfully identified, most studies have been insufficiently powered because GWAS require a large sample size (although some studies used multivariate approaches to increase statistical power[26,58]). In addition, these studies typically did not consider the effects of demographic and behavioural phenotypes. In addition to GWAS, both twin and family-based studies have reported that GMV[45–49,57] and FC[27–43] are heritable. These studies typically achieve larger effect sizes than GWAS studies with smaller sample sizes, although the possibility of inflated effect sizes due to shared environments is a concern. These studies also typically ignore demographic and behavioural phenotypes by treating them as covariates. Importantly, none of the previous studies described here directly investigated the effects of intergenerational transmission from parents to children.

In recent years, some studies directly investigated intergenerational transmission of brain information using parent-child dyads. Although extended pedigree studies with sufficient sample sizes could answer such a question, it is logistically more difficult to recruit participants for pedigree studies than for studies with a parent-child design. Thus, parent-child

design studies play a complementary role in the investigation of intergenerational
transmission. For example, some studies found that, during specific tasks, brain activity of
parent-child dyads was synchronised[60,61]. Other, more relevant studies to the current studies
reported that patterns of FC [59] and GMV [62] were similar between parent-child dyads. However,
these studies did not quantitatively and comprehensively investigate the effects of different
brain networks, and did not compare FC and GMV. It is also unclear which factors affect brain
similarities, including age, sex, hormonal level, and behavioural traits (although Yamagata et
al. investigated the effects of sex on GMV-based similarity using ROI-based analyses [62]). This
is partly because investigating such questions require a large neuroimaging dataset of parent-
child dyads with rich behavioural phenotypes. Overall, the present study is the first to
investigate whether, to what extent, and how brains of parent-child dyads are similar.

For both function and structure, the brain regions that contributed to similarity
tended to be broadly distributed across the entire brain when we used whole-brain
information (Figure 2c–2f), as in previous FC-based individual identification studies[12,67].
Although this tendency was retained when we only used the edges or regions within sub-
networks, we observed slightly different contrasts between functional and structural
information (Figure 3b and Table 2). Specifically, for function, the medial frontal and
frontoparietal areas, which are known to be involved in networks related to higher cognitive
function, revealed higher accuracies than structural information. In contrast, for structure,
visual, subcortical, and cerebellum networks exhibited higher accuracies than function. This
contrast is interesting because the prefrontal cortex is one of the last regions of the brain to
reach maturation, exhibiting development until approximately 25 years of age, whereas
subcortical regions reach maturation earlier[72]. Note that, for structural brain information, only
a small number of studies using GMV are comparable to the current study[73,74], and none of
them investigated each network's contribution in detail.

We confirmed that our results were not solely driven by structural or functional
similarity alone. The neural similarity between FC and GMV exhibited a weak correlation
(Supplementary Figure 6). In addition, combining FC and GMV led to the highest accuracy for
many brain regions (Table 1). These results indicate that FC and GMV are distinct and contain
complementary information. Previous studies reported that function and structure contain
similar information, but also distinct information[68]. One study reported that information
obtained from function and structure contain complementary information for the
identification of siblings, but did not investigate the contributions of distinct anatomical brain

locations[75]. The current results are not only consistent with those of previous studies but also
demonstrated qualitative differences between function and structure by exhaustively
investigating the contributions of each distinct anatomical brain location. It is noteworthy that
GMV was able to obtain similar accuracy, even with short scans that are routinely acquired as
initial scans in MRI protocols.

We also found that several demographic and behavioural factors had effects on the
brain similarity of parent-child dyads. First, we found that age had an effect on accuracies of
GMV, but not FC. It is known that both FC and GMV change through adolescence[76,77]; thus,
this difference suggests that the functional and structural development of the brain are not
qualitatively equal, at least from the perspective of parent-child similarity. Second, when we
split children into males and females, we found age effects on the brains of female children,
both for FC and GMV. This finding suggests that the developmental trajectory of the brain
qualitatively differs between females and males. In addition, female children were more
similar to their parents than male children, particularly at age 13. This finding is intriguing
because previous studies also reported that female children are more similar to their mothers
both behaviourally[78] and neurally[62]. Third, testosterone affected brain similarity in parent-
child dyads. Interestingly, children with high levels of testosterone exhibited greater similarity
than other children, despite female children having greater accuracy than male children. Note
that there were no significant differences in levels of testosterone between males and females
both at age 11 (two-sample t-test, $t(49) = -0.64$, $P = 0.53$, Hedge's g = -0.19) and 13 (two-
sample t-test, $t(44) = 0.81$, $P = 0.42$, Hedge's g = 0.25). It is known that levels of testosterone
increase through adolescence, especially in male children[79]. Thus, different results may have
been obtained if older children were tested. Fourth, questionnaire-based development scores
had effects on similarity. Children with higher developmental problem scores were more
similar to their parents than other children. Although this result is somewhat counterintuitive,
it suggests that similarity of the brain does not merely represent behavioural maturity
assessed by questionnaire. Overall, the current results provide the first detailed picture of the
signature of intergenerational transmission in the brain.

Recent developments in cognitive neuroscience have made it possible to investigate
individual differences, an issue that has not been deeply investigated because of the absence
of adequate datasets, analytical techniques, and computational resources. The current results
shed light on the importance of investigating family-level differences, in addition to individual-
level differences. Families are not a neutral environment for identity development, but deeply

affect individuals from adolescence, strongly influencing the development of a person's
identity[80]. Given that the current results revealed that parental brains are similar to the brains
of their children, we propose that future studies should investigate the relationships between
family-level behavioural indices and family-level brain information to enable more reliable
predictions of children's behaviour and development. We believe our dataset will help to
extend research in this direction[64].

The current findings indicate several potentially interesting questions for future
research. The first and perhaps most important question is whether parent-child brain
similarity changes across children's development. Recent studies proposed a brain-based
quantitative approach for investigating the trajectory of development[81,82]. Although the
current study revealed that children at age 11 and age 13 are different, it may be valuable to
investigate whether the trajectory of the parent-child brain similarity affects the various risks
faced by adolescents, including psychiatric disorders and criminal behaviour, using much
longer-term longitudinal datasets. Second, insights may be gained by using genetic
information to confirm the extent to which genetic factors contribute to neural similarity. We
examined biological parent-child relationships in the current study. Thus, it may be valuable
to test whether non-biological parents and children show the same level of brain similarity.
Third, we used GMV as a structural brain measurement because a number of previous studies
investigated the relationship between GMV and various traits. However, the human brain also
exhibits individual differences in white matter microstructure. Diffusion tensor imaging
provides measures of white matter integrity in the brain, and can provide useful data, but, like
GMV, produces different information to FC[83]. Indeed, previous studies have shown that
thickness correlations partially reflect underlying fibre connections but contain exclusive
information[84]. Future studies should use other types of structural information, such as
structural connectivity obtained by diffusion tensor imaging. Fourth, because our dataset
mostly consisted of mothers (81 mothers among 84 parents), it may be valuable to test
whether we can also identify children' brains from fathers. Although we confirmed that we
were able to successfully conduct analyses both male and female children, future studies
should investigate the effects of parents' sex on performance.

In the present study, we sought to address a critical question in social science:
whether, to what extent, and how parents and children are similar. Our analytical framework
and the richness of our dataset made it possible to ask the question from the neurobiological
perspective. The results revealed that parents' and their children's brains exhibit a high degree

of similarity, and that various factors, including age, sex, hormones, and development score
have effects on similarity. These results provide a comprehensive picture of the
neurobiological substrates of parent-child similarity, and show the usability of our dataset for
investigating the neurobiological substrates of intergenerational transmission.

Acknowledgments

This study is the result of the projects "Science of personalized value development through
adolescence: integration of brain, real-world, and life-course approaches" (16H06396,
16H06398, 16H06399) and "Adolescent Mind & Self-Regulation" (23118001, 23118002) of
Grants-in-Aid for Scientific Research on Innovative Areas from the Japan Society for the
Promotion of Science. We thank Hiroshi Imamizu, Ph.D., Ryu Ohata, Ph.D., Ayumu Yamashita,
Ph.D., and Aurelio Cortese, Ph.D. for reviewing the draft and helpful discussions. We thank
Okito Yamashita, Ph.D., and Jun-ichiro Hirayama, Ph.D. for helpful discussions. We thank Kana
Inoue and Kaori Tachi for their support with data curation. We thank Giuseppe Lisi, Ph.D. for
his support with code sharing. We thank Corey Horien for his support with code and data
sharing.

Author Contributions

N.O., S.A., N.Y., K.M., D.K., S. Kawakami, K.S., S. Koike, K.E., S.Y., A.N., K.K., and S.C.T. recruited
participants for the study and collected their behavioural and imaging data. N.O. inspected
imaging data. N.Y. provided codes for preprocessing resting state fMRI data. S. Koike, and K.K.,
interpreted the results and provided technical advice. S.C.T. supervised the study. Y.T.
conceptualised the study, conducted the analysis, visualised the data, and wrote the
manuscript. All authors reviewed the manuscript.

Data and code availability statement

475 The data that support the findings of the current study may be available from the
476 corresponding author upon reasonable request (http://value.umin.jp/data-resource.html).
477 The code supporting the findings of this study will be published after acceptance.

Methods

Overview of the dataset

The Tokyo TEEN Cohort (TTC) study, which was launched in 2012, is a large-scale longitudinal general population-based survey to elucidate puberty development during adolescence, particularly the acquisition processes of self-regulation and willingness to face challenges, by focusing on the interaction between biological, psychological, and social factors[63,64]. This study was conducted as part of the population-neuroscience component of the TTC (pn-TTC) study, in which 301 early adolescents were recruited from the general population. Subjects of the pn-TTC study were subsampled from a larger subject group of the TTC study, and it was confirmed that the pn-TTC subsample was representative of the TTC study population. Written informed consent was obtained from each subject and the subject's primary parent before participation. All protocols were approved by the research ethics committees of the Graduate School of Medicine and Faculty of Medicine at the University of Tokyo, Tokyo Metropolitan Institute of Medical Science, and the Graduate University for Advanced Studies. All research was performed in accordance with relevant guidelines/regulations. The detailed methods for subject recruitment are described elsewhere[63,64]. The dataset is publicly shared upon request (http://value.umin.jp/data-resource.html).

We excluded subjects who exhibited anomalies in fMRI or T1w images. We also excluded parents who did not have either fMRI or T1w images, and children who did not have either fMRI or T1w images at age 11 and age 13. After this screening process, 84 dyads were included in the final analysis (39 female children; 81 mothers; age = 11.59 ± 0.66 for children at age 11, 13.63 ± 0.62 for children at age 13, and 43.35 ± 0.62 for parents, mean ± s.t.d).

MRI parameters

Subjects were instructed to lie supine on the bed of the MRI scanner. MRI scanning was performed on a Philips Achieva 3T system (Philips Medical Systems, Best, The Netherlands) using an eight-channel receiver head coil. Each subject underwent resting-state fMRI and T1-weighted (T1w) three-dimensional magnetisation-prepared rapid gradient echo (3D-MPRAGE) sequences.

Sagittal T1w images were acquired using the 3D-MPRAGE sequence with the

following parameters: repetition time (TR) = 7.0 ms, echo time (TE) = 3.2 ms, minimum

inversion time = 875.8 ms, flip angle = 9°, matrix = 256 × 256, field of view (FOV) = 256 mm ×

240 mm × 200 mm, voxel size = 1 mm × 1 mm × 1 mm, slice thickness = 1 mm, number of

slices = 200. The acquisition time was approximately 10 min 42 sec.

Resting-state fMRI images were acquired using a gradient-echo echo-planar imaging

(EPI) sequence with the following parameters: TR / TE, 2500 ms / 30 ms; flip angle, 80°; matrix,

64 × 64; FOV, 212 mm × 199 mm × 159 mm; voxel size, 3.31 mm × 3.31 mm; slice thickness,

3.20 mm; slice gap, 0.8 mm. Each brain volume consisted of 40 axial slices and each functional

run contained 250 image volumes preceded by four dummy volumes, resulting in a total scan

time of 10 min 40 sec. Subjects were instructed to stay awake, to keep their minds as clear as

possible, and to keep their eyes on a fixation point at the centre of the screen through a mirror

during scanning.

Information extraction

We used Statistical Parametric Mapping 8 (SPM8: Wellcome Department of Cognitive

Neurology, http:/ /www.fil.ion.ucl.ac.uk/spm/software/) in MATLAB (MathWorks, Natick,

Massachusetts) for preprocessing and statistical analyses.

Preprocessing of structural *MRI:* T1w images were segmented into three tissue

classes (grey matter [GM], white matter [WM], and cerebrospinal fluid [CSF]) using a

segmentation approach implemented in SPM8. The segmented images (only GM) were then

normalised into standardised Montreal Neurological Institute (MNI) space by applying a

deformation field in SPM8. The GMV of each ROI was extracted and averaged within that ROI.

We used a functional atlas defining 268 ROIs that cover the entire brain (functional atlas from

Finn et al.[12], which used the method developed by Shen et al.[65]) (this atlas can be downloaded

from https://www.nitrc.org/frs/?group_id=51), enabling us to obtain a vector with a size of

268 for each subject. Note that, although an alternative method for inter-subject registration

called Diffeomorphic Anatomical Registration Exponentiated Lie algebra (DARTEL) exists, we

did not employ it because our goal was not to conduct comparisons at the group-level. Future

studies should investigate whether employing another segmentation and normalisation

method can improve accuracy.

Preprocessing of resting-state fMRI: Preprocessing of resting-state fMRI included

slice-timing correction, realignment, co-registration, normalisation to MNI space, and spatial

smoothing with an isotropic Gaussian kernel of 6 mm full-width at half-maximum. To avoid the effects of head motion artefacts, we calculated framewise displacement (FD). FD is defined as the mean relative displacement between two consecutive volumes for each of the six motion parameters. We conducted a "scrubbing" procedure by removing volumes with FD > 0.5 mm, along with the previous volume and two subsequent volumes, as proposed by Power et al.[66] The average grey matter time-course for each ROI was calculated, then temporally filtered using a first-order Butterworth filter with a pass band between 0.01 Hz and 0.08 Hz. The time-course of each ROI was linearly regressed by the temporal fluctuations in white matter, cerebrospinal fluid, and the entire brain, as well as six head motion parameters. The time-course of white matter and cerebrospinal fluid were filtered using a first-order Butterworth filter with a pass band between 0.01 Hz and 0.08 Hz, and a white matter mask was eroded by one voxel to consider a partial volume effect. All parameters were determined in accord with a previous study[19]. For each subject, an FC matrix between all ROIs was then calculated by evaluating pair-wise temporal Pearson's correlations of blood-oxygenation level dependent time courses, based only on the remaining images after the scrubbing step above. We used the same 268 ROIs that were used for GMV. Because FC matrices are symmetrical, values on only one side of the diagonal were kept, resulting in 35,778 unique edges (268 × 267/2). We then regressed the motion and total grey matter volume and mean FD out from data matrices.

Motion index: In addition to resting-state fMRI and structural MRI information, we performed the same analyses using motion estimates during resting-state fMRI to investigate the effects of motion artefacts[12]. We first specified 20 bins to span {0:0.05:1} to calculate discrete motion distribution vectors for each parent and child based on FD over an entire scan. These motion distribution vectors were then used in the same way as the FC or GMV vectors.

Similarity analysis

We modified a connectome fingerprinting approach by Fin et al[12]. They used two datasets consisting of the same individual but different task sessions, called "source" and "target" dataset. They correlated the connectivity vector from one participant in the source dataset to the vectors of all participants in the target dataset and identified the maximum correlation. If the two vectors showing the strongest correlations came from the same individual, the resulting binary accuracy was 100%, whereas binary accuracy was 0% otherwise. Although these studies successfully identified brain networks that contributed to the individual

identification, their method treats second-ranked and worst-ranked cases as equally failed

cases, thus discarding some information that might be useful for improving the statistical

power. In addition, the chance rate depends on the number of samples, thus making the

interpretation difficult, especially comparing different datasets with different sample sizes. To

overcome these issues, we modified the method as follows. We confirmed that the proposed

method is more sensitive than conventional methods (Supplementary Figure 5) and the

chance rate was always 50%, irrespective of the sample size.

For each parent-child dyad, we first calculated the similarity of their FC and/or GMV

patterns based on their Pearson's correlation. We next assessed whether the similarity of the

parent-child dyad (child and their own parent) was larger than a stranger-child dyad (child and

another child's parent). We then calculated the winning rate of the similarity between parent-

child dyad, denoted as "accuracy". We repeated this procedure across all children and

averaged accuracies.

Intuitively, the obtained statistics can be considered as a "pairwise classification

accuracy" calculated by the following procedure:

1. Select a child randomly from all children in the sample.

2. Select two parents, including the child's own parent and a randomly selected parent in the

sample.

3. If the Pearson's correlation coefficient between parent-child dyad is higher than that of

stranger-child dyad, the result is recorded as a correct parent-child identification.

4. In contrast, if the Pearson's correlation coefficient between parent-child dyad is smaller

than that of stranger-child dyad, it is recorded as a failed identification.

5. Repeat this procedure and calculate accuracy across repetition.

By increasing the number of repetitions, this approach converges to the accuracy

obtained by the main analysis. The chance rate of this approach is always 50%.

We performed 1,000-times bootstrapping to estimate the 95% confidence interval

of accuracy, by randomly subsampling 90% of the subjects in each iteration. To determine

whether accuracy was achieved at above-chance levels, we used 1,000-times permutation

testing to generate a null distribution by randomly shuffling the parent-child mapping.

To determine the role of specific edges/regions in the performance, we quantified

highly unique and highly consistent edges/regions using a differential power (DP) measure

and a group consistency measure (ϕ) described in detail elsewhere[12]. DP provides an

estimate, for each given edge/region, of the likelihood that within parent-child dyad similarity (between a parent and their parent) is higher than stranger-child similarity (between a parent and another parent's child). Specifically, we computed the edge/region product vector (φ_i) from two sets of FC/GMV vectors $[X_i^{Child}], [X_i^{Parent}]$,

$$\varphi_i(f) = X_i^{Child}(f) * X_i^{Parent}(f), f = 1, 2, \ldots, M$$

where i indexes dyad, f indexes edge/region, and M is the total number of edges/regions in the entire FC/GMV vector. We can calculate φ_i between vectors of a child and another child's parent

$$\varphi_{ij}(f) = X_i^{Child}(f) * X_j^{Parent}(f), i \neq j$$

To compute the DP for all the dyads in a given dataset, we calculate an empirical probability

$$P_i(f) = \frac{P(|\varphi_{ji}(f) > \varphi_{ii}(f)| + |\varphi_{ij}(f) > \varphi_{ii}(f)|)}{2(N)}, N = number\ of\ dyads$$

A low $P_i(f)$ indicates a more discriminative edge/region. We can finally calculate DP of an edge/region across all children in a sample:

$$DP(e) = \sum_i \{-\ln(P_i(e))\}$$

If the parent-child dyad product was higher than the stranger-child product across all children in a sample, this corresponds to a high DP value, and the edge/region is helpful.

The group consistency measure, φ is simply the mean of φ_i for a given edge/region across all children. Edges/regions with high φ values are therefore high across all pairs of children and parents, and thus are not helpful.

For the analyses in Figure 2, we used whole-brain FC or GMV. For the analyses of data shown in Figure 3, we split the whole brain into 10 sub-networks and conducted the same analyses using FC or GMV within each sub-network. The definition of sub-networks was obtained from Horien et al.[67].

Testosterone and Child Behavior Checklist

We investigated the effects of testosterone and Child Behavior Checklist (CBCL) scores on the brain similarity of parent-child dyads. The detailed methods for data collection are described

elsewhere[63,64]. In the main analyses, we excluded dyads if either parent or child did not have
a score. We also excluded children who had an extremely high testosterone measurement
(more than mean + 1.5 s.t.d. Mean values of excluded children were 41.65 pg/mL [N=4] and
92.63 pg/mL [N=7] for children at age 11 and 13, respectively). After exclusion, the number of
dyads was 51 for testosterone at age 11 (4.01 ± 3.63 pg/mL, mean ± s.t.d), and 46 for
testosterone at age 13 (15.96 ± 17.83 pg/mL, mean ± s.t.d), 82 for CBCL at age 13 (6.55 ± 6.16,
mean ± s.t.d). There was no exclusion for CBCL age at 11 (10.48 ± 9.10, mean ± s.t.d).

1. *Testosterone:* The adolescents collected their salivary samples at home early in the
morning. In advance, both the adolescents and their primary parents were informed of
how to collect the adolescents' saliva using sample tubes. The adolescents tried it under
the guidance of the survey staff for practice. They were instructed not to collect the saliva
within a week after a tooth extraction or immediately after dental treatment to avoid
contamination with blood. They were also asked not to eat food after brushing their teeth
on the night before the saliva collection. They were instructed to rinse their mouth soon
after getting up and to make sure they were at their normal body temperature, and not to
have breakfast and not to brush teeth before the collection. Furthermore, they were asked
to wait for 20 min after the rinse and then to collect 4.5 ml of their saliva by passive drool
in sterilized tubes (1.5 ml/tube * 3 tubes) made of polypropylene (Nalgene™ General
Long-Term Storage Cryogenic Tubes, Thermo Fisher SCIENTIFIC, U.S.A.) within 60 min.
Salivary samples were collected in only one day, since high correlation among morning
salivary testosterone levels across days in adolescents was reported[85]. Salivary samples
were kept in household refrigerator freezers, delivered frozen to our laboratory, where the
weights were measured and tubes stored at minus 80 degrees C until the testosterone
levels were measured. The concentration of salivary testosterone was measured once by
liquid chromatography- tandem mass spectrometry (LC–MS/MS), which has become the
current standard[86]. All testosterone measurements were then square-eroot transformed
to better approximate a normal distribution prior to quantitative analyses.

2. *CBCL:* CBCL is a parental-report questionnaire used to screen children for behavioural
problems: there are 20 competence items and 120 items on behavioural and emotional
problems. The CBCL includes the following eight empirically-based syndrome scales: 1)
Aggressive Behavior, 2) Anxious/Depressed, 3) Attention Problems, 4) Rule-Breaking
Behavior, 5) Somatic Complaints, 6) Social Problems, 7) Thought Problems, and 8)
Withdrawn/Depressed, as well as summary scores reflecting "Internalization" and
"Externalization." We used the average scores of "Internalization" and "Externalization" in

the main results.

References

1.   de Graaf, P. M. & Kalmijn, M. Trends in the Intergenerational Transmission of Cultural
and Economic Status. *Acta Sociol.* **44**, 51–66 (2001).
2.   Deary, I. J., Spinath, F. M. & Bates, T. C. Genetics of intelligence. *Eur. J. Hum. Genet.*
14, 690–700 (2006).
3.   Grønhøj, A. & Thøgersen, J. Like father, like son? Intergenerational transmission of
values, attitudes, and behaviours in the environmental domain. *J. Environ. Psychol.*
29, 414–421 (2009).
4.   Kalmuss, D. The intergenerational transmission of marital aggression. *J. Marriage*
Fam. 11–19 (1984).
5.   Corak, M. Income inequality, equality of opportunity, and intergenerational mobility.
J. Econ. Perspect. **27**, 79–102 (2013).
6.   Demirkan, A. *et al.* Genetic risk profiles for depression and anxiety in adult and elderly
cohorts. *Mol. Psychiatry* **16**, 773–783 (2011).
7.   Miller, R. B. & Glass, J. Parent-child attitude similarity across the life course. *J.*
Marriage Fam. 991–997 (1989).
8.   Ho, T. C., Sanders, S. J., Gotlib, I. H. & Hoeft, F. Intergenerational Neuroimaging of
Human Brain Circuitry. *Trends Neurosci.* **39**, 644–648 (2016).
9.   Flint, J., Timpson, N. & Munafò, M. Assessing the utility of intermediate phenotypes
for genetic mapping of psychiatric disease. *Trends Neurosci.* **37**, 733–741 (2014).
10. Dubois, J. & Adolphs, R. Building a Science of Individual Differences from fMRI. *Trends*
Cogn. Sci. **20**, 1–19 (2016).
11. Biswal, B. B. *et al.* Toward discovery science of human brain function. *Proc. Natl.*
Acad. Sci. U. S. A. **107**, 4734–4739 (2010).
12. Finn, E. S. *et al.* Functional connectome fingerprinting: identifying individuals using
patterns of brain connectivity. *Nat. Neurosci.* **18**, 1–11 (2015).
13. Rosenberg, M. D. *et al.* A neuromarker of sustained attention from whole-brain
functional connectivity. *Nat. Neurosci.* **19**, 165–71 (2016).
14. Takagi, Y., Hirayama, J. ichiro & Tanaka, S. C. State-unspecific patterns of whole-brain
functional connectivity from resting and multiple task states predict stable individual
traits. *Neuroimage* **201**, 116036 (2019).
15. van den Heuvel, M. P., Stam, C. J., Kahn, R. S. & Hulshoff Pol, H. E. Efficiency of
functional brain networks and intellectual performance. *J. Neurosci.* **29**, 7619–24

698		(2009).
699	16.	Fox, M. D. & Greicius, M. Clinical applications of resting state functional connectivity.
700		*Front. Syst. Neurosci.* **4**, 19 (2010).
701	17.	Takagi, Y. *et al.* A neural marker of obsessive-compulsive disorder from whole-brain
702		functional connectivity. *Sci. Rep.* **7**, 7538 (2017).
703	18.	Takagi, Y. *et al.* A common brain network among state, trait, and pathological anxiety
704		from whole-brain functional connectivity. *Neuroimage* **172**, 506–516 (2018).
705	19.	Yahata, N. *et al.* A small number of abnormal brain connections predicts adult autism
706		spectrum disorder. *Nat. Commun.* **7**, (2016).
707	20.	Kanai, R. & Rees, G. The structural basis of inter-individual differences in human
708		behaviour and cognition. *Nat. Rev. Neurosci.* **12**, 231–242 (2011).
709	21.	Busch, V., Schuierer, G., Bogdahn, U. & May, A. Changes in grey matter induced by
710		training Newly honed juggling skills show up as a transient feature on a brain-imaging
711		scan . *Nature* **427**, 311–312 (2004).
712	22.	Duncan, J. *et al.* A neural basis for general intelligence. *Science (80-.).* **289**, 457–460
713		(2000).
714	23.	Okada, N. *et al.* Smaller anterior subgenual cingulate volume mediates the effect of
715		girls' early sexual maturation on negative psychobehavioral outcome. *Neuroimage*
716		**209**, 116478 (2020).
717	24.	Kasai, K. *et al.* Progressive decrease of left superior temporal gyrus gray matter
718		volume in patients with first-episode schizophrenia. *Am. J. Psychiatry* **160**, 156–164
719		(2003).
720	25.	Elliott, L. T. *et al.* Genome-wide association studies of brain imaging phenotypes in UK
721		Biobank. *Nature* **562**, 210–216 (2018).
722	26.	Feng, J. *et al.* Partitioning heritability analyses unveil the genetic architecture of
723		human brain multidimensional functional connectivity patterns. *Hum. Brain Mapp.*
724		(2020).
725	27.	Ge, T., Holmes, A. J., Buckner, R. L., Smoller, J. W. & Sabuncu, M. R. Heritability
726		analysis with repeat measurements and its application to resting-state functional
727		connectivity. *Proc. Natl. Acad. Sci.* **114**, 5521–5526 (2017).
728	28.	Moodie, C. A., Wisner, K. M. & MacDonald III, A. W. Characteristics of canonical
729		intrinsic connectivity networks across tasks and monozygotic twin pairs. *Hum. Brain*
730		*Mapp.* **35**, 5532–5549 (2014).
731	29.	Yang, Z. *et al.* Genetic and environmental contributions to functional connectivity
732		architecture of the human brain. *Cereb. cortex* **26**, 2341–2352 (2016).

30. Sinclair, B. *et al.* Heritability of the network architecture of intrinsic brain functional
connectivity. *Neuroimage* **121**, 243–252 (2015).

31. Xu, J. *et al.* Heritability of the effective connectivity in the resting-state default mode
network. *Cereb. cortex* **27**, 5626–5634 (2017).

32. Fu, Y. *et al.* Genetic influences on resting-state functional networks: A twin study.
Hum. Brain Mapp. **36**, 3959–3972 (2015).

33. van den Heuvel, M. P. *et al.* Genetic control of functional brain network efficiency in
children. *Eur. Neuropsychopharmacol.* **23**, 19–23 (2013).

34. Achterberg, M. *et al.* Distinctive heritability patterns of subcortical-prefrontal cortex
resting state connectivity in childhood: A twin study. *Neuroimage* **175**, 138–149
(2018).

35. Gao, W. *et al.* Intersubject variability of and genetic effects on the brain's functional
connectivity during infancy. *J. Neurosci.* **34**, 11288–11296 (2014).

36. Teeuw, J. *et al.* Genetic and environmental influences on functional connectivity
within and between canonical cortical resting-state networks throughout adolescent
development in boys and girls. *Neuroimage* **202**, 116073 (2019).

37. Glahn, D. C. *et al.* Genetic control over the resting brain. *Proc. Natl. Acad. Sci.* **107**,
1223–1228 (2010).

38. Sudre, G. *et al.* Estimating the heritability of structural and functional brain
connectivity in families affected by attention-deficit/hyperactivity disorder. *JAMA*
psychiatry **74**, 76–84 (2017).

39. Korgaonkar, M. S., Ram, K., Williams, L. M., Gatt, J. M. & Grieve, S. M. Establishing the
resting state default mode network derived from functional magnetic resonance
imaging tasks as an endophenotype: A twins study. *Hum. Brain Mapp.* **35**, 3893–3902
(2014).

40. Fornito, A. *et al.* Genetic influences on cost-efficient organization of human cortical
functional networks. *J. Neurosci.* **31**, 3261–3270 (2011).

41. Meda, S. a *et al.* Multivariate analysis reveals genetic associations of the resting
default mode network in psychotic bipolar disorder and schizophrenia. *Proc. Natl.*
Acad. Sci. U. S. A. **111**, E2066-75 (2014).

42. Adhikari, B. M. *et al.* Comparison of heritability estimates on resting state fMRI
connectivity phenotypes using the ENIGMA analysis pipeline. *Hum. Brain Mapp.* **39**,
4893–4902 (2018).

43. Colclough, G. L. *et al.* The heritability of multi-modal connectivity in human brain
activity. *Elife* **6**, 1–19 (2017).

44.   Bis, J. C., DeCarli, C., Smith, A. V & others. Common variants at 12q14 and 12q24 are
769   associated with hippocampal volume. *Nat Genet* **44**, 545–551 (2012).

770   45.   den Braber, A. *et al.* Heritability of subcortical brain measures: a perspective for
771   future genome-wide association studies. *Neuroimage* **83**, 98–102 (2013).

772   46.   Eyler, L. T. *et al.* Conceptual and data-based investigation of genetic influences and
773   brain asymmetry: a twin study of multiple structural phenotypes. *J. Cogn. Neurosci.*
774   **26**, 1100–1117 (2014).

775   47.   Blokland, G. A. M., de Zubicaray, G. I., McMahon, K. L. & Wright, M. J. Genetic and
776   environmental influences on neuroimaging phenotypes: a meta-analytical
777   perspective on twin imaging studies. *Twin Res. Hum. Genet.* **15**, 351–371 (2012).

778   48.   Kremen, W. S. *et al.* Genetic and environmental influences on the size of specific
779   brain regions in midlife: the VETSA MRI study. *Neuroimage* **49**, 1213–1223 (2010).

780   49.   Jansen, A. G., Mous, S. E., White, T., Posthuma, D. & Polderman, T. J. C. What twin
781   studies tell us about the heritability of brain development, morphology, and function:
782   a review. *Neuropsychol. Rev.* **25**, 27–46 (2015).

783   50.   Zhao, B. *et al.* Genome-wide association analysis of 19,629 individuals identifies
784   variants influencing regional brain volumes and refines their genetic co-architecture
785   with cognitive and mental health traits. *Nat. Genet.* **51**, 1637–1644 (2019).

786   51.   Ikram, M. A. *et al.* Common variants at 6q22 and 17q21 are associated with
787   intracranial volume. *Nat. Genet.* **44**, 539 (2012).

788   52.   Guadalupe, T. *et al.* Human subcortical brain asymmetries in 15,847 people
789   worldwide reveal effects of age and sex. *Brain Imaging Behav.* **11**, 1497–1514 (2017).

790   53.   Franke, B. *et al.* Genetic influences on schizophrenia and subcortical brain volumes:
791   large-scale proof of concept. *Nat. Neurosci.* **19**, 420–431 (2016).

792   54.   Hibar, D. P. *et al.* Novel genetic loci associated with hippocampal volume. *Nat.*
793   *Commun.* **8**, 1–12 (2017).

794   55.   Hibar, D. P. *et al.* Common genetic variants influence human subcortical brain
795   structures. *Nature* **520**, 224–229 (2015).

796   56.   der Meer, D. *et al.* Brain scans from 21,297 individuals reveal the genetic architecture
797   of hippocampal subfield volumes. *Mol. Psychiatry* 1–13 (2018).

798   57.   Wen, W. *et al.* Distinct genetic influences on cortical and subcortical brain structures.
799   *Sci. Rep.* **6**, 32760 (2016).

800   58.   Ge, T. *et al.* Multidimensional heritability analysis of neuroanatomical shape. *Nat.*
801   *Commun.* **7**, 1–10 (2016).

802   59.   Lee, T. H., Miernicki, M. E. & Telzer, E. H. Families that fire together smile together:

803   Resting state connectome similarity and daily emotional synchrony in parent-child

804   dyads. *Neuroimage* **152**, 31–37 (2017).

805 60. Quiñones-Camacho, L. E. *et al.* Parent–child neural synchrony: a novel approach to

806   elucidating dyadic correlates of preschool irritability. *J. Child Psychol. Psychiatry Allied*

807   *Discip.* (2019). doi:10.1111/jcpp.13165

808 61. Reindl, V., Gerloff, C., Scharke, W. & Konrad, K. Brain-to-brain synchrony in parent-

809   child dyads and the relationship with emotion regulation revealed by fNIRS-based

810   hyperscanning. *Neuroimage* **178**, 493–502 (2018).

811 62. Yamagata, B. *et al.* Female-specific intergenerational transmission patterns of the

812   human corticolimbic circuitry. *J. Neurosci.* **36**, 1254–1260 (2016).

813 63. Ando, S. *et al.* Cohort Profile: The Tokyo Teen Cohort study (TTC). *Int. J. Epidemiol.* **48**,

814   (2019).

815 64. Okada, N. *et al.* Population-neuroscience study of the Tokyo TEEN Cohort (pn-TTC):

816   Cohort longitudinal study to explore the neurobiological substrates of adolescent

817   psychological and behavioral development. *Psychiatry Clin. Neurosci.* **73**, 231–242

818   (2019).

819 65. Shen, X., Tokoglu, F., Papademetris, X. & Constable, R. T. Groupwise whole-brain

820   parcellation from resting-state fMRI data for network node identification.

821   *Neuroimage* **82**, 403–415 (2013).

822 66. Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L. & Petersen, S. E. Spurious but

823   systematic correlations in functional connectivity MRI networks arise from subject

824   motion. *Neuroimage* **59**, 2142–2154 (2012).

825 67. Horien, C., Shen, X., Scheinost, D. & Constable, R. T. The individual functional

826   connectome is unique and stable over months to years. *Neuroimage* **189**, 676–687

827   (2019).

828 68. Alexander-Bloch, A., Raznahan, A., Bullmore, E. & Giedd, J. The convergence of

829   maturational change and structural covariance in human cortical networks. *J.*

830   *Neurosci.* **33**, 2889–2899 (2013).

831 69. Sisk, C. L. & Zehr, J. L. Pubertal hormones organize the adolescent brain and behavior.

832   *Front. Neuroendocrinol.* **26**, 163–174 (2005).

833 70. Achenbach, T. M. *Child behavior checklist/4-18*. (VT: University of Vermont, 1991).

834 71. Petty, C. R. *et al.* The child behavior checklist broad-band scales predict subsequent

835   psychopathology: A 5-year follow-up. *J. Anxiety Disord.* **22**, 532–539 (2008).

836 72. Arain, M. *et al.* Maturation of the adolescent brain. *Neuropsychiatr. Dis. Treat.* **9**,

837   449–461 (2013).

838   73.   Valizadeh, S. A., Liem, F., Mérillat, S., Hänggi, J. & Jäncke, L. Identification of individual
839   subjects on the basis of their brain anatomical features. *Sci. Rep.* **8**, 1–9 (2018).

840   74.   Wachinger, C., Golland, P., Kremen, W., Fischl, B. & Reuter, M. BrainPrint: A
841   discriminative characterization of brain morphology. *Neuroimage* **109**, 232–248
842   (2015).

843   75.   Kumar, K., Toews, M., Chauvin, L., Colliot, O. & Desrosiers, C. Multi-modal brain
844   fingerprinting: A manifold approximation based framework. *Neuroimage* **183**, 212–
845   226 (2018).

846   76.   Gennatas, E. D. *et al.* Age-related effects and sex differences in gray matter density,
847   volume, mass, and cortical thickness from childhood to young adulthood. *J. Neurosci.*
848   **37**, 5065–5073 (2017).

849   77.   Váša, F. *et al.* Conservative and disruptive modes of adolescent change in human
850   brain functional connectivity. *Proc. Natl. Acad. Sci. U. S. A.* **117**, 3248–3253 (2020).

851   78.   Whitley, E., Gale, C. R., Deary, I. J., Kivimaki, M. & Batty, G. D. Association of maternal
852   and paternal IQ with offspring conduct, emotional, and attention problem scores:
853   Transgenerational evidence from the 1958 British birth cohort study. *Arch. Gen.*
854   *Psychiatry* **68**, 1032–1038 (2011).

855   79.   Harden, K. P., Kretsch, N., Tackett, J. L. & Tucker-Drob, E. M. Genetic and
856   environmental influences on testosterone in adolescents: Evidence for sex
857   differences. *Dev. Psychobiol.* **56**, 1278–1289 (2014).

858   80.   Grotevant, H. D. & Cooper, C. R. Individuation in family relationships. *Hum. Dev.* **29**,
859   82–100 (1986).

860   81.   Zuo, X. N. *et al.* Human Connectomics across the Life Span. *Trends Cogn. Sci.* **21**, 32–
861   45 (2017).

862   82.   Rosenberg, M. D., Casey, B. J. & Holmes, A. J. Prediction complements explanation in
863   understanding the developing brain. *Nat. Commun.* 1–13 (2018).
864   doi:10.1038/s41467-018-02887-9

865   83.   Honey, C. J. *et al.* Predicting human resting-state functional connectivity from
866   structural connectivity. *Proc. Natl. Acad. Sci. U. S. A.* **106**, 2035–2040 (2009).

867   84.   Gong, G., He, Y., Chen, Z. J. & Evans, A. C. Convergence and divergence of thickness
868   correlations with diffusion connections across the human cerebral cortex.
869   *Neuroimage* **59**, 1239–1248 (2012).

870   85.   Granger, D. A., Shirtcliff, E. A., Booth, A., Kivlighan, K. T. & Schwartz, E. B. The
871   'trouble' with salivary testosterone. *Psychoneuroendocrinology* **29**, 1229–1240
872   (2004).

873 86. Ketha, H., Kaur, S., Grebe, S. K. & Singh, R. J. Clinical applications of LC-MS sex steroid

874   assays: Evolution of methodologies in the 21st century. *Curr. Opin. Endocrinol.*

875   *Diabetes Obes.* **21**, 217–226 (2014).

Supplementary Figure 1: Overall patterns of edges/regions with high differential power (highly discriminative among subjects) and group consistency (highly similar among subjects) tended to be similar across different thresholds and across development (**(a)–(d)** child at age 11); **(e)–(h)** child at age 13). The figures show the results when the edges were thresholded at the 99.8th and 99.75th percentiles for FC, and regions were thresholded at the 85th, 80th percentiles for GMV. For each threshold, a circle plot (FC) and a bar graph (GMV) are shown, in which nodes are grouped according to anatomical location.

Supplementary Figure 2: Box plots of signal strength of each lobule for GMV. The red line indicates the

median. The bottom and top edges of the box indicate the 25th and 75th percentiles, respectively. The

crosses denote outliers, and the whiskers extend to the most extreme data points not considered outliers.

The brainstem (the right-most boxes in each figure) exhibited much lower signal strength compared with

the other lobules. Means ± s.e.m are shown.

Supplementary Figure 3: Accuracies using only low-movement subjects. We confirmed that qualitatively

similar results were obtained when we excluded children whose head movements were in the top 25%, at

both age 11 and age 13, resulting in the inclusion of 41.25% of the total sample.

Supplementary Figure 4: Between-network analyses using only the between-network edges. Statistical

significance was assessed by comparing the distributions for each network obtained through bootstrapping,

uncorrected.

Supplementary Figure 5: Accuracies obtained using conventional identification methods. Box plots of

accuracies using FC (top row) and GMV (bottom row) for children at age 11 (left column) and 13 (right

column) are shown. The bottom and top edges of the box indicate the 25th and 75th percentiles,

respectively. The crosses denote outliers, and the whiskers extend to the most extreme data points not

considered outliers. n.s. non-significant.

Supplementary Figure 6 For each sub-network (1–10: sub-networks shown in Figure 3a) and whole-brain

(ALL), we calculated Pearson's correlation coefficients between parent-child brain similarities defined by FC

and those defined by GMV across all parent-child dyads (red box). We found that they exhibited relatively

low correlations in general, suggesting that they contained independent information. Directly to the right

of these red boxes (grey box) are the results of the 1,000-times permutation testing.

Supplementary Figure 7: The number of top ranked networks for COMB (red bar), GMV (green bar), and

FC (blue bar) estimated via bootstrapping and null distribution (grey bar) estimated via permutation are

shown.