Examining the relationship between social cognition and neural synchrony during movies in children with and without autism

Lyons, KM[a,b,*], Stevenson, RA[a,b], Owen, AM[a,b,c], & Stojanoski, B[a,b]

a The Brain and Mind Institute, Western University, London ON, N6A 5B7, Canada.

b The Department of Psychology, Western University, London ON, N6A 5B7, Canada.

c The Department of Physiology and Pharmacology, Western University, London ON, N6A 5B7, Canada.

Corresponding author
Kathleen Lyons
Email: klyons8@uwo.ca
The Brain and Mind Institute, Department of Psychology,
The University of Western Ontario, London, Ontario, N6C 5B7, Canada

Abstract

Children who have been diagnosed with autism spectrum disorder (ASD) often show a

marked deficit in measures of social cognition. In autistic adults, measures of social

cognition have been shown to relate to differences in brain synchronization (as measured

by fMRI) when individuals are processing naturalistic stimuli, such as movies. However,

whether children with social impairments, with or without a diagnosis of ASD, differ in

their neural responses to movies has not yet been investigated. In the current study, neural

synchrony, measured using fMRI, was examined in three groups of children aged 7 to 12,

who differed with respect to scores on a measure of autistic traits associated with social

impairment and whether or not they had been diagnosed with ASD. While watching the

movie 'Despicable Me', those diagnosed with ASD had significantly less neural

synchrony in areas that have been previously shown to be associated with social

cognition (e.g. areas related to 'theory of mind'), and plot following (e.g. the lateral

prefrontal cortex), than those who did not have an ASD diagnosis. In contrast, two groups

who differed in their degree of social impairment, but did not have a diagnosis of ASD,

showed no significant differences in neural synchrony across the whole brain. These

results shed some light on how autistic traits may contribute to an individual's conscious

experience of the world, and how, for children with ASD, that experience may differ

markedly from that of those without ASD.

Keywords

Autism spectrum disorder, neural synchrony, fMRI, development, Theory of Mind, social

impairment

Introduction

Autism spectrum disorder (ASD)[1] is a complex developmental condition characterised by a variety of neurological and psychological features; however, the most prominent feature of ASD is a marked deficit in 'social cognition'. Social cognition refers to understanding what other people believe, how they will react in situations, and why they feel the way they do, and is a core element of successful human interactions. Autistic individuals perform poorly on tasks that assess social cognition, such as face perception (Spencer et al., 2011), perspective taking (Hamilton et al., 2009), and theory of mind (ToM), or the ability to attribute mental states to oneself and others (Pedreño et al., 2017). One of the most common tools to screen for deficits associated with ASD is the Social Responsiveness Scale, which measures aspects of social awareness, communication, and motivation (Constantino & Gruber, 2012).

The brains of autistic individuals often show differences when compared to those of typically-developing individuals. These include structural abnormalities (Barnea-Goraly et al., 2004; Brieber et al., 2007), functional differences during task-based fMRI (Bölte et al., 2008; Gilbert et al., 2008; Just et al., 2007; Mason et al., 2008; Solomon et al., 2009) and changes in resting-state functional connectivity (Cherkassky et al., 2006; Kana et al., 2015; Monk et al., 2009; Weng et al., 2010). Many of the brain regions that show differences in autistic individuals have been linked to ToM in healthy individuals, including the temporal parietal junction (Saxe & Kanwisher, 2003; Saxe & Wexler, 2005), the medial prefrontal cortex (Hartwright et al., 2013; Krause et al., 2012; Völlm et

---

[1]Abbreviations: ASD: Autism spectrum disorder; fMRI: functional magnetic resonance imaging; L-SRS: High SRS-2 score; SRS-2: Low SRS-2 score; H-SRS: Social responsiveness scale – revised; ToM: Theory of mind; Weschler Intelligence Scale for Children: WISC.

al., 2006), and the posterior superior temporal sulcus (Otsuka et al., 2009; Yang et al.,

2015).

Evidence has recently emerged that autistic adults process social information in

naturalistic, or 'real-life' contexts differently than typically-developing individuals.

Several studies have investigated social processing differences between those with and

without ASD by examining brain activity in response to watching movies (Bolton et al.,

2018; Byrge et al., 2015; Hasson et al., 2009; Salmi et al., 2013). Movie watching mimics

real-world experiences by requiring the viewer to integrate perceptual and cognitive

systems in order to follow the complexities of the plot. It is known that the brains of

healthy individuals become highly synchronized (or correlated) when viewing the same

movie (Hasson, Landesman, et al., 2008). This measure of synchronization across

different brains is termed *inter-subject correlation* and high levels of synchrony suggest

that individuals are experiencing the movie in much the same way. For example, Naci et

al. (2014) noted a high degree of synchrony in frontoparietal regions when healthy

individuals watched "Bang You're Dead!" by Alfred Hitchcock and this was shown to

relate to how suspenseful and engaging viewers found the movie. The brains of autistic

adults have been shown to be less synchronized than those of typically-developing adults

during movie watching, and synchrony across individuals tends to be more variable

(Bolton et al., 2018; Byrge et al., 2015; Hasson et al., 2009; Salmi et al., 2013). However,

this has not been examined in autistic children.

Richardson et al. (2018) have shown that in typically-developing children, those

with poorer social cognition have reduced synchrony during movie watching in areas

known to be involved with ToM, suggesting that lower synchrony in these areas may also

be a feature of autistic children. In the current study, this question was investigated in

three groups of children who differed with respect to their social impairment scores and

whether or not they had been diagnosed with ASD. Specifically, a data-driven approach

was used to examine differences in the degree of inter-subject correlation during movie

watching in children aged 7 to 12, who had either been diagnosed with ASD, or did not

have ASD but their scores on the Social Responsiveness Scale – revised (SRS-2)

indicated social impairments related to autistic traits, or did not have ASD and had typical

SRS-2 scores for their age.

On the basis of the existing literature, it was predicted that group differences

would emerge in inter-subject correlation within brain networks associated with social

cognition. Specifically, it was hypothesized that brain activity within both frontoparietal

(Naci et al., 2014), and the ToM networks (Richardson et al., 2018) would be less

synchronized in children without ASD with higher social impairment scores than those

with lower social impairment scores. Furthermore, it was hypothesized that the brains of

children with ASD would be the least synchronized of all, based on their known

impairments in many aspects of social cognition.

84
Methods

Dataset

Data was analyzed from the Healthy Brain Network Biobank collected by the

Child Mind Institute (described in Alexander et al., 2017), which is an ongoing initiative

to collect neuroimaging, medical, and behavioural data on 10,000 participants between

the ages of 5 to 21. The Chesapeake Institutional Review Board approved this study.

Detailed information on the dataset can be found at

http://fcon_1000.projects.nitrc.org/indi/cmi_healthy_brain_network/

Participants and data acquisition

94   The Healthy Brain Network Biobank used a community-referred recruitment

model to generate a heterogeneous and transdiagnostic sample. Briefly, recruitment

involved advertising the study to community members, educators, local care providers,

and parents who were on email lists or at events. Potential participants were screened,

and were excluded if there were safety concerns, impairments that would interfere with

the study procedure (such as being nonverbal or having an IQ of less than 66), and/or

medical concerns that could potentially impact brain related findings (for a full

description, see Alexander et al., 2017). The study protocol included, where possible, the

acquisition of T1 weighted anatomical MRI scans and functional MRI data acquired

while the participants watched a ten-minute clip of 'Despicable Me' (from 1:02:09 to

1:12:09). All MRI data was collected on a 3T Siemens scanner using a Siemens 32-

channel head coil. Functional images were acquired with a gradient-echo planar imaging

pulse sequence (TR =800 ms, TE =30 ms, Flip Angle =31 degrees, whole brain coverage

60 slices, resolution 2.4 x 2.4 mm^2). High-resolution T1-weighted MPRAGE structural

images were acquired in 224 sagittal (TR = 2500 ms, TE = 3.15 ms, resolution .8 x .8

mm^2).

110   From this database, participants were included in the current analysis if they were

between the ages of 7-12 and both anatomical and functional MRI data had been

successfully acquired. Everyone included in the current study had written consent

obtained from their legal guardians and written assent obtained from the participant.

Participants were not excluded based on their handedness. All participants also had scores

on the Social Responsiveness Scale Revised (SRS-2), which is a measure of social

reciprocity and communication associated with deficits in ASD (Constantino & Gruber,

2012). Specifically, the SRS-2 assesses deficits associated with social awareness, social

cognition, social communication, social motivation, and restrictive interests and repetitive

behavior, and is rated by parents or caregivers of the child. A score of 59 or below

suggests that the child does not have social impairments associated with ASD. A score

above 59 is suggestive of impairments in social functioning.

As part of this study, all participants completed a computerized version of the

Schedule for Affective Disorders and Schizophrenia - Children's version (KSADS) in

addition to the social responsiveness scale - revised (SRS-2). The KSADS is a semi-

structured diagnostic interview used to assess current and past psychopathology

according to the DSM-IV criteria, and is rated by a research clinician or social worker

(Alexander et al., 2017; Kaufman et al., 1997). Participants who were suspected to have

ASD were then assessed in person by a clinician. These participants were assessed using

the Autism Diagnostic Observation Schedule – 2nd edition (Lord, Rutter, DiLavore, Risi,

Gotham & Bishop, 2012) and the Autism Diagnostic Interview – Revised (Rutter, Le

Couteur & Lord, 2003) and those who met the relevant criteria were diagnosed with

ASD.

Participants were divided into three groups: The "Low SRS-2 score" (L-SRS)

group included those who had an SRS-2 score ≤ 59; the "High SRS-2 Score (H-SRS)"

group included participants who had an SRS-2 score of ≥ 60 (the ASD screener cut-off),

but were not diagnosed with ASD; and the Autism Spectrum Group (ASD) included

participants who were diagnosed with ASD by a clinician as part of the HBN protocol

(for details, see Table 1).

Because the groups differed with respect to sample size, age and sex, the L-SRS

and H-SRS groups were resampled to produce three demographically matched sub-

groups. Specifically, for each participant in the ASD group, an L-SRS and an H-SRS

individual who had the same sex and was closest in age (to the month) were selected for

inclusion where possible (see Table 1). This resulted in three groups of 28 participants,

ensuring sufficient power for acquiring reliable inter-subject correlation results (Pajula &

Tohka, 2016). The matched sample was used to statistically compare the groups in the

whole brain and network of interest analyses. All but one of the participants in the High

SRS-2 group were assessed in person by a clinician. All but one participant also

completed the Weschler Intelligence Scale for Children (WISC; Wechsler, 2014).

Table 1 Participant demographics. Means, standard deviations, and ranges of the ages,
SRS-2 scores, and the WISC full scale IQ, as well as the number of females and males
(F/M) are displayed for each group in the full and matched sample. The full sample of
participants was used to create the matched groups. The matched sample was then used
for all group comparisons (i.e. the whole brain analysis, the network of interest analysis,
and the percentage of synchronized cortex). Only the pairwise cluster-based analysis used
the full sample of participants.

		Group			Test of group differences
	Measure	L-SRS	H-SRS	ASD	
Full Sample:					
	N	64	34	28	
	Age	9.9 (1.7) 7.1 to 12.9	9.2 (1.6) 7.1 to 11.8	9.4 (1.5) 7.1 to 12.5	$F_{(2,123)} = 1.99$, $p = .141$
	Sex (F/M)	27/37	13/21	2/26	$X^2_{(2)} = 11.27$, $p = .003$
	SRS-2 scores	49.6(4.8)* 39 to 59	67.5 (6.0)* 60 to 84	76.6 (10.5)* 62 to 90	$F_{(2, 123)} = 180.98$, $p < .001$
	WISC full scale IQ	103 (17) 61 to 143	98 (16) 53 to 135	93 (18) 56 to 129	$F_{(2,122)} = 3.77$, $p = .026$
Matched Sample:					
	N	28	28	28	
	Mean Age	9.4(1.5) 7.0 to 12.6	9.6 (1.6) 7.2 to 11.8	9.4 (1.5) 7.1 to 12.5	$F_{(2,81)} = .155$, $P = .857$
	Sex (F/M)	3/25	7/21	2/26	$X^2_{(2)} = 4.08$, $p = .129$
	SRS-2	48.6 (4.8)* 41 to 58	67.2 (5.7)* 60 to 84	76.6 (10.5)* 62 to 90	$F_{(2,81)} = 101.79$, $p < .001$
	WISC full scale IQ	103 (18) 61 to 141	96 (15) 53 to 121	93 (18) 56 to 129	$F_{(2,80)} = 2.71$, $p = .073$

MRI pre-processing

For the current study, the MRI data were preprocessed and analyzed using the

Automatic Analysis (AA) toolbox (Cusack et al., 2015), SPM8, and in-house MATLAB

scripts. Pre-processing of functional data included motion correction (using six motion

parameters: left/right, anterior/posterior, superior/inferior, chin up/down, top of head

left/right, nose left/right), functional and structural scans were co-registered and

normalized to the Montreal Neurological Institute (MNI) template. Functional data were

then spatially smoothed using a Gaussian filter (8 mm kernel), and low-frequency noise

(e.g., drift) was removed by high-pass filtering with a threshold of 1/128 Hz. The data

was denoised using Bandpass filter regressors, with cerebrospinal fluid, white matter

signals, motion parameters, their lag-3 2^{nd}-order volterra expansion (Friston et al., 2000),

and "spikes" (based on mean signal variance across volumes) as nuisance regressors.

Exploratory whole brain synchronization

To determine the degree of synchronization separately for each group, the degree

of inter-subject correlation across the whole brain was calculated using a leave-one-out

approach using the matched sample. That is, the pre-processed time course of every voxel

was correlated (Pearson and then Fisher z-transformed) between each participant and the

mean time course of every voxel from the rest of the group (N-1). A one-sample t-test

was calculated on the resulting individual brain-wide correlation values. Multiple

comparisons were corrected with a false discovery rate (FDR) of .05 to generate group

maps of significantly correlated voxels. To identify where in the brain inter-subject

correlation differences existed between the three groups, t-tests were performed on the

correlation values at each voxel derived for all of the individuals within each group.

Multiple comparisons were corrected with an FDR of .05.

Network of interest inter-subject correlation

The degree of synchronization within eight previously defined functional

networks was calculated. To address our specific hypotheses, a map for the ToM network

was used (Dufour et al., 2013) as well as the frontoparietal network from the Yeo et al.,

(2011) parcellation. Six additional networks (Visual, Dorsal Attention, Ventral Attention,

Somatomotor, Limbic, Default Mode Network) from Yeo et al. (2011) were also included

in an exploratory analysis to examine potential differences in other areas of the brain. The

8 network parcellations are displayed in Figure 1. Similar to the whole brain inter-subject

correlation analysis, the intra-group inter-subject correlation for each of these eight

networks was calculated using a leave-one-out approach using the matched sample.

Specifically, the time course of each network (based on the average time course of each

voxel within the network) for each participant was correlated with the average time

course of each network for the remaining participants in the group, minus that participant

(N-1). Finally, we used a general linear model to determine if group membership was a

significant predictor of intra-group synchronization across the 8 networks. The model

included inter-subject correlation values as the predicted variable and group as the

predictor variable. This was done separately for each network. The networks that showed

a significant effect of group were followed up with Welch t-tests (all results were FDR

corrected to .05).

To better understand the results from the intra-group analysis, the degree of inter-

group inter-subject correlation was then calculated, by taking the mean time course for

each individual in one group and correlating it with the mean of the two other groups.

This generated a correlation value that reflected how similar each participant's time

course was to the two other groups. For instance, we calculated how correlated each ASD

participant was to the mean of the other two groups. Finally, we calculated three separate

general linear models to determine if participants correlated significantly more with their

own group than the mean of the other two groups. The networks that showed a significant

effect of group were followed up with Welch t-tests (all results were FDR corrected to

.05).

Percent synchronization across the cortex

The percent of significant voxels across the cortex was calculated, for descriptive

purposes, to quantify the number of synchronized voxels common across all individuals

in each of the three matched sample groups. To calculate the total percentage of cortex

that was synchronized, the number of voxels that were significant per group were divided

by the total number of voxels in the brain. To calculate the total percentage of each

network that was synchronized, the number of voxels that were significant per group

were divided by the total number of voxels in the network of interest.

229
Figure 1 Network of interests used to parcellate the brain. a) ToM network parcellation. The ToM
network (displayed in red) included regions in the dorsal, ventral, and lateral medial prefrontal cortex,
bilateral temporal parietal junction, precuneus, and right superior temporal sulcus (Dufour et al.,
2013). b) The seven network parcellation from Yeo et al. (2011) contained 1) the frontoparietal
network (displayed in orange) which includes the lateral prefrontal cortex, medial cingulate,
intraparietal sulcus, and inferior temporal gyrus, 2) the visual network (displayed in dark blue) which
encompasses the visual cortex, 3) the somatomotor network (displayed in light blue) which includes
the motor cortex, premotor cortex, and postcentral gyrus, 4) the dorsal attention network (displayed in
dark green) which includes the frontal eye fields, precentral ventral frontal cortex, middle temporal
area, and intraparietal sulcus, 5) the ventral attention network (displayed in light green) which includes
the dorsal anterior prefrontal cortex, and anterior and posterior cingulate, 6) the limbic network
(displayed in mustard) which includes the temporal pole and orbital frontal cortex, 7) the default mode
network (displayed in red) which includes the dorsal medial prefrontal cortex, temporal parietal
junction, postcentral gyrus, precuneus, the superior temporal sulcus, the posterior cingulate cortex, and
retrosplenial cortex.

Cluster-based inter-subject correlation analysis

To explore the relationship between SRS-2 scores (as a continuous variable) and

neural synchrony, pairwise correlations were calculated between each participant and that

of every other participant in the ToM and frontoparietal networks. This was done by

calculating the mean time course (i.e. the mean activation across each voxel in the

network for each time point) in both networks for each participant, and then correlating it

with every other participant's mean time course. Because SRS-2 scores were skewed

(upwards) in the ASD and H-SRS groups, this analysis included all participants (N =

126), rather than the smaller matched groups. These pairwise correlations were then

plotted in a matrix by ranking each participant by their SRS-2 score (from low to high)

for descriptive purposes. Finally, a clustering analysis was conducted to determine

whether groups of participants could be identified based solely on their neural

synchronization, rather than group membership or SRS-2 scores. To do this, a k-means

clustering algorithm was used to group together participants using the time series of

neural activity in the ToM and frontoparietal networks. The MATLAB evalclusters

function was used to identify the optimal number of clusters based on the variance in the

data using the Calinski-Harabasz Index computed over 1000 iterations to minimize the

fitting parameter. Based on the groupings generated from this cluster analysis, a logistic

regression analysis was computed to investigate which factors (SRS-2 total and subscale

scores, age, sex, and social impairment group membership) best predicted the cluster-

generated groupings.

267
268
269

270 **Results**
271
There was a total of 267 eligible participants who met the inclusion criteria (see

Methods). Of this sample, 141 participants were removed because of excessive motion,

defined as large "spikes", or significant fluctuations in signal intensity (greater than 3

standard deviations of the mean), in at least 25% of the data.

There was a significant difference between the three groups in terms of SRS-2

scores ($F_{(2,81)}$ = 101.76, p < .001) and post-hoc t-tests showed that the H-SRS group had

significantly higher scores than the L-SRS group ($t_{(51.9)}$ = 12.96, p < .001) and had

significantly lower scores than the ASD group ($t_{(42.54)}$ = 4.12, p < .001). There were no

significant differences between the groups on the WISC full scale IQ scores ($F_{(2,80)}$ =

2.71, p = .073), or any of the WISC subscales except for working memory; ($F_{(2,80)}$ = 3.29,

p = .042). The ASD group had significantly lower working memory scores compared to

the L-SRS group ($t_{(52.10)}$ = 2.35, p = .023) but not the H-SRS group ($t_{(50.24)}$ = 1.05, p =

.30).

Differences in correlated motion within each group were examined, in order to

ensure that this did not inflate the inter-subject correlation results. Correlated motion was

calculated separately for each group, by taking each participant's 6 motion parameters for

each frame and correlating the time course with that of the mean of the rest of the group

(N-1). No significant differences were found between the groups in their degree of

correlated motion ($F_{(2,81)}$ = .181, p = .835).

Exploratory whole brain synchronization

Whole brain synchronization was characterized in the three groups. All groups

showed significant synchronization in the auditory and visual areas (Figure 2a). In fact,

synchronization in these areas was stronger than in any other brain areas, replicating

previous inter-subject correlation findings during movie watching (Hasson et al., 2008).

The H-SRS and L-SRS groups also showed significant inter-subject correlation in areas

associated with ToM and executive processing, including parts of the right and left

temporal parietal junction, the precuneus, the intraparietal sulcus, the superior parietal

lobe, and portions of the medial and lateral prefrontal cortex. In contrast, the ASD group

had very little significant inter-subject correlation outside of visual and auditory areas

(see Figure 2a, bottom row).

Next, whole brain contrasts were conducted (Figure 2b) to examine whether the

magnitude of synchronization differed between the three groups. When the L-SRS group

was contrasted to the H-SRS group, only tiny areas of difference were observed after

multiple comparisons corrections, in the inferior temporal gyrus (MNI coordinates $_{X, Y, Z}$

= -46, -37, -17, $t_{(54)} = 4.61$, $p_{corrected} = .030$), and white matter (see Figure 2b, top row).

The L-SRS group showed significantly greater inter-subject correlation than the ASD

group (Figure 2b, middle row) in the bilateral temporal parietal junction (MNI

coordinates (left) = -57, -61, 30, $t_{(54)} = 3.97$, $p_{corrected} = .011$, MNI coordinates (right) = 49,

-67, 31, $t_{(54)} = 5.07$, $p_{corrected} = .002$), precuneus (MNI coordinates = 4, -51, 41, $t_{(54)} = 4.14$,

$p_{corrected} = .009$), right superior temporal sulcus (MNI coordinates =60, -11, -16, $t_{(54)} =$

4.47, $p_{corrected} = .005$), right hippocampus (MNI coordinates = 32, -14, -19, $t_{(54)} = 3.52$,

$p_{corrected} = .026$), and in regions of the lateral (MNI coordinates = 39, 54, -9, $t_{(54)} = 3.61$,

$p_{corrected} = .022$), and the right medial prefrontal cortex (MNI coordinates = 22, 42, 37, $t_{(54)}$

= 3.89, $p_{corrected} = .014$). The H-SRS group had significantly greater synchronization than

the ASD group in the precuneus (MNI coordinates = -3, -55, 45, $t_{(54)} = 4.00$, $p_{corrected} =$

.017), right hippocampus (MNI coordinates = 28, -5, -21, $t_{(54)}$ = 3.37, $p_{corrected}$ = .043), and

in regions of the lateral (MNI coordinates = 46, 44, 12, $t_{(54)}$ = 6.30, $p_{corrected}$ < .001), and

medial prefrontal cortex (MNI coordinates = -5, 65, -7, $t_{(54)}$ = 4.27, $p_{corrected}$ = .012)

(Figure 2b, bottom row).

Figure 2 Exploratory whole brain inter-subject correlation analysis. **a)** Voxels displayed in red
showed significant inter-subject correlation during movie watching in the L-SRS group. Voxels
displayed in violet showed significant inter-subject correlation in the H-SRS group. Voxels displayed
in blue showed significant inter-subject correlation in the ASD group. All p values were FDR
corrected to an alpha of .05. **b)** Whole brain contrasts were calculated by conducting one-tailed t-tests
on the inter-subject correlation values between each group (p values corrected to an FDR of .05).
Voxels displayed in yellow showed significantly greater inter-subject correlation values based on this
contrast, voxels displayed in red show the ToM network parcellation. c) Voxels displayed in yellow
showed significantly greater inter-subject correlation values based on the same contrast displayed in b,
overlaid on top of the Yeo et al. (2011) 7-network parcellation. (Frontoparietal = orange, Visual =
dark blue, Somatomotor = light blue, Dorsal attention = dark green, Ventral attention = light green,
Limbic = mustard, Default mode = red).
334
335   *Network based synchronization*

336   Group differences in the magnitude of intra-group synchronization revealed a

main effect of group in the ToM ($F_{(2,81)} = 4.94$, $p = .009$) and the limbic ($F_{(2,81)} = 3.93$, p

$= .023$) networks (Figure 3), but not in any of the others examined, including the

frontoparietal network ($F_{(2,81)} = 2.02$, $p = .140$, Cohen's d ranged from .037 to .476).

Post-hoc analyses of neural synchronization revealed that the ASD group had

significantly lower inter-subject correlation values compared to the L-SRS group within

the ToM ($t_{(50.11)} = 3.50$, $p_{corrected} = .006$, Cohen's d = .934) and limbic networks ($t_{(50.00)} =$

2.48, $p_{corrected} = .044$, Cohen's d = .664). They also had significantly lower inter-subject

correlation values compared to the H-SRS group in the limbic network ($t_{(50.21)} = 2.18$,

$p_{corrected} = .044$, Cohen's d = .631), although differences in inter-subject correlation just

failed to meet the corrected alpha level in the ToM network ($t_{(52.33)} = 2.36$, $p_{corrected} =$

.0504, Cohen's d = .584). Moreover, no significant differences in inter-subject correlation

were observed between the L-SRS and H-SRS groups within the ToM ($t_{(45.72)} = .488$,

$p_{corrected} = .628$, Cohen's d = .130) or limbic networks ($t_{(45.21)} = .417$, $p_{corrected} = .628$,

Cohen's d = .111).

Figure 3 Intra-group network of interest analysis. Mean inter-subject correlation, based on the leave one out correlation analysis conducted separately for each group, is displayed as dots for each participant in the eight networks. Boxplots indicate the median inter-subject correlation value and interquartile range for each group (blue = L-SRS, yellow = H-SRS, red = ASD). The ASD group had significantly lower inter-subject correlation in the limbic and ToM networks compared to the L-SRS group. The ASD group also had significantly lower inter-subject correlation in the limbic network compared to the H-SRS group, while in the ToM network this difference narrowly missed statistical significance (corrected p value = .0504). The groups did not differ significantly in any of the six other networks.

An inter-group inter-subject correlation network analysis was performed to

investigate whether individuals in one group had significantly greater neural

synchronization with their own group than that of the other two groups. The results

revealed that the degree of inter-subject correlation was not significantly different

between any of the groups in any of the examined networks, including the frontoparietal

and ToM networks.

Percent synchronization across the cortex

When looking at the percentage of synchronized voxels across the whole brain,

the ASD group had nearly one-third less (38%) than the L-SRS (56%) and H-SRS (52%)

groups (see Figure 4). The percentage of significant voxels in each of the eight networks

of interest was also calculated (see Figure 4). The difference in percentage across the

whole brain between the groups was not accounted for by less synchronization in any one

network; rather, the ASD group had fewer synchronized voxels in every network,

including in the ToM and frontoparietal networks.

376
377 **Figure 4** Percentage of correlated voxels. The percent of significant voxels across the cortex was
378 calculated, for descriptive purposes, to quantify the number of synchronized voxels common across all
379 individuals in each of the three matched sample groups. This was calculated by dividing the number of
380 voxels with significant inter-subject correlation by the total number of voxels in the whole brain or
381 network for each group separately (blue = L-SRS, yellow = H-SRS, red = ASD).
382
383
384
385

388   *Cluster-based inter-subject correlation analysis*

389   To explore whether SRS-2 scores predicted inter-subject correlation values when

used as a continuous measure (instead of a categorical variable), pairwise inter-subject

correlations were calculated between each participant (N=126) in the frontoparietal and

ToM networks. The entire sample was used so that the SRS-2 scores were normally

distributed and to increase statistical power. Pairwise correlations were conducted to

reduce any influence the groupings may have had on the mean time course originally

used to calculate inter-subject correlation. For instance, if those with low SRS-2 scores

and those with high SRS-2 scores both correlated with their own group to a similar

degree, but the pattern of activations was different, using these groupings would

obfuscate any differences. For descriptive purposes, the matrix of pairwise correlation

values was plotted by ranking each participant by their SRS-2 score, from low to high

(see Figure 5a). A k-means clustering analysis was conducted on the pairwise correlations

to explore potential factors that predicted groups of participants who have the most

similar degree of synchrony in these two networks. The best fit was achieved by dividing

the data into two clusters in both the frontoparietal and ToM networks; cluster 1 included

individuals with similar neural responses to the movie (large positive correlations) and

cluster 2 included individuals with unrelated neural responses to the movie (Figure b).

Moreover, there was also a large overlap between the participants who were in cluster 1

in the ToM and frontoparietal networks. Specifically, of the 58 participants who had high

similarity in the ToM network (cluster 1), 45 of them also had high similarity in the

frontoparietal network.

Figure 5 Pairwise inter-subject correlations. Yellow squares indicate a higher positive correlation (i.e. high similarity in time series), blue squares indicated a low or negative correlation (i.e. low similarity in time series). **a)** Pairwise correlations in time series in the frontoparietal and ToM networks between each pair of participants are ordered by SRS-2 scores (from low to high). **b)** Pairwise correlations in time series are ordered based on the K-means analysis in the frontoparietal and ToM networks. Black boxes show cluster 1 (the high similarity group) for each network.

Logistic regression was run to determine whether the probability of being in

cluster 1 versus cluster 2 could be predicted by age, sex, full scale IQ, SRS-2 total and

subscales, or group membership (i.e., L-SRS, H-SRS and ASD). None of these factors

significantly predicted cluster membership in the frontoparietal network. However, in the

ToM network, group membership significantly predicted cluster membership. Cluster 1

comprised 35 participants (60%) in the L-SRS group, 17 individuals (29%) from the H-

SRS groups, and 6 individuals (10%) who were diagnosed with ASD. In contrast, cluster

2 consisted of 29 individuals (45%) from the L-SRS group, 17 individuals (20%) from

the H-SRS group, and 22 individuals (35%) diagnosed with ASD. There were

significantly more participants from the ASD group in cluster 2 than in cluster 1 in the

ToM network ($X^2_{(1)} = 7.5$, p = .006), while there was no significant difference in the

number of H-SRS participants between the two clusters ($X^2_{(1)} = .11$, p = .73), and

although there were more L-SRS participants in cluster 1, this difference did not reach

significance ($X^2_{(1)} = 3.25$, p = .072).

432
433   **Discussion**
434
In the current study, a group of ASD participants had significantly less neural

synchronization when watching a movie compared to the L-SRS and H-SRS groups

across the whole brain, including the ToM and limbic networks, as well as the lateral and

medial prefrontal cortex. These regions have been shown previously to be associated with

elements of 'plot following' during movie watching (Hasson, Furman, et al., 2008;

Hasson, Landesman, et al., 2008; Naci et al., 2014; Nguyen et al., 2019), suggesting that

the children in the ASD group were experiencing the movie qualitatively differently than

the participants in the other two groups. These results, in particular the fact that the ToM

network was less synchronized in the ASD group, are intriguing given that regions within

this network are associated with social cognition (Dufour et al., 2013; Mills et al., 2014;

Richardson et al., 2018; Rilling et al., 2004), which is known to be affected in ASD

(Hamilton et al., 2009; Pedreño et al., 2017; Spencer et al., 2011). While aspects of social

cognition are usually discussed in the context of inter-personal relationships, they are also

essential components of movie-watching, allowing one to become immersed in the plot

by taking the perspective of the characters appropriately, understanding their motives,

and following their verbal and nonverbal communication cues. Yeshurun et al. (2017)

have reported previously that manipulating an individual's understanding of a plot

reduces neural synchrony in ToM regions, including the precuneus, temporal parietal

junction, and medial prefrontal cortex. Thus, these findings support the idea that autistic

children process social stimuli in a distinct way, as they have different neural responses in

the ToM network during a movie, when compared to children without ASD.

It is also interesting that participants in the ASD group had significantly less

synchrony in the lateral prefrontal cortex, a region within the frontoparietal network,

when compared to those in the other two groups. Understanding a complex narrative

(such as a movie's plot) requires a viewer to remember previous events, pay attention to

what is currently happening, make predictions about the future consequences of current

events, and integrate this information over time, all of which depends on frontoparietal

executive processing (Naci et al., 2014). In previous studies, reduced synchrony in this

network has been associated with 'losing the plot' during deep sedation (Naci et al.,

2018), and in patients with severe brain damage (Naci et al., 2014). Thus, this decrease in

inter-subject correlation in the lateral prefrontal cortex may suggest that participants in

the ASD group are also failing to grasp elements of the plot in the way that the other

participants do.

Despite finding that inter-subject correlation was reduced in prefrontal regions

using a whole brain analysis, no differences in the degree of inter-subject correlation

were found in the frontoparietal network when a network of interest analysis was used.

One potential reason is that the parcellation used for the frontoparietal network was based

on adult data and may not accurately capture this network in children. Previous work has

shown that the frontoparietal network continues to develop into early adulthood (Baum et

al., 2017; Peters et al., 2016), and so the parcellation masks from Yeo et al. (2011) may

have led us to average neural activity from regions that are not yet fully integrated in

children.

While not part of our hypotheses, it is interesting that the ASD group showed less

inter-subject correlation in the right hippocampus in the whole brain analysis as well as in

the limbic network, when examined using the parcellation by Yeo et al. (2011). Similar

findings have been reported in autistic adults watching movies (Byrge et al., 2015).

Moreover, Chen et al. (2017) found that, in healthy adults, the degree of inter-subject

correlation within the hippocampus during movie watching predicted events that were

later recalled, although this has not been examined during development. Nevertheless,

long-term memory deficits have been reported in ASD; specifically, autistic individuals

perform worse on episodic, but not semantic, memory tasks (Crane & Goddard, 2008;

Lind, 2010)

Contrary to our hypothesis, no meaningful differences in neural synchrony were

found between the L-SRS and H-SRS groups. This contrasts with the results of

Richardson et al., (2018) who found that social cognition in typically-developing children

was related to the degree of inter-subject correlation within the ToM network during

movie-watching. One potential reason for this difference is that Richardson et al., (2018)

calculated inter-subject correlation based on how similar each child's time course was to

a group of adults watching the same movie, whereas in the current study, inter-subject

correlation was calculated by correlating each participant's time course to the mean of

their own group. Moreover, the measure of social cognition used by Richardson et al.

(2018) focused specifically on comprehension of a social narrative, which has many

things in common with how people follow the plot of a movie. It is perhaps not surprising

then, that the two things correlated. In the current study, a measure of social impairment

was used – the SRS-2, which measures an individual's motivation to engage in social

interactions, their use of social communication, and their ability to understand social cues

(Constantino & Gruber, 2012). Thus, while the H-SRS and L-SRS groups differed in

terms of their social impairments as measured by the SRS-2 scale, these mechanisms may

be unrelated, or only moderately related, to those that are involved in plot following.

Moreover, it is also possible that creating categorical groups based on the SRS-2 scores

may have obscured subtle differences in individuals with differing levels of social

impairment. To investigate this possibility, the exploratory pairwise correlation analysis

was conducted, which found that SRS-2 scores as a continuous measure did not predict

whether participants had similar patterns of neural activity in the ToM or frontoparietal

networks. Taken together, these results suggest that it is only when social impairment is

in the clinical range, as is seen in ASD, that differences in conscious processing of

naturalistic stimuli emerge.

As a group, autistic participants had less inter-subject correlation compared to

those without ASD, but these differences did not apply uniformly to each individual. The

clustering analysis indicated that the majority of ASD participants had low similarity in

their time courses compared to all other participants. However, six out of 28 of those

diagnosed with ASD clustered with the 'high similarity' group (comprising about 10% of

the group) according to their synchronization in the ToM network. Using a similar

clustering analysis, Byrge et al., (2015) found that in a sample of 17 high functioning

autistic adults, five showed idiosyncratic patterns of inter-subject correlation compared to

typically-developing individuals, while the other 12 clustered with the control group.

Moreover, they found that these five individuals were significantly worse than the control

group and the other 12 ASD participants, when asked to explain elements of a movie plot.

Together, these findings suggest that lower synchronization during movie-watching may

be common, but not a uniform characteristic of either autistic children or adults. Indeed,

heterogeneity in clinical features, cognitive profiles, and differing genetic and

environmental risk factors has plagued research in ASD (Betancur, 2011; Jeste &

Geschwind, 2014; Lenroot & Yeung, 2013). For example, within the neuroimaging

literature, some studies have reported underconnectivity across the brains of autistic

individuals (Cherkassky et al., 2006; Di Martino et al., 2014; von dem Hagen et al.,

2013), while others find hyperconnectivity (Supekar et al., 2013; Uddin et al., 2010,

2013).

Finally, it is important to keep in mind the exploratory nature of the current study

when interpreting these findings. This is a step towards a better understanding of how

children with and without ASD process naturalistic stimuli, but replication and further

investigation is needed to better understand the nature of the differences observed. For

instance, one potential mechanism underlying our results could be that participants in the

ASD group had more variable neural responses to the movie. However, it would be

valuable for future studies to directly examine if more variable neural responses to

movies are driving reduced neural synchronization in those diagnosed with ASD.

Additionally, a major limitation of this study is that no memory test, or measure of how

well the movie clip was understood, was collected. A behavioral measure of movie

comprehension may help to explain the nature of the neural differences observed in this

study. It is possible that individuals were attending to different features of the movie,

which has been shown to influence the degree of neural synchrony (Nguyen et al., 2019),

although previous work has confirmed that movies similar to 'Despicable me' maintain

the viewers' attention (Hasson, Landesman, et al., 2008; Naci et al., 2015). It is also

unlikely that participants were asleep during the movie, as most of the visual network was

synchronized across the three groups during the movie, which is not observed when

individuals are sedated (Naci et al., 2018).

Conclusion

552   In sum, the current results suggest that autistic children, as a group, process

movies in a unique way compared to those without ASD. Interestingly, a minority of

these children had time courses that were highly correlated with a group of children

without ASD in the ToM network. Future research should investigate factors that underlie

this heterogeneity, as this may be one avenue to better understand how autistic

individuals process the world around them.

Author contributions

Conceptualization, writing–reviewing and editing, K.M.L., R.A.S, A.M.O, B.S.;

Methodology, Formal analysis, K.M.L., B.S.; writing–original draft preparation, K.M.L.,

B.S., A.M.O; funding acquisition, A.M.O., B.S

Acknowledgements

The authors thank the Child and Mind Institute for designing and collecting data for the

Healthy Brain Biobank.

Funding

This research was funded by the Canada Excellence Research Chairs Program (#215063),

the Canadian Institutes of Health Research (#408004), and the Natural Sciences and

Engineering Research Council of Canada (#390057).

Conflict of interest

The authors have no conflicts of interest to declare.

References

Alexander, L. M., Escalera, J., Ai, L., Andreotti, C., Febre, K., Mangone, A., ... & Litke,

S. (2017). The Healthy Brain Network Biobank: An open resource for

transdiagnostic research in pediatric mental health and learning disorders.

Scientific data, *4*, 170181. https://doi.org/10.1038/sdata.2017.181.

Barnea-Goraly, N., Kwon, H., Menon, V., Eliez, S., Lotspeich, L., & Reiss, A. L. (2004).

White matter structure in autism: Preliminary evidence from diffusion tensor

imaging. *Biological Psychiatry*, *55*(3), 323–326.

https://doi.org/10.1016/j.biopsych.2003.10.022

Baum, G. L., Ciric, R., Roalf, D. R., Betzel, R. F., Moore, T. M., Shinohara, R. T., Kahn,

A. E., Vandekar, S. N., Rupert, P. E., Quarmley, M., Cook, P. A., Elliott, M. A.,

Ruparel, K., Gur, R. E., Gur, R. C., Bassett, D. S., & Satterthwaite, T. D. (2017).

Modular Segregation of Structural Brain Networks Supports the Development of

Executive Function in Youth. *Current Biology*, *27*(11), 1561-1572.e8.

https://doi.org/10.1016/j.cub.2017.04.051

Betancur, C. (2011). Etiological heterogeneity in autism spectrum disorders: More than

100 genetic and genomic disorders and still counting. *Brain Research*, *1380*, 42–

77. https://doi.org/10.1016/j.brainres.2010.11.078

Bölte, S., Hubl, D., Dierks, T., Holtmann, M., & Poustka, F. (2008). An fMRI-study of

locally oriented perception in autism: Altered early visual processing of the block

design test. *Journal of Neural Transmission*, *115*(3), 545–552.

https://doi.org/10.1007/s00702-007-0850-1

Bolton, T. A. W., Jochaut, D., Giraud, A.-L., & Ville, D. V. D. (2018). Brain dynamics in

ASD during movie-watching show idiosyncratic functional integration and

segregation. *Human Brain Mapping*, *39*(6), 2391–2404.

https://doi.org/10.1002/hbm.24009

Brieber, S., Neufang, S., Bruning, N., Kamp-Becker, I., Remschmidt, H., Herpertz-

Dahlmann, B., Fink, G. R., & Konrad, K. (2007). Structural brain abnormalities in

adolescents with autism spectrum disorder and patients with attention

deficit/hyperactivity disorder. *Journal of Child Psychology and Psychiatry*,

48(12), 1251–1258. https://doi.org/10.1111/j.1469-7610.2007.01799.x

Byrge, L., Dubois, J., Tyszka, J. M., Adolphs, R., & Kennedy, D. P. (2015). Idiosyncratic

Brain Activation Patterns Are Associated with Poor Social Comprehension in

Autism. *Journal of Neuroscience*, *35*(14), 5837–5850.

https://doi.org/10.1523/JNEUROSCI.5182-14.2015

Chen, J., Leong, Y. C., Honey, C. J., Yong, C. H., Norman, K. A., & Hasson, U. (2017).

Shared memories reveal shared structure in neural activity across individuals.

Nature Neuroscience, *20*(1), 115–125. https://doi.org/10.1038/nn.4450

Cherkassky, V. L., Kana, R. K., Keller, T. A., & Just, M. A. (2006). Functional

connectivity in a baseline resting-state network in autism. *NeuroReport*, *17*(16),

1687–1690. https://doi.org/10.1097/01.wnr.0000239956.45448.4c

Constantino, J. N., & Gruber, C. P. (2012). *Social Responsiveness Scale–Second Edition*

(SRS-2). Western Psychological Services.

Crane, L., & Goddard, L. (2008). Episodic and Semantic Autobiographical Memory in

Adults with Autism Spectrum Disorders. *Journal of Autism and Developmental*

Disorders, *38*(3), 498–506. https://doi.org/10.1007/s10803-007-0420-2

Cusack, R., Vicente-Grabovetsky, A., Mitchell, D. J., Wild, C. J., Auer, T., Linke, A. C.,

& Peelle, J. E. (2015). Automatic analysis (aa): Efficient neuroimaging workflows

and parallel processing using Matlab and XML. *Frontiers in Neuroinformatics*, *8*.

https://doi.org/10.3389/fninf.2014.00090

Di Martino, A., Yan, C.-G., Li, Q., Denio, E., Castellanos, F. X., Alaerts, K., Anderson, J.

S., Assaf, M., Bookheimer, S. Y., Dapretto, M., Deen, B., Delmonte, S., Dinstein,

I., Ertl-Wagner, B., Fair, D. A., Gallagher, L., Kennedy, D. P., Keown, C. L.,

Keysers, C., … Milham, M. P. (2014). The autism brain imaging data exchange:

Towards a large-scale evaluation of the intrinsic brain architecture in autism.

Molecular Psychiatry, *19*(6), 659–667. https://doi.org/10.1038/mp.2013.78

Dufour, N., Redcay, E., Young, L., Mavros, P. L., Moran, J. M., Triantafyllou, C.,

Gabrieli, J. D. E., & Saxe, R. (2013). Similar Brain Activation during False Belief

Tasks in a Large Sample of Adults with and without Autism. *PLoS ONE*, *8*(9).

https://doi.org/10.1371/journal.pone.0075468

Friston, K. J., Mechelli, A., Turner, R., & Price, C. J. (2000). Nonlinear responses in

fMRI: the Balloon model, Volterra kernels, and other hemodynamics.

NeuroImage, 12(4), 466-477. https://doi.org/10.1006/nimg.2000.0630

Gilbert, S. J., Bird, G., Brindley, R., Frith, C. D., & Burgess, P. W. (2008). Atypical

recruitment of medial prefrontal cortex in autism spectrum disorders: An fMRI

study of two executive function tasks. *Neuropsychologia*, *46*(9), 2281–2291.

https://doi.org/10.1016/j.neuropsychologia.2008.03.025

Hamilton, A. F., Brindley, R., & Frith, U. (2009). Visual perspective taking impairment

in children with autistic spectrum disorder. *Cognition*, *113*(1), 37–44.

https://doi.org/10.1016/j.cognition.2009.07.007

Hartwright, C. E., Apperly, I. A., & Hansen, P. C. (2013). Representation, Control, or

Reasoning? Distinct Functions for Theory of Mind within the Medial Prefrontal

Cortex. *Journal of Cognitive Neuroscience*, *26*(4), 683–698.

https://doi.org/10.1162/jocn_a_00520

Hasson, U., Avidan, G., Gelbard, H., Vallines, I., Harel, M., Minshew, N., & Behrmann,

M. (2009). Shared and idiosyncratic cortical activation patterns in autism revealed

under continuous real-life viewing conditions. *Autism Research*, *2*(4), 220–231.

https://doi.org/10.1002/aur.89

Hasson, U., Furman, O., Clark, D., Dudai, Y., & Davachi, L. (2008). Enhanced

Intersubject Correlations during Movie Viewing Correlate with Successful

Episodic Encoding. *Neuron*, *57*(3), 452–462.

https://doi.org/10.1016/j.neuron.2007.12.009

Hasson, U., Landesman, O., Knappmeyer, B., Vallines, I., Rubin, N., & Heeger, D. J.

(2008). Neurocinematics: The Neuroscience of Film. *Projections*, *2*(1), 1–26.

https://doi.org/10.3167/proj.2008.020102

Jeste, S. S., & Geschwind, D. H. (2014). Disentangling the heterogeneity of autism

spectrum disorder through genetic findings. *Nature Reviews. Neurology*, *10*(2),

74–81. https://doi.org/10.1038/nrneurol.2013.278

Just, M. A., Cherkassky, V. L., Keller, T. A., Kana, R. K., & Minshew, N. J. (2007).

Functional and Anatomical Cortical Underconnectivity in Autism: Evidence from

an fMRI Study of an Executive Function Task and Corpus Callosum

Morphometry. *Cerebral Cortex*, *17*(4), 951–961.

https://doi.org/10.1093/cercor/bhl006

Kana, R. K., Maximo, J. O., Williams, D. L., Keller, T. A., Schipul, S. E., Cherkassky, V.

L., Minshew, N. J., & Just, M. A. (2015). Aberrant functioning of the theory-of-

mind network in children and adolescents with autism. *Molecular Autism*, *6*(1),

59. https://doi.org/10.1186/s13229-015-0052-x

Kaufman, J., Birmaher, B., Brent, D., Rao, U. M. A., Flynn, C., Moreci, P., ... & Ryan, N.

(1997). Schedule for affective disorders and schizophrenia for school-age

children-resent and lifetime version (K-SADS-PL): Initial reliability and validity

data. *Journal of the American Academy of Child & Adolescent Psychiatry, 36(7),*

980-988. https://doi.org/10.1097/00004583-199707000-00021.

Krause, L., Enticott, P. G., Zangen, A., & Fitzgerald, P. B. (2012). The role of medial

prefrontal cortex in theory of mind: A deep rTMS study. *Behavioural Brain*

Research, *228*(1), 87–90. https://doi.org/10.1016/j.bbr.2011.11.037

Lenroot, R. K., & Yeung, P. K. (2013). Heterogeneity within Autism Spectrum

Disorders: What have We Learned from Neuroimaging Studies? *Frontiers in*

Human Neuroscience, 7, 733. https://doi.org/10.3389/fnhum.2013.00733

Lind, S. E. (2010). Memory and the self in autism: A review and theoretical framework.

Autism, *14*(5), 430–456. https://doi.org/10.1177/1362361309358700

Lord, C., Rutter, M., DiLavore, P., Risi, S., Gotham, K., & Bishop, S. (2012). Autism

diagnostic observation schedule–2nd edition (ADOS-2). *Los Angeles, CA:*

Western Psychological Corporation.

Mason, R. A., Williams, D. L., Kana, R. K., Minshew, N., & Just, M. A. (2008). Theory

of Mind disruption and recruitment of the right hemisphere during narrative

comprehension in autism. *Neuropsychologia*, *46*(1), 269–280.

https://doi.org/10.1016/j.neuropsychologia.2007.07.018

Mills, K. L., Lalonde, F., Clasen, L. S., Giedd, J. N., & Blakemore, S.J. (2014).

Developmental changes in the structure of the social brain in late childhood and

adolescence. *Social Cognitive and Affective Neuroscience*, *9*(1), 123–131.

https://doi.org/10.1093/scan/nss113

Monk, C. S., Peltier, S. J., Wiggins, J. L., Weng, S.-J., Carrasco, M., Risi, S., & Lord, C.

(2009). Abnormalities of intrinsic functional connectivity in autism spectrum

disorders,. *NeuroImage*, *47*(2), 764–772.

https://doi.org/10.1016/j.neuroimage.2009.04.069

Naci, L., Cusack, R., Anello, M., & Owen, A. M. (2014). A common neural code for

similar conscious experiences in different individuals. *Proceedings of the*

National Academy of Sciences, *111*(39), 14277–14282.

https://doi.org/10.1073/pnas.1407007111

Naci, L., Haugg, A., MacDonald, A., Anello, M., Houldin, E., Naqshbandi, S., Gonzalez-

Lara, L. E., Arango, M., Harle, C., Cusack, R., & Owen, A. M. (2018). Functional

diversity of brain networks supports consciousness and verbal intelligence.

Scientific Reports, *8*. https://doi.org/10.1038/s41598-018-31525-z

Nguyen, M., Vanderwal, T., & Hasson, U. (2019). Shared understanding of narratives is

correlated with shared neural responses. *NeuroImage*, *184*, 161–170.

https://doi.org/10.1016/j.neuroimage.2018.09.010

Otsuka, Y., Osaka, N., Ikeda, T., & Osaka, M. (2009). Individual differences in the

theory of mind and superior temporal sulcus. *Neuroscience Letters*, *463*(2), 150–

153. https://doi.org/10.1016/j.neulet.2009.07.064

Pajula, J., & Tohka, J. (2016). How Many Is Enough? Effect of Sample Size in Inter-

Subject Correlation Analysis of fMRI. *Computational Intelligence and*

Neuroscience. https://doi.org/10.1155/2016/2094601

Pedreño, C., Pousa, E., Navarro, J. B., Pàmias, M., & Obiols, J. E. (2017). Exploring the

Components of Advanced Theory of Mind in Autism Spectrum Disorder. *Journal*

of Autism and Developmental Disorders, *47*(8), 2401–2409.

https://doi.org/10.1007/s10803-017-3156-7

Peters, S., Van Duijvenvoorde, A. C. K., Koolschijn, P. C. M. P., & Crone, E. A. (2016).

Longitudinal development of frontoparietal activity during feedback learning:

Contributions of age, performance, working memory and cortical thickness.

Developmental Cognitive Neuroscience, *19*, 211–222.

https://doi.org/10.1016/j.dcn.2016.04.004

Richardson, H., Lisandrelli, G., Riobueno-Naylor, A., & Saxe, R. (2018). Development

of the social brain from age three to twelve years. *Nature Communications*, *9*(1),

1027. https://doi.org/10.1038/s41467-018-03399-2

Rilling, J. K., Sanfey, A. G., Aronson, J. A., Nystrom, L. E., & Cohen, J. D. (2004). The

neural correlates of theory of mind within interpersonal interactions. *NeuroImage*,

22(4), 1694–1703. https://doi.org/10.1016/j.neuroimage.2004.04.015

Rutter, M., Le Couteur, A., & Lord, C. (2003). Autism diagnostic interview-revised. *Los*

Angeles, CA: Western Psychological Services, 29, 30.

Salmi, J., Roine, U., Glerean, E., Lahnakoski, J., Nieminen-von Wendt, T., Tani, P.,

Leppämäki, S., Nummenmaa, L., Jääskeläinen, I. P., Carlson, S., Rintahaka, P., &

Sams, M. (2013). The brains of high functioning autistic individuals do not

synchronize with those of others. *NeuroImage: Clinical*, *3*, 489–497.

https://doi.org/10.1016/j.nicl.2013.10.011

Saxe, R, & Kanwisher, N. (2003). People thinking about thinking people: The role of the

temporo-parietal junction in "theory of mind." *NeuroImage*, *19*(4), 1835–1842.

https://doi.org/10.1016/S1053-8119(03)00230-1

Saxe, R., & Wexler, A. (2005). Making sense of another mind: The role of the right

temporo-parietal junction. *Neuropsychologia*, *43*(10), 1391–1399.

https://doi.org/10.1016/j.neuropsychologia.2005.02.013

Solomon, M., Ozonoff, S. J., Ursu, S., Ravizza, S., Cummings, N., Ly, S., & Carter, C. S.

(2009). The neural substrates of cognitive control deficits in autism spectrum

disorders. *Neuropsychologia*, *47*(12), 2515–2526.

https://doi.org/10.1016/j.neuropsychologia.2009.04.019

Spencer, M. D., Holt, R. J., Chura, L. R., Suckling, J., Calder, A. J., Bullmore, E. T., &

Baron-Cohen, S. (2011). A novel functional brain imaging endophenotype of

autism: The neural response to facial expression of emotion. *Translational*

Psychiatry, *1*(7), e19. https://doi.org/10.1038/tp.2011.18

Supekar, K., Uddin, L. Q., Khouzam, A., Phillips, J., Gaillard, W. D., Kenworthy, L. E.,

Yerys, B. E., Vaidya, C. J., & Menon, V. (2013). Brain Hyperconnectivity in

Children with Autism and its Links to Social Deficits. *Cell Reports*, *5*(3), 738–

747. https://doi.org/10.1016/j.celrep.2013.10.001

Yeo, T. H., Krienen, F. M., Sepulcre, J., Sabuncu, M. R., Lashkari, D., Hollinshead, M.,

Roffman, J. L., Smoller, J. W., Zöllei, L., Polimeni, J. R., Fischl, B., Liu, H., &

Buckner, R. L. (2011). The organization of the human cerebral cortex estimated

by intrinsic functional connectivity. *Journal of Neurophysiology*, *106*(3), 1125–

1165. https://doi.org/10.1152/jn.00338.2011

Uddin, L. Q., Supekar, K., & Menon, V. (2010). Typical and atypical development of

functional human brain networks: Insights from resting-state fMRI. *Frontiers in*

Systems Neuroscience, *4,* 21. https://doi.org/10.3389/fnsys.2010.00021

Uddin, L. Q., Supekar, K., & Menon, V. (2013). Reconceptualizing functional brain

connectivity in autism from a developmental perspective. *Frontiers in Human*

Neuroscience, *7,* 458. https://doi.org/10.3389/fnhum.2013.00458

Völlm, B. A., Taylor, A. N. W., Richardson, P., Corcoran, R., Stirling, J., McKie, S.,

Deakin, J. F. W., & Elliott, R. (2006). Neuronal correlates of theory of mind and

empathy: A functional magnetic resonance imaging study in a nonverbal task.

NeuroImage, *29*(1), 90–98. https://doi.org/10.1016/j.neuroimage.2005.07.022

von dem Hagen, E. A. H., Stoyanova, R. S., Baron-Cohen, S., & Calder, A. J. (2013).

Reduced functional connectivity within and between 'social' resting state

networks in autism spectrum conditions. *Social Cognitive and Affective*

Neuroscience, *8*(6), 694–701. https://doi.org/10.1093/scan/nss053

Wechsler, D. (2014). The Wechsler intelligence scale for children—fifth edition. *San*

Antonio, TX: The Psychological Corporation.

Weng, S.-J., Wiggins, J. L., Peltier, S. J., Carrasco, M., Risi, S., Lord, C., & Monk, C. S.

(2010). Alterations of resting state functional connectivity in the default network

in adolescents with autism spectrum disorders. *Brain Research*, *1313*, 202–214.

https://doi.org/10.1016/j.brainres.2009.11.057

Yang, D. Y.-J., Rosenblau, G., Keifer, C., & Pelphrey, K. A. (2015). An integrative

neural model of social perception, action observation, and theory of mind.

Neuroscience & Biobehavioral Reviews, *51*, 263–275.

https://doi.org/10.1016/j.neubiorev.2015.01.020

Yeshurun, Y., Swanson, S., Simony, E., Chen, J., Lazaridi, C., Honey, C. J., & Hasson,

U. (2017). Same Story, Different Story: The Neural Representation of Interpretive

Frameworks. *Psychological Science*, *28*(3), 307–319.

https://doi.org/10.1177/0956797616682029