A New *tet*(X6) Tigecycline Resistance Determinant Co-carried with *mcr-1* by A Single Plasmid

Yongchang Xu1, Lizhang Liu1, Youjun Feng1,2,3*

1 Department of Pathogen Biology & Microbiology and Department of General Intensive Care Unit of the Second Affiliated Hospital, Zhejiang University School of Medicine, Hangzhou, Zhejiang 310058, China

2 Non-coding RNA and Drug Discovery Key Laboratory of Sichuan Province, Chengdu Medical College, Chengdu, Sichuan 610500, China

3 College of Animal Sciences, Zhejiang University, Hangzhou, Zhejiang 310058, China

* To whom correspondence should be addressed: fengyj@zju.edu.cn (Y Feng)

Abstract

Tigecycline and colistin are last-resort defenses used in anti-infection therapies of carbapenem-resistant bacterial pathogens. The successive emergence of plasmid-borne *tet*(X) tigecycline resistance mechanism and mobile colistin resistance (*mcr*) determinant, renders them clinically ineffective, posing a risky challenge to global public health. Here, we report, for the first time, that co-transfer of *tet*(X6) and *mcr-1* gives co-resistance to both classes of antibiotics by a single plasmid in an epidemic ST95 lineage of virulent *Escherichia coli*. Genomic analysis suggested that transposal transfer of *mcr-1* proceeds into the plasmid pMS8345A, in which a new variant *tet*(X6) is neighbored with Class I integron. The structure-guided mutagenesis finely revealed the genetic determinants of Tet(X6) in the context of phenotypic tigecycline resistance. The combined evidence *in vitro* and *in vivo* demonstrated its enzymatic action of Tet(X6) in the destruction of tigecycline. The presence of Tet(X6) (and/or MCR-1) robustly prevents the accumulation of reactive oxygen species (ROS) induced by tigecycline (and/or colistin). Unlike that *mcr-1* exerts fitness cost in *E. coli*, *tet*(X6) does not. Also, the co-expression of *tet*(X6) and *mcr-1* gives no synergistic delayed growth of the recipient *E. coli*. Therefore, spread of such a plasmid is of great concern in the context of "one health" comprising environmental/animal/human sectors, and heightened efforts are required to monitor its dissemination.

Keywords: Tigecycline resistance; Tet(X); Tet(X6); Colistin resistance; *mcr-1*; Co-transfer

Running title: Co-carriage of *tet*(X6) and *mcr-1*

Importance

We report that *tet*(X6), a new tigecycline resistance gene, is co-transferred with the other resistance gene *mcr-1* by a single plasmid. Not only have we finely mapped genetic determinants of *tet*(X6), but also revealed its biochemical action of tigecycline destruction. Crosstalk of fitness cost is addressed between Tet(X6) and MCR-1. Together, it constitutes mechanistic basis for Tet(X) resistance enzyme, and heightens the need to prevent and even control its dissemination.

Introduction

Antimicrobial resistance is an increasingly-devastating challenge in the context of "one health" that covers the environmental, animals and human sectors. Colistin is one of cationic antimicrobial polypeptides (CAMP) with an initial target of the surface-anchored lipid A moieties on the Gram-negative bacterium [1]. In contrast, tigecycline is the third-generation of tetracycline-type antibiotic, which interferes the machinery of protein synthesis of both Gram-negative, and Gram-positive bacteria [2]. In general, both colistin and tigecycline are an ultimate line of antibiotics to combat against lethal infections with carbapenem-resistant pathogens [3, 4]. Unfortunately, the emergence and global distribution of MCR family of mobile colistin resistance (*mcr-1* [5, 6] to *mcr-9* [7]) has potentially threatened the renewed interest of colistin in clinical therapies [8]. The majority of transferable colistin resistance depends on the surface lipid A remodeling by the MCR enzymes via the "ping-pong" action [9].

In addition to the two well-known actions, efflux and ribosome protection [10, 11], antibiotic degradation also constitutes in part the mechanism of tigecycline resistance [12-14]. The Tet(X) enzyme is a class of flavin-requiring monooxygenase [15, 16], which possesses the ability of modifying tetracycline and its derivatives (such as the glycylcycline, tigecycline) [13]. In general, Tet(X) inactivates tigecycline to give 11a-Hydroxytigecycline, rendering the carrier host bacterium insusceptible to tigecycline [13]. In particular, soil functional metagenomics performed by Forsburg *et al.* [17] revealed a number of new tetracycline destructases (10 in total, namely Tet47 to Tet56). Among them, *tet56* is only one exclusively from human pathogen *Legionella longbeachae*, indicating its potential spread from environments to clinical sector [17]. Subsequent structural studies suggested that these Tet tetracycline destructases accommodate antibiotics in diverse orientation, which highlights their architectural plasticity [12]. Indeed, the discovery of an inhibitor blocking the entry of flavin adenine dinucleotide cofactor into Tet(50) enzyme also paved a new way to reversing tigecycline resistance [12].

Very recently, two new variants [*tet*(X4) and *tet*(X5)] of *tetX*-type determinants that encode a tigecycline-inactivating enzyme, was found to spread by distinct plasmids in *Escherichia coli* from China [18, 19]. Though the limited distribution of *tet*(X4) thus far [20], it constitutes an expanding family of TetX resistance enzyme (Tet(X) [21, 22] to Tet(X5) [19]), and raises the possibility of rendering tigecycline (and even the newly-FDA approved eravacycline, the fourth-generation of tetracycline derivatives [23]) clinically ineffective. Worrisomely, the co-transfer of *mcr-1* and *tet*(X) probably promotes the emergence of a deadly superbug with the co-resistance to polymyxin and

tigecycline. However, this requires further epidemiological evidence.

Here, we report that this is the case. It underscores an urgent need to monitor and evaluate a potential risk for the convergence of Tet(X) tigecycline resistance to MCR colistin resistance by a single highly-transmissible plasmid in an epidemic ST95 lineage of virulent *E. coli* [24].

Results

Discovery of *tet*(X6), a new variant of Tet(X) resistance enzyme

To address this hypothesis, we systematically screened the whole NCBI nucleotide database, in which each of the six known *tet*(X) variants [*tet*(X) to *tet*(X5)] acts as a query. Among them, it gives six hits with the significant score (97%-99% identity) upon using the *tet*(X3) of *Pseudomonas aeruginosa* (1137bp, Acc. no.: AB097942) as a searching probe. The resultant hits correspond to four contigs of uncultured bacterium and two plasmids. These contigs include TE_6F_Contig_7 (3328bp, Acc. no.: KU547125), TG_6F_Contig_3 (3323bp, Acc. no.: KU547130), TG_7F_Contig_3 (2575bp, Acc. no.: KU547185); and TE_7F_Contig_3 (2588bp, Acc. no.: KU547176). The matched plasmids refer to pMS8345A (241,162bp; Acc. no.: CP025402) [24] and p15C38-2 (150,745bp; Acc.no.: LC501585), in which the gene of MS8345_A00031 exhibits 97.1% identity and 100% coverage when compared to *tet*(X3) (**Fig. S1**). This *tet*(X3)-like gene, MS8345_A00031, is thereafter renamed *tet*(X6) (Acc. no.: BK011183) in this study (**Fig. 1A**). Strikingly, we found that this plasmid coharbors the *mcr-1* gene encoding a phosphoethanolamine (PEA)-lipid A transferase [25]. Because that the plasmid is detected in an epidemic clone of ST95 Extraintestinal Pathogenic *E. coli* (ExPEC) [24], this single plasmid pMS8345A possesses the potential to confer its recipient host *E. coli* the co-resistance to both tigecycline and colistin. It is unusual, but not without any precedent. In fact, *mcr-1* has ever coexist with *bla*NDM in a single isolate [26] and even a single plasmid [27].

Characterization of a *tet*(X6)-harboring plasmid

In fact, this *tet*(X6)-positive pMS8345A (Acc. no.: CP025402) attracting our attention in the *in silico* search, is an IncHI2-type large plasmid (~241kb) that was initially discovered by Beatson and coworkers [24] from a virulent lineage of *E. coli* with multiple drug resistance (MDR) via the routine screen of colistin resistance (**Fig. 1A**). Generally, this plasmid has an average GC% value of 46.29%, and is predicted to contain 548 putative open reading frames (ORFs). Importantly, pMS8345A is found to possess a series of Integrative and Conjugative Elements (ICEs) using the web-based tool of oriTfinder (**Fig. 1A**) [28]. Although that pMS8345A is not accessible in China right now, we applied its surrogate one, pDJB-3 (**Fig. 1A**), in the conjugation assays. Consistent with its carriage of T4SS alone (**Fig. 1A**), pDJB-3 cannot survive in the conjugation trials. In contrast, pHNSHP45-2 has the ability of conjugation [29]. This is because that it is fulfilled with all the four essential modules for self-transmission, namely *oriT* region, relaxase gene, type IV coupling protein (T4CP) gene and type IV secretion system (T4SS) (**Fig. 1B**). A similar scenario is also seen in the plasmid pM8345A. Taken together, this highlights the potential transferability of pM8345A.

The hallmark of pMS8345A lies in two disconnected/unique resistance regions, one of which refers to the *tet*(X6)-bearing MDR region of appropriate ~40kb long (**Fig. 1B**), and the other denotes the *mcr-1*-containing region (**Fig. 1C**). Surprisingly, 11 kinds of antimicrobial resistance (AMR) genes have been recruited and integrated into this unusual MDR region (**Fig. 1B**), giving multiple drug resistance. In addition to *tet*(X6) of tigecycline resistance, these AMR loci are listed as follows: i) *arr-2* (rifampicin resistance), ii) *ere*(A) encoding erythromycin esterase; iii) two copies of *aadA* family genes: *aadA1* and *aadA22* (aminoglycoside resistance); iv) an extended-spectrum β-lactamase gene $bla_{CTX-M-1}$; v) *tmrB* producing tunicamycin resistance; vi) an operon of *strA-strB* conferring streptomycin resistance; vii) bla_{TEM-1B} encoding β-lactamase; viii) *mph*(A), a macrolide 2'-phosphotransferase I-expressing gene; ix) two copies of *sul1* genes of sulfonamide resistance; and x) *dfrA7*, the trimethoprim resistance gene (**Fig. 1B**). However, the colistin resistance arises from the "*mcr-1*-IS*Apl1*" transposon alone (**Fig. 1C**). Therefore, the occupation of MDR (including, but not limited to the co-resistance to tigecycline and polymyxin, two last-resort anti-infection options) by pMS8345A is a serious risk in the clinic sector, once it further successfully enters and spreads into a number of epidemic virulent clones.

Genomic analyses of *tet*(X6)-containing MDR region

The linear genome alignment of MDR plasmids revealed that pMS8345A displays high level of similarity to at least three other resistance plasmids (**Fig. 1A**). Namely, they include i) the *mcr-1*-harboring plasmid pHNSHP45-2 with 99.75% identity and 87% query coverage (Acc. no.: KU341381), ii) and a tetracycline resistance plasmid of *Yersinia pseudotuberculosis*, pYps.F1 with 100% identity and 74% query coverage (Acc. no.: LT221036), and iii) a typical IncHI2-group, *mcr-1*-carrying plasmid pDJB-3 with 99.74% identity and 63% query coverage (Acc. no.: MK574666) we isolated and sequenced in this study (**Fig. 1A**). Unlike that the *mcr-1*-lacking plasmid pYps.F1 exists in *Y. pseudotuberculosis*, all the other three *mcr-1*-harboring plasmids disseminate in different clones of *E. coli* with varied sequence types (**Fig. 1A**). In brief, i) pMS8345A is detected in ST95, a globally-distributed clone having the relevance to bacterial bloodstream infections and neonatal meningitis [24]; ii) pHNSHP45-2 is recovered from an intensive pig farm [29]; and iii) pDJB-3 is recently determined by our group to occur in ST165 in an pig farm, a rare sequence type (**Fig. 1A**).

Among them, the organization of MDR differs greatly (**Fig. 1**). Unlike the pMS8345A having both *tet*(X6)-positive MDR (**Fig. 1B**), and the *mcr-1*-containing cassette (**Fig. 1C**), the plasmid pDJB-3 has *mcr-1*, but not

MDR region (**Fig. 1A**). In contrast, the plasmid pYps.F1 carries the MDR region, but not *mcr-1* (**Fig. 1A**). Genetic dissection indicated that three class 1 integrons are located in the MDR region. Given that a pool of gene cassettes can be integrated, the majority of which encode resistance to antibiotics [30], Class 1 integron facilitates the global spread of AMRs [31]. In particular, the MDR region of pMS8345A seems structurally unusual (**Fig. 1B**). First, it comprises a cluster of transposons and insert sequences (*IS*) with its boundary of two integrases (one is *int1* on the forward strand, and the other denotes a truncated *int1* on the reverse strand, **Fig. 1B**); Second, the integrase-encoding locus *int1* is adjacent to an integron-associated recombination site *attI*, and then followed by several *attC* sites (**Fig. 1B**); Third, the occurrence of Tn*2* and Tn*As1* (two copies of Tn*3* family transposons) in the pMS8345A MDR region suggests the association with its mobility (**Fig. 1B**); Fourth, the multiple *IS* elements located within the MDR region (namely *bla*$_{CTX-M-1}$ carried on an IS*Ecp1*, the operon of *strA-strB* adjacent to an IS*26*-IS*1133* structure [32], the *aac(3)-IIa/tmrB* plus *bla*$_{TEM-1B}$ genes neighbored by IS*26*), facilitate the formation of transposons via the recombination events (**Fig. 1B**); Fifth, circular gene cassettes [such as *arr-2/ere(A)/aadA1* and *sul1/tet*(X6)] are presumably integrated by site-specific recombination between *attI* and *attC*, a process mediated by the integron integrase (**Fig. 1B**). In general consistency with scenarios seen in class 1 integron (**Fig. 1B**), the fact that the GC content (37.8%) of *tet*(X6) is far less than the average GC% (46.29%) of the pMS8345A, hints its acquisition through gene horizontal transfer. Sequence alignment unveils that the *tet*(X6)-positive MDR region in pMS8345A is highly similar to the MDR region in p15C38-2 (**Fig. 1B**). In brief, *tet*(X6) and *sul1* are downstream of a class 1 integron carrying the *aadA* family of resistance genes (*aadA22* in pMS8345A and *aadA12* in p15C38-2). p15C38-2 harbors a Tn*3*-like transposon, Tn*As2* with 85% identity.

Functional insights into Tet(X6) tigecycline resistance

In addition to the pMS8345A plasmid, the *tet*(X6)-based *in silico* search unveils two more *tet*(X6)-containing contigs (**Fig. S2**), namely TG_7F_Contig_3 (2575bp, Acc. no.: KU547185) and TE_7F_Contig_3 (2588bp, Acc. no.: KU547176). More intriguingly, the two contigs are derived from uncultivated bacterium from latrine, in EI Salvador, 2012 (**Fig. 2**). It seems likely that *tet*(X6) appears earlier than that of *tet*(X4) initially detected in a contig of *K. pneumoniae* (4069bp, Acc. no.: NQBP01000050) from Thailand, 2015 [20]. further database mining suggests a number of *tet*(X) homologs [designated Tet(X7) to Tet(X13)] that are similar to *tet*(X6) at the identity ranging 91.27% to 97.62%. Phylogenetic analysis of these Tet(X) enzymes illustrates an ongoingly-expanding family of Tet(X) resistance determinants

(**Fig. 2**), raising a possible ancestor shared amongst these *tet*(X) variants.

Given that i) the statement by He *et al*. [18] that TetX3 of *Pseudomonas* (Acc. no.: AB097942) is not active, is argued by our recent study [20]; and ii) as a new variant, Tet(X6) displays 96.03% identity to the *Pseudomonas* Tet(X3) **(Fig. S3)**, integrative evidences are extremely needed for the functional assignment of Tet(X6) in the context of tigecycline resistance. Therefore, we cloned *tet*(X6) into an arabinose-inducible pBAD24 expression vector and test its function in the strain MG1655 of *E. coli*. As predicted, the presence of *tet*(X6) can restore the growth of its recipient strain on LBA agar plates with tigecycline (16 to 32μg/ml, **Fig. S4A**). This level of resistance is almost as same as *tet*(X3) does, but slightly lower than that of Tet(X4) (**Fig. S4A**). As predicted, structural modeling of Tet(X6) presents a substrate-loading channel (**Figs 3A-B**). Similar to the scenario with Tet(X4) [20], it consists of a tigecycline substrate-binding motif (**Figs 3C** and **E**) and a FAD cofactor-occupied cavity (**Figs 3D** and **F**). As for Tet(X6), the substrate-binding requires the cooperation of five residues E182, R203, H224, G226 and M365 (**Table 1** and **Fig. 3E**). Similarly, the FAD cofactor is surrounded with the following six residues E36, R37, R107, D301, P308 and Q312 in Tet(X6) (**Table 1** and **Fig. 3F**). Except that the substitution of H224T occurs in Tet(X1) [and/or H234Y in Tet(X5), in **Fig. S3**], all the aforementioned residues are relatively-conserved across the newly-proposed family of Tet(X) tigecycline-inactivating enzymes (**Table 1**). Among them, a number of residues have been functionally verified. In the case of Tet(X4), two of 5 substrate-binding cavity (H231 and M372) and three of 6 FAD-interactive residues (E43, R114, and D308) somewhat play roles in the phenotypic tigecycline resistance [20]. As expected, structure-guided alanine substitution of Tet(X6) revealed that i) two of 5 tigecycline-binding residues (H224A and M365A) partially determine its phenotypic resistance to tigecycline (**Fig. 3G**); and ii) three of 6 FAD-surrounding residues (E36A, R107A, and D301A) give differential level of impact on its resultant tigecycline resistance (**Fig. 3H**). Evidently, this result represents a functional proof for Tet(X6) as a new member of the expanding family of Tet(X) enzymes that have a role in the action of tigecycline degradation.

Action of inactivation of tigecycline by Tet(X6)

To further elucidate biochemical mechanism of Tet(X6)-catalyzed tigecycline destruction (**Fig. 4**), we integrated an *in vivo* approach of microbial bioassay (**Fig. 4A**) with the *in vitro* system of enzymatic reaction (**Figs 4B-D**). This tigecycline bioassay we developed is dependent on the indicator strain DH5α of *E. coli* in that it was verified to be tigecycline susceptibility (**Fig. 4A**). As predicted, a zone of bacterial inhibition was clearly seen to surround a paper disk of the blank control, on which 2.5μg/ml tigecycline is spotted (**Fig.**

4A). A similar scenario was also observed in the negative control, the supernatant from *E. coli* MG1655 with the empty vector pBAD24 alone (**Fig. 4A and S1 Table**). In contrast, the zone of tigecycline inhibition disappears around the paper disk containing supernatants of *E. coli* MG1655 expressing either *tet*(X6) or its homologous gene *tet*(X3) (**Fig. 4A**). This highlighted an *in vivo* role of *tet*(X6) [*tet*(X3)] in the context of inactivating tigecycline.

Subsequently, we attempted to produce the recombinant forms of Tet(X6) and its homolog Tet(X3) aiming at the *in vitro* enzymatic activity. Of note, Tet(X6) and Tet(X3) protein consistently gives yellow in solution (**Fig. 4B**), hinting a possibility of being occupied with a FAD cofactor. Gel filtration analysis indicated that both Tet(X6) and Tet(X3) display the solution structure of being a monomer (**Fig. 4B**). This is generally in an agreement with the apparent molecular mass (~36kDa) seen in the SDS-PAGE (**Fig. 4B**). The identification of polypeptide fingerprint with liquid chromatography (LC)/mass spectrometry allowed us to further dissect catalytic action of Tet(X6) [Tet(X3)] using the *in vitro* reconstituted system of tigecycline oxygenation (**Fig. S5A-B**). As expected, LC/MS-based detection of the substrate tigecycline showed a unique peak at the position of 586.2 m/z (**Fig. 4C**). In particular, the reaction mixture of Tet(X6) [Tet(X3)] consistently gave two distinct peaks in the spectrum of LC/MS (**Figs 4D and S6**). Namely, they included a peak of substrate tigecycline (586.2 m/z), and an additional peak assigned to its oxygenated product of tigecycline at the position of 602.2 m/z (**Figs 4D and S6**). This finding is consistent with an observation with the newly-identified Tet(X4) [18]. As Forsberg and coworkers described [17], a similar scenario was also seen with the inactivation of tetracycline by either Tet(X3) or Tet(X6) (**Fig. S7**), which is evidenced by the fact liquid culture of *tet*(X3) [and/or *tet*(X6)]-bearing *E. coli* gives dark (**Fig. S7**). Unlike that the negative control strain MG1655 with empty vector pBAD24 alone displays a big inhibition circle, E-test of tigecycline unveiled that expression of *tet*(X3) [or *tet*(X6)] renders the recipient strains significantly antagonistic to the tigecycline challenge (**Fig. S8**). Thereafter, we formulated a working model that Tet(X6) exploits a FAD cofactor to oxygenate/destruct the last-line antibiotic tigecycline (**Fig. 4E**). It seems likely this chemical reaction proceeds via a 'ping-pong' action, and this hypothesis requires further experimental evidence.

Variation in the *mcr-1*-containing cassettes

Sequence analyses of *mcr-1*-bearing elements from the three IncHI2-type plasmids (pMS8345A, pSA186_MCR1, and pDJB-3) reveals the core structure of "IS*Apl1-mcr-1*" (**Fig. 1C**). Unlike the plasmid of pDJB-3 containing a cassette of "IS*Apl1-mcr-1-pap2*-IS*Apl1*", the pSA186_MCR1 plasmid possesses an inactivated *pap2* inserted with an inverted copy of

ISApl1 (**Fig. 1C**). As a member of the IS30 family, the ISApl1 can transpose into its target by formatting a synaptic complex between an inverted repeat (IR) in the transposon circle and an IR-like sequence in the target [33]. After the initial formation of this composite transposon, one or both copies of ISApl1 might be lost. As such, this loss might improve the stability of mcr-1 in a diverse range of plasmids and then intensify its spread of mcr-1 [34]. Therefore, we favor to believe that pMS8345A having "ISApl1-mcr-1" alone (**Fig. 1C**) proceeds the loss of its downstream ISApl1 following the transposition. Not surprisingly, the E. coli strain carrying mcr-1 gives the minimum inhibitory concentration (MIC) at 4.0µg/ml. In addition, functional expression of a single mcr-1 confers the polymyxin-susceptible recipient strain of E. coli MG1655 to be viable on the LBA agar plate with colistin of up to 16µg/ml (**Fig. S4B**). Evidently, these data proved that both tet(X6) and mcr-1 functionally cooccurs by a single plasmid in an epidemic ST95 clone of pathogenic E. coli.

Physiological alteration by Tet(X6) and MCR-1

To address physiological consequence of Tet(X6) and MCR-1, we separately examined the pool of intracellular reactive oxygen species (ROS) and various growth curve-based metabolic fitness in an array of different E. coli strains (**Figs 5** and **6**). As illustrated in the assay of fluorescence activated cell sorting (FACS), the cytosolic ROS level in the M1655 with empty vector alone was relatively low (**Fig. 5A**). Similar scenarios were also seen in derivatives of the MG1655 strain, regardless of the presence of mcr-1 (**Fig. 5D**), tet(X6) (**Fig. 5E**), and even both (**Fig. 5F**). As predicted, the level of intracellular ROS was increased greatly upon its exposure to either colistin (**Figs 5B and J**) or tigecycline (**Figs 5C and J**). The expression of mcr-1 effectively prevented the colistin-stimulated ROS formation (**Figs 5G and J**). Similarly, the presence of tet(X6) robustly interfered the ROS production triggered by tigecycline (**Figs 5H and J**). In fact, the addition of both colistin and tigecycline only gave slight increment of ROS accumulation in the MG1655 strain coharboring mcr-1 and tet(X6) (**Figs 5I-K**). Therefore, Tet(X6) attenuates the tigecycline-induced ROS generation as does MCR-1 in response to colistin (**Fig. 5**).

As expected, the presence of plasmid-borne mcr-1 can cause the delayed growth of its recipient host E. coli MG1655, whereas the empty vector not (**Figs 6A-B**). In agreement with earlier observations by our research group and others [35-39], this underscored that MCR-1 exerts significantly fitness cost in E. coli. In contrast, the expression of tet(X6) fails to trigger any detectable deficiency in bacterial growth (**Fig. 6C**), indicating that Tet(X6)-causing metabolic burden/disorder is minimal. To further probe whether or not the crosstalk between Tet(X6) and MCR-1 occurs in E. coli, we

developed an engineered strain coharboring derivatives of two compatible plasmids (one refers to a low copy number, T7 promoter-driven pWSK129 [40], and the other denotes an arabinose-inducible pBAD24 with ampicillin resistance [41]). In fact, the two resistance enzymes MCR-1 and Tet(X6) are produced by pWSK129*::mcr-1*, and pBAD24*::tet*(X6), respectively (**S1 Table**). Evidently, the coexistence of *tet*(X6) and *mcr-1* failed to augment the inhibition of bacterial growth (**Fig. 6D**).

As recently performed with MCR-3/4 [4], we also integrated an approach of LIVE/DEAD cell staining (**Fig. 7**). Unlike that the negative control, MG1655 strains with vector alone are almost fulfilled with alive cells (**Figs 7A-B**), the *mcr-1*-producing strains contained around 30% dead cells (**Figs 7C-D**, and **I**). Consistent with no obvious effect exerted by Tet(X6) on growth curves (**Fig. 6B**), our confocal microscopy assays revealed that low level of DAED/LIVE ratio is assigned to the *tet*(X6)-carrying MG1655 (**Figs 7E-F**, and **I**). Not surprisingly, the co-expression of *mcr-1* and *tet*(X6) cannot promotes the ratio of DEAD/LIVE cells (**Figs 7G-H**, and **I**), in relative to the *mcr-1*-positive strains (**Figs 7C-D**). Taken together, these data suggested that no synergism can be assigned to the fitness cost caused by the lipid A modifier MCR-1 and the tigecycline-inactivating enzyme Tet(X6). Unlike the fact that MCR-1 modifies bacterial lipid A, the initial target of the cationic antimicrobial peptide colistin receptor [6, 25], Tet(X6) presumably hydrolyzes the tetracycline class antibiotics, like tigecycline (**Fig. 4**) [15, 20]. The former action results in bacterial surface remodeling by MCR-1 through an addition of PEA moiety to 1(4')-phosphate position of lipid A [8, 9]. Consequently, this might in part shape metabolic flux of the recipient microbe to balance *mcr-1* expression and bacterial survival stressed with colistin, giving the phenotypic fitness cost [35]. In contrast, it seems likely that the destruction of tigecycline by the flavin-dependent Tet(X) enzyme exerts minor effects on metabolic process (or claims few metabolic requirement). While such explanation for the limited fitness cost by Tet(X6) needed experimental evidence.

Discussion

The pMS8345A, a large IncHI2-type MDR-plasmid is firstly identified by Beatson and coworkers [24] to coexist with a big ColV-like virulence plasmid in the ST95 virulent lineage of *E. coli*. This alarms us that the spread of such pathogen might herald an era of post-antibiotic where we stand. The data we report here furthers our understanding tigecycline resistance mechanism of TetX family enzymes (**Fig. 4E**). To the best of knowledge, it is a first report addressing a case of the co-transfer of *tet*(X6) and *mcr-1* by a single plasmid. Since that no known mobile elements are adjacent to *tet*(X6), we hypothesize that the transposon of "IS*Apl1*-*mcr-1*-*pap2*-IS*Apl1*" mediates the transfer of *mcr-1* into this plasmid. The discovery of Tet(X6), a new member of TetX family, allows us to engineer an array of TetX-expressing bacteria, which paves a way to the development of bioremediation strategy for the environmental tetracycline contamination in the agricultural/industrial productions.

Also, the co-carriage of *tet*(X6) and *mcr-1* on a single IncHI2-type plasmid is far different from the observation by Sun *et al.* [19] that the co-existence of *tet*(X4) and *mcr-1* is mediated by two distinct plasmids in an *E. coli* clone. Unlike that the mobility of *tet*(X4) relies on IS*CR2*-mediated transposition [18], the gain of *tet*(X6) transferability is not clear (**Fig. 1B**). Not surprisingly, the IncHI2-type plasmid carries *mcr-1* along with *tet*(X6) here, because that it has ever been found to act as a vehicle of global *mcr-1* dissemination [8, 42]. As expected, two types of different antibiotics (colistin and tigecycline) consistently stimulate the formation of hydroxyl radicals in *E. coli* (esp. ROS, in **Fig. 5**), which might constitute additional example and/or evidence for an improved postuate of "efficient antibiotic killing associated with bacterial metabolic state (ATP [43-45] and ROS [46-49]). As for a given version of MCR resistance determinants [36, 38, 39], the recipient bacterial host has been demonstrated to give fitness cost exemplified with the delayed growth prior to the entry into log-phase. It is reasonable that the presence of either *mcr-1* or *tet*(X6) also cause metabolic fitness to some extent. Consistent with earlier observations of MCR-like members by Yang *et al.* [35] and Zhang *et al.* [36, 37, 39], we confirmed the obvious fitness cost caused by MCR-1 (**Figs 6-7**). This is in part (if not all) explained by the fact that bacterial membrane integrity is altered by MCR-1-mediated lipid A remodeling [9]. In contrast, the expression of *tet*(X6) does not lead to metabolic burden (**Figs 6-7**), which is because that Tet(X6) destructs the antibiotic of glycyl-cycline tigecycline, rather than the ribosome target [10]. It is unusual, but not without any precedent. Retrospectively, a similar scenario has ever been assigned to the other resistance enzyme β-lactamase-encoding gene bla_{TEM1b} (i.e., no fitness cost is correlated with it) [35]. Therefore, it is not surprising that no synergistic fitness arises from the co-carriage of *tet*(X6) and *mcr-1* in *E. coli*. Although a

number of new *tet*(X) variants [*tet*(X6) to *tet*(X13)] have been proposed in this study (**Fig. 2**), most of versions apart from *tet*(X6) await experimental demonstration in the near future. This is because that rare case of cryptic version might occur naturally, such as the prototypical *tet*(X) [we called *tet*(X0)] preexisting in an obligate anaerobe *Bacteroides fragilis* [21]. Given that i) tigecycline and colistin both are one of few alternative options to combat against carbapenem-resistant Enterobacteriaceae and Acinetobacter species, ii) both MCR-1 [5]and Tet(X4) [19] have accordingly rendered colistin and tigecycline ineffective in the therapy of mice with MDR infection, the cooccurrence and co-transfer of *tet*(X6) and *mcr-1* by a single plasmid amongst epidemic pathogens is a risky challenge to public health and clinical therapies.

Taken together, it is plausible and urgent to introduce *mcr* variants along with, but not only limited to, *tet*(X) variants in the routine national (and/or international) investigation in the context of "one health" (environmental/animal/human sectors). Along with major findings of other research group [12], functional definition of Tet(X3) and its homologue Tet(X6) here might supplement additional insights into Tet(X) tigecycline resistance, and even benefit the development of anti-Tet(X) resistance enzyme inhibitors.

Materials and Methods

Sequencing, assembly and annotation of plasmids

The plasmid pDJB-3 was isolated from the colistin-resistant *E. coli* DJB-3 of swine origin, verified with *mcr-1*-specific PCR, and then subjected to genome sequencing with the Hiseq X ten PE150 sequencer platform (Illumina, USA). As a result, the DNA library of pDJB-3 plasmid prepared by KAPA Hyper Prep Kit (Roche, Basel, Switzerland) gave a pool of 150 bp paired-end reads that are destined to be assembled into a contig by the SPAdes Genome Assembler (version 3.11.0). A BLASTN search was conducted to probe whether or not the resultant *mcr-1*-containing contig has a best-hit plasmid candidate. Together with Sanger sequencing, PCR was applied to close all the suspected gaps.

The resultant plasmid genome was annotated through the prediction of open reading frames (ORFs) with RAST (rapid annotation using subsystem technology, http://rast.nmpdr.org). PlasmidFinder 1.3 (https://cge.cbs.dtu.dk/services/PlasmidFinder-1.3/) was used to type the plasmid incompatibility, and ResFinder 3.1 (https://cge.cbs.dtu.dk/services/ResFinder/) was applied to screen possible antimicrobial resistance genes. The plasmid map was given with GenomeVx (http://wolfe.ucd.ie/GenomeVx/), and its linear alignment was proceeded with Easyfig [50].

Plasmid conjugation experiments

As recently described by Sun *et al*. [27], the experiments of plasmid conjugation were performed, in which the rifampin-resistant *E. coli* recipient strain EC600 (and/or strain DJB-3) acted as a donor. In brief, overnight cultures were re-grew in LB broth, donor and recipient strains were mixed at the logarithmic phase and spotted on a filter membrane, and then incubated at 37°C overnight. Subsequently, bacteria were washed from filter membrane and spotted on LB agar plate containing 400 μg/ml rifampin and 4 μg/ml colistin for selection of transconjugants. The suspected transformants were validated with PCR assays.

Molecular and microbial manipulations

With all the known *tet*(X) variants [*tet*(X0)-*tet*(X5)] as queries, BLASTN was carried out. In particular, a *tet*(X3)-based search returned a plasmid pMS8345A with significant hit, leading to the discovery of new variant of *tet*(X6). Then, the *tet*(X6) was synthesized *in vitro*, and cloned into pABD24, giving pABD24*::tet*(X6) (**Table S1**). Following the verification of its identity with direct DNA sequencing, this recombinant plasmid was introduced into the MG1655 strain of *E. coli* to assess its role *in vivo*. The generation of all the

point-mutants of *tet*(X6) were based on pBAD24*::tet*(X6) (**Table S1**) using

Site-directed mutagenesis kit (Vazyme Biotech), along with an array of specific

primers (**Table S2**). To test relationship of MCR-1 with Tet(X6), *mcr-1* was

cloned into pWSK, giving pWSK*::mcr-1*, compatible with pBAD*::tet*(X6) within

a single *E. coli* colony (**Table S1**).

As recently described with *tet*(X4) [20], the ability of *tet*(X6) and *tet*(X3)

in phenotypic tigecycline resistance was evaluated with LBA agar plates

containing tigecycline in a series of dilution. The strain expressing *tet*(X4) is

used as a positive control. In addition, the MCR-1 colistin resistance was also

judged as we earlier conducted with *mcr-1* [51] with little change. All the

examined *E. coli* strains were cultivated at 37°C overnight. Overnight cultures

were standardized to OD_{600} 0.05, inoculated (1:10; v/v) into 96-well

glass-bottomed plates in fresh LB broth ± arabinose or lactose (0, 0.02%, and

0.2%, w/v), and shaken (180 r.p.m) at 37°C. Of note, arabinose acted as an

inducer of pBAD24, and lactose was used to trigger expression of

pWSK129-based MCR-1. As we very recently reported [36, 38, 39], bacterial

growth curves were automatically plotted with spectrophotometer (Spectrum

lab S32A) to evaluate the fitness cost caused by *tet*(X6) and *mcr-1*. During the

total period of 20 hours, the value of optical absorbance (i.e., OD600) was

consistently recorded at an interval of 1 hour.

Bioassay for tigecycline destruction

The hydrolytic activity of Tet(X6) [and/or Tet(X3)] enzyme was

determined as Balouiri *et al*. described [52] with little change. In brief, the strain

of MG1655 harboring pBAD24::*tet*(X6) [or pBAD24::*tet*(X3)] was cultivated

overnight on LB agar plates supplemented with 0.1% arabinose. As a result,

bacterial colonies stripped, were suspended with 0.5ml of LB broth containing

0.1% arabinose and 2.5mg/ml tigecycline, whose optical density at 600nm

(OD600) was adjusted to about 2.0. Then, the suspension cultures were

proceeded to 8h stationary growth at 37°C. Following centrifugation

(13,600rpm, 20min) and filtration (at 0.22μm cut-off), bacterial supernatants

were prepared. The *E. coli* DH5α here referred to an indicator strain of

tigecycline susceptibility. Of note, the overnight culture of *E. coli* DH5α (~100μl)

was spread on a LB agar plate, which is centered with a paper disk of 6mm

diameter. To visualize the inhibition zones, the supernatant of interest (~20μl)

was spotted the paper disk on the aforementioned bioassay plates, and

incubated at 37°C for 16h. The negative control denotes the supernatant from

E. coli MG1655 bearing the empty vector pBAD24 (**Table S1**), and the blank

control referred to the tigecycline-containing LB broth.

Expression, purification and identification of Tet(X) enzymes

1   To produce the proteins of Tet(X6) and its homologue Tet(X3), the

strains of *E. coli* BL21 carrying pET21*::tet*(X6) [and pET21*::tet*(X3)] were

engineered (**Table S1**) for the inducible expression via the addition of 0.5 mM

isopropyl β-D-1-thiogalactopyranoside (IPTG). Bacterial lysates obtained by a

French Press (JN-Mini, China), were subjected to 1h of centrifugation at

16,800 rpm at 4°C, and the resultant supernatants were incubated with

pre-equilibrated Ni-NTA agarose beads on ice for 3 hours. Following the

removal of protein contaminants, the Tet(X6) [and/or Tet(X3)] protein was

eluted from the Ni-NTA agarose beads using the elution buffer [20mM Tris-HCl

(pH 8.0), 150mM NaCl, 20mM immidazole, and 5%glycerol], and concentrated

with a 30kDa cut-off ultra-filter (Millipore, USA). Subsequently, gel filtration was

performed to probe solution structure of Tet(X6) [Tet(X3)], using a Superdex

200/300GL size exclusion column (GE Healthcare). The purity of protein

pooled from the target peak was judged with SDS-PAGE (15%), and its identity

was validated with liquid chromatography (LC) mass spectrometry.

Enzymatic assay for Tet(X6) *in vitro*

To confirm the enzymatic activity of Tet(X6) enzyme, the *in vitro* reaction

system was established as recently described by Sun *et al*. [19] with little

change. The components of this assay (50µl in total) consisted of 20mM Tris

(pH7.5), 150mM NaCl, 1mM NADPH, 4mg/ml tigecycline, and 2mg/ml purified

Tet(X6) [and/orTet(X3)] enzyme. Following the maintenance (~12h) of

enzymatic reaction at room temperature, the resultant reaction mixture was

subjected to further analysis of liquid chromatography mass spectrometry

(LC/MS) using an Agilent 6460 triple quadrupole mass spectrometer (Agilent

Technologies, USA) [18]. As for LC/MS here, it was carried out as follows: i)

Nitrogen acted as the sheath gas and drying gas, the nebulizer pressure was

set to 45 psi, and the flow rate of drying gas was 5 liter/min. The flow rate and

temperature of the sheath gas were 11 liter/min and 350°C, respectively; ii)

Chromatographic separation proceeded on a Zorbax SB C8 column

(150 × 2.1mm, 3.5µm); iii) Mass spectrometric detection was completed using

an electrospray ionization (ESI) source in positive mode. Scan range was

100~1000amu; and iv) The resultant data was processed with Agilent Mass

Hunter Workstation.

Flow cytometry

Mig-log phase cultures (OD600, 0.5) were prepared for the detection of

intra-cellular reactive oxygen species (ROS). The oxidant sensor dye,

DCFH2-DA (sigma) was mixed with bacterial strains and kept for 0.5h.

Accordingly, the antibiotic (colistin and/or tigecycline) was added. Then,

bacterial samples (10^5-10^6) diluted with 0.85% saline were subjected to the

analysis of flow cytometry [37, 39]. The resultant FACS data was recorded with

a BD FACSVerse flow cytometer through counting 10,000 cells at a flow rate of 35 ml/min (and/or 14 ml/min). In particular, DCFH florescence was excited with a 488 nm argon laser and emission was detected with the FL1 emission filter at 525 nm using FL1 photomultiplier tub.

Confocal microscopy

As recently described [35], confocal microscopy was conducted to examine the potential effects on bacterial viability exerted by resistance enzymes [MCR-1 and/or Tet(X6)]. Prior to confocal microscopy assays, mid-log phase cultures were processed with the LIVE/DEAD BacLight™ Bacterial Viability Kit (Cat. No. L7012) [35]. Namely, the three strains tested here included i) *E. coli* MG1655 (*mcr-1*/pWSK), ii) MG1655 [*tet*(X6)/pBAD], and iii) MG1655 [*tet*(X6)/pBAD and *mcr-1*/pWSK). Of note, lactose is an inducer of *mcr-1* expression, and arabinose acts as an activator for Tet(X6) enzyme production. After the removal of supernatants, bacterial biofilms were stained with 3% LIVE/DEAD kit solution, and maintained at room temperature in the dark for 15 minutes. Photographs were captured by the confocal laser scanning microscopy (Zeiss LSM 800) with a 63 × oil immersion lens and analyzed using COMSTAT image analysis software. The Tukey–Kramer multiple comparison post hoc test was applied to judge the COMSTAT data. Statistical significance was set at p< 0.01 with T-test.

Bioinformatics

Multiple sequence alignments of Tet(X) variants at the levels of both amino acids and nucleic acids proceeded with ClustalOmega (https://www.ebi.ac.uk/Tools/msa/clustalo). Consequently, the phylogeny of Tet(X) was generated with TreeView (https://www.treeview.co.uk/). Tet(X6) was structurally modeled using Swiss-Model (https://swissmodel.expasy.org/interactive) [53], in which the structural template detected refers to Tet(X2) (PDB: 2Y6Q) [16, 54]. Both GMQE (global model quality estimation) and QMEAN (a global and local absolute quality estimate on the modeled structure) was applied to judge the quality of the modeled structure. Finally, structural presentation and cavity illustration of TetX6 was given with PyMol (https://pymol.org/2).

Accession number

Nucleotide sequence data of *tet(X6)* reported here is available in the Third-Party Annotation Section of the DDBJ/ENA/GenBank databases under the accession number TPA: BK011183. The full genome sequence of the *mcr-1*-harboring plasmid pDJB-3 of swine origin is accessed under the accession number: MK574666.

Acknowledgements

We would like to thank Dr. Yuanyuan Zhang for the technical assistance in flow cytometer assays with BD FACSVerse (Shared Management Platform for Large Instrument, College of Animal Sciences, Zhejiang University). This work was supported by National Key R&D Program of China (2017YFD0500202, YF) and National Natural Science Foundation of China (31830001, 31570027 & 81772142, YF). Dr. Feng is a recipient of the national "Young 1000 Talents" Award of China.

Author contributions

YF designed and supervised this study; YF, YX and LL conducted experiments and analyzed data; YF and BS contributed regents and interpreted data; YF drafted this manuscript.

Competing Interests

We declare that no conflict of interest is present.

References

1. Li J, Nation RL, Turnidge JD, Milne RW, Coulthard K, Rayner CR, et al. Colistin: the re-emerging antibiotic for multidrug-resistant Gram-negative bacterial infections. The Lancet Infectious diseases. 2006;6(9):589-601. doi: 10.1016/S1473-3099(06)70580-1. PubMed PMID: 16931410.

2. Doan TL, Fung HB, Mehta D, Riska PF. Tigecycline: a glycylcycline antimicrobial agent. Clinical therapeutics. 2006;28(8):1079-106. doi: 10.1016/j.clinthera.2006.08.011. PubMed PMID: 16982286.

3. Noskin GA. Tigecycline: a new glycylcycline for treatment of serious infections. Clinical infectious diseases : an official publication of the Infectious Diseases Society of America. 2005;41 Suppl 5:S303-14. doi: 10.1086/431672. PubMed PMID: 16080069.

4. Feng Y. Transferability of MCR-1/2 polymyxin resistance: Complex dissemination and genetic mechanism. ACS infectious diseases. 2018;4(3):291-300. doi: 10.1021/acsinfecdis.7b00201. PubMed PMID: 29397687.

5. Liu YY, Wang Y, Walsh TR, Yi LX, Zhang R, Spencer J, et al. Emergence of plasmid-mediated colistin resistance mechanism MCR-1 in animals and human beings in China: a microbiological and molecular biological study. The Lancet Infectious diseases. 2016;16(2):161-8. doi: 10.1016/S1473-3099(15)00424-7. PubMed PMID: 26603172.

6. Gao R, Hu Y, Li Z, Sun J, Wang Q, Lin J, et al. Dissemination and mechanism for the MCR-1 colistin resistance. PLoS pathogens. 2016;12(11):e1005957. doi: 10.1371/journal.ppat.1005957. PubMed PMID: 27893854; PubMed Central PMCID: PMC5125707.

7. Carroll LM, Gaballa A, Guldimann C, Sullivan G, Henderson LO, Wiedmann M. Identification of novel mobilized colistin resistance gene *mcr-9* in a multidrug-resistant, colistin-susceptible *Salmonella enterica* serotype Typhimurium isolate. mBio. 2019;10(3). doi: 10.1128/mBio.00853-19. PubMed PMID: 31064835; PubMed Central PMCID: PMC6509194.

8. Sun J, Zhang H, Liu YH, Feng Y. Towards understanding MCR-like colistin resistance. Trends in microbiology. 2018;26(9):794-808. doi: 10.1016/j.tim.2018.02.006. PubMed PMID: 29525421.

9. Zhang H, Srinivas S, Xu Y, Wei W, Feng Y. Genetic and biochemical mechanisms for bacterial lipid A modifiers associated with polymyxin resistance. Trends in biochemical sciences. 2019. doi: 10.1016/j.tibs.2019.06.002. PubMed PMID: 31279652.

10. Nguyen F, Starosta AL, Arenz S, Sohmen D, Donhofer A, Wilson DN. Tetracycline antibiotics and resistance mechanisms. Biol Chem. 2014;395(5):559-75. Epub 2014/02/06. doi: 10.1515/hsz-2013-0292. PubMed PMID: 24497223.

11. Li W, Atkinson GC, Thakor NS, Allas U, Lu CC, Chan KY, et al. Mechanism of

tetracycline resistance by ribosomal protection protein Tet(O). Nature communications. 2013;4:1477. Epub 2013/02/14. doi: 10.1038/ncomms2470. PubMed PMID: 23403578; PubMed Central PMCID: PMCPMC3576927.

12. Park J, Gasparrini AJ, Reck MR, Symister CT, Elliott JL, Vogel JP, et al. Plasticity, dynamics, and inhibition of emerging tetracycline resistance enzymes. Nature chemical biology. 2017;13(7):730-6. doi: 10.1038/nchembio.2376. PubMed PMID: 28481346; PubMed Central PMCID: PMC5478473.

13. Moore IF, Hughes DW, Wright GD. Tigecycline is modified by the flavin-dependent monooxygenase TetX. Biochemistry. 2005;44(35):11829-35. doi: 10.1021/bi0506066. PubMed PMID: 16128584.

14. Thaker M, Spanogiannopoulos P, Wright GD. The tetracycline resistome. Cellular and molecular life sciences : CMLS. 2010;67(3):419-31. doi: 10.1007/s00018-009-0172-6. PubMed PMID: 19862477.

15. Yang W, Moore IF, Koteva KP, Bareich DC, Hughes DW, Wright GD. TetX is a flavin-dependent monooxygenase conferring resistance to tetracycline antibiotics. The Journal of biological chemistry. 2004;279(50):52346-52. doi: 10.1074/jbc.M409573200. PubMed PMID: 15452119.

16. Volkers G, Palm GJ, Weiss MS, Wright GD, Hinrichs W. Structural basis for a new tetracycline resistance mechanism relying on the TetX monooxygenase. FEBS letters. 2011;585(7):1061-6. doi: 10.1016/j.febslet.2011.03.012. PubMed PMID: 21402075.

17. Forsberg KJ, Patel S, Wencewicz TA, Dantas G. The tetracycline destructases: A novel family of tetracycline-inactivating enzymes. Chemistry & biology. 2015;22(7):888-97. doi: 10.1016/j.chembiol.2015.05.017. PubMed PMID: 26097034; PubMed Central PMCID: PMC4515146.

18. He T, Wang R, Liu D, Walsh TR, Zhang R, Lv Y, et al. Emergence of plasmid-mediated high-level tigecycline resistance genes in animals and humans. Nature microbiology. 2019;4(9):1450-6. doi: 10.1038/s41564-019-0445-2. PubMed PMID: 31133751.

19. Sun J, Chen C, Cui CY, Zhang Y, Liu X, Cui ZH, et al. Plasmid-encoded *tet*(X) genes that confer high-level tigecycline resistance in *Escherichia coli*. Nature microbiology. 2019;4(9):1457-64. doi: 10.1038/s41564-019-0496-4. PubMed PMID: 31235960.

20. Xu Y, Liu L, Sun J, Feng Y. Limited distribution and mechanism of the TetX4 tetracycline resistance enzyme. Sci Bull (Beijing). 2019:doi: 10.1016/j.scib.2019.08.024 doi: doi: 10.1016/j.scib.2019.08.024.

21. Guiney DG, Jr., Hasegawa P, Davis CE. Expression in *Escherichia coli* of cryptic tetracycline resistance genes from *Bacteroides* R plasmids. Plasmid. 1984;11(3):248-52. PubMed PMID: 6379711.

22. Speer BS, Bedzyk L, Salyers AA. Evidence that a novel tetracycline resistance gene found on two *Bacteroides* transposons encodes an NADP-requiring oxidoreductase. Journal of bacteriology. 1991;173(1):176-83. doi:

10.1128/jb.173.1.176-183.1991. PubMed PMID: 1846135; PubMed Central PMCID: PMC207172.

23. Thakare R, Dasgupta A, Chopra S. Eravacycline for the treatment of patients with bacterial infections. Drugs of today. 2018;54(4):245-54. doi: 10.1358/dot.2018.54.4.2800623. PubMed PMID: 29869646.

24. Forde BM, Zowawi HM, Harris PNA, Roberts L, Ibrahim E, Shaikh N, et al. Discovery of *mcr-1*-mediated colistin resistance in a highly virulent *Escherichia coli* lineage. mSphere. 2018;3(5): pii: e00486-18. doi: 10.1128/mSphere.00486-18. PubMed PMID: 30305321; PubMed Central PMCID: PMC6180223.

25. Xu Y, Lin J, Cui T, Srinivas S, Feng Y. Mechanistic insights into transferable polymyxin resistance among gut bacteria. The Journal of biological chemistry. 2018;293(12):4350-65. doi: 10.1074/jbc.RA117.000924. PubMed PMID: 29462787; PubMed Central PMCID: PMC5868246.

26. Mao J, Liu W, Wang W, Sun J, Lei S, Feng Y. Antibiotic exposure elicits the emergence of colistin- and carbapenem-resistant *Escherichia coli* coharboring MCR-1 and NDM-5 in a patient. Virulence. 2018;9(1):1001-7. doi: 10.1080/21505594.2018.1486140. PubMed PMID: 30047824; PubMed Central PMCID: PMC6067848.

27. Sun J, Yang RS, Zhang Q, Feng Y, Fang LX, Xia J, et al. Co-transfer of bla_{NDM-5} and *mcr-1* by an IncX3-X4 hybrid plasmid in *Escherichia coli*. Nature microbiology. 2016;1:16176. doi: 10.1038/nmicrobiol.2016.176. PubMed PMID: 27668643.

28. Li X, Xie Y, Liu M, Tai C, Sun J, Deng Z, et al. oriTfinder: a web-based tool for the identification of origin of transfers in DNA sequences of bacterial mobile genetic elements. Nucleic Acids Res. 2018;46(W1):W229-W34. Epub 2018/05/08. doi: 10.1093/nar/gky352. PubMed PMID: 29733379; PubMed Central PMCID: PMCPMC6030822.

29. Zhi C, Lv L, Yu LF, Doi Y, Liu JH. Dissemination of the mcr-1 colistin resistance gene. The Lancet Infectious diseases. 2016;16(3):292-3. Epub 2016/03/15. doi: 10.1016/S1473-3099(16)00063-3. PubMed PMID: 26973307.

30. Gillings MR. Integrons: past, present, and future. Microbiol Mol Biol Rev. 2014;78(2):257-77. Epub 2014/05/23. doi: 10.1128/MMBR.00056-13. PubMed PMID: 24847022; PubMed Central PMCID: PMCPMC4054258.

31. Zhang AN, Li LG, Ma L, Gillings MR, Tiedje JM, Zhang T. Conserved phylogenetic distribution and limited antibiotic resistance of class 1 integrons revealed by assessing the bacterial genome and plasmid collection. Microbiome. 2018;6(1):130. Epub 2018/07/23. doi: 10.1186/s40168-018-0516-2. PubMed PMID: 30031405; PubMed Central PMCID: PMCPMC6054849.

32. Chiou CS, Jones AL. Nucleotide sequence analysis of a transposon (Tn5393) carrying streptomycin resistance genes in Erwinia amylovora and other gram-negative bacteria. J Bacteriol. 1993;175(3):732-40. Epub 1993/02/01. doi: 10.1128/jb.175.3.732-740.1993. PubMed PMID: 8380801; PubMed Central

PMCID: PMCPMC196212.

33. Wang R, van Dorp L, Shaw LP, Bradley P, Wang Q, Wang X, et al. The global distribution and spread of the mobilized colistin resistance gene mcr-1. Nat Commun. 2018;9(1):1179. Epub 2018/03/23. doi: 10.1038/s41467-018-03205-z. PubMed PMID: 29563494; PubMed Central PMCID: PMCPMC5862964.

34. Snesrud E, He S, Chandler M, Dekker JP, Hickman AB, McGann P, et al. A Model for Transposition of the Colistin Resistance Gene mcr-1 by ISApl1. Antimicrobial agents and chemotherapy. 2016;60(11):6973-6. Epub 2016/09/14. doi: 10.1128/AAC.01457-16. PubMed PMID: 27620479; PubMed Central PMCID: PMCPMC5075121.

35. Yang Q, Li M, Spiller OB, Andrey DO, Hinchliffe P, Li H, et al. Balancing *mcr-1* expression and bacterial survival is a delicate equilibrium between essential cellular defence mechanisms. Nature communications. 2017;8(1):2054. doi: 10.1038/s41467-017-02149-0. PubMed PMID: 29233990; PubMed Central PMCID: PMC5727292.

36. Zhang H, Wei W, Huang M, Umar Z, Feng Y. Definition of a family of nonmobile colistin resistance (NMCR-1) determinants suggests aquatic reservoirs for MCR-4. Advanced science. 2019;6(11):1900038. doi: 10.1002/advs.201900038. PubMed PMID: 31179218; PubMed Central PMCID: PMC6548957.

37. Zhang H, Hou M, Xu Y, Srinivas S, Huang M, Liu L, et al. Action and mechanism of the colistin resistance enzyme MCR-4. Communications biology. 2019;2:36. doi: 10.1038/s42003-018-0278-1. PubMed PMID: 30701201; PubMed Central PMCID: PMC6347640.

38. Xu Y, Zhong LL, Srinivas S, Sun J, Huang M, Paterson DL, et al. Spread of MCR-3 colistin resistance in China: An epidemiological, genomic and mechanistic study. EBioMedicine. 2018;34:139-57. doi: 10.1016/j.ebiom.2018.07.027. PubMed PMID: 30061009; PubMed Central PMCID: PMC6116419.

39. Zhang H, Zong Z, Lei S, Srinivas S, Sun J, Feng Y, et al. A genomic, evolutionary, and mechanistic study of MCR-5 action suggests functional unification across the MCR family of colistin resistance. Advanced science. 2019;6(11):1900034. doi: 10.1002/advs.201900034. PubMed PMID: 31179217; PubMed Central PMCID: PMC6548960.

40. Wang RF, Kushner SR. Construction of versatile low-copy-number vectors for cloning, sequencing and gene expression in *Escherichia coli*. Gene. 1991;100:195-9. PubMed PMID: 2055470.

41. Khlebnikov A, Risa O, Skaug T, Carrier TA, Keasling JD. Regulatable arabinose-inducible gene expression system with consistent control in all cells of a culture. Journal of bacteriology. 2000;182(24):7029-34. doi: 10.1128/jb.182.24.7029-7034.2000. PubMed PMID: 11092865; PubMed Central PMCID: PMC94830.

42. Wang Q, Li Z, Lin J, Wang X, Deng X, Feng Y. Complex dissemination of the

diversified *mcr-1*-harbouring plasmids in *Escherichia coli* of different sequence types. Oncotarget. 2016;7(50):82112-22. doi: 10.18632/oncotarget.12621. PubMed PMID: 27741523; PubMed Central PMCID: PMC5347679.

43. Stokes JM, Lopatkin AJ, Lobritz MA, Collins JJ. Bacterial metabolism and antibiotic efficacy. Cell Metab. 2019;30(2):251-9. Epub 2019/07/08. doi: 10.1016/j.cmet.2019.06.009. PubMed PMID: 31279676.

44. Lopatkin AJ, Stokes JM, Zheng EJ, Yang JH, Takahashi MK, You L, et al. Bacterial metabolic state more accurately predicts antibiotic lethality than growth rate. Nature microbiology. 2019. Epub 2019/08/28. doi: 10.1038/s41564-019-0536-0. PubMed PMID: 31451773.

45. Belenky P, Ye JD, Porter CB, Cohen NR, Lobritz MA, Ferrante T, et al. Bactericidal antibiotics induce toxic metabolic perturbations that lead to cellular damage. Cell Rep. 2015;13(5):968-80. Epub 2015/11/14. doi: 10.1016/j.celrep.2015.09.059. PubMed PMID: 26565910; PubMed Central PMCID: PMCPMC4648786.

46. Lobritz MA, Belenky P, Porter CB, Gutierrez A, Yang JH, Schwarz EG, et al. Antibiotic efficacy is linked to bacterial cellular respiration. Proceedings of the National Academy of Sciences of the United States of America. 2015;112(27):8173-80. Epub 2015/06/24. doi: 10.1073/pnas.1509743112. PubMed PMID: 26100898; PubMed Central PMCID: PMCPMC4500273.

47. Kohanski MA, Dwyer DJ, Hayete B, Lawrence CA, Collins JJ. A common mechanism of cellular death induced by bactericidal antibiotics. Cell. 2007;130(5):797-810. Epub 2007/09/07. doi: 10.1016/j.cell.2007.06.049. PubMed PMID: 17803904.

48. Brynildsen MP, Winkler JA, Spina CS, MacDonald IC, Collins JJ. Potentiating antibacterial activity by predictably enhancing endogenous microbial ROS production. Nat Biotechnol. 2013;31(2):160-5. Epub 2013/01/08. doi: 10.1038/nbt.2458. PubMed PMID: 23292609; PubMed Central PMCID: PMCPMC3568245.

49. Dwyer DJ, Belenky PA, Yang JH, MacDonald IC, Martell JD, Takahashi N, et al. Antibiotics induce redox-related physiological alterations as part of their lethality. Proceedings of the National Academy of Sciences of the United States of America. 2014;111(20):E2100-9. Epub 2014/05/08. doi: 10.1073/pnas.1401876111. PubMed PMID: 24803433; PubMed Central PMCID: PMCPMC4034191.

50. Sullivan MJ, Petty NK, Beatson SA. Easyfig: a genome comparison visualizer. Bioinformatics. 2011;27(7):1009-10. doi: 10.1093/bioinformatics/btr039. PubMed PMID: 21278367; PubMed Central PMCID: PMC3065679.

51. Ye H, Li Y, Li Z, Gao R, Zhang H, Wen R, et al. Diversified *mcr-1*-harbouring plasmid reservoirs confer resistance to colistin in human gut microbiota. mBio. 2016;7(2):e00177. doi: 10.1128/mBio.00177-16. PubMed PMID: 27048797; PubMed Central PMCID: PMC4817250.

52. Balouiri M, Sadiki M, Ibnsouda SK. Methods for in vitro evaluating

antimicrobial activity: A review. Journal of pharmaceutical analysis. 2016;6(2):71-9. doi: 10.1016/j.jpha.2015.11.005. PubMed PMID: 29403965; PubMed Central PMCID: PMC5762448.

53. Biasini M, Bienert S, Waterhouse A, Arnold K, Studer G, Schmidt T, et al. SWISS-MODEL: modelling protein tertiary and quaternary structure using evolutionary information. Nucleic acids research. 2014;42(Web Server issue):W252-8. doi: 10.1093/nar/gku340. PubMed PMID: 24782522; PubMed Central PMCID: PMC4086089.

54. Walkiewicz K, Benitez Cardenas AS, Sun C, Bacorn C, Saxer G, Shamoo Y. Small changes in enzyme function can lead to surprisingly large fitness effects during adaptive evolution of antibiotic resistance. Proceedings of the National Academy of Sciences of the United States of America. 2012;109(52):21408-13. doi: 10.1073/pnas.1209335110. PubMed PMID: 23236139; PubMed Central PMCID: PMC3535585.

Supporting information

Supporting tables

Supporting figures

Fig. S1 Nucleotide acid sequence analysis of *tet*(X3) and *tet*(X6)

To determine the similarity between *tet*(X3) and *tet*(X6), their nucleotide acid sequences were subjected to Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/) , and resultant form of sequence is given with ESPript 3.0 (http://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi). Identical residues are in white letters with black background, and different residues are black letters with white background. The identity between *tet*(X3) and *tet*(X6) is 97.1%, with 31 substitution of 1137 residues in total.

Fig. S2 Evidence that an intact *tet*(X6) is only harbored on one plasmid and two contigs thus far

Sequence alignment was conducted as described in **Fig. S1**.

An initial codon "ATG" and the stop codon "TAA" are underlined.

Fig. S3 Sequence alignment of the Tet(X6) enzyme with other five homologs (X0 to X5)

Multiple sequence alignment was conducted with Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/), generating its output with ESPript 3.0 (http://espript.ibcp.fr/ESPript/cgi-bin/ESPript.cgi). The putative substrate-loading cavity is composed of FAD-interactive residues (with red triangles) and Tigecycline-binding residues (with blue triangles). Identical residues are indicated with white letters in red background.

Fig. S4 Co-occurrence of *tet*(X6) and *mcr-1* gives co-resistance to colistin and tigecycline

A. Contrasting the level of tigecycline resistance by Tet(X) variants

The *E. coli* MG1655 carrying different variants of *tet*(X) [X3, X4, and X6] were maintained at 37°C on LBA agar plates containing tigecycline in series of dilution.

B. The MCR-1 confers phenotypic colistin resistance in the *E. coli* MG1655

The derivatives of *E. coli* MG1655 bear *mcr-1* or *tet*(X6) alone (or both) were spotted on LBA agar plates containing colistin in series of dilution, and maintained at 37°C for overnight.

Fig. S5 MS identity of the purified two proteins Tet(X3) and Tet(X6)
A. MS-based identification of polypeptide fragments from the purified Tet(X3) protein
B. MS-based determination of the purified TetX6 protein
The underlined letters denote the polypeptides that match Tet(X3) [and/or Tet(X6)] protein.

Fig. S6 Use of LC/MS to identify the oxygenated product of tigecycline by Tet(X3) enzyme
In the spectrum, the peak of 586.2 (m/z) refers to tigecycline, whereas the other peak of 602.2 (m/z) denotes an oxygenated product of tigecycline. Chemical structures were given with ChemDraw.

Fig. S7 Visualization for destruction of tetracycline by Tet(X3) [and Tet(X6)] enzyme
Unlike the fact that the blank and the negative control (liquid culture of *E. coli* with empty vector alone) display yellow, bacterial culture of Tet(X3) [and/or Tet(X6)]-expressing *E. coli* gives dark. This indicates that the expression of Tet(X3) [and Tet(X6)] leads to the oxygenation of tetracycline, which is fully consistent with the observation of other tetracycline-inactivating enzymes by Forsberg and coworkers [17].
0.1% arabinose is used to induce the expression of pBAD24-borne *tet*(X3) [and *tet*(X6)].
Designation: Blank, the LB liquid medium containing tetracycline; Vec, pBAD24.

Fig. S8 Use of E-test to evaluate phenotypic growth of MG1655 strains expressing *tet*(X3)/*tet*(X6)
Semi-solid medium was poured, which was supplemented with the MG1655 strains with the empty vector alone, or the plasmid-borne *tet*(X3) [and/or *tet*(X6)], accordingly. E-test strip is featuring with a gradient concentration of tigecycline. The cut-off value of tigecycline resistance is indicated with an arrow. A representative result of three independent experiments is given.

  Figure legends

Figure 1 Genetic context-based analyses for co-transfer of *tet*(X6) and *mcr-1*
by a single plasmid
A. Co-linear genome alignment of the *tet*(X6), and *mcr-1*-coharboring plasmid
pMS8345A with other three IncHI2-type mega plasmids (pDJB-3, pYps.F1, and
pHNSHP45-2)

The backbones of the four resistance plasmids are of high similarity. Except
that pYps.F1 (Acc. no.: LT221036) of *Yersinia pseudotuberculosis* lacks *mcr-1*,
all the other three resistance plasmids here carry this gene.
pDJB-3 (Acc. no.: MK574666) is a new variant of the *mcr-1*-containing plasmid
we reported here. Unlike that pDJB-3 lacking multidrug resistance region
(MDR), the remaining three plasmids possess different MDRs.
ICEs are colored: *oriT* in brown, relaxase gene in orange, T4SS in green, and
T4CP in purple.

B. Fine dissection of the *tet*(X6)-positive MDR regions in pMS8345A and
p15C38-2

As for this unique MDR region in pMS8345A, three Class 1 integrons are
circled by dashed lines, 14 kinds of antibiotic resistance determinants like
addA1 (with red arrows), are integrated, and no less than 10 different mobile
elements such as IS*Ecp1* and Tn*As1* (indicated with yellow arrows) are
determined. In particular, a new *tet*(X) variant of tigecycline resistance
enzymes, named *tet*(X6) (in bold red letters), is adjacent to *sul1* that conferring
resistance to sulfonamide. Additionally, five *attL* sites (highlighted in purple)
are detected.

C. Genetic context of the *mcr-1*-centering transposal region in pMS8345A,
pHNSHP45-2, and pDJB-3

Colored arrows indicate ORFs and the shaded region depicts sequence
similarity (70%-100%). The resistance genes are highlighted in red and mobile
elements are highlighted in yellow.

Designations: pDJB-3 (156.5kb, IncHI2/*E. coli*, China/2017, Acc. no.:
MK574666); pYps.F1 (200.66kb, IncH1/*Y. pseudotuberculosis*, France/2017,
Acc. no.: LT221036); pHNSHP45-2 (251.5kb, IncHI2/*E. coli*, China/2013, Acc.
no.: KU341381); p15C38-2 (150.745kb, IncA/C2, *E. coli*/Japan/2003, Acc. no.:
LC501585); and pMS8345A (241.162kb, IncHI2/*E. coli*, Qutar/2017, Acc. no.:
CP025402).

Figure 2 Phylogeny of Tet(X6)

The protein sequences of TetX variants were selected from NCBI database, and the accession numbers were labeled accordingly on the graph. Clustal Omega (https://www.ebi.ac.uk/Tools/msa/clustalo/) was applied to generate the phylogenetic tree, whose final output is built with TreeView (https://www.treeview.co.uk/).

In addition to Tet(X3) (colored pink), all the eight newly-named TetX variants (X6 to X13) are labeled in red. Besides the pMS8345A plasmid producing TetX6, two uncultured bacterium contigs (TE_7F_Contig_3 and TG_7F_Contig_3) from Latrine, EI Salvador, were found to carry an intact version of *tet*(X6).

Figure 3 Structural and functional analysis of the substrate tigecycline-loading tunnel in Tet(X6)

Surface structure of Tet(X6) with tigecycline-loading tunnel in the front view (**A**) and the rotated view (**B**).

The substrate-loading tunnel in open state is indicated with an arrow.

C. Sectional view of tigecycline-occupied cavity in Tet(X6) protein

D. Sectional view of FAD-binding cavity in Tet(X6) enzyme

E. An enlarged view of tigecycline-binding cavity in Tet(X6)

F. Structural snapshot of the FAD cofactor-binding motif in Tet(X6)

The critical residues are labeled. The molecule of tigecycline is colored in cherry red, and the FAD molecule is labeled in purple.

G. Use of site-directed mutagenesis to assay the five tigecycline-binding residues in Tet(X6)

H. Structure-guided alanine substitution analyses of FAD-interacting residues in Tet(X6)

Negative control is the *E. coli* MG1655 alone, and positive control refers to the MG1655 expressing the wild-type of *tet*(X6) (**Table S1**). The pBAD24-driven expression of Tet(X6) and its derivatives is triggered by the addition of 0.2% arabinose. Three independent assays were conducted, each of which is indicated with a dot.

Figure 4 Action and mechanism for Tet(X6) tigecycline resistance

A. The bioassay for Tet(X6) [and Tet(X3)]-mediated destruction of tigecycline

The presence of tigecycline on paper disks leads to the appearance of a transparent zone of microbial inhibition. In contrast, the inhibition zone is absent upon the destruction of tigecycline by Tet(X6) [and/or Tet(X3)].

B. Gel filtration analysis of the purified Tet(X6) resistance enzyme

To address the solution structure of Tet(X6) and Tet(X3), the method of size

exclusion chromatography was applied. The protein samples were loaded into

Superdex 200/300GL size exclusion column (GE Healthcare, USA). The

eluted protein of interest (indicated with an arrow) is visualized in the

right-hand inside gel (15% SDS-PAGE). The almost-identical elution volume

(~16ml) of Tet(X6) and Tet(X3) is generally consistent with the apparent size of

molecular weight ~36kDa in the inside PAGE gel, validating the monomeric

form. Of note, the yellow color of Tet(X6) [Tet(X3)] solution is given in inside gel

on the left hand, indicating the presence of a FAD cofactor-bound protein form.

LC/MS identification of tigecycline (**C**) and its oxygenated product by Tet(X6)

enzyme (**D**)

Compared with the peak of tigecycline at 586.2 (m/z), the oxygenated

derivative of tigecycline is reflected by a unique peak of 602.2 (m/z) in that it is

added with an oxygen atom.

E. A chemical reaction model that Tet(X6) destructs/inactivates tigecycline

It was adapted appropriately from Xu *et al.* [20] with permission.

FAD-Tet(X6) denotes Tet(X6) enzyme in the form of binding FAD cofactor.

Chemical structures were given with ChemDraw.

Designations: blank, the LBA media with only tigecycline; vec, the vector of

pET21a; LC/MS, liquid chromatography mass spectrometry; FAD, flavin

adenine dinucleotide; NADP$^+$, the oxidized form of nicotinamide adenine

dinucleotide phosphate; NADPH, the reduced form of nicotinamide adenine

dinucleotide phosphate.

Figure 5 Use of flow cytometry to measure intracellular ROS level induced by

colistin and/or tigecycline

A. FACS analysis of basal level of ROS in the negative strain *E. coli* MG1655

with empty vector alone

Intracellular ROS level is boosted upon the addition of either colistin (**B**) or

tigecycline (**C**) into the MG1655 strain

Cytosolic ROS level of the MG1655 strains expressing *mcr-1* (**D**) or *tet*(X6) (**E**)

F. Determination of ROS level in the MG1655 carrying both *mcr-1* and *tet*(X6)

G. Colistin cannot stimulate the ROS production in the MG1655 strain

expressing *mcr-1*

H. Tigecycline cannot activates the ROS production in the MG1655 strain expressing *tet*(X6)

I. The mixture of tigecycline and colistin cannot significantly alter the cytosolic ROS accumulation in the MG1655 strain coharboring *mcr-1* and *tet*(X6)

J. Flow cytometry-based determination of relative level of intracellular ROS in different *E. coli* MG1655 strains in response to colistin and/or tigecycline

Along with Tukey-Kramer multiple comparisons post-test, the data is given using one-way analysis of variance (ANOVA). *p-value is less than 0.001.

Abbreviations: FACS, fluorescence activated cell sorting; ROS, reactivated oxygen species.

K. A scheme for colistin/tigecycline-induced accumulation of ROS, and its interfered formation by MCR-1/Tet(X6)

Blue asterisk denotes colistin, and green square refers to tigecycline. The oxygenated form of tigecycline is indicated with green square attached with a red dot. The reactive oxygen species (ROS) are illustrated with the symbols of lightning.

Figure 6 Use of growth curves to monitor metabolic fitness caused by *mcr-1* and *tet*(X6) in *E. coli*

A. The empty vector exerts no effect on bacterial growth of the *E. coli* MG1655

B. Expression of *tet*(X6) causes slightly the delayed growth of the *E. coli* MG1655

C. Expression of *mcr-1* gives significantly metabolic fitness in its recipient host *E. coli* MG1655 slightly the delayed growth of the *E. coli* MG1655

D. Co-expression of *mcr-1* and *tet*(X6) cannot lead to synergistic fitness cost of the recipient strain

The data was given in means ± SD from three independent plotting of growth curves.

Figure 7 Confocal microscopy-based visualization for bacterial fitness cost caused by functional expression of *tet*(X6) (and/or *mcr-1*) in *E. coli*

A&B Bacterial viability of the negative control strains, *E. coli* MG1655 with the empty vector alone is not affected by the presence of the inducer arabinose and/or lactose

C&D The ratio of DEAD/LIVE cells suggests that arabinose-induced expression of *mcr-1* gives appreciable level of fitness cost in *E. coli*

E&F Regardless of the inducer lactose added, the MG1655 strains harboring *tetX6* is indistinguishable, when compared with those of negative control strains

G. The *mcr-1*, and *tet*(X6)-coharboring MG1655 does not display the phenotypic fitness cost, without the addition of neither arabinose nor lactose

H. The co-expression of MCR-1 and Tet(X6) cannot exert synergistic effect of metabolic fitness

I. Bacterial viability-based measurement of bacterial fitness costs in various MG1655 strains bearing either *mcr-1* or *tet*(X6) (or both)

Bacterial viability was determined through cell staining with LIVE/DEAD kit, which is followed by imaging with confocal laser scanning microscopy. The color of green and red separately denotes alive and dead cells.

Data was collected from four independent trials, and evaluated using one-way analysis of variance (ANOVA) followed by Tukey–Kramer multiple comparisons post hoc test. **p-value is less than 0.01. ns, no significance.

1   **Table 1** Substrate-loading cavities across TetX-type enzymes

Enzymes	Substrate-loading cavities									
	FAD-binding sites						Tetracycline-recognizable sites			
Tet(X0)	E46	R47	R117	D311	P318	Q322	Q192	R213	H234	G236	M375
Tet(X1)	E36	R37	R107	D302	P309	Q313	Q182	R203	T224	G226	M366
Tet(X2)	E46	R47	R117	D311	P318	Q322	Q192	R213	H234	G236	M375
Tet(X3)	E36	R37	R107	D301	P308	Q312	Q182	R203	H224	G226	M365
Tet(X4)	E43	R44	R114	D308	P315	Q319	Q189	R210	H231	G233	M372
Tet(X5)	E46	R47	R117	D310	P317	Q321	Q192	R213	Y234	G236	M373
Tet(X6)	E36	R37	R107	D301	P308	Q312	Q182	R203	H224	G226	M365
Tet(X7)	E36	R37	R107	D301	P308	Q312	Q182	R203	H224	G226	M365
Tet(X8)	E36	R37	R107	D301	P308	Q312	Q182	R203	H224	G226	M365
Tet(X9)	E36	R37	R107	D301	P308	Q312	Q182	R203	H224	G226	M365
Tet(X10)	E45	R46	R116	D310	P317	Q321	Q191	R212	H233	G235	M374
Tet(X11)	E36	R37	R107	D301	P308	Q312	Q182	R203	H224	G226	M365
Tet(X12)	E45	R46	R116	D310	P317	Q321	Q191	R212	H233	G235	M374
Tet(X13)	E46	R47	R117	D310	P317	Q321	Q192	R213	H234	G236	M373
Tet(X2/4)-P	E46	R47	R117	D310	P317	Q321	Q192	R213	H234	G236	M373

3   The substituted residues are indicated in yellow background.

A

pDJB-3, *E. coli*
(156.5kb, MK574666)

pYps.F1, *Y. pseudotuberculosis*
(200.66kb, LT221036)

pHNSHP45-2, *E. coli*
(251.493kb, KU341381)

pMS8345A, *E. coli*
(241.162kb, CP025402)

B

MDR region
(~40kb)

75976   116262

Resistance gene
Mobile element
oriT
Relaxase
T4SS
T4CP

p15C38-2
(LC501585)

C

pHNSHP45-2
(KU341381)

pDJB-3
(MK574666)

Tet(X2/4)-P (*Riemerella anatipestifer*; AKO71464)

Tet(X4) (*E. coli*, 2017; MK134376)

Tet(X2) (*Bacteroides*, 2001; AJ311171)

Tet(X0) (*Bacteroides*, 1991; M37699)

Tet(X1) (*Bacteroides*, 2001; AJ311171)

Tet(X5) (*Acinetobacter baumannii*, 2018; MK134375)

Tet(X13) (*Flavobacterium marinum*, 2019; WP_091095665)

Tet(X8) (TG_0402_Contig_27/Soils, El Salvador, 2012; KU548516)

Tet(X6) (TE_7F_Contig_3/Latrine, El Salvador, 2012; KU547176)

Tet(X6) (pMS8345A/*E.coli*, 2017; NZ_CP025402)

Tet(X6) (TG_7F_Contig_3/Latrine, El Salvador, 2012; KU547185)

Tet(X9) (TG_6F_Contig_3/Latrine, El Salvador, 2012; KU547130)

Tet(X3) (*P. aeruginosa*, 2002; AB097942)

Tet(X10) (*Acinetobacter indicus*, 2019; WP_075168333)

Tet(X11) (*E. coli*, 2014; KHO55827)

Tet(X12) (2019; WP_064754130)

Tet(X7) (TE_6F_Contig_7/Latrine, El Salvador, 2012; KU547125)

0.05

A Tigecycline-binding domain

Helix bridge

FAD-binding domain

B Tigecycline-binding domain

Helix bridge

FAD-binding domain

C Sectional view of tigecycline-occupied cavity

D Sectional view of FAD-binding cavity

E H224 G226 Q182 R203 M365

Tigecycline-interactive cavity

F R107 P308 Q312 D301 R37 E36

FAD-centering cavity

G Tigecycline-binding sites

Level of tigecycline resistance (μg/ml)

MG1655 TetX6 Q182A R203A H224A G226A M365A

H FAD-interacting residues

Level of tigecycline resistance (μg/ml)

MG1655 TetX6 E36A R37A R107A D301A P308A Q312A

A MG1655+Vec

C MG1655+MCR-1

E MG1655+Tet(X6)

G MG1655+MCR-1/Tet(X6)

No arabinose or lactose

B MG1655+Vec

D MG1655+MCR-1

F MG1655+Tet(X6)

H MG1655+MCR-1/Tet(X6)

plus arabinose or lactose

I

Ratio of dead/live bacteria

p< 0.01

p< 0.01

ns

Vec   MCR-1   Tet(X6)   MCR-1/Tet(X6)