Genomic islands of differentiation in a rapid avian radiation have been driven by recent selective sweeps

Hussein A. Hejase[1*], Ayelet Salman-Minkov[2], Leonardo Campagna[3,4], Melissa J. Hubisz[5], Irby J. Lovette[3,4], Ilan Gronau[2], Adam Siepel[1*]

[1]Simons Center for Quantitative Biology, Cold Spring Harbor Laboratory, Cold Spring Harbor, NY, USA

[2]Efi Arazi School of Computer Science, The Herzliya Interdisciplinary Center (IDC), Herzliya, Israel

[3]Fuller Evolutionary Biology Program, Cornell Lab of Ornithology, Ithaca, NY, USA

[4]Department of Ecology and Evolutionary Biology, Cornell University, Ithaca, NY, USA

[5]Department of Biological Statistics and Computational Biology, Cornell University, Ithaca, NY, USA

* To whom correspondence should be addressed: hijazi@cshl.edu, asiepel@cshl.edu

Abstract

Numerous studies of emerging species have identified genomic "islands" of elevated differentiation against a background of relative homogeneity. The causes of these islands remain unclear, however, with some signs pointing toward "speciation genes" that locally restrict gene flow and others suggesting selective sweeps that have occurred within nascent species after speciation. Here, we examine this question through the lens of recently obtained genome sequence data for five species of southern capuchino seedeaters, finch-like birds from South America that have undergone a species radiation during the last ~50,000 generations. By applying newly developed statistical methods for ancestral recombination graph inference and machine-learning methods for the prediction of selective sweeps, we show that the striking islands of differentiation in these birds appear to be generally associated with relatively recent, species-specific selective sweeps, most of which are predicted to be "soft" sweeps acting on standing genetic variation. Many of these sweeps coincide with genes associated with melanin-based variation in plumage, suggesting a prominent role for sexual selection. At the same time, a few loci also exhibit indications of possible selection against gene flow. These observations shed new light on the complex manner in which natural selection shapes genome sequences during speciation.

Introduction

The question of how new species arise is one of the oldest and thorniest puzzles in evolutionary biology, having occupied investigators at least since Darwin and Wallace[1–3]. Indeed, formulating a genetic basis for the process of speciation was a key component of the neodarwinian synthesis[1,4,5]. By the middle of the last century, theories of speciation had come to rely heavily on the notion of allopatry, or geographical or ecological barriers to interbreeding[6]. However, it has since become clear that separate species may also emerge in the absence of such barriers. Claims of such sympatric speciation can be controversial and difficult to verify[7,8], but now appear to be well supported in numerous cases[8–16].

In recent decades, inexpensive, large-scale DNA sequencing has allowed molecular genetic studies of speciation to be extended from traditional laboratory model systems to natural populations. Particularly useful are natural systems that consist of populations in the early stages of genetic separation—that is, groups of organisms "caught in the act" of speciation[1], often in sympatry. These systems allow genomic regions with relevant differences to stand out clearly against a background of relatively low genetic differentiation. Examples include *Heliconius* butterflies[17,18], *Anopheles* mosquitos[19,20], pea aphids[21], stick insects[22], sunflowers[23], monkeyflowers[24], house mice[25,26], threespine stickleback[27] and cichlid[28] fish, and various birds, including carrion and hooded crows[29–31], flycatchers[32], and blue- and golden-winged warblers[33]. In addition to enabling investigation of the genetic architecture of reproductive barriers, these systems allow many other questions to be addressed, such as what is the timeline of speciation, and what are the roles of chromosomal rearrangements and sex chromosomes[1]? In some cases, they also allow identification of particular mutations underlying important differences between incipient species.

A number of such studies have focused, in particular, on the intriguing observation that recently separated species often exhibit local genomic regions of pronounced interspecies differentiation against a background of relative homogeneity—a phenomenon dubbed "islands of differentiation" or, sometimes, "islands of speciation"[1,17,19–21,23,25,26,34]. Early on, these islands were widely believed to reflect "speciation genes" (sometimes called "barrier loci") that contribute in some way to reduced gene flow, presumably through inviability, sterility, or reduced fitness of hybrids[19,35]. Later, however, it was shown that a similar local elevation of genetic differentiation could be generated instead by reductions of genetic diversity within one or both nascent species during and following speciation, stemming from selective sweeps and possibly background selection[36–38]. In other words, the observed "islands" may not reflect differential gene flow, but instead, differential influences of selection on linked sites. This ambiguity follows from the

widespread use of relative measures of genetic differentiation such as F_{ST}, which can be elevated either by increases in interspecies divergence or decreases in intraspecies variation. Some recent studies have considered competing hypotheses for the origins of islands of differentiation and found mixed evidence regarding their primary causes[30,31,34].

Recently, we collected whole-genome sequence data for another natural system: southern capuchino seedeaters of the genus *Sporophila*, a group of passerine birds native to South America. These birds are similar to Darwin's finches (also members of the family Thraupidae) in that they appear to have recently separated into numerous distinct and largely genetically separate species. The southern capuchino species, however, are morphologically and ecologically indistinct and differ primarily in male plumage, suggesting that sexual selection may have been important in driving speciation[39–41]. These species have adjacent and, in some cases, overlapping ranges[42], and can be syntopic at breeding sites. In addition, they include many pairs at similar levels of genetic differentiation, allowing for numerous quasi-independent pairwise comparisons.

Our data set includes nine species of southern capuchinos, eight of which are estimated to have emerged <50,000 generations ago from a large, highly polymorphic ancestral species[43]. We focus here on the five of these eight species that were most broadly sampled (12 diploid individuals per species), including *Sporophila pileata* (pil), *S. melanogaster* (mel), *S. nigrorufa* (nig), *S. palustris* (pal), and *S. hypoxantha* (hyp). The combination of a recent radiation and a large ancestral population for these species leads to modest levels of genetic differentiation and a great deal of shared polymorphism, reflecting widespread incomplete lineage sorting. Nevertheless, these genomes include numerous striking islands of differentiation, as defined by peaks in pairwise F_{ST}, and these islands are strongly enriched for genes in the melanogenesis pathway, suggesting an association with melanin-based variation in plumage[42]. Campagna *et al.* focused in particular on 25 peaks with elevated F_{ST}, which represent candidate targets of selection during or after speciation. Interestingly, many of these islands appear recurrently across different pairwise comparisons. In addition, 10 of the 25 peaks are located on the *Z* chromosome.

In this paper, we re-examine these islands of differentiation in the southern capuchinos, as well as broader patterns of diversity across their genomes. Rather than focusing on the use of summary statistics in windows across the genome, as in most previous studies, we base our analysis on the underlying genealogies that precisely describe the relationships among the sequenced genomes. In particular, we make use of methods we recently developed for sequence-based statistical inference of the ancestral recombination graph (ARG), which describes both genealogical relationships and changes to those relationships along the genome due to historical

recombination events[43,44]. Furthermore, we combine ARG inference with the latest machine-learning methods[45] to identify selective sweeps of various ages and types, and systematically compare these predictions with the previously identified islands of differentiation. As we discuss in detail below, we find multiple lines of evidence for a strong relationship between selective sweeps—particularly recent, species-specific "soft" sweeps—and the observed islands. By contrast, we find much less evidence for an influence from selection against gene flow, with a few notable exceptions. Altogether, our study provides new insights into the processes leading to genetic differentiation between species in an ideal model system for rapid speciation.

Results

Genealogical Patterns in F_{ST} Peaks

We began by reanalyzing the 25 F_{ST} peaks (**Supplementary Table S1 and Supplementary Figure S1**) identified by Campagna et al.[42], with the goal of gaining further insight into their evolutionary causes. As noted above, these peaks are presumed to reflect the speciation process in some manner, but the specific mechanisms that generated them have not been explored.

We applied the *ARGweaver* program to the 60 individual genome sequences across 1.06 Gb of sequenced scaffolds (see **Methods**). We then summarized the sampled genealogies both within the peaks and in flanking "background" regions using several features, including the expected time to most recent common ancestry (TMRCA) within each species—which should be reduced in peaks in the presence of local sweeps but unaffected by selection against gene flow between species during speciation—and the cross-coalescence time between species—which should be increased in peaks by selection against gene flow but unaffected by local sweeps (**Figure 1**). We also calculated the standard summary statistics F_{ST} and d_{XY} in a 10kb sliding window. Notice that the within-species TMRCA and the cross-coalescence time are closely related to the conventional summary statistics π_{Within} (a.k.a. π_W) and d_{XY} (a.k.a. $\pi_{Between}$, π_B, or π_{XY}), respectively, which have been previously used to distinguish among different contributing factors to F_{ST} peaks[34,36,46]. However, the ARG-based statistics are potentially more informative, by reflecting a comprehensive, model-based inference that considers correlated, fully resolved genealogies. Notably, these ARG-based statistics also provide a natural means for accommodating background selection and mutation rate variation (see **Methods** and **Discussion**).

We first compared the windowed F_{ST} and d_{XY} statistics within the F_{ST} peaks. If the F_{ST} peaks are driven by selection against gene flow, then these F_{ST} and d_{XY} estimates should be positively correlated, with larger values indicating deeper, and smaller values indicating shallower, cross-species coalescence (**Figure 1**)[34,36]. By contrast, if the F_{ST} peaks are driven by recent, species-specific sweeps, then no correlation is expected, because F_{ST} will be determined by the strength and ages of the sweeps, and d_{XY} will be largely consistent with the background. We found that the correlation between these statistics was quite low overall (Pearson's r = 0.197; **Supplementary Figure S2**), more consistent with a "recent sweeps" model than an "early selection against gene flow" model.

Next we turned to the *ARGweaver*-inferred genealogies and compared the within-species TMRCAs in the F_{ST} peaks and the background regions. If the F_{ST} peaks are driven by selective sweeps, we would expect them to exhibit a reduced within-species TMRCA relative to the background, reflecting recent increases in the frequency of selected alleles and corresponding "bursts" of coalescence. Indeed, the inferred genealogies at several F_{ST} peaks, such as the one on scaffold 404 near the *SLC45A2* gene (**Figure 2A**), display this qualitative behavior, with dense clusters of recent coalescence events suggesting selective sweeps (see additional examples in **Supplementary Figure S3**). Accordingly, a measure of the within-species TMRCA called TMRCAH (the TMRCA for the first 50% of all lineages from a given species to find a common ancestor; see **Methods**), is depressed at these F_{ST} peaks (**Figure 2B**). At the same time, the cross-coalescence time does not tend to be elevated at these peaks, as would be expected if they were driven by selection against gene flow during speciation (**Figure 2C**).

To see whether these trends held more generally, we devised statistical tests for (1) enrichment of particular species within clades of these trees, and (2) a corresponding reduction in the within-species TMRCA. The first test identifies instances of species differentiation, and the second identifies likely species-specific sweeps (see **Methods** and **Supplementary Text**). In 23 of the 25 (92%) F_{ST} peaks, we observed a significant enrichment for at least one species based on test 1, with 7 peaks (28%) showing significant enrichment for more than one species (**Table 1**). Overall, we detected 35 instances of species-enriched clades in these peaks, with *S. nigrorufa* showing the largest contribution (14 peaks) and *S. hypoxantha* showing the smallest (2 peaks). Of these 35 instances, 18 (64%) in 16 different peaks had significantly reduced within-species TMRCAs compared to the background, based on test 2. Together, these observations suggest a substantial contribution from recent selective sweeps to differentiation between southern capuchino species in these genomic islands.

We further examined the remaining 17 cases of species enrichment that did not exhibit reduced TMRCAs. These could represent cases in which the F_{ST} peaks are driven by early selection against gene flow rather than recent selective sweeps, although they could equally well reflect weaker or older sweeps that failed to meet our criteria for significance. We devised a test for elevated cross-coalescence times (test 3) between species, as expected in the case of selection against gene flow at the time of speciation within the F_{ST} peaks (see **Methods**). We found that most F_{ST} peaks had either no significant difference in cross-coalescence time with respect to flanking regions, or reduced—rather than elevated—cross-coalescence times (**Supplementary Table S4** and **Supplementary Text**). Nonetheless, four of the peaks did exhibit elevations of cross-coalescence times consistent with selection against gene flow (**Table 1**). The genomic signatures at these peaks are subtle and would likely be missed by summary statistics such as d_{XY}, but the genealogical evidence is compelling, particularly in the peaks on scaffolds 252 and 1635 (**Supplementary Figures S4**-**S6**). Notably, the region near the peak on scaffold 252, which is upstream of the gene encoding the Agouti-signaling protein (ASIP), contains multiple signatures of divergence. These include an apparent recent species-specific sweep in *S. melanogaster*, and a modestly recent sweep in *S. nigrorufa*, alongside deep separation for *S. palustris* and *S. pileata*, which indicates possible selection against gene flow in these species (**Supplementary Text**). Altogether, the observed patterns suggest possible selection against gene flow at a few loci of interest but its overall contribution to islands of differentiation appears to be limited compared to that of recent selective sweeps.

Prediction of Selective Sweeps by Machine Learning

The previous analysis suggests that selective sweeps have played a major role in driving the islands of differentiation observed in these recently emerged southern capuchino species. To characterize these sweeps in more detail, we developed machine-learning methods to predict individual sweeps of various kinds across the available genome sequences. We adopted a prediction strategy based on supervised learning, using a diverse collection of standard population genomic summary statistics as features and a linear Support Vector Machine (SVM) for classification (**Figure 3** and **Methods**). Similar approaches are becoming increasingly widely used in population genomics, and have been shown to be particularly effective in integrating multiple disparate signals to address challenging prediction problems[47,48,45,49]. As in most previous applications, we trained our classifiers from simulated data, which has the advantage of providing essentially limitless training data with error-free annotation of selection histories, but the potential disadvantage of being sensitive to the choice of simulation parameters (see **Discussion**). To

make our method as robust as possible to this choice, we used a forward simulation scheme based on the SLiM simulator[50], making use of a previously inferred demographic model[51] and allowing for ranges of values of key parameters, including selection coefficients, mutation and recombination rates, effective population sizes, times of selection onset, and positions of beneficial mutations (see **Methods** and **Supplementary Text**).

We were interested in several properties of the selective sweeps that contributed to the diversification of these species, including whether they tended to be "soft" or "hard" sweeps; to be recent or ancestral; or to be "complete" or "partial" sweeps[52,53]. To make these complex and intertwined questions more tractable, we sought first to address them individually, in sequence, and then later considered more complex cases combining multiple factors. Moreover, we focused first on the simple case of a pairwise analysis of present-day species, and simulations involving an ancestral population that diverged into two species, taking advantage of the fact that the five species under study are believed to have diverged from one another at roughly the same time, leading to a high degree of symmetry in the pairwise comparisons. In all cases, we simulated 48 haploid genomes (24 per species) consisting of 9000 regions of length 50 kb, using 8000 regions for training and the remaining 1000 for testing. Each genomic region was analyzed in five nonoverlapping 10 kb windows for signs of selection in the middle 10-kb window (see **Methods**).

Soft Sweeps have been Common in the Diversification of Southern Capuchinos

We focused first on the distinction between soft and hard sweeps, training a three-way classifier to distinguish species-specific soft and hard sweeps from each other and from neutral regions (see **Methods**). On simulated test sets, the classifier displayed an overall accuracy of ~93%, with most errors coming from the misclassification of soft sweeps as hard sweeps (~9% of soft sweeps) or of hard sweeps as soft sweeps (~6% of hard sweeps). (**Supplementary Table S5** and **Supplementary Figure S7**). We applied the method to the real data for *S. melanogaster* and *S. nigrorufa*, focusing at first on four scaffolds (252, 412, 404, and 1717) that contain top F_{ST} peaks and harbor known pigmentation-related genes (**Supplementary Table S6** and **Figure 4A**). The classifier predicted a total of 154 soft and 33 hard sweeps across these four scaffolds, including high-confidence (probability > 0.95) predictions of 28 soft and 3 hard sweeps. We observed similar patterns in three other pairwise analyses, selected based on their high levels of differentiation[42] (**Supplementary Figure S8-10** and **Supplementary Table S6**). In general, we identified large numbers of apparent species-specific sweeps, many of which coincided with F_{ST} peaks or otherwise occurred nearby genes involved in the regulation of melanogenesis, consistent with the hypothesis that selective sweeps have been a major force behind the

differentiation of these species. Furthermore, we consistently observed much larger numbers of predicted soft sweeps than hard sweeps. For example, in a broader analysis of sweeps in *S. melanogaster* (relative to *S. nigrorufa*) in the 19 scaffolds that contain F_{ST} peaks for this species, we observed 520 predicted soft sweeps and only 85 predicted hard sweeps, of which 86 and 8 were predicted with probability greater than 0.95, respectively (**Supplementary Table S7**).

To validate these predictions, we made use of three orthogonal analyses. First, we examined many individual predictions in detail, considering the local trees inferred by *ARGweaver* at both predicted hard and soft sweeps (**Figure 4B**). We found, in numerous cases, that the hard and soft sweeps had distinct genealogical features, with the hard sweeps displaying evidence of single derived haplotypes at high frequency, corresponding to unusually large and young clades, and the soft sweeps showing signatures of multiple potentially beneficial haplotypes all at elevated frequencies (**Supplementary Figure S11**). Second, we carried out a more systematic analysis based on two measures derived from the inferred ARGs: (1) the time to most recent common ancestry (TMRCA) for the species experiencing the sweep, and (2) a related statistic called the relative TMRCA half-life (RTH), which is expected to be reduced in partial sweeps[43] (**Figure 4C** and **Supplementary Figure S12–13**). We found that both hard and soft sweeps exhibited reductions of both statistics relative to flanking neutral regions. Interestingly, hard sweeps show a lower RTH than soft sweeps, likely because of their recent origin on a single haplotype background. Finally, to validate our ability to differentiate between soft and hard sweeps, we examined two haplotype homozygosity statistics, H_{12} and $\frac{H_2}{H_1}$, which are sensitive to differences in frequency between the two most prevalent haplotypes[54] (see **Methods**). We found, as expected, that the predicted soft sweeps tended to have larger values of $\frac{H_2}{H_1}$ and smaller values of H_{12}, while the hard sweeps showed the opposite pattern (**Figure 4C** and **Supplementary Figure S14**). Simulated hard and soft sweeps showed qualitatively similar behavior, although, unsurprisingly, they separated more cleanly than the empirical ones. Our results suggest that the empirical data is indeed dominated by the soft sweeps, but the classifier has perhaps erred on the side of miscalling some soft sweeps as hard sweeps (consistent with tests of our classifier on simulated data).

Many of the Soft Sweeps are also Recent and Partial

We expanded our simulation scheme to consider other distinguishing characteristics of selective sweeps. First, focusing on the case of soft sweeps, we asked whether these sweeps tended to occur recently, after a pair of species diverged, or whether they may instead be

ancestral, meaning that they began in the common ancestor of the two species. In simulation experiments, we found that we had some statistical power to distinguish these cases, with somewhat better classification accuracy for recent sweeps (~98%) than ancestral ones (~80%; see **Supplementary Table S8** and **Supplementary Figure S15**). In particular, the ancestral sweeps and neutral regions tended to be difficult to distinguish, probably owing to loss in older sweeps of the characteristic pattern of extended homozygosity[54]. We applied our classifier to the real data and did obtain predictions of 4–11 times as many recent, species-specific sweeps (per-species) as ancestral sweeps (see **Supplementary Table S9**), although this enrichment likely in part reflects differences in power.

Second, we attempted to ask whether species-specific soft sweeps tended to be "complete," meaning that the favored allele has been driven completely to fixation, or "partial," meaning that it has not fixed. We had limited power to address this question (see **Supplementary Table S10** and **Supplementary Figure S16**) but found some evidence suggesting that a substantial fraction of soft sweeps are partial sweeps (see **Supplementary Table S11**). This observation is consistent with the fact that the reduction in RTH observed in predicted soft sweeps is more pronounced than the reduction in TMRCA (**Figure 4C**).

An Expanded Analysis of all Five Species Further Supports Abundant Species-Specific Soft Sweeps

Having found that soft sweeps—often recent—have likely been common in these southern capuchino species, we extended our pairwise analyses to consider all five species simultaneously, allowing for a sweep to occur in any one of them. Our goal was to produce a comprehensive set of predictions encompassing all species. Based on simulated data, our multi-way classifier showed good accuracy in this setting (93–95% accuracy; **Supplementary Table S12** and **Supplementary Figure S17**). An analysis of the nineteen scaffolds containing F_{ST} peaks identified numerous species-specific soft sweeps (**Supplementary Figure S18** and **Supplementary Table S13**), consistent with our previous pairwise analyses. Furthermore, the inferred local trees associated with these predicted sweeps indicated reduced RTH statistics (**Figure 5** and **Supplementary Figures S19**). Together, these findings further support the prevalence of soft sweeps.

To gain insight into the functional relevance of the predicted sweeps, we examined the Gene Ontology (GO) biological process of nearby genes, focusing on predictions of species-specific soft sweeps with probability >0.95. We identified only two significantly enriched GO processes: "melanin biosynthetic process from tyrosine" (p = 0.023 after FDR correction) and "pigmentation"

($p = 0.022$ after FDR correction). Notably, the predicted sweeps were mostly located in non-coding regions near the genes in question, and likely contain cis-regulatory elements that control gene expression. Prominent examples include predictions near the genes *ASIP* (induces melanocytes to synthesize pheomelanin instead of eumelanin), *KITL* (stimulates melanocyte proliferation) (**Figure 5**), and *SLC45A2* (transports substances needed for melanin synthesis).

Discussion

In this article, we have presented an analysis of the genome sequences of 60 individuals representing five species of southern capuchino seedeaters. These birds serve as an excellent model for recent speciation in the presence of gene flow. Furthermore, because they differ primarily in male plumage and appear to be at least partially sympatric, they are an ideal system for studying the genetic effects of sexual selection in speciation[42,51]. Our analysis focused in large part on previously identified genomic islands of differentiation[42], defined by local peaks in F_{ST}. However, we substantially extended the examination of these regions, making use of state-of-the-art tools and techniques including ancestral recombination graph (ARG) inference and prediction of selective sweeps by machine learning.

We were particularly interested in comparing two competing models for the formation of genomic islands of differentiation: one in which selection acts during speciation to reduce gene flow in these genomic regions, and another in which selection acts to reduce within-species diversity through species-specific selective sweeps[35,36]. Notably, the "selective sweeps" model is a distillation of a larger family of models that allows for early sweeps, late sweeps, and adaptive introgression[34,36,46]. Nevertheless, we view these two general paradigms—early barriers to gene flow vs. recent sweeps—as representing a fundamental distinction between plausible models for the formation of islands, with the timing and specific nature of the sweeps being less essential.

At the same time, a binary choice between models is clearly an oversimplification—not only does it mask a diversity of scenarios within each paradigm, but it also obscures the possibility that both paradigms could be simultaneously at play. Importantly, the ARG-based statistics allow us to gain insight into the possible contributions of each model at each island of differentiation. For example, in the island upstream of the *ASIP* gene, we find evidence of a recent near-complete sweep in *S. melanogaster* alongside a deep separation between *S. pileata* and the other four species (**Supplementary Figures S4–S5** and **Supplementary Text**). Thus, both models likely contributed to differentiation in the regulatory sequence of this gene, but at different times and in different species. Notably, the distinction between the two paradigmatic models may not be absolute, since loci that experienced early barriers to gene flow could later undergo selective

sweeps, and loci that underwent species-specific sweeps could lead to reduced hybrid fitness resulting in barriers to gene flow. Nevertheless, when we contrast these two extreme scenarios, we find much stronger support overall for F_{ST} peaks being associated with recent sweeps than with older barriers to gene flow.

It is worth noting that, while we have focused on selective sweeps as an alternative to barriers to gene flow, previous models have allowed for a reduction in within-species diversity owing to background selection (BGS) as well as selective sweeps[34,36,46]. We focused on selective sweeps because they are far more plausible as an explanation for dramatic reductions in within-species diversity that have occurred recently and in a species-specific manner. At the same time, we took care to use indicators of selective sweeps that should be fairly robust to BGS, to avoid mistaking signatures of background selection for sweeps. In particular, our ARG-based analysis made use of measures based on the degree of "clustering" of coalescence events in local trees, which are designed to be sensitive to sweeps to the exclusion of BGS[43] (see **Methods**). Similarly, haplotype statistics such as H_{12} and $\frac{H_2}{H_1}$ should be elevated by sweeps but not by BGS. Thus, our conclusions about the prominence of sweeps should be relatively unaffected by BGS.

Consistent with other recent studies, we found machine-learning methods based on combinations of traditional population genetic summary statistics to be quite powerful in detecting selective sweeps. Notably, we went beyond previous work by characterizing sweeps not only as "soft" or "hard", but also as ancient or recent, and partial or complete. Overall, we found abundant evidence for sweeps in these capuchino species, and indications that these sweeps are enriched for soft sweeps and recent population-specific sweeps. These conclusions were further supported by semi-orthogonal evidence from ARG inference and haplotype statistics. At the same time, it was clear from our simulation experiments that some features of sweeps were much more difficult than others to discern from sequence data. For example, we had limited power to identify ancient sweeps and partial sweeps. We also attempted to distinguish between sweeps that were specific to a single species and sweeps that occurred in parallel at the same locus in multiple species, but found that our results were somewhat ambiguous, with true parallel sweeps being mixed together with cases of apparent shared sweeps or adaptive introgression (**Supplementary Table S14 and Supplementary Figures S20–S22**). In addition, our current machine-learning methods are still coarse-grained, in that they identify broad windows containing sweeps rather than specific causal variants, and they do not make direct use of informative ARG-derived features in classification. An important area for future work will be to develop improved high-resolution ARG-aware machine-learning predictors of sweeps[49], taking advantage of the latest advances in genealogy and ARG inference[44,55,56]. Finally, as in all studies that predict sweeps using machine-learning

classifiers trained from simulated data, our methods are sensitive to biases stemming from our choices of parameters for simulation[45,49]. We attempted to mitigate this issue by simulating from a broad family of models, and by validating our predictions with independent methods where possible. Nevertheless, training bias and limitations in both power and resolution have undoubtedly influenced our predictions.

Regardless of these caveats, the prediction of abundant soft sweeps—largely recent and population-specific—makes sense from first principles and previous findings. These capuchino species appear to have emerged from a large, highly polymorphic ancestral population, which would have provided high levels of standing variation to serve as the substrate for soft sweeps (**Supplementary Text**). Furthermore, these speciation events appear to have been quite recent, which might favor soft sweeps over hard sweeps, because they can occur more rapidly[57]. It has further been argued that standing genetic variants may be more likely than novel ones to drive speciation[58], because novel mutations tend to be deleterious if not neutral[59], whereas older variants have already been filtered and shaped by selection in their ecological context[60]. Notably, other recent empirical studies have similarly found a predominance of soft selective sweeps in both *Drosophila melanogaster*[61,62] and humans[63]. The recent speciation event would also allow limited time for haplotypes under selection to reach fixation, which would favor partial sweeps over complete sweeps. Finally, the likely role of sexual selection in driving many of these sweeps is consistent with selected alleles for features such as plumage having been present in the ancestral population, likely at low frequency, and having swept to high frequency relatively recently, in a species-specific manner. Sexual selection would also be consistent with our observations suggesting that many loci have experienced sweeps in parallel in different species.

Overall, the genetic changes underlying any speciation event are complex and unlikely to be fully explained by any simple model. They are undoubtedly highly context-dependent, depending on diverse features such as population size and structure, rates and patterns of gene flow, degrees of sympatry, differences in local environments, and the prominence of sexual selection. Nevertheless, our analysis of these capuchino species demonstrates that, at least under the right circumstances, prominent genomic islands of differentiation can be explained largely through recent, population-specific selective sweeps. Furthermore, we have shown that the analysis of these islands can benefit substantially by going beyond summary statistics such as F_{ST} and d_{XY} and making full use of the ancestral recombination graph (ARG), as well as machine-learning methods for prediction of selective sweeps. Together, our observations help to fill in the picture of how selection, recombination, and drift act together to shape the genomes of distinct species, with broad implications across the tree of life.

Methods

Whole genome data set. Our analysis made use of the reference genome previously assembled for S. *hypoxantha*[42]. The assembly covers 1.17 Gb of sequence partitioned into 5,120 scaffolds. The population-level sequencing data from that paper consisted of 72 individuals from nine capuchino seedeater species. We applied stringent filtering by masking sites with the following quality parameters: *QD<2, FS>40.0, MQ<20.0,* or *HaplotypeScore>*12.0. Variants were subsequently filtered if they satisfied one of the following conditions: (1) had more than two alleles, (2) had a minor allele frequency smaller than 0.1, (3) had mean sequencing depth smaller than 2, (4) had mean sequencing depth greater than 50, (5) had genotypes for less that 80% of the sequenced individuals (57 out of 72). This filtering pipeline produced 11,530,110 SNPs genotyped for 72 individuals of the nine southern capuchino species. Since four of the nine species—S. *ruficollis*, S. *hypochroma*, S. *cinnamomea*, and the outgroup S. *bouvreuil*—were represented by only three individuals, we focused our analysis on the five species for which there are 12 individuals: *S. hypoxantha, S. melanogaster*, *S. nigrorufa, S. palustris*, and *S. pileata*.

F_{ST} peaks. We considered the 25 F_{ST} peaks defined in ref. [42] (**Supplementary Table S1**). These regions were determined as stretches of two or more non-overlapping 25 kb windows with average F_{ST} above 0.2 in at least one of the ten pairwise comparisons and at least one variant site with F_{ST} above 0.85. Windows with less than ten variant sites were not considered (see ref. [42] for more details).

ARG blocks. To facilitate effective execution of *ARGweaver*, we filtered and partitioned the sequence scaffolds according to the following process. First, due to the low confidence in local genealogy inference near the edges of the aligned sequences, only scaffolds of length at least 120 kb were considered. This filtered out roughly 110 Mb of sequence, resulting in 1.06 Gb of sequence analyzed by *ARGweaver*. Among the remaining 595 scaffolds, those shorter than 2 Mb were analyzed as is, and longer scaffolds were split for computational efficiency into blocks of similar length with an overlap of 100 kb. This resulted in 1,003 genomic blocks (ARG blocks) which were analyzed via separate parallel runs of *ARGweaver*.

Executing *ARGweaver*. We applied *ARGweaver* on probabilistic genotypes specified in the previously generated sequence VCF file[42]. Genotypes based on sequence coverage less than 5 or greater than 30 were masked using an individual mask file, as well as sites with no genotype that had at least 80% certainty (`--mask-uncertain 0.8`). We used a site compression of 5

when possible (`-c 5`), and reduced compression in ARG blocks where the frequency of variant sites was too high. To make the analysis more robust to mutation rate variation, we supplied *ARGweaver* with local mutation rates estimated using two individuals sequenced by Campagna *et al.* (2017) from two species not analyzed in this study: L20.3 from *S. hypochroma* and L15.3 from *S. bouvreuil* (individuals from these two species were not used in our subsequent analyses). The absolute divergence between these two individuals was computed in 100 kb windows with 10 kb overlaps using probabilistic genotypes given in the sequence VCF file. Windows with fewer than 20,000 unfiltered bases were given the genome-wide average rate. The mutation rate at each genomic position was set by scaling the assumed average mutation rate of 10^{-9}/bp/gen by the absolute divergence in the 100 kb window centered at that position divided by the average divergence across all windows. In addition, we used the following options for setting the effective population size, recombination rate, time discretization, random genotype phasing, and MCMC sampling: `--popsize 1000000 -r 1e-9 --ntimes 20 --maxtime 1e7 --delta 0.005 --randomize-phase 1.0 --unphased --sample-phase 10 --resample-window-iters 1 --resample-window 10000 -n 1000`. *ARGweaver* was executed on each ARG block for 1000 MCMC iterations, and the ARG sampled in the last iteration was used to extract local trees for all subsequent analyses. In one of the ARG blocks that does not overlap an F_{ST} peak (the fourth ARG block for scaffold 329), *ARGweaver* was halted after 700 iterations due to time constraints. While *ARGweaver* was executed on the entire data set, consisting of 72 diploid individuals, all of our downstream analyses made use of the 60 individuals from the five focal species of this study (**Supplementary Text**).

A simple genealogical indicator of selective sweeps. The main genealogical signature associated with a selective sweep is a "burst" of coalescence events, which results in a large clade with recent origin. Motivated by this fundamental observation, we developed a simple ARG-based test for the association of a given F_{ST} peak with a selective sweep. First, we examined the local tree inferred by *ARGweaver* for the site with the largest F_{ST} value within an F_{ST} peak. In this tree, we chose a specific clade that was most likely associated with a recent species-specific sweep by looking for a clade that was: (1) sufficiently large, (2) relatively young, (3) enriched for haploid samples from one of the species for which the highest F_{ST} value was obtained in that site. The age and size of this target clade were recorded and compared to a background distribution computed using clades of identical size observed in a set of 10,000 local genealogies sampled uniformly at random along the same scaffold outside of the F_{ST} peak. If a sampled tree did not contain a clade of the target size, it was discarded, and if it contained multiple such clades, the

age of the youngest one was recorded. A *p*-value was computed for the age of the target clade according to its rank (order) in the background distribution. A low *p*-value implies that this clade is significantly younger than expected for clades of the same size, and thus suggests a recent selective sweep in the species enriched in that clade. This test is applied in the results shown in **Figure 2** and **Supplementary Figures S3** and **S4**.

ARG-based measures associated with (species-specific) selective sweeps. To apply the approach described above to cases where the target clade or its size are unknown, we used measures based on the time to the most recent common ancestor (TMRCA) within a given species. For a given local tree and a species of interest, we define the TMRCAH of that species as the age of the youngest clade that contains at least half of the haploid samples from that species (*n*=12). Requiring only half the samples of the species allows us to consider partial sweeps and provides robustness to the inherent uncertainty in the inferred local trees. Values of TMRCAH are plotted in **Figure 2** and **Supplementary Figure S4**. To account for variation in age estimates stemming from mutation rate variation and/or background selection, we used relative versions of TMRCAH, or relative time to coalescence of half the lineages (RTH). In different parts of our analysis we used slightly different normalization approaches for RTH that aimed to capture slightly different features of selective sweeps (see also **Validation of inferred sweeps using ARG-based measures**, below). In the comprehensive analysis of ARG-based signatures in the F_{ST} peaks (test 2 in **Table 1** and **Supplementary Table S3**), we used a measure termed RTH', obtained by dividing the TMRCAH of a given species by the age of the subtree that contained at least half the total number of samples (*n*=60). This variant is designed to capture sweeps even if they were near complete in one species or shared with other species, as long as they did not affect more than half of our sampled genomes. Furthermore, it was found to effectively reduce the variation in ages caused by mutation rate variation and background selection. To detect partial sweeps that were not necessarily associated with a certain species, we computed for every tree a relative time for the first coalescence of 12 lineages (RT12) from possibly different species. This relative age was obtained by taking the age of the youngest subtree with at least *n*=12 leaves, and dividing it by the age of the subtree that contained at least half the total number of samples.

Species enrichment score. A species enrichment score was computed to indicate separation of a given species from all others in a given local tree (test 1 in **Table 1**). For a subtree (clade) with *n* leaves out of which *k* are mapped to a certain species, we computed the probability of this observation under a hypergeometric distribution, and defined the associated enrichment score as

$-\log_{10}(P_{K=24,N=120}(X{\geq}k|n))$. The enrichment score associated with a certain species in a given local tree is then defined as the maximum enrichment score for that species over all subtrees, to reflect the fact that we are looking for the existence of a clade separating that species from the others.

Inter-species cross coalescence times. To detect possible selection against gene flow, we looked for a delay in times of inter-species cross coalescence within F_{ST} peaks relative to times of cross coalescence observed in nearby genomic regions. For a simple measure (used in **Figure 2** and **Supplementary Figure S4**), we use the average of all cross-coalescence events in a given local tree. A certain coalescence event may be counted multiple times toward this distribution if it is the most recent common ancestor of multiple pairs of samples from the two species. Thus, this measure could be viewed as the ARG-based version of d_{XY}. In the detailed analysis of F_{ST} peaks (test 3 in **Table 1** and **Supplementary Table S4**), we used a slightly different version that considered only the ten youngest cross-coalescence events, which would be the most affected by changes in the amount of gene flow in the recent past. As with the other age statistics (RTH' and RT12), we account for the effects of mutation rate variation and background selection by dividing these ten ages with the age of the youngest subtree that contains at least half of the total number of haploid samples.

Genomic control for enrichment scores, RTH', and RT12. The detailed analysis of F_{ST} peaks (**Table 1** and **Supplementary Tables S2** and **S3**) relies on detection of significant deviations of genealogical signatures within the peaks, as compared to the genomic background. First, to account for the uncertainty stemming from the fact that our genealogical measures are based on a single sampled ARG, we averaged values of enrichment scores, RTH', and RT12 across non-overlapping 20 kb windows. The windows were determined by tiling each scaffold, and masking out the 50 kb near each edge (due to inference uncertainty in the edges of every ARG block). In every 20 kb window, we computed an average value for each measure by using 40 representative local trees from that window, equally spaced 500 bp apart starting 250 bp from the edge of the window. Second, for every peak and every species, we computed the maximum (windowed) species enrichment scores and the minimum (windowed) RTH' for that species, taking into consideration all 20 kb windows that overlap the peak (**Supplementary Tables S2 and S3**). Similarly, a minimal (windowed) RT12 was recorded for every F_{ST} peak. To assess significance thresholds for these eleven statistics, we considered their distributions in 39,699 windows that belong to one of the 576 analyzed sequence scaffolds that do not contain F_{ST} peaks. For species enrichment scores and RTH', we set a significance threshold with empirical p-value of 0.0001, by

requiring that only three of these control segments exceed the threshold (tests 1 and 2 in Table 1). For RT12, we set a slightly more relaxed threshold with empirical *p*-value of 0.001, by requiring that only 39 of these segments exceed the threshold. To account for the use of extreme values (maximum or minimum) in long genomic blocks, we examined the distribution of values in a collection of 1,376 non-overlapping 500 kb blocks from the 576 scaffolds that do not contain F_{ST} peaks. The block length was selected to be conservative relative to the length distribution of F_{ST} peaks (mean length 243 kb and only four peaks longer than 500 kb; **Supplementary Table S1**).

Genomic control for cross coalescence times. We applied a slightly different control scheme for cross-coalescence times to enable detection of subtle changes in their distribution. First, rather than computing average values across 20 kb windows, we computed for each species-pair the distribution of normalized cross coalescence times in every F_{ST} peak. For peaks longer than 200 kb, we focused on the 200 kb region that contains the highest species enrichment scores. To account for possible fluctuations in coalescence times between sequence scaffolds, we used a local control set comprised of the 200,000 sites flanking the F_{ST} peak. When possible, we selected 100 kb from each side of the peak, unless the peak was close to the edge of a sequence scaffold. The peak on scaffold 308 was discarded from this analysis, because there are less than 200 kb flanking the peak on the sequence scaffold. The difference between the distribution of normalized cross coalescence times in an F_{ST} peak and that recorded in the flanking region is quantified using percentile differences. The percentile difference between two distributions is computed by taking the median value of one distribution, examining the percentile of this value in the second distribution, and taking the difference in absolute value from ½. This is done in both directions (F_{ST} peak versus flanking region and vice versa), and the difference associated with every peak is taken to be the maximum of the two. See **Supplementary Table S4** for the quantile differences computed for all F_{ST} peaks and all species pairs.

Contribution of recent selective sweeps and selection against gene flow. Evidence for a recent species-specific selective sweep is given by a high enrichment score and a low RTH' for a given species. F_{ST} peaks associated with such a signal are indicated by gray cells in Table 1 marked with a red circle (tests 1&2). Evidence for selection against gene flow is more subtle. A delay in inter-species cross coalescence times could be a result of selection against gene flow, but it could also be caused by other factors, such as random fluctuations and even selective sweeps (see **Supplementary Text**). We thus used the following combination of all three types of statistics to suggest cases of selection against gene flow in our data set:
 (1) The species has a significantly high enrichment score in the peak.

(2) The species does not have a significantly low RTH' in the peak.

(3) The species has elevated cross-coalescence times with some other species (quantile difference > 0.1).

These conditions imply a deep clade enriched for a given species, suggesting that selection against gene flow acted to delay cross-coalescence times. Thus, if all three conditions hold, we mark the cell in Table 1 associated with the given F_{ST} peak and species with a blue circle (test 3).

Summary statistics used as features for prediction of different models for selective sweeps. A series of sequence-based species-level summary statistics were used as possible covariates for a machine learning approach to distinguish between different models of selection (including neutral drift). All statistics were collected along five consecutive 10 kb windows with the objective of informing on possible sweeps induced by a positively selected mutation in the third (middle) window. Some of the summary statistics corresponded to standard measures of diversity, such as the number of segregating sites, π (ref. [64]), Tajima's D (ref. [65]), θ_W (ref. [66]), θ_H (ref. [64]), the number of distinct haplotypes (ref. [62]), Z_{nS} (ref. [67]), and maximum value of ω (ref. [68]). For each of these statistics, we computed an average value for each of the five 10 kb windows for each analyzed species separately and jointly for all species together. We also extracted statistics based on the distribution of linkage disequilibrium (LD) of each of the five windows with the middle window—where the site under selection is assumed to be located. LD was computed for every variant site in the middle window with all other variant sites in all five windows. Then, for each of the five windows, we recorded the distribution of LD values relative to the middle window, and from this distribution we extracted three statistics based on the three values for percentiles 75, 90, and 95. Each LD-based statistic was recorded for each analyzed species separately. Finally, each summary statistic was normalized by dividing the value recorded for a given window by the sum of values across all five windows. As a result, the average value of each statistic across the five windows was 0.2 (see **Figure 3**).

Simulated datasets used for training and testing the selective sweep classifier. Training and testing data sets were generated using SLiM[50] by simulating 9,000 regions of length 50 kb for each model we considered: (e.g., "neutral", "soft sweep", "hard sweep", etc.). Of these regions, 8,000 were used for training and 1,000 were used for testing (see below). The number of simulated species and sampled sequences was set to match the size of the analyzed data set. Thus, in the species-pair analyses, a total of 48 haploid sequences were sampled at the end of the simulation (24 per species), and in the expanded analysis the entire set of five species was

simulated and 120 haploid sequences were sampled. Simulations used a demographic model based on one inferred previously from ddRAD data[51]. In the expanded analysis we used the complete model, and in the species-pair analyses we used the appropriate two-species derivatives of this model (**Supplementary Text**). In non-neutral simulations, selection was applied to a single focal site located in the middle 10 kb window. We explored a range of values for the main parameters corresponding to demography and selection: (1) mutation rate, (2) recombination rate, (3) effective population sizes, (4) selection coefficient, (5) onset time of selection, and (6) position of beneficial mutation. Furthermore, for soft sweeps (partial or complete), we also varied the initial derived allele frequency at which a mutation switches from evolving under drift to becoming beneficial, and for partial sweeps we varied the target derived allele frequency at which a mutation switches back to being neutral (**Supplementary Text**). For each simulated region, we recorded the set of covariate statistics (see above). If one of these statistics fell outside of the range of values observed in the genomic data, the simulated 50 kb segment was excluded from the set and replaced by another segment (to maintain a total of 9,000 simulated segments per class). The comparison to genomic data was done based on the 19 scaffolds that contain F_{ST} peaks and the specific species being analyzed by the classifier. Thus, slightly different filtering was applied for each of the four species-pair analyses in each classification task (see below).

Training a linear SVM to classify different modes of selection. A linear support vector machine (SVM) was applied to the simulated training data sets to learn a classification model for each task separately. SVM is a supervised learning method widely used in classification and regression analysis. Classification into two categories (e.g., "neutral" vs. "non-neutral") is achieved by finding the hyperplane that maximizes the separation of the data from the two different classes, using either a linear or a non-linear decision boundary (**Figure 3**). Since most of our prediction tasks had more than two categories (e.g. "neutral", "soft sweep", and "hard sweep"), we used a one-vs-rest approach. Thus, a separate classifier was trained for each class, with the 8,000 regions simulated for that class as positive training examples, and regions simulated under other models as negative training examples. We used the C-Support Vector Classification tool (*sklearn.svm.SVC*) from the Python *sklearn* package to train and test the classifier. Default operating parameters were used in all cases. Classification into more than two categories was obtained by combining the results from multiple binary classifiers using the *predict_proba* function in *sklearn*. This function applies a logistic function to the individual score produced by each classifier, thus transforming it into a value reflecting the probability that the

binary classifier assigns to its "positive" class. It then divides the probability obtained (separately) for each class by the sum of all probabilities, to obtain a normalized prediction probability for each class. The resulting multi-way classifier was then tested using 1,000 regions simulated for each class. The results of these tests were summarized in a confusion matrix describing the frequency of different types of prediction errors when the predicted class is set as the one with the highest probability. We also computed a receiver operating characteristic (ROC) curve for every pair of classes, to provide a more complete summary of the behavior of different types of errors.

Species-pair analyses and expanded analysis of all five species. We started by examining different types of selective sweeps using a series of classification tasks, each focused on two types of sweeps and two species. The pairs of sweep categories considered were (1) soft vs. hard sweeps, (2) recent vs. ancestral sweeps, (3) partial vs. complete sweeps, and (4) species-specific vs. parallel sweeps (see **Supplementary Text** for more details). In each case, we aimed to distinguish the two types of sweeps from each other and from neutrally evolving regions. Each of these four classification tasks was applied to four species-pairs that overall exhibited the highest F_{ST} values genome-wide: (1) S. *melanogaster* vs. S. *nigrorufa*, (2) S. *nigrorufa* vs. S. *pileata*, (3) S. *pileata* vs. S. *palustris*, and (4) S. *hypoxantha* vs. S. *melanogaster*. Note that for each species-pair, simulated segments were filtered using slightly different criteria (see above). Based on the outcome of the four classification tasks in all four species pairs, we designed an expanded analysis of all five capuchino species. In particular, in this analysis we allowed any one of the five species to undergo a species-specific complete soft sweep.

Application of machine learning to genomic data. We analyzed the 19 scaffolds that contain F_{ST} peaks using each of our trained classifiers. Each scaffold was scanned by a sliding 50 kb window with a step size of 10 kb. Summary statistics were extracted, as described above, for each 50 kb window using the genome samples appropriate for the specific classification task at hand—24 genomes in each species-pair analyses and all 60 genomes in the expanded analysis. These summary statistics were then provided as input to the trained binary classifiers and their classification outputs were combined to provide a normalized probability for each class (see above). The middle 10 kb of the 50 kb window was then assigned the class with the highest score, and Manhattan plots were used to show the distribution of class assignments and their normalized probabilities across each scaffold (e.g., Figure 4A).

High-confidence prediction of sweeps. For high-confidence predictions, we considered classifications associated with a normalized probability 0.95 or higher, and we further confirmed

their reliability through *p*-values empirically computed on a set of 10,000 regions simulated under neutral evolution. Each trained classifier was applied to these regions and the distribution of prediction probabilities was recorded for each of the non-neutral classes, as a null distribution. Given a genomic window predicted to be in a non-neutral class, the *p*-value associated with this prediction was taken to be the order (rank) of the probability assigned by the classifier to this prediction in the null distribution described above. Finally, the set of *p*-values for each non-neutral class of each classifier were corrected using the false discovery rate (FDR), and we confirmed that calls with prediction probability above 0.95 were associated with an FDR-corrected *p*-value below 0.05.

Validation of inferred sweeps using ARG-based measures. The inferred soft and hard sweeps were validated using summary statistics extracted from the ARG inferred by *ARGweaver*. In particular, we explored two measures relating to the within-species TMRCA, which are all expected to be reduced in species-specific selective sweeps (see also **ARG-based measures associated with sweeps** above): (1) TMRCA - the age of the youngest clade that contains all 24 haploid samples from a given species and (2) RTH - the age of the youngest clade that contains at least half of the haploid samples from a given species (TMRCAH), divided by the TMRCA of that species. The distribution of these values was recorded for regions classified by the linear SVM into each of the three selection categories—soft sweeps, hard sweeps, and neutral—to examine observed differences. As additional validation, we used representative local trees extracted from the ARG in regions predicted to undergo sweeps by either the species-pair analyses or the expanded five species analysis. Within a given 10 kb window classified as a sweep in a given species, we extract as a representative tree the local tree with the lowest RTH.

Validation of soft versus hard sweeps using homozygosity statistics. To differentiate soft from hard sweeps, we used the haplotype homozygosity statistics, H_1, H_2, and H_{12} (ref. [61]). The haplotype homozygosity, H_1, is defined as the probability of observing a completely homozygous haplotype (50 kb long), which is computed by taking the sum of squares of all haplotype frequencies. Similarly, H_2 is the haplotype homozygosity computed by ignoring the most frequent haplotype, and H_{12} is the haplotype homozygosity computed by combining the two most frequent haplotypes. In hard sweeps, one haplotype rises to much higher frequency than all others, so H_1 and H_{12} are both expected to be high, but the ratio $\frac{H_2}{H_1}$ is expected to be low. However, in soft sweeps, the ratio $\frac{H_2}{H_1}$ is expected to be much higher, and H_{12} is typically lower (depending on how

"soft" the sweep is). Thus, the two statistics $\frac{H_2}{H_1}$ and H_{12} provide effective separation between the two types of sweeps, as demonstrated in ref. [61].

Acknowledgments

The authors would like to acknowledge Noah Dukler for help with figure preparation. This research was supported by US National Science Foundation grants (NSFDEB) 1555769, 1555754, US-Israel Binational Science Foundation grant BSF2015523, and US National Institutes of Health grant R35-GM127070. The content is solely the responsibility of the authors and does not necessarily represent the official views of the US National Institutes of Health or the US National Science Foundation. The bird illustrations are original drawings done by Jillian Ditner.

Competing Interests

The authors declare no competing interests.

Contributions

Designed the research: H.A.H., L.C., I.G., A.S.. Performed machine learning modeling and simulations: H.A.H.. Performed the ARG analysis: H.A.H., A.S-M., M.J.H., I.G.. Analyzed the data: H.A.H., A.S-M., I.G.. Wrote the paper: H.A.H., I.G., A.S.. Edited and approved paper: H.A.H., A.S-M., L.C., I.J.L., I.G., A.S.

References

1. Wolf, J. B. W. & Ellegren, H. Making sense of genomic islands of differentiation in light of speciation. *Nat. Rev. Genet.* **18**, 87–100 (2017).
2. Darwin, C. & Wallace, A. On the tendency of species to form varieties; and on the perpetuation of varieties and species by natural means of selection. *J. Proc. Linn. Soc. Lond. Zool.* **3**, 45–62 (1858).
3. Coyne, J. & Orr, H. *Studying speciation*. (Sinauer Associates, Inc Sunderland, MA, 2004).
4. Dobzhansky, T. G. *Genetics and the Origin of Species*. (Columbia Univ. Press, 1937).
5. Mayr, E. & Provine, W. B. *The Evolutionary Synthesis: Perspectives on the Unification of Biology*. (Harvard Univ. Press, 1998).
6. Mayr, E. *Systematics and the Origin of Species*. (Columbia Univ. Press, 1942).
7. Wang, X., He, Z., Shi, S. & Wu, C.-I. Genes and speciation – Is it time to abandon the Biological Species Concept? *Natl. Sci. Rev.* nwz220 (2019) doi:10.1093/nsr/nwz220.
8. Savolainen, V. *et al.* Sympatric speciation in palms on an oceanic island. *Nature* **441**, 210–213 (2006).
9. Schliewen, U. K., Tautz, D. & Pääbo, S. Sympatric speciation suggested by monophyly of crater lake cichlids. *Nature* **368**, 629–632 (1994).
10. Filchak, K. E., Roethele, J. B. & Feder, J. L. Natural selection and sympatric divergence in the apple maggot Rhagoletis pomonella. *Nature* **407**, 739–742 (2000).
11. Sorenson, M. D., Sefc, K. M. & Payne, R. B. Speciation by host switch in brood parasitic indigobirds. *Nature* **424**, 928–931 (2003).
12. Barluenga, M., Stölting, K. N., Salzburger, W., Muschick, M. & Meyer, A. Sympatric speciation in Nicaraguan crater lake cichlid fish. *Nature* **439**, 719–723 (2006).
13. Poelstra, J. W., Richards, E. J. & Martin, C. H. Speciation in sympatry with ongoing secondary gene flow and a potential olfactory trigger in a radiation of Cameroon cichlids. *Mol. Ecol.* **27**, 4270–4288 (2018).
14. Schliewen, U. K., Kocher, T. D., McKaye, K. R., Seehausen, O. & Tautz, D. Evidence for sympatric speciation? *Nature* **444**, E12–E13 (2006).
15. Pagan, I. & Holmes, E. C. Long-Term Evolution of the Luteoviridae: Time Scale and Mode of Virus Speciation. *J. Virol.* **84**, 6177–6187 (2010).
16. Meyer, J. R. *et al.* Ecological speciation of bacteriophage lambda in allopatry and sympatry. *Science* **354**, 1301–1304 (2016).
17. Nadeau, N. J. *et al.* Genomic islands of divergence in hybridizing *Heliconius* butterflies identified by large-scale targeted sequencing. *Philos. Trans. R. Soc. B Biol. Sci.* **367**, 343–353 (2012).
18. Martin, S. H. *et al.* Genome-wide evidence for speciation with gene flow in *Heliconius* butterflies. *Genome Res.* **23**, 1817–1828 (2013).
19. Turner, T. L., Hahn, M. W. & Nuzhdin, S. V. Genomic Islands of Speciation in Anopheles gambiae. *PLoS Biol.* **3**, e285 (2005).
20. White, B. J., Cheng, C., Simard, F., Costantini, C. & Besansky, N. J. Genetic association of physically unlinked islands of genomic divergence in incipient species of *Anopheles gambiae*. *Mol. Ecol.* **19**, 925–939 (2010).
21. Via, S. & West, J. The genetic mosaic suggests a new role for hitchhiking in ecological speciation. *Mol. Ecol.* **17**, 4334–4345 (2008).

22. Soria-Carrasco, V. *et al.* Stick Insect Genomes Reveal Natural Selection's Role in Parallel Speciation. *Science* **344**, 738–742 (2014).

23. Renaut, S. *et al.* Genomic islands of divergence are not affected by geography of speciation in sunflowers. *Nat. Commun.* **4**, 1827 (2013).

24. Brandvain, Y., Kenney, A. M., Flagel, L., Coop, G. & Sweigart, A. L. Speciation and Introgression between Mimulus nasutus and Mimulus guttatus. *PLoS Genet.* **10**, e1004410 (2014).

25. Harr, B. Genomic islands of differentiation between house mouse subspecies. *Genome Res.* **16**, 730–737 (2006).

26. Geraldes, A., Basset, P., Smith, K. L. & Nachman, M. W. Higher differentiation among subspecies of the house mouse (Mus musculus) in genomic regions with low recombination. *Mol. Ecol.* **20**, 4722–4736 (2011).

27. Marques, D. A. *et al.* Genomics of Rapid Incipient Speciation in Sympatric Threespine Stickleback. *PLOS Genet.* **12**, e1005887 (2016).

28. Malinsky, M. *et al.* Genomic islands of speciation separate cichlid ecomorphs in an East African crater lake. *Science* **350**, 1493–1498 (2015).

29. Poelstra, J. W. *et al.* The genomic landscape underlying phenotypic integrity in the face of gene flow in crows. *Science* **344**, 1410–1414 (2014).

30. Vijay, N. *et al.* Evolution of heterogeneous genome differentiation across multiple contact zones in a crow species complex. *Nat. Commun.* **7**, 13195 (2016).

31. Knief, U. *et al.* Epistatic mutations under divergent selection govern phenotypic variation in the crow hybrid zone. *Nat. Ecol. Evol.* **3**, 570–576 (2019).

32. Uy, J. A. C. *et al.* Mutations in different pigmentation genes are associated with parallel melanism in island flycatchers. *Proc. R. Soc. B Biol. Sci.* **283**, 20160731 (2016).

33. Toews, D. P. L. *et al.* Plumage Genes and Little Else Distinguish the Genomes of Hybridizing Warblers. *Curr. Biol.* **26**, 2313–2318 (2016).

34. Irwin, D. E. *et al.* A comparison of genomic islands of differentiation across three young avian species pairs. *Mol. Ecol.* **27**, 4839–4855 (2018).

35. Wu, C.-I. The genic view of the process of speciation. *J. Evol. Biol.* **14**, 851–865 (2001).

36. Cruickshank, T. E. & Hahn, M. W. Reanalysis suggests that genomic islands of speciation are due to reduced diversity, not reduced gene flow. *Mol. Ecol.* **23**, 3133–3157 (2014).

37. Noor, M. A. F. & Bennett, S. M. Islands of speciation or mirages in the desert? Examining the role of restricted recombination in maintaining species. *Heredity* **103**, 439–444 (2009).

38. Pennisi, E. Disputed islands. *Science* **345**, 611–613 (2014).

39. Campagna, L. *et al.* Rapid phenotypic evolution during incipient speciation in a continental avian radiation. *Proc. R. Soc. B Biol. Sci.* **279**, 1847–1856 (2012).

40. Machado, É. & Silveira, L. F. Plumage variability and taxonomy of the Capped Seedeater Sporophila bouvreuil (Aves: Passeriformes: Emberizidae). *Zootaxa* **2781**, 49–62 (2011).

41. Di Giacomo, A. S. *et al.* Landscape associations of globally threatened grassland birds in the Aguapey river Important Bird Area, Corrientes, Argentina. *Bird Conserv. Int.* **20**, 62–73 (2010).

42. Campagna, L. *et al.* Repeated divergent selection on pigmentation genes in a rapid finch radiation. *Sci. Adv.* **3**, e1602404 (2017).

43. Rasmussen, M. D., Hubisz, M. J., Gronau, I. & Siepel, A. Genome-Wide Inference of

Ancestral Recombination Graphs. *PLoS Genet.* **10**, e1004342 (2014).

44. Hubisz, M. J., Williams, A. L. & Siepel, A. Mapping gene flow between ancient hominins through demography-aware inference of the ancestral recombination graph. *bioRxiv* (2019) doi:10.1101/687368.

45. Schrider, D. R. & Kern, A. D. Supervised Machine Learning for Population Genetics: A New Paradigm. *Trends Genet.* **34**, 301–312 (2018).

46. Irwin, D. E., Alcaide, M., Delmore, K. E., Irwin, J. H. & Owens, G. L. Recurrent selection explains parallel evolution of genomic regions of high relative but low absolute differentiation in a ring species. *Mol. Ecol.* **25**, 4488–4507 (2016).

47. Sheehan, S. & Song, Y. S. Deep Learning for Population Genetic Inference. *PLoS Comput. Biol.* **12**, e1004845 (2016).

48. Kern, A. D. & Schrider, D. R. diploS/HIC: An Updated Approach to Classifying Selective Sweeps. *G3 Genes Genomes Genet.* **8**, 1959–1970 (2018).

49. Hejase, H. A., Dukler, N. & Siepel, A. From Summary Statistics to Gene Trees: Methods for Inferring Positive Selection. *Trends Genet.* S0168952519302690 (2020) doi:10.1016/j.tig.2019.12.008.

50. Haller, B. C. & Messer, P. W. SLiM 2: Flexible, Interactive Forward Genetic Simulations. *Mol. Biol. Evol.* **34**, 230–240 (2017).

51. Campagna, L., Gronau, I., Silveira, L. F., Siepel, A. & Lovette, I. J. Distinguishing noise from signal in patterns of genomic divergence in a highly polymorphic avian radiation. *Mol. Ecol.* **24**, 4238–4251 (2015).

52. Nielsen, R., Hellmann, I., Hubisz, M., Bustamante, C. & Clark, A. G. Recent and ongoing selection in the human genome. *Nat. Rev. Genet.* **8**, 857–868 (2007).

53. Pritchard, J. K., Pickrell, J. K. & Coop, G. The Genetics of Human Adaptation: Hard Sweeps, Soft Sweeps, and Polygenic Adaptation. *Curr. Biol.* **20**, R208–R215 (2010).

54. Przeworski, M. The signature of positive selection at randomly chosen loci. *Genetics* **160**, 1179–1189 (2002).

55. Speidel, L., Forest, M., Shi, S. & Myers, S. R. A method for genome-wide genealogy estimation for thousands of samples. *Nat. Genet.* **51**, 1321–1329 (2019).

56. Kelleher, J. *et al.* Inferring whole-genome histories in large population datasets. *Nat. Genet.* **51**, 1330–1338 (2019).

57. Barrett, R. D. H. & Schluter, D. Adaptation from standing genetic variation. *Trends Ecol. Evol.* **23**, 38–44 (2008).

58. Marques, D. A., Meier, J. I. & Seehausen, O. A Combinatorial View on Speciation and Adaptive Radiation. *Trends Ecol. Evol.* **34**, 531–544 (2019).

59. Ohta, T. The Nearly Neutral Theory of Molecular Evolution. *Annu. Rev. Ecol. Syst.* **23**, 263–286 (1992).

60. Abbott, R. *et al.* Hybridization and speciation. *J. Evol. Biol.* **26**, 229–246 (2013).

61. Garud, N. R., Messer, P. W., Buzbas, E. O. & Petrov, D. A. Recent Selective Sweeps in North American Drosophila melanogaster Show Signatures of Soft Sweeps. *PLoS Genet.* **11**, e1005004 (2015).

62. Messer, P. W. & Petrov, D. A. Population genomics of rapid adaptation by soft selective sweeps. *Trends Ecol. Evol.* **28**, 659–669 (2013).

63. Schrider, D. R. & Kern, A. D. Soft Sweeps Are the Dominant Mode of Adaptation in the

Human Genome. *Mol. Biol. Evol.* **34**, 1863–1877 (2017).

64. Fay, J. C. & Wu, C.-I. Hitchhiking Under Positive Darwinian Selection. *Genetics* **155**, 1405–1413 (2000).
65. Tajima, F. Statistical Methodfor Testing the Neutral Mutation Hypothesis by DNA Polymorphism. *Genetics* **123**, 585–595 (1989).
66. Watterson, G. A. On the number of segregating sites in genetical models without recombination. *Theor. Popul. Biol.* **7**, 256–276 (1975).
67. Kelly, J. K. A Test of Neutrality Based on Interlocus Associations. *Genetics* **146**, 1197–1206 (1997).
68. Kim, Y. & Nielsen, R. Linkage Disequilibrium as a Signature of Selective Sweeps. *Genetics* **167**, 1513–1524 (2004).
69. Nei, M. & Li, W. H. Mathematical model for studying genetic variation in terms of restriction endonucleases. *Proc. Natl. Acad. Sci. U. S. A.* **76**, 5269–5273 (1979).

Figure 1: Two models for the formation of islands of differentiation (F_{ST} peaks): selection against gene flow during speciation vs. recent species-specific selective sweeps. (**A**) Representative genealogies for six individuals from two recently diverged species in an F_{ST} peak (*yellow*) and in a nearby neutrally evolving genomic region (*gray*). Under both models, neutral genealogies exhibit frequent incomplete lineage sorting (ILS) due to large ancestral population sizes and possible ongoing gene flow between species (*vertical dashed gray line*). ILS is reduced in the F_{ST} peaks either because the time back to the first cross-coalescence (marked by *X*) is elevated by selection against gene flow (*left*), or because the within-species time to most recent common ancestry (TMRCA; marked by *triangle*) is reduced by a selective sweep in one or both species (*right*). (**B**) Hypothetical distributions of various sequence-based and genealogy-based measures under the two models. Selection against gene flow (*blue*) will tend to induce a positive correlation between F_{ST} and d_{XY}, whereas species-specific sweeps (*red*) will not (*left*). The within-species TMRCA is expected to be reduced by selective sweeps, but not substantially by selection against gene flow (*middle*). The cross-coalescence time is expected to be elevated by selection against gene flow but not substantially by species-specific sweeps (*right*).

Figure 2: Comparison of inferred genealogies in an F_{ST} peak and in flanking neutral regions for the first 10 Mb of scaffold 404. (**A**) Average F_{ST} values in non-overlapping 10-kb windows between *S. nigrorufa* (nig) and *S. palustris* (pal), showing a pronounced peak near the pigmentation-related gene, *SLC45A2*. Local trees are shown within the peak (*top right*) and in a flanking neutral region (*top left*), representing the relationships among 120 haploid samples from five species (see *legend*). The tree within the F_{ST} peak has two very young clades (*gray boxes*) each of which is enriched for *S. nigrorufa* or *S. palustris* and has a more recent TMRCA than expected given its size (*p*=0.01, see **Methods**). By contrast, the flanking tree is characterized by frequent deep coalescence events and incomplete lineage sorting. (**B**) Times to most recent common ancestry (in millions of generations) for half of the haploid samples (TMRCAH) from *S. nigrorufa*. Values are reduced in the F_{ST} peak compared to flanking regions, suggesting species-specific selective sweeps. (**C**) Cross-coalescence times (CC) in millions of generations between *S. nigrorufa* and *S. palustris*, as computed from the inferred ARG. No elevated cross coalescence times are observed in the F_{ST} peak, as would be expected if selection against gene flow were predominant. The smoothing method used in panels **B** and **C** was local polynomial regression; the gray bands represent 95% confidence intervals.

Figure 3: Illustration of machine-learning pipeline for prediction of selective sweeps. Based on an estimated demographic history[51], SLiM[50] is used to simulate both neutral genomic regions, and regions containing sweeps. Summary statistics specific to each species are then extracted from each simulated region. The summary statistics used were π (ref. [69]), the number of segregating sites, Tajima's D[65], θ_W (ref. [66]), θ_H (ref. [64]), the number of distinct haplotypes[62], Z_{nS} (ref. [67]), the maximum value of ω (ref. [68]), and several statistics that summarize the linkage disequilibrium (LD) distribution (see **Methods**). All statistics were used as features in training a machine-learning model (a linear SVM) to discriminate a sweep from a neutral region. The model detects not only characteristic combinations of features, but also their patterns of variation across genomic windows. The learned model is applied to the empirical data to predict individual sweeps across genomic scaffolds. Finally, the predicted sweeps are analyzed using ARG-based measures (e.g. species-specific TMRCA and RTH), and local trees that map to those predictions are explored for recent clusters of coalescent events. Similar pipelines are used to distinguish between soft and hard sweeps, recent and ancient sweeps, and partial and complete sweeps (see **Methods**).

Figure 4: (**A**) Manhattan plots showing the prediction probabilities for soft and hard sweeps in *S. melanogaster* (mel) and *S. nigrorufa* (nig) across three scaffolds harboring top F_{ST} peaks and known pigmentation genes (labeled in *red*). (**B**) Local trees inferred by *ARGweaver* in 10 kb windows classified as soft or hard sweeps in *S. melanogaster* (mel) and *S. nigrorufa* (nig). The red arrow in panel A points to the classified window from which each local tree was extracted. Shown also is the RTH of the local tree and the prediction probability for soft or hard sweeps associated with the 10 kb window. Highlighted for each local tree are the two youngest clades containing at least three haploid samples from the target species. We also highlight the youngest clade that contains at least half the haploid samples for the target species (TMRCAH). (**C**) Cumulative distribution functions of *ARGweaver*-based estimates of species-specific TMRCA (*left*) and RTH (*middle*) for *S. melanogaster* in regions classified as hard sweeps (red), soft sweeps (blue), or other classes (green) (see **Methods**). These statistics are depleted by selective sweeps, owing to clusters of recent coalescent events. (*right*) Relationship between homozygosity-based statistics $\frac{H_2}{H_1}$ and H_{12} (see **Methods**) for simulated and real *S. melanogaster* data. Each dot in the plot corresponds to a 50 kb region, and its color indicates its predicted (or simulated) class.

Figure 5: Predictions of the soft sweep classifier expanded to all five species and corresponding genealogical signatures. (**A**) Prediction probabilities for soft sweeps in each of the five species (top; see legend for species color code) in a region containing an F_{ST} peak (dashed vertical lines) upstream of the *KITL* gene (red) on scaffold 412. A high-confidence prediction for a soft sweep in S. *nigrorufa* (prediction probability 0.94) is inferred roughly 130 kb upstream of the *KITL* gene. The 50 kb region surrounding this predicted peak (gray bar) exhibits genealogical signatures of a sweep in two ARG-based statistics: (1) high enrichment scores for S. *nigrorufa* (middle) indicating that local trees in this region contain clades enriched for this species, and (2) low RTH (bottom) indicating that the clades enriched for S. *nigrorufa* are young. (**B**) A local tree inferred by *ARGweaver* in the region containing the inferred soft sweep (position 3.43 Mb). Tree tips are colored based on species label (see legend) and color of internal branches represents average over all offspring branches. The figure focuses on the last million generations, with deeper branches faded out and shown in half scale. The tree contains a young clade (~105,000 generations) with 15 haploid samples, all from S. *nigrorufa* (out of the total 24 samples). This clade contains two major subclades (gray background), which could possibly correspond to two haplotype backgrounds for the inferred soft sweep.

Table 1: Summary of genealogical signatures of speciation. Results are for the 25 F_{ST} peaks previously identified in pairwise analyses of *S. hypoxantha* (hyp), *S. melanogaster* (mel), *S. nigrorufa* (nig), *S. palustris* (pal), and *S. pileata* (pil)[42]. Associated local trees were extracted from the *ARGweaver*-inferred ancestral recombination graph and examined for evidence of species separation (test 1: gray cells, $p<0.0001$), young clades indicating recent species-specific selective sweeps (test 2: red circles, $p<0.0001$), and deep separation indicating selection against gene flow (test 3: blue circles, quantile difference > 0.1). See **Methods** and **Supplementary Tables S2-S4** for details.

peak id	zebra finch chromosome	length (kb)	nearby genes	hyp	mel	nig	pal	pil
Scaffold59	15	85		(gray)				
Scaffold118	2	25				🔴		
Scaffold252	20	90	*ASIP*		🔴	🔵	🔵	🔵
Scaffold257a	Z	370				🔴		
Scaffold257b	Z	500	*TYRP1*		🔴			
Scaffold257c	Z	840	*MLANA*				🔴	
Scaffold257d	Z	265			🔴			
Scaffold257e	Z	45						
Scaffold263	Z	535	*MYO5A*			(gray)		
Scaffold308	-	140			(gray)	🔴		(gray)
Scaffold404a	Z	765	*SCL45A2*			🔴	🔴	(gray)
Scaffold404b	Z	195				🔴		
Scaffold412	1A	205	*KITL*			🔴		(gray)
Scaffold430	1	85			(gray)			🔴
Scaffold567	2	260				🔴		🔴
Scaffold579	1	670				🔴		
Scaffold637a	Z	300				(gray)		
Scaffold637b	Z	40						
Scaffold762	11	35				🔴		🔴
Scaffold766	4	65					🔴	
Scaffold791	1	50	*DCT* or *TYR*				🔴	
Scaffold1635	6	30			🔵			
Scaffold1717	4	30	*CAMK2D*	(gray)	🔵			
Scaffold1954	5	75					🔵	
Scaffold3622	1	385			🔴	(gray)		