Plasmodium oocysts respond with dormancy to crowding and nutritional stress

Tibebu Habtewold*, Aayushi A. Sharma, Claudia A.S. Wyer, Ellen K.G. Masters, Nikolai Windbichler and George K. Christophides*

Department of Life Sciences, Imperial College London, London, UK

*For correspondence: t.habtewold@imperial.ac.uk, g.christophides@imperial.ac.uk

Abstract

Malaria parasites develop and grow as oocysts in the mosquito for several days before being able to infect another human. During this time, mosquitoes take regular bloodmeals to replenish their nutrient and energy reserves needed for flight and reproduction. We hypothesized that supplemental bloodmeals are critical for oocyst growth and that experimental infection protocols, typically involving a single bloodmeal, cause nutritional stress to developing oocysts. Therefore, enumerating oocysts independently of their growth and differentiation state may lead to erroneous conclusions regarding the efficacy of malaria transmission blocking interventions. We tested this hypothesis in *Anopheles coluzzii* mosquitoes infected with human and rodent parasites *Plasmodium falciparum* and *Plasmodium berghei*, respectively. We find that oocyst growth rates decrease at late developmental stages as infection intensities increase; an effect exacerbated at very high infection intensities. Oocyst growth and differentiation can be restored by supplemental bloodmeals even at high infection intensities. We show that high infection intensities as well as starvation conditions reduce RNA Polymerase III activity in oocysts unless supplemental bloodmeals are provided. Our data suggest that oocysts respond to crowding and nutritional stress by employing a dormancy-like strategy and urge development of alternative methods to assess the efficacy of transmission blocking interventions.

Main text

The lifecycle of the malaria parasite *Plasmodium* inside a mosquito vector begins with gametocytes ingested by a female mosquito during a bloodmeal. In the mosquito midgut lumen, gametes fuse to form zygotes that develop into motile ookinetes. Ookinetes traverse the midgut-epithelium and transform into oocysts. Multiple rounds of endomitotic replications inside an oocyst result in production of thousands of sporozoites, a process known as sporogony. Upon oocyst rupture, sporozoites migrate to the salivary gland for transmission to humans via a mosquito bite.

Current malaria control efforts focus on development of transmission blocking interventions including transmission blocking vaccines and drugs as well as genetically engineered mosquitoes expressing antimalarial effectors[1,2]. Experimental mosquito infections with *Plasmodium* parasites, and microscopic enumeration of oocysts is the gold standard for evaluation of transmission reduction, whereby mosquitoes are offered an infectious bloodmeal and maintained on sugar solution until oocysts are counted[3]. Infections often lead to high oocyst loads and consequently massive proliferation of sporozoites, causing depletion of nutrients in the vector[4-7]. For example, ultra-high intensity infections of *Anopheles stephensi* mosquitoes with the rodent parasite *P. berghei* leads to reduced number of sporozoites reaching the salivary glands and high mosquito mortality compared with lower infection intensities[8]. High parasite infections have been linked to marked reduction in free amino acids in the mosquito[6], which together with lipids are chiefly acquired from bloodmeals and are essential for most physiological processes and reproduction[9]. Free amino acids are also critical for parasite development; e.g. Isoleucine (Ile), a major component of *P. falciparum* proteins, is exclusively provided by the mosquito through digestion of blood serum[10].

Under nutritionally limiting conditions, asexual *P. falciparum* is shown to enter a dormancy-like state, restricting protein synthesis to halt cell proliferation and inhibiting catabolism to maintain viability[11,12]. Since *Plasmodium* parasites lack the target of rapamycin (TOR), the key regulator of the canonical nutrient-sensing pathway in most eukaryotes, they use the mitochondrial association factor 1 (MAF1) to regulate this metabolic switch[13]. MAF1 acts as a repressor of RNA polymerase III (Pol III) that controls expression of highly abundant non-coding RNAs and is thus essential for cell maintenance, growth and proliferation[14,15]. Under nutritional stress, *Plasmodium* MAF1 binds to Pol III switching off tRNA transcription. When conditions become favorable, MAF1 is phosphorylated and reverse translocated to the cytoplasm, freeing Pol III[13]. MAF1 mutant *P. falciparum* exhibits increased tyrosine tRNA ($tRNA^{Tyr}$) expression and fails to recover from dormancy after long periods of incubation in Ile-deficient medium.

To date, mostly indirect evidence exists for a *Plasmodium* oocyst strategy to cope with nutritional stress. *P. berghei* oocyst size and sporozoite output are markedly reduced in *Anopheles gambiae*

silenced for Lipophorin (Lp), the main lipid transporter of insects[4,16], and as mentioned previously
very high *P. berghei* infections result in reduced number of sporozoites in *A. stephensi* salivary
glands[8]. These reports and unpublished observations we and others have made over the years led
us to hypothesize that experimental infections may cause a nutritional stress to the developing
oocyst, affecting sporogonic development and the measurable infection outcome. Here we
investigated and corroborated this hypothesis using *P. berghei* and *P. falciparum* infections of *A.*
coluzzii (also known as M-form *A. gambiae*). Our results stress the need for improved assays in
studies of malaria transmission, especially those assessing the efficacy of transmission blocking
interventions.

Oocyst intensity affects *Plasmodium* DNA content

We optimized a quantitative PCR (qPCR) assay that detects *Plasmodium* mitochondrial
cytochrome-b (*Cyt-b*) gene in DNA extracted from as few as 50 merozoites spiked into *A. coluzzii*
midgut homogenates (**Fig. S1**). The qPCR amplification-curves were distinct from background
noise in uninfected midguts (**Fig. S2**). Using this assay, we found that *PbCyt-b* abundance (used as
proxy for oocyst growth) in the midgut of mosquitoes infected 7 days earlier with *P. berghei* was
positively and significantly (P<0.0001) correlated with oocyst intensity recorded by microscopic
enumeration prior to midgut homogenization (**Fig. 1a**). The results also showed that infection
intensity does not influence *PbCyt-b* DNA abundance per capita oocyst (PCO; **Fig. 1b, c**). This
pattern changes at later stages of development, as no correlation was detected between the *PbCyt-b*
abundance and oocyst intensity at 14 days post infection (dpi; **Fig. 1d**). Interestingly, the oocyst
intensity was negatively correlated (P=0.001) with *PbCyt-b* abundance PCO (**Fig. 1e**). When
midguts were grouped into two categories based on oocyst intensity, *PbCyt-b* abundance PCO was
markedly lower (P=0.007) in those harboring <15 oocysts than those with ≥15 oocysts (**Fig. 1f**).

The impact of oocyst intensity on the abundance of parasite DNA PCO was further examined in
mosquitoes where the immune factor *LRIM1* was silenced with dsRNA injection 3 days prior to *P.*
berghei infection. These mosquitoes harbored over 10 times more oocysts than control *dsLacZ*
injected mosquitoes (**Fig. S3**). At 14 dpi, *LRIM1*-silenced mosquitoes yielded oocysts of
significantly reduced (P<0.0001) diameter compared to controls (**Fig. 2a**). A significantly decreased
abundance of *PbCyt-b* PCO in *LRIM1*-silenced mosquitoes compared to controls was observed both
pre-sporulation (7 dpi; P<0.0001) and sporulation (14 dpi; P<0.003) stages (**Fig. 2b, c**). These data
suggested that high infection intensity negatively affects oocyst growth. Therefore, with standard
post-infection mosquito husbandry, where mosquitoes obtain a single bloodmeal and are then

maintained on sugar throughout the course of infection, nutrient starvation may have detrimental
impacts on the developing oocyst.

Supplemental bloodmeals accelerate oocyst growth

To assess the impact of post-infection mosquito blood feeding on oocyst growth, parasite DNA
abundance was compared between *P. berghei* oocysts in mosquitoes maintained under standard
husbandry regime and mosquitoes provided a supplemental, uninfected bloodmeal 7 dpi. The
results revealed that the supplemental bloodmeal significantly increased (P=0.029) the oocyst size
compared to control (**Fig. 3a, b**). In these mosquitoes, oocysts contained on average significantly
more DNA (P=0.003) than oocysts in control mosquitoes (**Fig. 3c**). The supplemental bloodmeal
significantly increased (P<0.0001) the correlation between *P. berghei* oocyst intensity and DNA
abundance per midgut (**Fig. 3d**). It also abolished the negative correlation between oocyst intensity
and abundance of parasite DNA PCO (**Fig. 3d**), which was observed in mosquitoes under standard
husbandry regime (see **Fig. 1e**), revealing that oocyst growth ensues when mosquitoes receive a
supplemental bloodmeal even when infection intensities are high.

We further investigated the effect of mosquito starvation on oocyst growth by augmenting oocyst
numbers through silencing *LRIM1* either alone or in combination with *Lp*, thus creating extreme
crowding and starvation conditions[4]. The results revealed a remarkable variation of *P. berghei*
oocyst size in midguts of both *LRIM1* and *LRIM1+Lp* silenced mosquitoes maintained under a
standard husbandry regime, particularly in the latter mosquito cohort (**Fig. 3f left panels**). This
phenotype was reversed when mosquitoes were provided a supplemental bloodmeal 7 dpi, as shown
by a significant increase of median oocyst size in both *LRIM1* (P<0.0001) and *LRIM1+Lp*
(P<0.0001) silenced mosquitoes (**Fig. 3f right panels** and **Fig. 4b**).

Analogous and indeed more pronounced differences were recorded for *P. falciparum* oocysts in
mosquitoes that received a supplemental bloodmeal at day-3 pi compared to control mosquitoes that
received only a single, infected bloodmeal (**Fig. 3h-j**). The supplemental bloodmeal was provided
at day-3 pi due to the faster bloodmeal digestion and oocyst development in these mosquitoes which
are maintained at 27°C instead of 21°C, while dissection was performed at day-9 pi.

Oocysts enter a dormancy-like state under starvation conditions

Female mosquitoes naturally lay their eggs before taking another bloodmeal to replenish energy
reserves and embark on a new gonotrophic cycle. Follicular resorption that is mediated by juvenile

hormone is employed to increase energy resources in case of nutrient limitation or when oviposition

is not possible[17], as is the case of most experimental infection protocols. This process is presumed

to also provide nutrient supplies to the developing oocyst[8,18]. We reproduced the natural infection

conditions by allowing mosquitoes infected with *P. falciparum* to lay their eggs at day 2 pi, before

they were provided supplemental bloodmeals at either day 3 or day 6 pi. The results showed a

dramatic difference reflected by significantly decreased oocyst sizes recorded at day 9 pi in

mosquitoes deprived of either of supplemental bloodmeals (both $P<0.0001$; **Fig. 4a, b**). This

finding was consistent with lower parasite DNA abundance PCO in bloodmeal-deprived mosquitoes

compared to those provided a supplemental bloodmeal at 3 ($P<0.005$) or 6 ($P<0.002$) dpi (**Fig. 4c**).

No significant differences in oocyst size or parasite DNA abundance PCO were recorded between

mosquitoes provided supplemental bloodmeals at 3 or 6 dpi, suggesting that oocyst growth can

quickly recover after a bloodmeal.

Oocysts retained their propensity to grow even when a supplemental bloodmeal was provided as

late as 9 dpi ($P<0.0001$; **Fig. 4d, e**). In this experiment, midgut dissections and oocyst observations

were carried out 13 dpi. Interestingly, the distribution of oocyst sizes in mosquitoes deprived of

supplemental bloodmeal was positive-skewed (**Fig. 4f**), and several oocysts in these mosquitoes

were as big as the biggest oocysts in mosquitoes provided a supplemental bloodmeal 9 dpi. This

implies that, under limiting nutrient conditions, the oocyst population employs an effective resource

management strategy whereby some oocysts are let to grow whereas most remain dormant. When

faced with resource stress, *Plasmodium* is indeed thought to induce growth restriction mediated by

extracellular vesicles, limiting growth to some parasite cells and ensuring transmission[19].

Furthermore, the distribution of oocyst sizes in mosquitoes receiving a supplemental bloodmeal 9

dpi was negative-skewed, and several oocysts were as small as the smallest oocysts in mosquitoes

deprived of supplemental bloodmeals, suggesting that not all oocysts effectively exit dormancy

following a long starvation period.

We investigated the abundance of *P. falciparum* sporozoites in salivary glands 14 dpi by

quantifying *PfCyt-b* in the mosquito thorax and head regions (**Fig. 4g**). The results revealed that

deprivation of supplemental bloodmeals combined with oviposition at 2 dpi resulted in significantly

lower levels of sporozoites compared to mosquitoes provided supplemental bloodmeals at 3

($P=0.009$) or 6 ($P=0.019$) dpi. Nevertheless, most of these mosquitoes (70%) did harbor detectable

sporozoite loads in their salivary glands, like those provided supplemental bloodmeals (81% at 3

dpi and 86% at 6 dpi), indicating that some oocysts, presumably the fully-grown ones, were fully

differentiated and produced sporozoites. Despite having much larger oocysts, mosquitoes provided

a supplemental bloodmeal 9 dpi, showed equal sporozoite loads compared to those deprived of

supplemental blood meals, suggesting that more time was needed for full oocyst differentiation.

The decelerated oocyst growth in nutrient limiting conditions suggested that oocysts might employ

a dormancy-like strategy to overcome starvation. To examine this, parasite *pre-tRNA^{Tyr}* abundance

was quantified as a proxy for Pol III activity in oocysts. Measuring immature instead of mature

tRNA^{Tyr} served to overcome the long half-life of mature tRNA that might affect real-time

quantification of gene expression[13]. *Pre-tRNA^{Tyr}* expression was calibrated to parasite DNA

abundance PCO. The results revealed that the levels of *P. falciparum pre-tRNA^{Tyr}* expression levels

in oocysts 7 dpi were significantly lower (P=0.008) in mosquitoes that received a single, infectious

bloodmeal compared to those that additionally received a supplemental bloodmeal 3 dpi (**Fig. 4h**).

Similarly, *P. berghei tRNA^{Tyr}* expression in *LRIM1* (P<0.0001) and *LRIM1+Lp* silenced (P<0.0001)

conditions measured in oocysts 14 dpi were significantly lower in mosquitoes under standard

husbandry regimes compared to those that received a supplemental bloodmeal 7 dpi (**Fig. 4i**). These

data together suggest that oocysts enter a dormancy-like state in the mosquito vector when faced

with nutrient depletion caused by bloodmeal deprivation, high infection intensities or disruption of

nutrient shuttling.

Discussion

Plasmodium spends a large part of its life as an oocyst within the *Anopheles* mosquito, where it

faces a variety of challenges that can adversely affect its growth, differentiation and transmission

potential[20]. A key challenge is the high nutritional demand to produce thousands of sporozoites. We

demonstrate that under starvation conditions, when an oocyst-harboring mosquito is deprived of

supplemental bloodmeals, infection intensity plays a major role in oocyst growth, as measured by

both DNA content and oocyst size. Mosquitoes harboring many oocysts, a condition that can occur

naturally in field infections and invariably in experimental infections and which can be further

elevated by inactivating the mosquito immune response, exhibit substantial variability in oocyst

growth. In such conditions of crowding and nutritional stress, expression of parasite *pre-tRNA^{Tyr}* is

significantly reduced, indicating that the oocyst enters a dormancy-like state.

Our findings bear important implications not only for understanding the mosquito-parasite

interactions underpinning malaria transmission but also in evaluating the efficacy of transmission

blocking interventions. The current gold standard in malaria transmission studies is microscopic

enumeration of oocysts present in the mosquito midgut. It has been previously shown that the

number of oocysts correlates well with the number of gametocytes ingested by the mosquito and

can therefore be used as a measure of the efficacy of interventions acting on gametocytes, gametes

and/or ookinetes[21-23]. However, this measurement does not consider any knock-on or additional

impacts of the intervention on the oocyst. Although this additional impact is unlikely to reverse any

detrimental effect on previous developmental stages, our data suggest that reduction in oocyst

numbers may translate to a lesser or no impact on the sporozoite load, especially when the effect is

small and oocyst competition for nutrients is reduced. Thus, a reduction in oocyst intensity alone is

an insufficient measurement and may not accurately reflect the efficacy of a transmission blocking

intervention.

At high intra-population densities, endoparasites are known to evoke competition for nutrients when

these are of limited supply[24]. This phenomenon is ubiquitous in nature and often leads to negative

density-dependent processes with significant effects on parasite fitness such as proliferation and

ability to avoid host immune defenses[25]. The density-dependent effect on *Plasmodium* sporogony

reported here is not related to mosquito immune responses known to reduce the number of oocysts

but to the rate of oocyst growth through endoreplication. The effect is indeed exacerbated upon

silencing the mosquito complement-like system due to the significant increase of the number of

ookinetes that successfully develop to oocysts.

The negative density-dependent effect on the oocyst is exacerbated in standard laboratory infections

where infected mosquitoes are often not provided supplemental bloodmeals, leading to acute

depletion of nutrient reserves[4-7,26]. In nature, a female mosquito acquires a bloodmeal every 2-3

days[27,28], which helps it meet the high energy and nutrient demand for flight and reproduction.

These additional bloodmeals are also thought to be important for the proliferating oocyst[18].

Plasmodium is auxotrophic for most amino acids; of these, free Isoleucine exclusively derives from

bloodmeal digestion, as neither the parasite nor the mosquito can synthesize it *de novo*[10,29,30].

Bloodmeal-derived fatty acids are also critical for mosquito fitness and reproduction, chiefly

derived from hemoglobin metabolism in the gut cells and distributed across mosquito tissues by Lp.

Silencing the *Lp* gene abolishes egg development and compromises growth of both *P. berghei* and

P. falciparum oocysts[4,16]; albeit studies showing that mature late-stage *P. falciparum* oocysts

synthesize lipids *de novo* via the type two fatty acid biosynthesis pathway (FAS-II), in contrast to *P.*

berghei and asexual stages that scavenge fatty acids exclusively from the host[31]. We show that

silencing *Lp* together with the complement-like immune pathway exacerbates the negative effects

of crowding on oocyst growth but cannot further reduce *pre-tRNA^{Tyr}* expression, indicating that the

dormancy-like state is likely to be regulated by amino acid sensing[12].

Better understanding of the oocyst response to crowding and nutritional stress could assist

development of novel interventions to limit malaria transmission. For example, a temporally tuned

targeting of pathways involved in mosquito amino acid or lipid metabolism using genetic engineering could prolong completion of the parasite extrinsic incubation period (EIP), delaying oocyst growth and differentiation and sporozoite release. In nature, only a small proportion of mosquitoes can survive through the *P. falciparum* EIP[32,33]; hence, even a small delay in oocyst growth and differentiation could have a drastic negative impact on malaria transmission.

Importantly, our findings highlight a major drawback of most standard laboratory infection protocols that can lead to erroneous conclusions when evaluating the transmission-blocking efficacy of an intervention. First, an intervention that inhibits or delays oocyst growth and/or differentiation without or by weakly affecting oocyst counts may be rejected owing to a decision merely based on reduction of oocyst numbers. This pertains to interventions targeting both mosquito or parasite molecules and pathways, e.g. mosquito *Lp* as described earlier[16,34] and parasite *misfit* gene of which the knockout produces oocysts arrested at early mitotic stages but continuing to grow in size[35]. Second, inducing a dormancy-like state, a universally resilient life stage, may make oocysts resistant to transmission blocking interventions. Third, the current standard protocol may lead to false positive conclusions in favor of an intervention reducing oocyst numbers whilst it has no real impact on malaria transmission owing to the density-dependent effects described here. We conclude that provision of supplemental bloodmeals to mitigate any parasite density-dependent effects on oocyst growth and development and assessment of the actual parasite load in the oocyst that escape the intervention through a molecular diagnostic method are profoundly important.

Methods

Mosquito infections

A. coluzzii of the N'gousso strain were reared following standard protocols and infected with *P. berghei ANKA507m6cl1* that constitutively expresses GFP[36] by direct feeding on infected mice or with *P. falciparum* gametocytes from a laboratory culture (NF54 strain) using standard membrane feeding. Engorged mosquitoes were provided 10% sucrose and maintained at 21°C for *P. berghei* infections and 27°C for *P. falciparum* infections until midgut dissection. Supplemental bloodmeals on human blood was provided via membrane feeding.

Midgut preparations and oocyst enumeration

Gut dissections were performed in ice-chilled PBS. For *P. berghei* infection, guts were mounted on glass slides and oocysts were counted with fluorescence microscopy. For *P. falciparum* infection,

guts were stained with 0.5% Mercurochrome for 10 min on ice and washed twice in ice chilled PBS

for 30 min, and oocysts were enumerated using light microscopy. After oocyst enumeration, the

foregut and hindgut were removed, and midguts were individually transferred to 1.5 ml screw-top

tubes with 350 µl RLT Plus buffer from Qiagen AllPrep DNA/RNA Mini Kit and kept at -20°C

until DNA and RNA extraction.

Genomic DNA and RNA extraction

Midguts were homogenized using PRECELLYS® 24 (Bertin Technologies) before simultaneous

extraction of genomic DNA and total RNA using the AllPrep DNA/RNA Mini Kit. DNA and RNA

concentrations were determined using Nanodrop (Thermo Scientific), and samples were kept at -

20°C until cDNA synthesis and qPCR analysis. To measure the abundance of *P. falciparum*

sporozoites in salivary glands, mosquitoes were killed with 75% ethanol and placed on dry ice and

their abdomens were removed. Genomic DNA was extracted from the remaining thorax and head

regions of each mosquito using Qiagen DNeasy 96 Blood and Tissue kit.

Quantitative PCR analysis

SYBR Green based qPCR was used to quantify *Plasmodium Cyt-b* mitochondrial DNA. Assay

sensitivity and specificity were determined with serial dilutions of *P. falciparum* merozoites from a

synchronized asexual culture after heat inactivation, quantified and collected using a BD FACSAria

III cell sorter. QPCR reactions were performed in 20 µl volumes according to the manufacturer's

instructions and previously described primers for *P. berghei*[37] and *P. falciparum*[38]. *Pre-*

tRNA^{Tyr} expression levels were determined with the Qiagen Quantinova SYBR Green PCR kit using

the stem-loop qPCR technique and qPCR primers described previously[13].

DNA and RNA abundances were measured using the standard curve-based qPCR method. Curves

for each target gene and the reference gene (*A. gambiae* S7 ribosomal protein gene[39]) were

constructed after serial dilution of nucleic acid templates. Ct-values were standardized using their

respective standard curves, and the target gene Ct value was normalized to that of the reference

gene. *Pre-tRNA^{Tyr}* PCO was standardized to *Cyt-b* abundance.

Data analysis

GraphPad Prism v.8.4.0 software package was used for data analysis and presentation. Spearman's

rank-order correlation analysis was used to measure the relationship between oocyst intensity ($\log_{10}$

transformed) and relative DNA abundance per single midgut or oocyst. All other statistical analyses
were performed using the Kolmogorov-Smirnov test.

Acknowledgements

297 We thank Jane Srivastava and Jessica E. Rowley for assisting with parasite cell sorting and Maria
298 G. Inghilterra for gametocyte culturing. The work was funded by the Bill and Melinda Gates
299 Foundation grant OPP1158151 to N.W. and G.K.C and the Wellcome Trust Investigator Award
300 107983/Z/15/Z to G.K.C.

Author contributions

303 T.H. and G.K.C. contributed conceptualization; T.H., A.A.S., C.A.S.W. and E.K.G.M. contributed
304 methodology; T.H. and A.A.S. conducted investigation; T.H. and G.K.C. performed formal
305 analysis; T.H. and A.A.S. wrote original draft; T.H. and G.K.C. wrote the paper; T.H., N.W. and
306 G.K.C. provided supervision; G.K.C. conducted project administration; N.W. and G.K.C. acquired
307 funding.

Competing interests

310 The authors declare no competing interests.

References

1   Gantz, V. M. *et al.* Highly efficient Cas9-mediated gene drive for population modification
of the malaria vector mosquito *Anopheles stephensi*. *Proc Natl Acad Sci U S A* **112**, E6736-
6743, doi:10.1073/pnas.1521077112 (2015).

2   Sinden, R. E. Developing transmission-blocking strategies for malaria control. *PLoS Pathog*
13, e1006336-e1006336, doi:10.1371/journal.ppat.1006336 (2017).

3   Churcher, T. S., Blagborough, A. M. & Sinden, R. E. Measuring the blockade of malaria
transmission: analyzing the results of mosquito feeding assays. *Malaria journal* **11**, P18
(2012).

4   Costa, G. *et al.* Non-competitive resource exploitation within mosquito shapes within-host
malaria infectivity and virulence. *Nature Communications* **9**, 3474, doi:10.1038/s41467-
018-05893-z (2018).

5   Ferguson, H. M. & Read, A. F. Genetic and environmental determinants of malaria parasite
virulence in mosquitoes. *Proc Biol Sci* **269**, 1217-1224, doi:10.1098/rspb.2002.2023 (2002).

6   Maier, W. A., Becker-Feldman, H. & Seitz, H. M. Pathology of malaria-infected
mosquitoes. *Parasitology Today* **3**, 216-218, doi:10.1016/0169-4758(87)90063-9 (1987).

7   Takken, W. *et al.* Larval nutrition differentially affects adult fitness and *Plasmodium*
development in the malaria vectors *Anopheles gambiae* and *Anopheles stephensi*. *Parasites*
& Vectors **6**, 345 (2013).

8   Pollitt, L. C. *et al.* Costs of crowding for the transmission of malaria parasites. *Evolutionary*
applications **6**, 617-629, doi:10.1111/eva.12048 (2013).

9   Rivera-Pérez, C., Clifton, M. E. & Noriega, F. G. How micronutrients influence the
physiology of mosquitoes. *Curr Opin Insect Sci* **23**, 112-117,
doi:10.1016/j.cois.2017.07.002 (2017).

10   Divo, A. A., Geary, T. G., Davis, N. L. & Jensen, J. B. Nutritional requirements of
Plasmodium falciparum in culture. I. Exogenously supplied dialyzable components
necessary for continuous growth. *The Journal of protozoology* **32**, 59-64,
doi:10.1111/j.1550-7408.1985.tb03013.x (1985).

11   Babbitt, S. E. *et al.* *Plasmodium falciparum* responds to amino acid starvation by entering
into a hibernatory state. **109**, E3278-E3287 (2012).

12   Mancio-Silva, L. *et al.* Nutrient sensing modulates malaria parasite virulence. *Nature* **547**,
213, doi:10.1038/nature23009 (2017).

13   McLean, K. J. & Jacobs-Lorena, M. *Plasmodium falciparum* Maf1 confers survival upon
amino acid starvation. *MBio* **8**, e02317-02316 (2017).

14   Boguta, M. Maf1, a general negative regulator of RNA polymerase III in yeast. *J*
Biochimica et Biophysica Acta -Gene Regulatory Mechanisms **1829**, 376-384 (2013).

15   Moir, R. D. & Willis, I. Regulation of pol III transcription by nutrient and stress signaling
pathways. *J Biochimica et Biophysica Acta -Gene Regulatory Mechanisms* **1829**, 361-375
(2013).

16   Mendes, A. M. *et al.* Conserved mosquito/parasite interactions affect development of
Plasmodium falciparum in Africa. *PLoS Pathog* **4**, e1000069,
doi:10.1371/journal.ppat.1000069 (2008).

17   Clifton, M. E. & Noriega, F. G. Nutrient limitation results in juvenile hormone-mediated
resorption of previtellogenic ovarian follicles in mosquitoes. *J Insect Physiol* **57**, 1274-1281,
doi:10.1016/j.jinsphys.2011.06.002 (2011).

18   Hurd, H., Hogg, J. C. & Renshaw, M. Interactions between bloodfeeding, fecundity and
infection in mosquitoes. *Parasitology Today* **11**, 411-416 (1995).

19   Mantel, P. Y. & Marti, M. The role of extracellular vesicles in *Plasmodium* and other
protozoan parasites. *Cell Microbiol* **16**, 344-354, doi:10.1111/cmi.12259 (2014).

20   Gamage-Mendis, A. C. *et al.* Infectivity of *Plasmodium vivax* and *P. falciparum* to
Anopheles tessellatus; relationship between oocyst and sporozoite development.
Transactions of the Royal Society of Tropical Medicine and Hygiene **87**, 3-6,
doi:10.1016/0035-9203(93)90396-8 (1993).

21   Churcher, T. S. *et al.* Measuring the blockade of malaria transmission--an analysis of the
Standard Membrane Feeding Assay. *International journal for parasitology* **42**, 1037-1044,
doi:10.1016/j.ijpara.2012.09.002 (2012).

22   Da, D. F. *et al.* Experimental study of the relationship between *Plasmodium* gametocyte
density and infection success in mosquitoes; implications for the evaluation of malaria
transmission-reducing interventions. *Exp Parasitol* **149**, 74-83,
doi:10.1016/j.exppara.2014.12.010 (2015).

23   Stone, W. J. R. *et al.* The relevance and applicability of oocyst prevalence as a read-out for
mosquito feeding assays. *Scientific Reports* **3**, 3418, doi:10.1038/srep03418 (2013).

24   Anderson, R. The regulation of host population growth by parasitic species. *Parasitology*
76, 119-157 (1978).

25   Churcher, T. S., Filipe, J. A. & Basanez, M. G. Density dependence and the control of
helminth parasites. *Journal of Animal Ecology* **75**, 1313-1320 (2006).

26   Nyasembe, V. O. *et al.* *Plasmodium falciparum* infection increases *Anopheles gambiae*
attraction to nectar sources and sugar uptake. *Current Biology* **24**, 217-221 (2014).

27   Ponnudurai, T. *et al.* Sporozoite load of mosquitoes infected with *Plasmodium falciparum.*
Transactions of the Royal Society of Tropical Medicine and Hygiene **83**, 67-70 (1989).

28   Beier, J. C. Malaria parasite development in mosquitoes. *Annual Review of Entomology* **43**,
519-543 (1998).

29   Liu, J., Istvan, E. S., Gluzman, I. Y., Gross, J. & Goldberg, D. *Plasmodium falciparum*
ensures its amino acid supply with multiple acquisition pathways and redundant proteolytic
enzyme systems. *Proceedings of the National Academy of Sciences* **103**, 8840-8845 (2006).

30   Mack, S. R., Samuels, S. & Vanderberg, J. P. Hemolymph of *Anopheles stephensi* from
uninfected and *Plasmodium berghei*-infected mosquitoes. 2. Free amino acids. *The Journal*
of parasitology **65**, 130-136 (1979).

31   van Schaijk, B. C. *et al.* Type II fatty acid biosynthesis is essential for *Plasmodium*
falciparum sporozoite development in the midgut of *Anopheles* mosquitoes. *Eukaryot Cell*
13, 550-559, doi:10.1128/EC.00264-13 (2014).

32   Charlwood, J. *et al.* Survival and infection probabilities of anthropophagic anophelines from
an area of high prevalence of *Plasmodium falciparum* in humans. *Bulletin of Entomological*
Research **87**, 445-453 (1997).

33   Ryan, S. J., Ben-Horin, T. & Johnson, L. R. Malaria control and senescence: the importance
of accounting for the pace and shape of aging in wild mosquitoes. *Ecosphere* **6**, 1-13 (2015).

34   Rono, M. K., Whitten, M. M., Oulad-Abdelghani, M., Levashina, E. A. & Marois, E. The
major yolk protein vitellogenin interferes with the anti-plasmodium response in the malaria
mosquito *Anopheles gambiae*. *PLoS Biol* **8**, e1000434, doi:10.1371/journal.pbio.1000434
(2010).

35   Bushell, E. S. *et al.* Paternal effect of the nuclear formin-like protein MISFIT on
Plasmodium development in the mosquito vector. *PLoS Pathog* **5**, e1000539,
doi:10.1371/journal.ppat.1000539 (2009).

36   Janse, C. J. *et al.* High efficiency transfection of *Plasmodium berghei* facilitates novel
selection procedures. *Mol Biochem Parasitol* **145**, 60-70,
doi:10.1016/j.molbiopara.2005.09.007 (2006).

37   Shimizu, S., Osada, Y., Kanazawa, T., Tanaka, Y. & Arai, M. J. M. j. Suppressive effect of
azithromycin on *Plasmodium berghei* mosquito stage development and apicoplast
replication. **9**, 73 (2010).

38   Farrugia, C. *et al.* Cytochrome b gene quantitative PCR for diagnosing *Plasmodium*
falciparum infection in travelers. **49**, 2191-2195 (2011).

39   Meister, S. *et al.* Immune signaling pathways regulating bacterial and malaria parasite
infection of the mosquito *Anopheles gambiae. Proc Natl Acad Sci U S A* **102**, 11420-11425,
doi:10.1073/pnas.0504950102 (2005).

 Figures

Fig 1. *P. berghei Cyt-b* **DNA abundance is oocyst density dependent.**

a, Correlation between oocyst density and abundance of *PbCyt-b* mitochondrial DNA 7 dpi (r = 0.24; P < 0.0001). **b**, Correlation between oocyst density and *PbCyt-b* DNA abundance PCO 7 dpi. **c**, *PbCyt-b* DNA abundance PCO in low and high oocyst density midguts 7 dpi. **d**, Relationship between oocyst intensity and *PbCyt-b* DNA abundance 14 dpi. **e**, Correlation between oocyst density and *PbCyt-b* DNA abundance PCO 14 dpi (r = -0.25; P = 0.001). **f**, *PbCyt-b* DNA abundance PCO in low and high oocyst density midguts 14 dpi. In scatterplots, each dot represents a single midgut and lines show linear regression fit. Horizontal lines in boxplots indicate median. Data were generated from at least two independent experiments.

Fig 2. Effect of crowding on *P. berghei* oocyst growth.

a, Oocyst diameter (dia.) in *LRIM1* and *LacZ* dsRNA injected mosquitoes. Data were generated from two independent biological replicates. **b-c**, *PbCyt-b* DNA abundance in *LRIM1* and *LacZ* dsRNA injected mosquitoes 7 (b) and 14 (c) dpi. Data were generated from three independent replicates. Horizontal lines in dot plots and boxplots indicate median.

Fig 3. Effect of supplemental bloodmeals on oocyst growth.

a, *P. berghei* oocyst diameter (dia.) in mosquitoes provided no supplemental bloodmeal after the time of infection (non-blood-fed; NBF) or an additional supplemental bloodmeal 7 dpi (BF7) measured 14 dpi. **b**, Representative microscopic pictures of *P. berghei* oocysts in NBF and BF7 mosquitoes taken 14 dpi. **c**, *PbCyt-b* DNA abundance in NBF and BF7 mosquitoes 14 dpi. **d**, Correlation between *P. berghei* oocyst density and *PbCyt-b* DNA abundance in the midgut of NBF and BF7 mosquitoes (r = 0.35, P<0.0001). **e**, Correlation between *P. berghei* oocyst density and *PbCyt-b* DNA abundance PCO. **f-g**, Representative microscopic pictures (**f**) and diameter (**g**) of *P. berghei* oocysts on midguts of mosquitoes silenced for *LRIM1* and *LRIM1+Lp* and maintained under a standard husbandry regime (NBF) or provided a supplemental bloodmeal 7 dpi (BF7). For oocyst diameter, oocysts were randomly selected from 20 midguts from two independent infection experiments. **h**, *P. falciparum* oocyst diameter (dia.) in mosquitoes provided no supplemental bloodmeal after the time of infection (NBF) or an additional supplemental bloodmeal 3 dpi (BF3) measured 9 dpi. **i**, Representative microscopic pictures of *P. falciparum* oocysts in NBF and BF3. **j**, *PfCyt-b* DNA abundance in NBF and BF3 mosquitoes. Horizontal lines in dot plots and boxplots indicate median. Data were generated from at least two independent experiments.

Fig 4. Effect of supplemental bloodmeal deprivation on oocyst development.

a, *P. falciparum* oocyst diameter (dia.) in mosquitoes provided no supplemental bloodmeal after the time of infection (non-blood-fed; NBF) or additional supplemental bloodmeals at 3 (BF3) or 6 (BF6) dpi measured 9 dpi. **b**, Representative microscopic pictures of *P. falciparum* oocysts in NBF, BF3 and BF6 mosquitoes taken 9 dpi. **c**, *PfCyt-b* DNA abundance in NBF, BF3 and BF6 mosquitoes 9 dpi. **d**, *P. falciparum* oocyst diameter (dia.) in mosquitoes provided no supplemental bloodmeal after the time of infection (NBF) or additional supplemental bloodmeals 9 dpi (BF9) measured 13 dpi. **e**, Representative microscopic pictures of oocysts in NBF and BF9 mosquitoes taken 13 dpi. **f**, Oocyst size distribution fitting the Poisson distribution model in NBF and BF9 mosquitoes 13 dpi. Solid vertical lines show median and dotted vertical lines show mean. **g**, Sporozoite load in NBF, BF3, BF6 and BF9 mosquitoes at 14 dpi. **h**, Relative expression levels of *P. falciparum* pre-tRNA^{Tyr} in NBF and BF3 mosquitoes measured 7 dpi. **i**, Relative expression levels of *P. berghei* pre-tRNA^{Tyr} in *LRIM1* and *LRIM1+Lp* silenced mosquitoes in NBF and BF7 mosquitoes at 14 dpi. Horizontal lines in dot plots and boxplots indicate median. Data were generated from at least two independent biological replicates.

    Supplementary Figures

Fig S1. Efficiency of quantitative PCR in determining *PfCyt-b* DNA abundance.

Merozoite cells from *P. falciparum* asexual cultures were purified and quantified using FACS
sorter. Single midgut homogenates from primiparous *A. coluzzii* mosquitoes were spiked in
merozoite serial dilutions. SYBR-based qPCR was performed following DNA extraction. The
standards curve represent average of 15 independent qPCR assays. Arrow indicates the limit of
assay detection (LoD). Note that each merozoite cell is thought to contain up to 20 mitochondria.
484
485
486

Fig S2. *PfCyt-b* DNA qPCR amplification curves in midguts with single or no oocysts.

Midguts of *A. coluzzii* mosquitoes fed on *P. falciparum* gametocytemic blood were dissected at 7
dpi and the number of oocysts were determined. DNA was extracted from 6 midguts showing
mono-oocyst infections and 1 with no oocysts. Abundances of *PfCyt-b* DNA in midguts containing
a single oocyst in relation to mosquito *S7* gene are 3.24, 6.14, 17.86, 40.37, 0.99, 0.92 and 0.002
from left to right, respectively.
495

Fig S3. *P. berghei* oocyst intensity following silencing of *LRIM1*.

Oocysts present on mosquito midguts were counted at day 14 pi. Injection of *LacZ* dsRNA was used as control. Horizontal lines in boxplots show median. Data were generated from three independent biological replicates. Kolmogorov-Smirnov test was applied to compare oocyst counts between *dsLacZ*-injected and *LRIM1* silenced mosquitoes.