Coastal development and human recreation disturbance limit Australian Pied
Oystercatcher *Haematopus longirostris* population sizes on E Australian beaches

STEPHEN TOTTERMAN

Empire Vale, NSW, 2478, Australia
Email: stephen@totterman.net.au

SUMMARY

Australian Pied Oystercatchers *Haematopus longirostris* and their habitat were surveyed on
72 beaches and 674 km of coastline, from Fraser Island, Queensland to near the New South
Wales–Victoria state border, in 2015–2018. A grand total of 232 individual birds (the sum of
mean beach counts) and 41 oystercatcher territories were counted. Regression models for
mean oystercatcher count density indicated a positive response to the abundance of the surf
clam *Donax deltoides,* a positive New South Wales Far North Coast regional effect and a
negative response to the proportion of urban beach. Models for oystercatcher territory density
indicated a positive Far North Coast effect and a negative response to pedestrian access
density. This report upgrades the coastal development and human recreation disturbance
threats for the species.

Keywords: beach-nesting bird; bird-habitat model; species distribution model

INTRODUCTION

The Australian Pied Oystercatcher *Haematopus longirostris* inhabits estuaries and ocean
beaches along the Australian coast, with major populations in the southern states of
Tasmania, South Australia and Victoria. The species is non-migratory and many breeding
pairs remain on their territories throughout the year. It is also long-lived and breeding
population sizes are usually limited by territorial behaviour and not by breeding success.
Delayed age of first breeding is normal, resulting in a surplus of non-territorial adults in
populations (see the conservation assessment by Taylor *et al.* 2014).

The Australian Pied Oystercatcher has been listed as Endangered in the state of New South
Wales (NSW Government 2016) because the estimated population size is low and projected
or continuing to decrease. Proposed conservation threats for the species in NSW include
habitat loss, human recreation disturbance, a reduction in food resources and low breeding
success, often because of depredation of eggs and chicks by the European Red Fox *Vulpes*
vulpes (NSW Scientific Committee 2010).

Microtidal, high energy ocean beaches (surf beaches) are common along the SE Australian
coast (Short 2007). These are essentially linear habitats with a narrow intertidal zone.
Australian Pied Oystercatchers typically nest on the low front dune (Wellman *et al.* 2000).
Spatial constraints and territorial behaviour mean that beach-resident oystercatcher
population sizes are smaller than those counted in large estuaries (see Taylor *et al.* 2014) and
are dispersed along the coast.

Commonly known as the 'pipi', *Donax deltoides* is a large surf clam that inhabits the
intertidal and shallow subtidal zones on ocean beaches from Fraser Island, in the state of
Queensland, to the Murray River, South Australia (Ferguson *et al.* 2018). Pipis are the major
prey for Australian Pied Oystercatchers on ocean beaches in NSW and previous studies have
reported that oystercatcher abundance is positively correlated with pipi biomass (Owner &
Rohweder 2003, Harrison 2009).

Pipis are harvested by humans for food and as bait for recreational fishing. Commercial pipi
fishing has been suggested to negatively impact stocks and thereby oystercatcher populations
(Owner & Rohweder 2003, Harrison 2009). However, surf clams are fast-growing and short-
lived, and natural variability in recruitment can also result in temporary stock declines
(Murray-Jones 1999). Only hand gathering of pipis is allowed in NSW. The commercial
fishery is small, with 52 licenses in 2018 and a FY 2015–16 catch of 176 t (Ferguson *et al.*
2018). There is no commercial pipi fishing in Queensland.

The feeding ecology of the Eurasian Oystercatcher *Haematopus ostralegus* and shellfishing
impacts have been thoroughly studied in Europe (*e.g.* Goss-Custard 1996). However, readers
are cautioned that the ecology and conservation of Australian Pied Oystercatchers on ocean
beaches is unlike that for Eurasian Oystercatchers in estuaries (Owner 1997, Taylor & Taylor
2005, Harrison 2009, Taylor *et al.* 2014, Totterman 2018).

Spatial constraints amplify human recreation disturbance for shorebirds on ocean beaches.
For example, mean beach widths, measured from the base of the front dune to the middle of
the low tide swash zone for 11 beaches on the Far North Coast of NSW were 47–98 m
(Totterman 2019a). These beach widths are similar to reported flight initiation distances for
Australian Pied Oystercatchers of 39–83 m for a human stimulus (reviewed in Weston *et al.*
2012) and 129 m for a human and dog stimulus (Harrison 2009). People also disperse from
beach access points and the mean along shore distance walked for 18 Far North Coast
beaches was 444 m (Totterman 2019b).

This study was motivated by concerns that previous studies (Owner & Rohweder 2003,
Harrison 2009) had overstated the importance of pipis for oystercatchers on ocean beaches
and overlooked the coastal development and human recreation disturbance threats. The
objective was to sample a larger number of beaches along the E Australian coast and re-
examine oystercatcher-habitat associations.

STUDY AREA AND METHODS

Sampling design

The sampling space for this study extended from the Fraser Island, Queensland S to the
NSW–Victoria border (Fig. 1). 117 exposed beaches with a SE–E aspect and > 1.5 km long,
where pipis could be abundant (Owner 1997), were identified for the sampling frame.
Beaches were defined by natural breaks, *e.g.* headlands and rivers. Some short beaches were
combined where the break was a minor obstacle, *e.g.* a short, low rocky interval that could

easily be traversed at all tides. Sampling prioritised longer, less-developed beaches, where

oystercatchers could be abundant (Owner 1997, Harrison 2009), for the purpose of estimating

bird-habitat models (Albert *et al.* 2010). Selection of high pipi and/or oystercatcher

abundance beaches was also guided by other studies and reports (Gray 2016a,b, Totterman

2019a, NSW Department of Planning, Industry and Environment (DPIE) 2019a).

The oystercatcher breeding season on the Far North Coast of NSW is Aug–Dec and late

attempts generally are replacement clutches (Wellman *et al.* 2000). Beaches were sampled in

Jul–Oct, early in the breeding season, when competition for territories and territory

occupancy was assumed to be maximum. Sampling started with beaches in the N, progressed

'down' the coast and included a few beaches on the return trip. Sampling in any one year

selected non-adjacent beaches to reduce spatial autocorrelation in oystercatcher counts that

could result from short-distance movements.

Counts

Beaches were surveyed with a handheld Global Positioning System satellite receiver (Garmin

GPS 76), marking 4–13 (mode 7) regularly spaced along shore waypoints (*i.e.* coordinates

saved in the device) that were used to measure beach length and distances for pipi sampling

and oystercatcher territory mapping.

Oystercatchers were usually counted at low tide and early in the morning, when there were

fewer people on beaches that could disturb birds and displace them along shore. A four-

wheel-drive (4WD) vehicle was used to count longer beaches, where permitted. Otherwise,

and for shorter beaches, a bicycle was used. Some of the shortest beaches, and those with a

steep face and soft sand, were counted on foot.

Oystercatchers present in any estuaries, lakes and associated sand spits adjacent to a beach

were noted but not added to the beach count. For example, coastal lakes are common along

the NSW South Coast, many of these were inhabited by oystercatchers and those birds can be

found roosting and sometimes nesting near the beach. However, those birds forage and spend

most of the time in the lakes.

'Adult'-plumage oystercatchers were counted including any second year immature plumage

birds which superficially look the same as adults (Marchant & Higgins 1993). More precise

age classification was not always possible (with variable observation distances, light, for

birds in flight, etc.) and beaches had to be counted quickly to avoid along shore movements

of birds. Any birds that were marked with bands or alphanumeric leg flags were noted to

assist in the identification of local residents.

Territory mapping (spot mapping) can easily be applied to beach-resident oystercatchers

because they are large, conspicuous birds and territories on ocean beaches are practically one-

dimensional (Totterman 2018). The minimum territory mapping effort was three counts per

beach and on non-consecutive days, to reduce serial correlation. Beaches with higher

oystercatcher counts also tended to have higher territory counts. Effort was increased to ≥

five counts for beaches with mean oystercatcher count densities > 0.5 birds/km, based on

breeding season densities of 0.3–1.9 birds/km for beaches on the Far North Coast of NSW
(Owner & Rohweder 2003, Harrison 2009). Additional, opportunistic counts were also made
for some beaches where pipi sampling continued beyond the initial counting period.
Territories were identified as adult 'pairs' that were present on the same stretch of beach on
≥ 50% of counts (*e.g.* 2/3 or 3/5). These are rapid, low-cost estimates and not all pairs that
occupy a stretch of beach may breed. Territories are compared with independent counts of
breeding pairs in the Results.

Pipi abundance

A new 'feet digging' method (Totterman 2019c) was used to efficiently sample pipis and
enable precise beach-scale estimates of pipi abundance and mean pipi length. The method
was developed from a recreational fishing technique that involves twisting one's feet into the
thixotropic sand to dislodge buried clams which are then recovered by hand. Several plots are
sampled across the swash zone in one 5-min sampling unit and the process is replicated at
several locations along the beach. Mean feet digging pipi counts have been shown to
correlate very strongly with quadrat-based mean transect pipi densities ($r = 0.98$) and pipi
length-frequency distributions were similar to those from quadrat-based samples except that
feet digging is not effective for pipis < 16 mm (Totterman 2019c). These young pipis are not
always present or abundant and can be ignored because individual mass increases
exponentially with shell length (Murray-Jones 1999) and large clams contribute more to
oystercatcher diets (Totterman 2018).

Variance increases strongly with mean pipi counts (Totterman 2019c). A constant precision
stopping rule was used to prevent oversampling of low pipi abundance beaches and
undersampling of high abundance beaches. A first pass sampled pipis at 20 locations,
regularly spaced along shore from a random starting distance and cycling through the tidal
stages high, ebb, low and flow (*i.e.* four tidal stages × five locations per tidal stage × 5
minutes per location). If the SE of the mean pipi count was > 2 pipis/5 min, then further
sampling passes, again with 20 locations in each pass, were performed until the SE was ≤ 2
pipis/5 mins or the cumulative sample size was 100. Stratified sampling was applied to three
long beaches, Teewah, Bennetts and Stockton (Table 1), which each had a distinct low pipi
abundance section at the southern end.

Pipi lengths were measured with a caliper to the nearest mm (maximum shell length). A
maximum sample size stopping rule was used to save time on high pipi abundance beaches.
All pipis were measured in the first sampling pass (see above). If pipi abundance was low and
the mean count from the first pass was precise (SE ≤ 2 pipis/5 mins), then mean length was
based on < 100 pipis. Otherwise, and if < 100 pipis were sampled in the first pass, pipis were
also measured in each subsequent pass until the sample size was ≥ 100, which was large
enough for a precise estimate of mean pipi length. After that, pipis were only counted.

Human recreation and other habitat variables

People, dogs, 4WDs, motorcycles, bicycles, horses and, on Fraser Island, aircraft were
counted simultaneous with oystercatchers. Dogs included dingos (naturalised dogs) that were

present in low densities on Fraser Island and in the Myall Lakes region of NSW. Counts of
people and dogs excluded those in 4WDs or in the surf. Motorcycle, bicycle, horse and
aircraft counts were small, with grand totals of three, 32, 42 and four respectively, and were
not included in the statistical analysis.

Coastal development variables and indirect measures of human recreation measured were:
urban length, length of public beach driving zones, whether or not beach driving permits were
required, length of dog exercise zones, length of beach adjoining any terrestrial protected
areas (National Parks and Nature Reserves), counts of pedestrian and, where public beach-
driving was permitted, 4WD beach access tracks and counts of beach front and near-beach
camping grounds and caravan parks. Access tracks include public and private, formal and
informal tracks. Tracks had to be frequently used and lead somewhere to be counted, *e.g.*
temporary dune driving tracks were not counted. 4WD tracks were also used by pedestrians
on many beaches but were not added to that count. There was no public beach driving for the
majority of beaches sampled and so the driving permits variable was not included in the
statistical analysis.

Human population density was calculated from the 1×1 km Australian Population Grid 2011
(Australian Bureau of Statistics 2011). The intersection of a line defined by GPS survey
waypoints with the Australian Population Grid 2011 identifies beach front grid cells and the
sum of these cells equals the 'beach front' human population.

Beach morphodynamic state (Short 2007) was observed during moderate surf conditions and
one handful of intertidal sand was sampled at each GPS waypoint. Sand grain size was
classified on the Wentworth scale by visual comparison to graded samples on a sand gauge.
Any shell fragments present were ignored. Beach morphodynamic state and sand grain size
within beaches were summarised as modal values. Sixty-two beaches sampled (86%) had
transverse bar and rip inner bars and 64 (89%) had medium sand (0.25–0.5 mm). These are
the most common conditions for ocean beaches in NSW (Short 2007). With such little
variation, these two variables were not included in the statistical analysis.

Statistical analysis

There were two response variables: mean oystercatcher count and territory count, and 19
input variables (Table 2). Input variables are all of those measured, while predictors are the
terms that are actually used in statistical modelling (Gelman & Hill 2007). Variables that
scale with beach length were converted to linear densities (*e.g.* birds/km) and proportions
(*e.g.* proportion of urban beach).

Exploratory data analysis was performed to detect patterns, multicollinearity and to reduce
the number of predictors. This involved three steps: 1) spatial patterns ('bubble maps',
correlograms); 2) bivariate associations and interactions (scatter plots), 3) variable clustering,
and; 4) variance inflation factors (to identify multicollinearity).

A spline correlogram (Bjørnstad & Falck 2001) for mean oystercatcher count density
indicated a spatial autocorrelation range of 63 km (95% CI 28–111 km). A radius of 50 km
was used for calculating binary spatial weights for spatial statistics.

Multiple regression models were fitted to oystercatcher counts because they allow for simple
and interpretable inference and the sample size was unbalanced and effectively small, with
high oystercatcher counts on only a few beaches. Repeated-measures models were not used
because mean oystercatcher counts were fairly precise, the number of counts per beach was
as small as three and estimating the variance among repeated counts was of no interest
(Murtaugh 2007).

Multicollinearity seriously confounds the interpretation of multiple regression results
(Graham 2003, Cade 2015). The first step to reducing multicollinearity was a cluster analysis
of bivariate correlations among continuous input variables. The dendrogram was cut at
Spearman $r^2 = 0.49$ ($|r| = 0.7$) and one representative variable was selected for each cluster
(Dormann *et al.* 2013). Next, variance inflation factors (VIFs) were computed for multiple
regression models with the reduced set of variables from the cluster analysis. Variables with
large VIFs were dropped backward-stepwise until all VIFs were < 3 (Zuur *et al.* 2010). To
avoid dropping useful predictors, variable selection in both of these analyses was made with
consideration to the strength of bivariate associations with oystercatcher count densities and
practical and *a priori* reasons (Dormann *et al.* 2013).

Generalised linear models (GLMs) were fitted to oystercatcher counts with a log-link and
log(beach length) as an offset, to effectively model count density. A negative binomial error
distribution was assumed for integerised mean oystercatcher counts (rounded to zero decimal
places) and a Poisson distribution was assumed for territory counts. GLM assumptions were
checked using overdispersion statistics, expected zeros (Warton 2005) mean-residual
variance plots (Ver Hoef & Boveng 2007) and diagnostic plots using Dunn-Smyth
randomised quantile residuals (Dunn & Smyth 1996).

Model selection was automated with three steps: 1) an exhaustive search of all subsets of
models, ranking them by small sample Akaike's Information Criterion (AICc); 2) selecting
the ΔAICc < 6 model set, and; 3) removal of models from the ΔAICc < 6 set that where more
complex versions of those having a lower AICc value (the 'nesting rule' of Richards 2008,
also see Arnold 2010).

Null models fitted to oystercatcher counts included the predictor mean pipi count with a
region covariate (Owner & Rohweder 2003). The pipis:region interaction was also included
in the exhaustive search of all subsets of models and marginality constraints were respected,
i.e. main effects were always included together with the interaction term. 'Nil models' for
oystercatcher counts were intercept-only models, *i.e.* with no predictor variables. The nil
model intercept is the logarithm of the grand mean oystercatcher count density.

Goodness of fit for oystercatcher count models was examined by plotting fitted versus
observed densities and calculating likelihood ratio pseudo-R^2 (Nagelkerke 1991) with

reference to nil models. Predictive performance (*e.g.* cross-validation accuracy) was not
evaluated because there were few high density sites for testing.

All GIS and statistical analyses were performed using the software R version 4.0.4 (R Core
Team 2021). Geographic analyses used the R packages *sp* version 1.4.5 (Bivand *et al.* 2013),
rgdal version 1.5-23 (Bivand *et al.* 2021) and *raster* version 3.4-5 (Hijmans 2020). Spatial
statistics used *ncf* version 1.2-9 (Bjørnstad 2020) and *spdep* version 1.1-5 (Bivand *et al.*
2013). Clustering of bivariate correlation coefficients was performed using *Hmisc* version
4.5-0 (Harrell 2021). Variance inflation factors were computed using *car* version 3.0-10 (Fox
& Weisberg 2019). Negative binomial models were fitted using *MASS* version 7.3-53
(Venables & Ripley 2002). Randomized quantile residuals were computed using *statmod*
version 1.4.35 (Dunn & Smyth 1996). Model searching, ranking and selection was performed
using the functions dredge and nested in *MuMIn* version 1.43.17 (Barton 2020). Spatial
autocorrelation in GLM residuals was checked using the lm.morantest function in *spdep*.
While this function is not intended for GLMs, it could be useful for relative comparisons
among models and there were no ready alternatives that take into account predictor variables.

RESULTS

Spatial patterns

72 of 117 beaches (62%) from the sampling frame and 674 of 942 km (72%) were sampled
including all of the longer beaches (except for Budgewoi on the Central Coast of NSW), all
of the known high pipi abundance beaches and all beach Key Management Sites for the
Australian Pied Oystercatcher (Fig. 1). The coastline was divided into five *ad hoc* contiguous
regions: SE Queensland (from Fraser Island S to the Queensland-NSW border); Far North
Coast (to Coffs Harbour); Mid-north Coast (to the Hunter River); Central Coast (to Broken
Bay), Sydney (to Port Hacking), and; South Coast (to the NSW-Victoria border). Sample
sizes were larger for regions where oystercatchers were more abundant: 20 of 28 beaches
from the Far North Coast; 27 of 30 from the Mid-north Coast, and; 18 of 37 from the South
Coast. Three of seven beaches from the sampling frame were sampled from the Central Coast
(where only one oystercatcher was counted), zero of four beaches from Sydney and four of
11 beaches from SE Queensland.

A grand total of 232 individual Australian Pied Oystercatchers (the sum of mean counts) and
41 territories were counted. Narrow count ranges (Table 1) indicate that there were no
substantial movements of birds in to and out of beaches between counts. Territory estimates
agreed reasonably well with independent breeding pair counts (Fig. 2).

The spatial pattern for mean oystercatcher count densities was strongly aggregated with a
'hot spot' in the Richmond River area of the Far North Coast (Figs. 3a,b). Mean pipi counts
showed a different spatial pattern, with a hot spot in the Mid-north Coast region (Fig. 3c).
Large pipis were available on all beaches. Mean pipi length increased from 37 mm on the
South Coast to 46 mm on the Far North Coast (Fig. 3d).

Bivariate associations

Mean oystercatcher count density was positively correlated with mean pipi count and the
proportion of protected beach (Spearman $r \geq 0.38$) and negatively correlated with the
proportion of urban beach, human population density, mean people count density, mean dog
count density, pedestrian access count density and the proportion of dogs permitted beach
(Spearman $r \leq -0.39$) (Table 3, Fig. 4).

Oystercatcher territory count density was positively correlated with mean pipi count, mean
pipi length and the proportion of protected beach (Spearman $r \geq 0.28$) and negatively
correlated with pedestrian access count density and human population density (Spearman $r \leq$
-0.30) (Table 3, Fig. 5).

Twelve beaches were sampled where commercial pipi fishing was observed or otherwise
known to occur and 60 beaches without. Mean pipi count, mean pipi length, mean
oystercatcher count density and mean oystercatcher territory density were greater for fished
beaches than for non-fished beaches (Figure 6).

Multicollinearity

Hierarchical clustering of continuous input variables indicated three groups of collinear
variables with Spearman $r^2 \geq 0.49$ (Fig. 7). The first cluster contained four coastal
development and recreation variables. Mean people and dog count densities were dropped
because those counts are imprecise, varying with the time of day, during the week, with
weather conditions, *etc*. Oystercatcher count densities were equally correlated with the
proportion of urban beach and human population density (Table 3) and those two variables
were examined separately in further analyses. The second cluster contained spatial co-
ordinates. Oystercatcher count densities were weakly correlated with UTM x and UTM y
(Spearman $r \leq 0.29$). UTM y was selected to account for possible latitudinal trends. The
third cluster contained three vehicle-based recreation variables. Oystercatcher count densities
were more strongly correlated with the proportion of 4WDs permitted beach (Spearman $r \geq$
0.22) than with either of the other two variables, which were dropped.

Next, large variance inflation factors resulted in UTM y (Spearman $r = 0.03$, VIF $= 4.7$)
being dropped from mean oystercatcher count models and UTM y (Spearman $r = 0.25$, VIF
$= 6.2$), camps and caravan parks count density (Spearman $r = -0.01$, VIF $= 5.1$), proportion
of protected beach (Spearman $r = 0.47$, VIF $= 3.1$) and proportion of dogs permitted beach
(Spearman $r = -0.39$, VIF $= 2.5$) being dropped from territory count models, in that order.
The last two variables were dropped because they contributed to multicollinearity for the
region predictor that was thought to be a necessary covariate for mean pipi count (Figs. 4a,
5a). After acting to reduce collinearity, the 19 input variables were reduced to 13 predictors
for mean oystercatcher count and 10 for territory count (Table 2).

Oystercatcher count models

Major features of the mean oystercatcher count ΔAICc < 6 model set were positive mean pipi
count (except in model seven) and Far North Coast (*region2*OTH = 0) effects and a negative
proportion of urban beach effect (or human population density in model six) (Table 4). There
was no support for the pipi abundance null model (ΔAICc = 23.6; Fig. 8a). Including the
proportion of urban beach improved the fit to lower densities (Figs. 8b,c).

Major features of the oystercatcher territory count ΔAICc < 6 model set were a positive Far
North Coast effect and a negative pedestrian access count density effect (Table 4). There was
no support for the pipi abundance null model (ΔAICc = 17.9; Fig 9a). Including pedestrian
access density improved the fit but there were some false positives (observed = 0 and fitted >
0) and underestimates from the selected models (Figs 9b,c).

DISCUSSION

The statistical analysis in this study was exploratory and descriptive because there was not
the detailed ecological knowledge for Australian Pied Oystercatchers on ocean beaches for
explanatory models (*e.g.* Goss-Custard 1996). Moreover, science can become 'stuck' in
paradigms and exploratory, hypothesis-generating studies are necessary for science to evolve
(Kuhn 1962, Smith 2000). With the usual caveat that 'association is not causation', the
oystercatcher-habitat models presented suggest some important variables for conservation
management and further research.

Oystercatcher counts in this study agreed with previous counts and estimates. The NSW
Wader Study Group counted 232 oystercatchers on NSW ocean beaches in Oct 1998 (Owner
& Rohweder 2003). There were 222 oystercatchers counted on NSW beaches in this study.
The NSW Government has estimated that there are < 200 Australian Pied Oystercatcher
breeding pairs in NSW (NSW DPIE 2019b), with 88 pairs at 15 Key Management Sites
including 24 pairs at four beach sites: South Ballina, Broadwater (which was defined for
management purposes as Airforce Beach N to Boundary Creek, at the S end of South Ballina
Beach, Table 1), Bombing Range and a number of beaches within Yuraygir National Park
(NSW DPIE 2019a). It is estimated that approximately one quarter (24/88 = 0.27) of
oystercatcher pairs are beach-residents from which it can be estimated that there are < 54
(0.27 × 200) beach-resident pairs in NSW. There were 40 oystercatcher territories counted on
NSW beaches in this study. A previous study counted 26 oystercatcher territories on beaches
between the Richmond River and Bonville Creek (8 km S of Coffs Harbour) in 2003 and 27
in 2005 (Harrison 2009). There were 27 territories counted on Far North Coast beaches (N of
Coffs Harbour) in this study. The agreement between these counts and estimates indicates
that the 68 NSW beaches sampled in this study represented almost the entire state population
of beach-resident oystercatchers.

Oystercatcher-habitat correlations were weaker than like correlations from the preceding
study that sampled 11 Far North Coast beaches, N of the Clarence River (Owner & Rohweder
2003). This disagreement was largely resolved by recomputing correlations for only Far
North Coast beaches (Table 5) however the strong oystercatcher-pipi correlation was then

defined by only two beaches: South Ballina and Bombing Range (Fig 4a). There were several
long and largely undeveloped Mid-north Coast beaches in this study with high mean pipi
counts but only moderate oystercatcher count densities. Established pipi fisheries in both of
these regions indicate that they both have a long history of high pipi abundance and, several
years after the 2003–2009 'pipi crash' (Harrison 2009, Totterman 2018), results from this
'snap shot' study are assumed to be representative of typical pipi abundances.

Automated information-theoretic model selection methods were effective at selecting small
sets of reasonable models in this study. However, the usual rule of 'garbage in, garbage out'
applies. Multicollinearity was particularly troublesome for territory count models, where
oystercatcher-habitat associations were weak. Of the 19 input variables in this study, only
four were frequent in the ΔAICc < 6 model sets for oystercatcher counts: mean pipi count,
proportion of urban beach, region and pedestrian access count density. The proportion of
urban beach and pedestrian access count density were not measured in previous
oystercatcher-habitat studies. Alternatively, mean people count density was positively
correlated with the proportion of urban beach in this study and could serve as a proxy for that
variable. Mean people count density was measured in one of the previous studies, however
the correlation with mean oystercatcher count density was weak (Owner & Rohweder 2003,
Table 4). In the other previous study, distance to towns (isolation) was suggested to be a
useful predictor for oystercatcher presence (Harrison 2009).

Mean oystercatcher count models in the ΔAICc < 6 set fitted observed count densities quite
well. These models indicated that oystercatchers aggregate at Far North Coast beaches with
high pipi abundance and that they avoid more urbanised beaches. Four of the seven models in
the mean oystercatcher count ΔAICc < 6 model set included the region predictor but not as a
covariate for mean pipi count. Model seven in that set did not include mean pipi count. These
models provide further evidence that high oystercatcher count densities on the Far North
Coast were a regional effect and, unlike what was proposed in previous studies (Owner &
Rohweder 2003, Harrison 2009), not simply a predator-prey numerical response. High
oystercatcher territory counts contributed to this Far North Coast effect and so local
population histories and recruitment are suggested to be important. Proportion of urban beach
(or human population density) substantially improved model fit for low mean oystercatcher
count density beaches. The proposed explanation is that coastal development and associated
human recreation disturbance displace shorebirds from otherwise suitable habitat (reviewed
by Hockin *et al.* 1992, also see Harrison 2009).

Oystercatcher territory count models in the ΔAICc < 6 set fitted observed count densities
rather poorly. The proposed explanation is that territorial birds are less responsive to habitat
than are non-territorial birds. For example, pipi abundance at South Ballina declined to near
zero during the 2003–2009 pipi crash but there remained eight territories and five breeding
pairs on the beach in 2009 (Totterman 2018, 2020). Nonetheless, models in the ΔAICc < 6 set
did indicate that territory density generally decreases towards zero on beaches with higher
pedestrian access count density. The proposed explanation is that higher access density can
result in shorter separation distances between access points and shorebird habitat zones,
higher intrusion rates into those zones and more frequent disturbance (Dowling & Weston

1999, Totterman 2019c). With the exception of Tallow Beach, where spot mapping suggested
one territory at Tallow Creek but where breeding is not known to occur, zero oystercatcher
territories were observed in this study when pedestrian access density exceeded 1.3 tracks/km
(Fig. 5d).

Beach access density is not only an issue for more urban beaches. Coastal tourism can also
reduce the breeding range, density and breeding success for shorebirds (reviewed by
Pienkowski 1993). More than two decades ago, it was proposed that tourism on Fraser Island
was negatively impacting beach-nesting birds (Fisher *et al.* 1998). Teewah Beach and 75
Mile Beach (Fraser Island) in this study were long, largely undeveloped (both are within
Great Sandy National Park) and had high mean pipis counts but also extensive beach front
camping zones with high count densities of pedestrian and 4WD tracks. Spot mapping
indicated only one oystercatcher territory for these two beaches with a combined length of
132 km. Moreover, mean oystercatcher counts on 75 Mile Beach have declined from 15 birds
between Hook Point and Indian Head in 1995 (Fisher *et al.* 1998) to three in this study (Table
1). Together with the negative proportion of urban beach effect for mean oystercatcher
counts, it is recommended that the coastal development and human recreation disturbance
threats for the Australian Pied Oystercatcher be upgraded.

No overall negative fishing effects for pipi stocks and oystercatcher abundance were found in
this study. Instead, mean pipi biomass and mean oystercatcher count and territory count
densities were higher on commercially fished beaches than on non-fished beaches. These
results suggest that both fishers and oystercatchers target beaches with higher pipi biomass.
The purposive sampling design in this study selected beaches where pipis and oystercatchers
could be abundant, including all known fished beaches. The sample of fished beaches is
assumed to be exhaustive and positive bias is assumed for pipi and oystercatcher means for
the non-fished beach sample (which would act to reduce, rather than increase, observed
differences). Together with uncertainty about the oystercatcher-pipi response, it is
recommended that the pipi harvesting threat for the Australian Pied Oystercatcher be
downgraded.

Coastal development and human recreation disturbance appeared to limit oystercatcher
population sizes on E Australian beaches in this study. With commercial interests for
developing coastal land and social interests for maintaining the *status quo* with regard to
access and recreation (James 2000, Priest *et al.* 2002), degradation and loss of beach-nesting
bird habitat is set to continue. Conservation efforts for Australian Pied Oystercatchers in
NSW, which have been almost exclusively focussed on breeding success (Wellman *et al.*
2000, NSW Office of Environment & Heritage 2011), will fail in the long term unless
important habitat for the species is recognised and protected from development and intense
human recreation (*e.g.* Totterman 2020).

ACKNOWLEDGEMENTS

REFERENCES

Albert, C.H., N.G. Yoccoz, T.C. Edwards Jr, C.H. Graham, N.E. Zimmermann & W. Thuiller. 2010. Sampling in ecology and evolution – bridging the gap between theory and practice. *Ecography* 33: 1028–1037. https://doi.org/10.1111/j.1600-0587.2010.06421.x

Arnold, T.W. 2010. Uninformative parameters and model selection using Akaike's Information Criterion. *Journal of Wildlife Management* 74: 1175–1178. DOI: 10.2193/2009-367

Australian Bureau of Statistics. 2011. *Australian Population Grid 2011*. Australian Government, ABS. Accessed 28 May 2017 at: https://www.abs.gov.au/ausstats/abs@.nsf/mf/1270.0.55.007

Barton, K. 2020. MuMIn: Multi-Model Inference. R package version 1.43.17. https://CRAN.R-project.org/package=MuMIn

Bivand, R.S, E. Pebesma &V. Gomez-Rubio. 2013. Applied spatial data analysis with R, Second edition. Springer, New York, U.S.A.

Bivand, R., T. Keitt & B. Rowlingson. 2021. rgdal: Bindings for the 'Geospatial' Data Abstraction Library. R package version 1.5-23. https://CRAN.R-project.org/package=rgdal

Bjørnstad, O.N & W. Falck. 2001. Nonparametric spatial covariance functions: Estimation and testing. *Environmental and Ecological Statistics* 8: 53–70.

Bjørnstad, O.N. 2020. ncf: Spatial Covariance Functions. R package version 1.2-9. https://CRAN.R-project.org/package=ncf

Burnham, K.P., D.R. Anderson & K.P. Huyvaert. 2011. AIC model selection and multimodel inference in behavioral ecology: some background, observations, and comparisons. *Behavioral Ecology and Sociobiology* 65: 23–35.

Cade, B.S. 2015. Model averaging and muddled multimodel inferences. *Ecology* 96: 2370–2382.

Dormann, C.F., J. Elith, S. Bacher, C. Buchmann , G. Carl, G. Carré, J.R. García Marquéz, B. Gruber, B. Lafourcade, P.J. Leitão, T. Münkemüller, C. McClean, P.E. Osborne, B. Reineking, B. Schröder, A.K. Skidmore, D. Zurell & S. Lautenbach. 2013. Collinearity: a review of methods to deal with it and a simulation study evaluating their performance. *Ecography* 36: 27–46. doi: 10.1111/j.1600-0587.2012.07348.x

Dowling, B. & M.A. Weston. 1999. Managing a breeding population of the Hooded Plover *Thinornis rubricollis* in a high-use recreational environment. *Bird Conservation International* 9: 255–270.

Dunn, P.K. & G.K. Smyth. 1996. Randomized quantile residuals. *Journal of Computational and Graphical Statistics* 5: 236–244.

Ferguson, G., D. Johnson & H. Gorfine. 2018. Pipi *Donax deltoides*. In: Status of Australian Fish Stocks Reports 2018 (C. Stewardson, J. Andrews, C. Ashby, M. Haddon, K. Hartmann, P. Hone, P. Horvat, S. Mayfield, A. Roelofs, K. Sainsbury, T. Saunders, J. Stewart, S. Nicol & B. Wise, Eds.). Fisheries Research and Development Corporation, Canberra, Australia. http://www.fish.gov.au/report/262-Pipi-2018

Fisher, F., M. Hockings & R. Hobson. 1998. Recreational impacts on waders on Fraser
Island. *Sunbird* 28: 1–11.

Fox, J. & S. Weisberg. 2019. An {R} Companion to Applied Regression, Third Edition.
Sage, Thousand Oaks, California, USA.

Fraser, N. & Lindsey, A. 2018. Some observations of Australian Pied Oystercatcher on
Worimi Conservation Lands. *Whistler* 12: 35–42.

Gelman, A. & J. Hill. 2007. Data Analysis Using Regression and Multilevel/Hierarchical
Models. Cambridge UniversityPress, Cambridge, UK.

Goss-Custard, J.D., Ed. 1996. *The oystercatcher: from individuals to populations.* Oxford
University Press, Oxford, UK.

Graham, M.H. 2003. Confronting multicollinearity in ecological multiple regression.
Ecology 84: 2809–2815.

Gray, C.A. 2016a. Effects of fishing and fishing closures on beach clams: experimental
evaluation across commercially fished and non-fished beaches before and during harvesting.
PLoS ONE 11: e0146122. https://doi.org/10.1371/journal.pone.0146122.

Gray, C.A. 2016b. Evaluation of fishery-dependent sampling strategies for monitoring a
small-scale beach clam fishery. *Fisheries Research* 177: 24–30.

Harrell, F.E. (with contributions from C. Dupont and many others). 2021. Hmisc: Harrell
Miscellaneous. R package version 4.5-0. https://CRAN.R-project.org/package=Hmisc

Harrison, A.E. 2009. *The ecology of two vulnerable shorebirds (Haematopus fuliginosus and*
H. longirostris) in sub-tropical northern NSW, Australia: implications for conservation and
management. PhD thesis, University of New England, Armidale, Australia.

Hijmans, R.J. 2020. raster: Geographic Data Analysis and Modeling. R package version 3.4-
5. https://CRAN.R-project.org/package=raster

Hockin, D., M. Ounsted, M. Gorman, D. Hill, V. Keller & M.A. Barker. 1992.
Examination of the effects of disturbance on birds with reference to its importance in
ecological assessments. *Journal of Environmental Management* 36: 253–286.

James, R. J. 2000. From beaches to beach environments: linking the ecology, human-use and
management of beaches in Australia. *Ocean and Coastal Management* 43: 495–514.

Kuhn, T.S. 1962. The Structure of Scientific Revolutions. First Edition. University of
Chicago Press, Chigao, USA.

Marchant S. & P.J. Higgins, Eds. 1993. *Haematopus longirostris* Pied Oystercatcher. PAges
716–726 in: *Handbook of Australian, New Zealand and Antarctic birds. Vol. 2, raptors to*
lapwings. Oxford University Press, Melbourne, Australia.

Murray-Jones, S. 1999. *Conservation and management in variable environments: the surf*
clam, Donax deltoides. PhD thesis, University of Wollongong, Australia.

Murtaugh, P.A. 2007. Simplicity and complexity in ecological data analysis. *Ecology* 88:
56–62.

Nagelkerke, N.J.D. 1991. A note on a general definition of the coefficient of determination.
Biometrika 78: 691–692.

NSW Department of Planning, Industry and Environment. 2019a. *Saving our species.*
Pied Oystercatcher (Haematopus longirostris) Key Management Sites. NSW Government,
DPIE. Accessed 26 Aug 2019 at:
https://www.environment.nsw.gov.au/savingourspeciesapp/project.aspx?ProfileID=10386

NSW Department of Planning, Industry and Environment. 2019b. Threatened species.
Pied Oystercatcher - profile. NSW Government, DPIE. Accessed 26 Aug 2019 at:
https://www.environment.nsw.gov.au/threatenedSpeciesApp/profile.aspx?id=10386

NSW Government. 2016. Biodiversity Conservation Act 2016, No. 63. Accessed 18 Feb
2019 at: https://www.legislation.nsw.gov.au/~/view/act/2016/63/

NSW Office of Environment & Heritage. 2011. NSW Threat abatement plan for predation
by the red fox (Vulpes vulpes). NSW Office of Environment and Heritage, Sydney. Accessed
13 May 2020 at:
https://www.environment.nsw.gov.au/resources/pestsweeds/110791FoxTAP2010.pdf

NSW Scientific Committee. 2010. Pied Oystercatcher *Haematopus longirostris* Vieillot
1817 - endangered species listing, final determination. Accessed 11 Nov 2018 at:
https://www.environment.nsw.gov.au/determinations/piedoystercatcherFD.htm

Owner, D. 1997. The ecology and management of the Pied Oystercatcher (*Haematopus*
longirostris) in Northern NSW. BSc thesis, Southern Cross University, Lismore, Australia.

Owner, D. & D.A. Rohweder. 2003. Distribution and habitat of Pied Oystercatchers
(*Haematopus longirostris*) inhabiting beaches in northern New South Wales. *Emu* 103: 163–
170.

Pienkowski, M.W. 1993. The impact of tourism on coastal breeding waders in western and
southern European overview. *Wader Study Group Bulletin* 68: 92–96.

Priest, B., P. Straw & M.A. Weston. 2002. Shorebird conservation in Australia. *Wingspan*
12 (Supplement): 1–15.

R Core Team. 2021. *R: A Language and Environment for Statistical Computing.* Version
4.04. Vienna: R Foundation for Statistical Computing. http://www.R-project.org

Richards, S.A. 2008. Dealing with overdispersed count data in applied ecology. *Journal of*
Applied Ecology 45: 218–227. https://doi.org/10.1111/j.1365-2664.2007.01377.x

Short, A.D. 2007. *Beaches of the New South Wales coast.* Second Edition. Sydney University
Press, Sydney, Australia.

Smith, J. 2000. Nice work — but is it science? *Nature* 408: 293.

Totterman, S.L. 2018. Response of Australian Pied Oystercatchers *Haematopus longirostris*
to increasing abundance of the beach bivalve prey *Donax deltoides*. Unpubl. preprint.
bioRxiv. https://doi.org/10.1101/444786

Totterman, S.L. 2019a. Seasonal zonation patterns of the sandy beach bivalve *Donax*
deltoides (Bivalvia: Donacidae) in subtropical eastern Australia. Unpubl. preprint. *bioRxiv*.
https://doi.org/10.1101/610576

Totterman, S. 2019b. Distances walked by beach users and protecting shorebird habitat from
human recreation disturbance. Unpubl. preprint. *bioRxi*v. https://doi.org/10.1101/783696

Totterman, S.L. 2019c. A "feet digging" swash zone sampling method for the sandy beach
bivalve *Donax deltoides* (Bivalvia: Donacidae). Unpubl. preprint. *bioRxiv*.
https://doi.org/10.1101/686196

Totterman, S. 2020. Taking fright: the decline of Australian Pied Oystercatchers
Haematopus longirostris at South Ballina Beach, New South Wales. Unpubl. prepint.
bioRxiv. https://doi.org/10.1101/2020.06.09.141796

Taylor, I.R. & S.G. Taylor. 2005. Foraging behaviour of Pied Oystercatchers in the presence
of kleptoparasitic Pacific Gulls. *Waterbirds 28*: 156–161.

Taylor, I.R., O.M.G. Newman, P. Park, B. Hansen, C.D.T. Minton & R. Jessop. 2014.
Conservation assessment of the Australian Pied Oystercatcher *Haematopus longirostris*.
International Wader Studies 20: 116–128.

Venables, W.N. & B.D. Ripley. 2002. *Modern Applied Statistics with S*. Fourth Edition.
Springer, New York, USA.

Ver Hoef, J.M. & Boveng, P.L. 2007. Quasi-poisson vs. negative binomial regression: how
should we model overdispersed count data? *Ecology* 88: 2766–2772.
https://doi.org/10.1890/07-0043.1

Warton, D.I. 2005. Many zeros does not mean zero inflation: comparing the goodness-of-fit
of parametric models to multivariate abundance data. *Environmetrics* 16: 275–289.

Wellman L., B. Moffatt, B. Totterman & N. Hing. 2000. A co-operative approach to
protecting threatened species - Pied Oystercatchers, South Ballina, Northern NSW. Pages 99–
104 in: *Proceedings of the NSW pest animal control conference, 25–27 Oct 2000* (S. Bulogh,
Ed.) . NSW Agriculture, Orange.

Weston, M.A., E.M. McLeod, D.T. Blumstein & P.-J. Guay. 2012. A review of flight
initiation distances and their application to managing disturbance to Australian birds. *Emu*
112: 269–286.

Zuur, A.F., E.N. Ieno & C.S. Elphick. 2010. A protocol for data exploration to avoid
common statistical problems. *Methods in Ecology and Evolution* 1: 3–14.
https://doi.org/10.1111/j.2041-210X.2009.00001.x

Table 1. Beaches that were sampled in this study and oystercatcher counts (beaches are ordered N to S by mid-point latitude). Regions are SE Queensland (SEQ), the Far North Coast of NSW (FNC), Mid-north Coast (MNC), Central Coast (CEC) and South Coast (SOC). Coded breeding records accompany beach territory counts: *n* pairs; habitat (Beach, Estuary, Lake); observation (C = Copulation; N = nest with eggs; Y = dependent young), and; M indicates marked nests (*e.g.* with fencing and signage). For example, 2BNM translates to 2 Beach Nests Marked. Several of these observations were of estuary/lake residents and not beach territories. Breeding pair count data sources were: 1) Harrison 2009 (br. season 2005); 2) Anon., pers. comm. 2015; 3) NSW DPIE unpubl. data (br. season 2016), and; 4) NSW DPIE unpubl. data (br. season 2018), and; 5) Fraser & Lindsey 2018 (br. seasons 2014–2017; there were only five reliable territories in that study and surely < five breeding pairs). Nest coordinates were used to assign breeding pairs from the NSW DPIE reports to beaches defined in this study. Beaches with commercial pipi fishing are indicated (Y).

Beach	Region	Sampled	S–N end	Lat. (°S)	Length (km)	Mean (range), *n*	Territories	Br. pairs	Fished
75 Mile (Fraser Island)	SEQ	Jul-16	11 km ramp (Hook Pt.) to Indian Head	25.33	84.5	3.3 (2–5), 3	1		
Teewah (Cooloola)	SEQ	Jul-17	Noosa River to Double Island Point	26.20	51.0	7 (5–9), 3	0		
Coolum	SEQ	Jul-16	Coolum Beach to Noosa Head	26.47	15.4	0, 3	0		
Warana	SEQ	Jul-16	Currimundi Lake to Point Cartwright	26.72	9.1	0, 3	0		
Kingscliff	FNC	Oct-18	Cudgen Creek to Fingal Head	28.23	7.0	0.7 (0–2), 3	0		
Bogangar	FNC	Jul-16	Cabarita Beach to Cudgen Creek	28.30	8.6	0.7 (0–2), 3	0		
Cudgera	FNC	Oct-18	Mooball Creek to Hastings Point	28.38	2.8	0, 3	0		
Wooyung	FNC	Jul-16	Brunswick River to Mooball Creek	28.46	16.6	0.7 (0–2), 3	0		
Tallow	FNC	Oct-17	Cocked Hat Rock to Cape Byron	28.68	7.4	3.7 (3–4), 3	1		
Seven Mile (Lennox Head)	FNC	Oct-17	Lennox Head to Jews Point	28.77	8.4	4 (4–4), 3	0		
Angels & Sharpes	FNC	Oct-18	Black Head to Whites Head	28.85	2.9	0, 3	0		
South Ballina	FNC	Aug-16	Broadwater Hd. to Richmond River	28.96	23.0	66 (60–78), 5	11	8[3]	Y
Airforce	FNC	Aug-15	Evans River to 'Coffee rocks'	29.09	5.8	4.8 (4–6), 5	2 (1BC)	2[3]	
Bombing Range	FNC	Aug-18	Jerusalem Creek to Goanna Headland	29.18	10.5	16.7 (13–19), 3	7	8[4]	
Ten Mile (Iluka)	FNC	Aug-16	Shark Bay to Black Rocks	29.31	12.5	8.7 (6–11), 3	1	1[3]	Y
Pippi (Yamba)	FNC	Sep-18	Green Point to Yamba Point	29.45	3.3	0.7 (0–2), 3	0		
Plumbago & Shelley (L. Arragan)	FNC	Sep-15	Red Cliff to Shelley Beach Head	29.55	4.6	2.7 (2–4), 3	1 (1BCM)	1[1]	
Sandon	FNC	Sep-17	Sandon River to Brooms Head	29.64	7.0	4.3 (3–5), 3	2 (2BNM)		
Illaroo	FNC	Aug-16	Illaroo to Sandon Bluffs	29.72	9.0	3.3 (2–6), 3	0		
Minnie Water Back	FNC	Sep-17	Diggers Camp to Minnie Water	29.79	2.8	0.7 (0–2), 3	0	0[1]	
Wooli	FNC	Sep-15	Wooli Wooli River to Wilsons Head	29.86	6.7	3.7 (2–6), 3	2 (1BY)	1[2]	
Station Creek	FNC	Sep-18	Corindi River to Station Creek	29.97	3.9	0.7 (0–2), 3	0 (1EYM)		

Hearns Lake & Woolgoolga Back	FNC	Sep-18	Sandy Beach to Woolgoolga Head	30.13	3.7	0, 3	0		
Moonee	MNC	Sep-17	Moonee Creek to Look At Me Now Hd.	30.19	4.4	0, 3	0		
Boambee	MNC	Oct-16	Boambee Creek to Corambirra Point	30.33	5.5	0.3 (0–1), 3	0	0[1]	
Bonville	MNC	Sep-18	Bundagen Headland to Bonville Creek	30.40	6.6	3 (2–4), 3	1		
North Beach (Mylestom)	MNC	Aug-16	Bellinger River to Bundagen Headland	30.47	8.4	0.7 (0–1), 3	0		
South Urunga (Hungry Head)	MNC	Sep-18	Wenonah Head to Bellinger River	30.53	4.9	0.7 (0–1), 3	0		
South Valla	MNC	Sep-15	Nambucca Heads to Deep Creek	30.62	4.8	0, 3	0		
Forster (Scotts Head)	MNC	Oct-16	Scotts Head to Nambucca River	30.70	10.8	0.7 (0–2), 3	0		
Grassy & Middle	MNC	Sep-15	Grassy Head to Scotts Head	30.78	3.4	0, 3	0		
South Smoky	MNC	Aug-17	Korogoro Creek to Smoky Cape	30.99	15.9	2 (2–2), 3	1		Y
Killick	MNC	Aug-16	Killick Creek to Hungry Hill	31.14	13.9	5.5 (4–6), 4	2		Y
Goolawah	MNC	Aug-17	Racecourse Head to Crescent Head	31.22	6.4	0, 3	0		Y
North Shore (Port Macquarie)	MNC	Oct-16	Hastings River to Queens Head	31.37	12.6	1.7 (1–3), 3	0		Y
Lighthouse (Port Macquarie)	MNC	Aug-17	Lake Cathie (open) to Tacking Point	31.51	10.1	1 (0–3), 3	0		Y
Grants (North Haven)	MNC	Aug-16	Camden Haven Inlet to Grants Head	31.62	3.1	0, 4	0		
Dunbogan	MNC	Aug-17	Diamond Head to Camden Head	31.68	8.9	1.3 (0–3), 3	0		Y
Crowdy Bay	MNC	Sep-15	Crowdy Head to Diamond Head	31.78	14.3	5.2 (4–7), 5	0		
Crowdy Head (Harrington)	MNC	Aug-17	Harrington Inlet to Crowdy Head	31.86	6.7	1.3 (0–4), 3	0		
Manning Point	MNC	Oct-16	Farquhar Inlet to Harrington Inlet	31.92	10.4	0, 3	0		
Old Bar	MNC	Sep-15	Wallabi Point (N) to Farquhar Inlet	31.98	5.4	0, 3	0		
Diamond	MNC	Sep-18	Diamond Beach to Wallabi Point (S)	32.03	5.0	0.3 (0–1), 3	0		
Nine Mile (Tuncurry)	MNC	Aug-16	Wallis Lake to Hallidays Point	32.12	12.0	7 (6–8), 3	0		
Seven Mile (Forster)	MNC	Aug-17	Booti Booti to Cape Hawke	32.28	9.6	8 (7–9), 3	0		Y
Sandbar (Smiths Lake)	MNC	Oct-16	Smiths Lake (Sandbar) to Bald Head	32.39	2.8	3.3 (2–6), 3	1 (1BN)		
Fiona & Submarine	MNC	Aug-17	Big Gibber Hd. to Yagon Headland	32.47	8.9	3.3 (2–4), 3	2 (1BC)		
Mungo (Myall Lakes)	MNC	Sep-16	Sandy Point to Big Gibber Headland	32.53	15.4	1.7 (0–5), 3	0		Y
Bennetts (Hawks Nest)	MNC	Sep-17	Yacaaba Head to Dark Point	32.64	14.3	7.7 (5–10), 3	1 (1BN)		
One Mile (Anna Bay)	MNC	Sep-16	Anna Bay	32.77	2.5	0, 3	0		
Stockton	MNC	Oct-15	Hunter River to Birubi Point	32.83	31.9	19.6 (14–24), 5	3 (1BC)	5[5]	Y
Nine Mile (Belmont)	CEC	Sep-18	Lake Entrance to Redhead Point	33.04	10.2	0.3 (0–1), 3	0		
Middle Camp	CEC	Sep-18	Catherine Hill Bay	33.15	1.7	0, 3	0		

Tuggerah	CEC	Sep-16	Tuggerah Entrance to Pelican Point	33.31	7.6	0, 3	0	
Bellambi & Woonona	SOC	Oct-15	Ocean pools to Flat Rock	34.36	2.1	0, 3	0	
Perkins (Windang)	SOC	Sep-16	Lake Illawarra to Red Point	34.52	6.6	0, 3	0	
Seven Mile (Shoalhaven)	SOC	Oct-15	Shoalhaven Hds. (open) to Crooked River	34.81	12.2	2 (1–4), 3	0	
Comerong Island	SOC	Sep-16	Crookhaven R. to Shoalhaven Hds. (open)	34.88	4.4	0.7 (0–2), 3	0 (1EN, 1ENM)	
Bherwerre	SOC	Sep-16	Sussex Inlet to Cave Head	35.17	6.8	7.6 (6–10), 5	2 (2BN)	Y
Cudmirrah	SOC	Sep-17	Cudmirrah to Sussex Inlet	35.19	3.6	0, 3	0	
Buckleys & Conjola	SOC	Sep-16	Narrawallee Inlet to Conjola Lake	35.28	4.6	1.3 (1–2), 3	0 (1ENM, 1LNM)	
Wairo	SOC	Sep-17	Tabourie Creek to Lagoon Head	35.43	4.7	1.7 (0–5), 3	0	
Durras Lake South	SOC	Sep-16	Durras to Durras Lake (open)	35.65	2.2	0, 3	1 (1LNM)	
Bengello (Moruya North)	SOC	Sep-16	Moruya River to Broulee Head	35.88	5.9	0, 3	0	
Pedro & Moruya South	SOC	Sep-17	Congo Point to Toragy Point	35.94	4.8	0, 3	1 (1ENM)	
Bingie	SOC	Oct-15	Coila Lake (open) to Bingie Bingie Point	36.03	3.8	0, 3	0	
Brou Lake	SOC	Oct-15	Lake Mummuga to Jemisons Point	36.14	6.5	5 (2–7), 3	0	
Wallaga	SOC	Sep-16	Wallaga Lake to Tilba Head	36.35	5.1	2.7 (1–5), 3	0	
Baragoot	SOC	Sep-17	Cuttagee Point to Baragoot Point	36.47	3.1	1.3 (0–2), 3	0	
Gillards & Middle	SOC	Oct-15	Cowdroys Head to Middle Beach	36.66	3.0	1 (0–2), 3	0 (1LN)	
Merimbula	SOC	Oct-15	Pambula Beach to Merimbula Lake	36.92	5.8	0.7 (0–2), 3	0	
Wonboyn	SOC	Oct-15	Greenglade to Bay Cliff	37.27	4.1	1.7 (1–2), 3	0	

582 Table 2. Variables used in the statistical analysis. 'Extensive variables' were converted to densities (*dens.*) and
583 proportions (*prop.*) by dividing by beach length. Predictors identified in the fifth column (B for mean
584 oystercatcher count models and T for oystercatcher territory count models) are those remaining after data
585 exploration and reduction (see Results).

Description	Measured	Scale	Predictor	Model
Response variables				
Oyc. mean count	*count.B*	length	*dens.B*	
Oyc. terr. count	*count.T*	length	*dens.T*	
Input variables				
Beach length (km)	*length*		*length*	B, T
Urban length (km)	*length.urban*	length	*prop.urban*	B, T
4WD length (km)	*length.4WD*	length	*prop.4WD*	B, T
Dogs length (km)	*length.dogs*	length	*prop.dogs*	B
Dogs off-leash length (km)	*length.offleash*	length	*prop.offleash*	B, T
Protected area length (km)	*length.protect*	length	*prop.protect*	B
Pedestrian access count	*access.people*	length	*dens.acc.people*	B, T
4WD access count	*access.4WD*	length	*dens.acc.4WD*	
Camping & c'van parks count	*count.camps*	length	*dens.camps*	B
People mean count	*count.people*	length	*dens.people*	
Dogs mean count	*count.dogs*	length	*dens.dogs*	
4WDs mean count	*count.4WDs*	length	*dens.4WDs*	
Human population (/km^2)	*pop1*	length×100	*dens.pop1* (/Ha)	B, T
Mean pipis count (/5 min.)	*pipis*		*pipis*	B, T
Pipi mean length (mm)	*pipis.length*		*pipis.length*	B, T
UTM x-coordinate (m)	*x*	1000	*x* (km)	
UTM y-coordinate (m)	*y*	1000	*y* (km)	
Commercial pipi fishing	*fished* (Y/N)		*fished*	B, T
Region	*region* (SEQ/FNC/ MNC/CEC/SOC)		*region2* (FNC/Other)	B, T

586 **Table 3.** Spearman rank-based correlation matrix for all continuous input variables (Table 2). Weak correlations ($|r| < 0.3$) are not presented.

	dens.B	length	prop.urban	prop.4WD	prop.dogs	prop.offleash	prop.protect	dens.access.people	dens.access.4WD	dens.camps	dens.people	dens.dogs	dens.4WD	dens.pop1	pipis	pipis.length	x	y
dens.T	0.6						0.4	−0.4								0.3		
dens.B			−0.5		−0.4		0.5	−0.5			−0.4	−0.5		−0.5	0.4			
length				0.5				−0.4	0.3				0.3		0.5	0.3		0.3
prop.urban				0.4	0.4	−0.6	0.6				0.8	0.8		0.8				
prop.4WD							−0.4	0.9				0.8			0.3	0.3		0.3
prop.dogs						0.5	−0.7	0.4			0.4	0.5		0.5				
prop.offleash							−0.6					0.4		0.3				
prop.protect								−0.6			−0.6	−0.7		−0.5				
dens.access.people									−0.3		0.7	0.5		0.6				
dens.access.4WD													0.8		0.3			
dens.camps																		
dens.people												0.8		0.7				
dens.dogs														0.7				
dens.4WD														0.3				
dens.pop1																		
pipis																		
pipis.length																	0.5	0.5
x																		1.0

Table 4. Summary of oystercatcher count generalised linear models sorted by small sample Akaike's Information Criterion (AICc). ΔAICc = 0 indicates the 'best approximating
model' of those in the model set; ΔAICc < 2 indicates strong support; ΔAICc = 2–7 indicates some support and ΔAICc > 14 indicates no support (Burnham *et al.* 2011). LR-R^2
is likelihood-ratio pseudo-R^2 (with reference to Nil models) and Moran's *I* is an index of residual spatial autocorrelation. *Ѳ* is the negative binomial overdispersion parameter
(variance = mean + mean^2/*Ѳ*; blank cells indicate a Poisson error distribution), *K* equals the number of parameters in the model (number of predictors plus intercept plus Ѳ).
Predictors are described in Table 2 (blank cells indicate that a model did not include that predictor). The *region2* predictor is a binary factor with reference Far North Coast
(OTH = other) and *region2*OTH and *pipis:region2*OTH are intercept and slope contrasts respectively. The *fished* predictor is a binary factor with reference N (Y = yes). The
parameter log(*length*) is an offset.

Predictor coefficients (and SEs)

ID	log(*length*)	intercept	pipis	Region2OTH	pipis:region2 OTH	prop.urban	dens.pop1	dens.acc.ppl	prop.offleash	prop.4WD	length	fishedY	*Ѳ*	*K*	ΔAIC	AIC weight	LR-R^2	Moran's *I*
Oystercatcher mean total count models																		
1[1]	1	−0.69 (0.36)	0.19 (0.06)	−0.57 (0.39)	−0.15 (0.06)	−3.73 (0.84)					−0.02 (0.01)		3.5	7	0.0	0.44	0.54	0.10
2	1	−0.75 (0.38)	0.17 (0.07)	−0.69 (0.41)	−0.13 (0.07)	−3.80 (0.89)							2.7	6	1.9	0.17	0.51	0.08
3	1	−0.13 (0.24)	0.05 (0.02)	−1.32 (0.25)		−3.95 (0.87)					−0.02 (0.01)	0.45 (0.26)	3.1	7	2.5	0.13	0.52	0.14
4	1	−0.03 (0.25)	0.06 (0.02)	−1.33 (0.26)		−4.28 (0.89)					−0.02 (0.01)		2.6	6	2.7	0.11	0.50	0.14
5	1	−0.16 (0.25)	0.05 (0.02)	−1.33 (0.27)		−4.24 (0.92)							2.3	5	2.9	0.10	0.48	0.11
6	1	−0.78 (0.36)	0.21 (0.06)	−0.40 (0.4)	−0.16 (0.06)		−0.44 (0.13)	−0.87 (0.45)			−0.02 (0.01)		3.6	8	5.1	0.034	0.52	0.09
7	1	−0.06 (0.25)		−1.20 (0.26)		−4.23 (0.91)						0.51 (0.28)	2.2	5	5.8	0.025	0.46	0.18
Null	1	−1.69 (0.37)	0.25 (0.08)	−0.23 (0.43)	−0.20 (0.08)								1.8	5	23.6		0.31	0.09
Nil	1	−1.13 (0.15)											0.8	2	42.7			0.30

[1]Model 1 prediction example:
FNC count = exp{log(*length*) − 0.69 + 0.19×*pipis* − 3.73×*prop.urban* − 0.02×*length*} = *length* × exp(−0.69) × exp(0.19×*pipis*) × …
OTH count = exp{log(*length*) + (−0.69−0.57) + (0.19−0.15)×*pipis* − 3.73×*prop.urban* − 0.02×*length*} = *length* × exp(−1.26) × exp(0.04×*pipis*) × …

ID	log(*length*)	intercept	pipis	Region2OTH	pipis:region2 OTH	prop.urban	dens.pop1	dens.acc.ppl	prop.offleash	prop.4WD	length	fishedY	*Ѳ*	*K*	ΔAIC	AIC weight	LR-R^2	Moran's *I*
Oystercatcher territory count models																		
1	1	−1.13 (0.30)	0.06 (0.02)	−2.36 (0.38)					−0.85 (0.26)					4	0.0	0.78	0.43	0.05
2	1	−0.57		−1.98					−0.86	−0.87		0.80		5	3.6	0.13	0.42	0.07

		(0.34)		(0.33)		(0.24)	(0.45)	(0.40)						
3	1	−0.71 (0.25)		−1.98 (0.33)		−0.93 (0.26)				3	4.4	0.087	0.36	0.13
Null	1	−2.91 (0.58)	0.22 (0.11)	−1.26 (0.77)	−0.16 (0.11)				1.1	5	17.9		0.24	0.09
Nil	1	−2.81 (0.28)							0.3	2	27.1			0.34

594 **Table 5.** Mean oystercatcher count density Pearson correlations compared with those from the preceding study that
595 sampled 11 Far North Coast beaches, N of the Clarence River (Owner 1997, for birds/km and Owner & Rohweder 2003,
596 for birds/km^2). Far North Coast (FNC) beaches are those between Coffs Harbour and the NSW-Queensland border (Table
597 1, Fig. 1).

Mean oystercatcher count density ~	Owner (1997) r	Owner & Rohweder (2003) r	This study – all beaches r	This study – NSW FNC r
Pipi mean density/count	0.78	0.83	0.26	0.76
Mean pipi length	0.70	0.66	0.22	0.44
Beach length	0.55	0.64	0.09	0.72
4WD mean count density	0.39	0.24	−0.02	0.04
People mean count density	−0.17	−0.03	−0.15	−0.23

599
Fig. 1. Map of the SE Australian coast showing the location of beaches (points plotted at beach centres) that were sampled in this study (Table 1). Colour-fill identifies regions: SE Queensland (magenta), Far North Coast (blue), Mid-north Coast (green), Central Coast (yellow) and South Coast (red). The background image shows human population densities (scale 0–1000 persons/km^2) from the Australian Population Grid 2011 (Australian Bureau of Statistics 2011). Text annotations identify states and larger coastal cities, towns and populated regions.

605
606 **Fig. 2.** Oystercatcher territory estimates compared to breeding pairs (Table 1). Point sizes are proportional to the number
607 of observations. A statistical comparison is not useful because the sample size (nine beaches) was small and there is
608 insufficient power to detect small deviations from the 1:1 line of agreement (dotted line).

Fig. 3. Oystercatcher and pipi spatial patterns along the E Australian coast. Point sizes are proportional to abundance/size and colour identifies regions: SE Queensland (magenta), Far North Coast (blue), Mid-north Coast (green), Central Coast (yellow) and South Coast (red). Note that mean pipi length is offset by 20 mm. Getis-Ord G^* and Moran's I statistics measure spatial concentration and similarity (autocorrelation) respectively for which all P-values were < 0.001.

Fig. 4. Mean oystercatcher count density correlations. Variables are described in Table 2 and Spearman rank-based correlations are in Table 3. Far North Coast beaches (Coffs Harbour to the NSW-Queensland border) are plotted as blue-filled circles and other regions are plotted as white-filled circles in (a). Oystercatcher-pipi lines in were fitted by ordinary least squares linear regression with *region2* (Far North Coast/Other) as a covariate, however the reported correlation statistic does not include that covariate. These notes also apply to Fig. 5.

Fig. 5. Oystercatcher territory density correlations.

619 **Fig. 6**. Pipi and oystercatcher contrasts for beaches with (squares) and without (circles) commercial pipi fishing. Individual
620 results are plotted with a small amount of random horizontal 'jitter'. Black points are means and error bars are 95%
621 confidence intervals from generalised linear models (a, c, d) and an unequal variance *t*-test with Welch's approximate
622 degrees of freedom (b). Contrasts (mean ratios/difference) are from the same models and *t*-test.

624
Fig. 7. Hierarchical clustering of continuous input variables using squared Spearman rank-based correlations (Table 3). Variables are described in Table 2. The horizontal line cuts the dendrogram at Spearman $r^2 = 0.49$ and indicates three clusters of collinear variables below the line: 1) *dens.people, dens.dogs, prop.urban, dens.pop1*; 2) *x, y,* and; 3) *dens.4WDs, prop.4WD, dens.acc.4WD*.

629 **Fig. 8.** Fitted versus observed mean oystercatcher count density for the null (a) minimum AICc (b) and minimum *K*
630 (simplest, model five) (c) models in Table 4. Far North Coast beaches (Coffs Harbour to the NSW-Queensland border) are
631 plotted as blue-filled circles and other regions are plotted as white-filled circles.

632 **Fig. 9.** Fitted versus observed oystercatcher territory density for the null (a) minimum AICc (b) and minimum *K* (simplest,
633 model four) (c) models in Table 4.