Type I interferon susceptibility distinguishes SARS-CoV-2 from SARS-CoV.

Kumari G. Lokugamage[1*], Adam Hage[1*], Maren de Vries[3], Ana M. Valero-Jimenez[3], Craig Schindewolf[1], Meike Dittmann[3], Ricardo Rajsbaum[1,2], Vineet D. Menachery[1,2]

[1]Department of Microbiology and Immunology, [2]Institute for Human Infection and Immunity, University of Texas Medical Branch, Galveston TX, USA

[3]Department of Microbiology, New York University School of Medicine, New York, NY 10016, USA

*Equal contributions

Corresponding Author: Vineet D. Menachery

Address: University of Texas Medical Branch, 301 University Blvd, Route #0610 Galveston, TX 77555

Email: Vimenach@utmb.edu

Article Summary: SARS-CoV-2 has similar replication kinetics to SARS-CoV, but demonstrates significant sensitivity to type I interferon treatment.

Running title: SARS-CoV-2 sensitive to type I IFN pretreatment

Keywords: Coronavirus, 2019-nCoV, SARS-CoV-2, COVID-19, SARS-CoV, type I interferon, IFN

Abstract

SARS-CoV-2, a novel coronavirus (CoV) that causes COVID-19, has recently emerged causing

an ongoing outbreak of viral pneumonia around the world. While distinct from SARS-CoV, both

group 2B CoVs share similar genome organization, origins to bat CoVs, and an arsenal of

immune antagonists. In this report, we evaluate type-I interferon (IFN-I) sensitivity of SARS-

CoV-2 relative to the original SARS-CoV. Our results indicate that while SARS-CoV-2

maintains similar viral replication to SARS-CoV, the novel CoV is much more sensitive to IFN-I.

In Vero and in Calu3 cells, SARS-CoV-2 is substantially attenuated in the context of IFN-I

pretreatment, while SARS-CoV is not. In line with these findings, SARS-CoV-2 fails to

counteract phosphorylation of STAT1 and expression of ISG proteins, while SARS-CoV is able

to suppress both. Comparing SARS-CoV-2 and influenza A virus in human airway epithelial

cultures (HAEC), we observe the absence of IFN-I stimulation by SARS-CoV-2 alone, but detect

failure to counteract STAT1 phosphorylation upon IFN-I pretreatment resulting in near ablation

of SARS-CoV-2 infection. Next, we evaluated IFN-I treatment post infection and found SARS-

CoV-2 was sensitive even after establishing infection. Finally, we examined homology between

SARS-CoV and SARS-CoV-2 in viral proteins shown to be interferon antagonists. The absence

of an equivalent open reading frame (ORF) 3b and changes to ORF6 suggest the two key IFN-I

antagonists may not maintain equivalent function in SARS-CoV-2. Together, the results identify

key differences in susceptibility to IFN-I responses between SARS-CoV and SARS-CoV-2 that

may help inform disease progression, treatment options, and animal model development.

Importance

With the ongoing outbreak of COVID-19, differences between SARS-CoV-2 and the original

SARS-CoV could be leveraged to inform disease progression and eventual treatment options.

In addition, these findings could have key implications for animal model development as well as

further research into how SARS-CoV-2 modulates the type I IFN response early during

infection.

Introduction

At the end of 2019, a cluster of patients in Hubei Province, China was diagnosed with a viral pneumonia of unknown origins. With community links to the Huanan seafood market in Wuhan, the disease cluster had echoes of the severe acute respiratory syndrome coronavirus (SARS-CoV) outbreak that emerged at the beginning of the century (1). The 2019 etiologic agent was identified as a novel coronavirus, 2019-nCoV, and subsequently renamed SARS-CoV-2 (2). The new virus has nearly 80% nucleotide identity to the original SARS-CoV and the corresponding CoV disease, COVID-19, has many of the hallmarks of SARS-CoV disease including fever, breathing difficulty, bilateral lung infiltration, and death in the most extreme cases (3, 4). In addition, the most severe SARS-CoV-2 disease corresponded to old age (>50 years old), health status, and healthcare workers, similar to both SARS- and MERS-CoV (5). Together, the results indicate SARS-CoV-2 infection and disease have strong similarity to the original SARS-CoV epidemic occurring nearly two decades earlier.

In the wake of the outbreak, major research efforts have sought to rapidly characterize the novel CoV to aid in treatment and control. Initial modeling studies predicted (6) and subsequent cell culture studies confirmed that spike protein of SARS-CoV-2 utilizes human angiotensin converting enzyme 2 (ACE2) for entry, the same receptor as SARS-CoV (7, 8). Extensive case studies indicated a similar range of disease onset and severe symptoms seen with SARS-CoV (5). Notably, less severe SARS-CoV-2 cases have also been observed and were not captured in the original SARS-CoV outbreak. Importantly, screening and treatment guidance has relied on previous CoV data generated with SARS-CoV and MERS-CoV. Treatments with both protease inhibitors and type-I interferon (IFN-I) have been employed (4); similarly, remdesivir, a drug targeting viral polymerases, has been reported to have efficacy against SARS-CoV-2 similar to findings with both SARS- and MERS-CoV (9-12). Importantly, several vaccine efforts have been initiated with a focus on the SARS-CoV-2 spike protein as the

major antigenic determinant (13). Together, the similarities with SARS-CoV have been useful in

responding to the newest CoV outbreak.

The host innate immune response is initiated when viral products are recognized by host

cell pattern recognition receptors, including Toll-like receptors (TLRs) and RIG-I-like receptors

(RLRs) (14, 15). This response ultimately results in production of IFN-I and other cytokines,

which together are essential for an effective antiviral response (16). IFN-I then triggers its own

signaling cascade via its receptor, in an autocrine or paracrine manner, which induces

phosphorylation of signal transducers and activators of transcription 1 (STAT1) and STAT2.

Together, STAT1, STAT2, and a third transcription factor, IRF9, form the Interferon Stimulated

Gene Factor 3 (ISGF3) complex, which is essential for induction of many IFN-stimulated genes

(ISGs), and ultimately elicit an effective antiviral response (17, 18). To establish productive

replication, viruses have developed different mechanisms to escape this antiviral response

targeting different parts of the IFN-I response machinery (19).

In this study, we further characterize SARS-CoV-2 and compare it to the original SARS-

CoV. Using Vero E6 cells, we demonstrate that SARS-CoV-2 maintains similar viral replication

kinetics as SARS-CoV following a low dose infection. In contrast, we find that SARS-CoV-2 is

significantly more sensitive to IFN-I pretreatment as compared to SARS-CoV. Infection of IFN-I

competent Calu3 2B4 cells resulted in reduced SARS-CoV-2 replication compared to SARS-

CoV. Similar to Vero cells, Calu3 cells pretreated with IFN-I had a greater reduction of

replication of SARS-CoV-2 compared to SARS-CoV. In human airway epithelial cultures, SARS-

CoV-2 showed robust replication and an absence of IFN-I stimulation contrasting influenza A

virus. However, pretreatment with IFN-I confirmed SARS-CoV-2 sensitivity and inability to

control IFN-I responses once initiated. These results suggest distinct changes between SARS-

CoV and SARS-CoV-2 in terms of IFN-I antagonism and we subsequently examined sequence

homology between the SARS-CoV and SARS-CoV-2 viral proteins that may be responsible for

these differences. Together, the results suggest SARS-CoV-2 lacks the same capacity to

control the IFN-I response as SARS-CoV.

Results

SARS-CoV-2 is sensitive to IFN-I pre-treatment

Our initial studies infected Vero E6 cells using a low multiplicity of infection (MOI) to

explore the viral replication kinetics of SARS-CoV-2 relative to SARS-CoV. Following infection,

we found that both SARS-CoV and SARS-CoV-2 replicate with similar kinetics, peaking 48

hours post infection (**Fig. 1A**). While SARS-CoV-2 titer was slightly lower than that of SARS-

CoV at 24 hours post infection, the results were not statistically different. By 48 hours,

replication of both viruses had plateaued and significant cytopathic effect (CPE) was observed

for both SARS-CoV and SARS-CoV-2 infections. Together, the results indicated that SARS-CoV

and SARS-CoV-2 replicate with similar replication kinetics in Vero E6 cells.

We next evaluated the susceptibility of SARS-CoV-2 to IFN-I pretreatment. Treatment

with IFN-I (recombinant IFN-α) has been attempted as an antiviral approach for a wide variety of

pathogens including hepatitis B and C viruses as well as HIV (20). During both the SARS and

MERS-CoV outbreaks, IFN-I has been employed with limited effect (21, 22). In this study, we

pretreated Vero E6 cells with 1000 U/mL of recombinant IFN-I (IFN-α) 18 hours prior to

infection. Vero E6 lack the capacity to produce IFN-I, but are able to respond to exogenous

treatment (23). Following pretreatment with IFN-I, SARS-CoV infection has a modest reduction

in viral titer of 1.5 $\log_{10}$ plaque forming units (PFU) as compared to untreated control 24 hours

post infection (**Fig. 1A**). However, by 48 hours, SARS-CoV has nearly equivalent viral yields as

the untreated conditions (7.2 $\log_{10}$PFU versus 7.5 $\log_{10}$PFU). In contrast, SARS-CoV-2 shows a

significant reduction in viral replication following IFN-I treatment. At both 24 and 48 hours post

infection, SARS-CoV-2 had massive 3-$\log_{10}$ (24 HPI) and 4-$\log_{10}$ (48 HPI) drops in viral titer as

compared to control untreated cells. Together, the results demonstrate a clear sensitivity to a

primed IFN-I response in SARS-CoV-2, which is not observed with SARS-CoV.

To explore differences in IFN-I antagonism between SARS-CoV and SARS-CoV-2, we

examined both STAT1 activation and IFN stimulated gene (ISG) expression following IFN-I

pretreatment and infection. Examining Vero cell protein lysates, we found that IFN-I treated cells

infected with SARS-CoV-2 induced phosphorylated STAT1 by 48 hours post infection (**Fig. 1B**).

SARS-CoV had no evidence of STAT1 phosphorylation in either IFN-I treated or untreated cells,

illustrating robust control over IFN-I signaling pathways. In contrast, SARS-CoV-2 is unable to

control signaling upon IFN-I treatment. Examining further, IFITM1, a known ISG (17), had

increased protein expression in the context of SARS-CoV-2 infection following IFN-I

pretreatment compared to SARS-CoV under the same conditions (**Fig. 1B**). Basal STAT1 levels

are reduced during SARS-CoV infection relative to uninfected control and to a lesser extent

during SARS-CoV-2, likely due to the mRNA targeting activity of non-structural protein 1 (NSP1)

(24). However, IFN-I treatment results in augmented protein levels for IFITM1 following SARS-

CoV-2 infection as compared to untreated SARS-CoV-2. In contrast, IFN-I treated SARS-CoV

had no significant increase in IFITM1 relative to control infection. Together, the STAT1

phosphorylation, ISG production, and viral protein levels indicate that SARS-CoV-2 lacks the

same capacity to modulate the IFN-I stimulated response as the original SARS-CoV.

SARS-CoV-2 attenuated in interferon competent cells.

While capable of responding to exogenous IFN-I, Vero cells lack the capacity to produce

IFN-I following infection which likely plays a role in supporting robust replication of a wide range

of viruses. To evaluate SARS-CoV-2 in a IFN-I responsive cell type, we infected Calu3 2B4

cells, a lung epithelial cell line sorted for ACE2 expression and previously used in coronavirus

and influenza research (25). Using an MOI of 1, we examined the viral replication kinetics of

SARS-CoV-2 relative to SARS-CoV in Calu3 cells. We found that both SARS-CoV and SARS-

CoV-2 replicate with similar overall kinetics, peaking 24 hours post infection (**Fig. 2A**). However,

SARS-CoV-2 replication is slightly attenuated relative to SARS-CoV at 24 hours post infection

(0.82 $\log_{10}$ reduction).   The attenuation in viral replication expands at 48 hours (1.4 $\log_{10}$

reduction) indicating a significant change in total viral titers between SARS-CoV and SARS-

CoV-2. Notably, no similar attenuation was observed in untreated Vero cells (**Fig. 1A**)

suggesting possible immune modulation of SARS-CoV-2 infection in the respiratory cell line,

due to differential sensitivity to secreted IFN-I during infection. We next evaluated the

susceptibility of SARS-CoV-2 to IFN-I pretreatment in Calu3 cells. When pretreating cells with

1000 U/mL of recombinant IFN-I 18 hours prior to infection, SARS-CoV infection has a modest

reduction in viral titer of ~0.8 $\log_{10}$ PFU as compared to untreated control at both 24 and 48

hours post infection (**Fig. 2A**). Similar to Vero cell results, SARS-CoV-2 shows a significant

reduction in viral replication following IFN-I treatment in Calu3 cells. At both 24 and 48 hours

post infection, SARS-CoV-2 had 2.65- $\log_{10}$ (24 HPI) and 2- $\log_{10}$ (48 HPI) drops in viral titer,

respectively, as compared to control untreated Calu3 cells. Together, the results demonstrate a

clear sensitivity to a primed IFN-I response in SARS-CoV-2, which is not observed with SARS-

CoV.

To further evaluate activation by IFN-I, we examined both STAT1 phosphorylation and

ISG expression following infection of Calu3 2B4 cells at 48 hours. Probing cell protein lysates,

we found that untreated cells infected with SARS-CoV or SARS-CoV-2 induced phosphorylated

STAT1 by 48 hours post infection (**Fig. 2B**). However, the level of STAT1 is markedly

diminished in SARS-CoV infection as compared to SARS-CoV-2 suggesting the original

epidemic strain disrupts the expression of the ISG. The diminished STAT1 levels in SARS-CoV

correspond to robust spike expression in untreated cells. In contrast, SARS-CoV-2 spike is

reduced as compared to SARS-CoV, consistent with lower replication observed in untreated

Calu3 cells (**Fig. 2A**). The noticeable IFITM1 expression levels observed in untreated Calu3

cells infected with SARS-CoV-2 that were not observed in Vero cells may signify higher ISG

levels in Calu3 cells during SARS-CoV-2 infection and may contribute to its attenuation relative

to SARS-CoV.

Following IFN-I pretreatment in Calu3 cells the differences in STAT1 activation and ISG

induction are less prominent compared the Vero experiments, likely due to an already induced

IFN-I response to the virus infection. However, we still found a reduction in STAT1

phosphorylation and ISG induction in IFN-I pretreated SARS-CoV versus SARS-CoV-2

infection. Total STAT1 levels were augmented in IFN-I treated cells following SARS-CoV

infection to values similar to control IFN-I treated cells. In contrast, SARS-CoV-2 STAT1 levels

remained amplified relative to SARS-CoV, but were similar to untreated SARS-CoV-2 infection.

Although marginal, IFITM1 expression was slightly increased in IFN-I pretreatment SARS-CoV-

2 relative to SARS-CoV. Importantly, the spike protein levels show a significant impact of IFN-I

on SARS-CoV infection consistent with titer. For SARS-CoV-2, the spike blot demonstrates the

massive attenuation in the presence of IFN-I. Overall, the results in Calu3 cells are consistent

with Vero cell findings and indicate a significant sensitivity of SARS-CoV-2 to IFN-I

pretreatment.

SARS-CoV-2 blocks IFN-I signaling in polarized human airway epithelial cultures.

Having established IFN-I sensitivity in Vero cells and Calu3 respiratory cell lines, we next sought

to evaluate SARS-CoV-2 in polarized human airway epithelial cultures (HAEC). These cultures

provide both the complexity of different cell types and the architecture of the human respiratory

epithelium. Previous work with SARS-CoV had already established its capacity to control the

IFN-I response in human airway cultures (26). Therefore, to examine the impact on overall level

of IFN-I signaling in HAECs, we compared SARS-CoV-2 infection to influenza A virus (H1N1)

infection, known to induce robust innate immune responses in these cultures. Briefly, HAEC

were pretreated with IFN-α on the basolateral side prior to and during infection (**Fig. 3A**). Apical

washes were collected at multiple time points and analyzed for viral infectious titers by plaque

assay. In addition, culture lysates at endpoint were examined for STAT1 phosphorylation by

western blot. Both influenza A and SARS-CoV-2 triggered different levels of immune stimulation

(**Fig. 3B**). Influenza A virus infection alone induced robust expression of both total and

phosphorylated STAT1 48 hours post infection. In stark contrast, SARS-CoV-2 infection alone

indicated no increase in STAT1 or phosphorylated STAT1. IFN-I pretreatment resulted in robust

induction of STAT1 and pSTAT1 in all cultures. Despite the distinct immune stimulation profiles

upon infection alone, both viruses robustly replicated in the HAEC, with influenza A virus

achieving higher viral yields as compared to SARS-CoV-2 (**Fig. 3C & 3D**). Pretreatment

reduced influenza infection ~1.5 $\log_{10}$ at both 24 and 48 hours post infection, consistent with

previous reports (27). In contrast, IFN-I pretreatment nearly ablated SARS-CoV-2 with 2-$\log_{10}$

and 4-$\log_{10}$ reduction in viral titers at 24 and 48 hours relative to untreated controls. Together,

the results highlight the capacity of SARS-CoV-2 to prevent IFN-I signaling, but also

demonstrate the sensitivity of SARS-CoV-2 to IFN-I pretreatment.

SARS-CoV-2 impacted by IFN-I post treatment. Having established that SARS-CoV-2 cannot

overcome a pre-induced IFN-I state, we next evaluated the impact of IFN-I treatment post

infection. Vero cells were infected with SARS-CoV or SARS-CoV-2 at an MOI 0.01 and

subsequently treated with 1000 U/mL IFN-α four hours post infection. For SARS-CoV, treatment

post infection had no significant impact on viral yields either 24 or 48 hours post infection,

consistent with findings from pretreatment experiments. In contrast, SARS-CoV-2 had a

substantial 2-$\log_{10}$ reduction in viral titers at 24 hours post infection relative to control. However,

by 48 hours, SARS-CoV-2 replication had achieved similar level to untreated controls indicating

that post treatment had only a transient impact in Vero cells. We subsequently performed the

post-treatment experiment utilizing Calu3 respiratory cells at an MOI 1. Similar to Vero cells,

SARS-CoV infection post treatment resulted in no significant changes at 24 hours and has a

modest decrease (~0.5 $\log_{10}$) at 48 hours, illustrating its resistance to IFN-I. In contrast, IFN-I

post-treatment of SARS-CoV-2 infection resulted in a >2-$\log_{10}$ reduction in viral titers at 24 hours

and expanded at 48 hours post (~4-$\log_{10}$ reduction). The results indicate that in Calu3 cells,

SARS-CoV-2 is unable to prevent inhibitory effects of IFN-I even after establishing initial

infection. Overall, the pre- and post-treatment data highlight distinct differences between

SARS-CoV and SARS-CoV-2 in modulation of IFN-I pathways.

Conservation of IFN-I antagonists across SARS-CoV and SARS-CoV-2

Previous work has established several key IFN-I antagonists in the SARS-CoV genome,

including NSP1, NSP3, ORF3b, ORF6, and others (28). Considering SARS-CoV-2's sensitivity

to IFN-I, we next sought to evaluate conservation of IFN-I antagonist proteins encoded by

SARS-CoV-2, SARS-CoV, and several bat SARS-like viruses including WIV16-CoV (29),

SHC014-CoV (30), and HKU3.1-CoV (31). Using sequence analysis, we found several changes

to SARS-CoV-2 that potentially contribute to IFN-I sensitivity (**Fig. 4**). For SARS-CoV structural

proteins, including the nucleocapsid (N) and matrix (M) protein, a high degree of sequence

homology (>90%AA identity) suggests that their reported IFN-I antagonism is likely maintained

in SARS-CoV-2 and other SARS-like viruses. Similarly, the ORF1ab poly-protein retains high

sequence identity in SARS-CoV-2 and several known IFN-I antagonists contained within the

poly-protein (NSP1, NSP7, NSP14-16) are highly conserved relative to SARS-CoV. One notable

exception is the large papain-like protease, NSP3, which is only 76% conserved between

SARS-CoV and SARS-CoV-2. However, SARS-CoV-2 does maintain a deubiquitinating domain

thought to confer IFN-I resistance (32). For SARS-CoV ORF3b, a 154 amino acid (AA) protein

known to antagonize IFN-I responses by blocking IRF3 phosphorylation (33), sequence

alignment indicates that the SARS-CoV-2 equivalent ORF3b contains a premature stop codon

resulting in a truncated 24 AA protein. Similarly, HKU3.1-CoV also has a premature termination

resulting in a predicted 39 AA protein. Both WIV16-CoV and SHC014-CoV, the most closely

related bat viruses to SARS-CoV, encode longer 114 AA truncated protein with >99% homology

with SARS-CoV ORF3b suggesting that IFN-I antagonism might be maintained in these specific

group 2B CoV strains. In addition, SARS-CoV ORF6 has been shown to be an IFN-I antagonist

that disrupts karyopherin-mediated transportation of transcription factors like STAT1 (33, 34). In

contrast to ORF3b, all five surveyed group 2B CoVs maintain ORF6; however, SARS-CoV-2

had only 69% homology with SARS-CoV while the other three group 2B bat CoVs had >90%

conservation. Importantly, SARS-CoV-2 has a two amino acid truncation in its ORF6; previous

work has found that alanine substitution in this C-terminus of SARS-CoV ORF6 resulted in

ablated antagonism (34). Together, the sequence homology analysis suggests that differences

in NSP3, ORF3b, and/or ORF6 may be key drivers of SARS-CoV-2 IFN-I susceptibility.

Discussion

With the ongoing outbreak of COVID-19 caused by SARS-CoV-2, viral characterization remains

a key factor in responding to the emergent novel virus. In this report, we describe differences in

the IFN-I sensitivity between SARS-CoV-2 and the original SARS-CoV. While both viruses

maintain similar replication in untreated Vero E6 cells, SARS-CoV-2 has a significant decrease

in viral protein and replication following IFN-I pretreatment. The decreased SARS-CoV-2

replication correlates with phosphorylation of STAT1 and augmented ISG expression largely

absent following SARS-CoV infection despite IFN-I pretreatment. Infection of IFN-I competent

Calu3 2B4 cells resulted in reduced SARS-CoV-2 replication relative to SARS-CoV; IFN-I

pretreatment also corresponded to increased sensitivity for SARS-CoV-2 in Calu3 cells.

However, SARS-CoV-2 fails to induce IFN-I pathways during infection of unprimed polarized

HAEC as compared to influenza A virus, suggesting a capacity to control IFN-I signaling. Yet,

pretreatment nearly ablates replication of SARS-CoV-2 highlighting robust sensitivity to IFN-I

pathways once activated. Finally, SARS-CoV-2, unlike SARS-CoV, responded to post-infection

IFN-I treatment with reduced viral yields in both Vero and Calu3 cells. Analysis of viral proteins

finds SARS-CoV-2 has several changes that potentially impact its capacity to modulate the IFN-

I response, including loss of an equivalent ORF3b and a short truncation of ORF6, both known

as IFN-I antagonists for SARS-CoV (33). Together, our results suggest SARS-CoV and SARS-

CoV-2 have differences in their ability to antagonize the IFN-I response once initiated and that

this may have major implications for COVID-19 disease and treatment.

With a similar genome organization and disease symptoms in humans, the SARS-CoV-2

outbreak has drawn insights from the closely related SARS-CoV. However, the differences in

sensitivity to IFN-I pretreatment illustrate a clear distinction between the two CoVs. Coupled with

a novel furin cleavage site (35), robust upper airway infection (8), and transmission prior to

symptomatic disease (36), the differences between SARS-CoV and SARS-CoV-2 could prove

important in disrupting the ongoing spread of COVID-19. For SARS-CoV, *in vitro* studies have

consistently found that wild-type SARS-CoV is indifferent to IFN-I pretreatment (37, 38).

Similarly, *in vivo* SARS-CoV studies have found that the loss of IFN-I signaling had no

significant impact on disease (39), suggesting that this virus is not sensitive to the antiviral

effects of IFN-I. However, more recent reports suggest that host genetic background may

majorly influence this finding (40). For SARS-CoV-2, our results suggest that IFN-I pretreatment

produces a 3 - 4 $\log_{10}$ drop in viral titer. This level of sensitivity is similar to MERS-CoV and

suggests that the novel CoV lacks the same capacity to escape a primed IFN-I response as

SARS-CoV (41, 42). Notably, the sensitivity to IFN-I does not completely ablate viral replication;

unlike SARS-CoV 2'O methyl-transferase mutants (37), SARS-CoV-2 is able to replicate to low,

detectable levels even in the presence of IFN-I. This finding could help explain positive test

results in patients with minimal symptoms and the range of disease observed. In addition, while

SARS-CoV-2 is sensitive to IFN-I pretreatment, both SARS-CoV and MERS-CoV employ

effective means to disrupt virus recognition and downstream signaling until late during infection

(25). While SARS-CoV-2 may employ a similar mechanism early during infection, STAT1

phosphorylation and reduced viral replication are observed in IFN-I primed and post-treatment

conditions indicating that the novel CoV does not block IFN-I signaling as effectively as the

original SARS-CoV.

For SARS-CoV-2, the sensitivity to IFN-I indicates a distinction from SARS-CoV and

suggests differential host innate immune modulation between the viruses. The distinct ORF3b

and truncation/changes in ORF6 could signal a reduced capacity of SARS-CoV-2 to interfere

with IFN-I responses. For SARS-CoV ORF6, the N-terminal domain has been shown to have a

clear role in its ability to disrupt karyopherin-mediated STAT1 transport (34); in turn, the loss or

reduction of ORF6 function for SARS-CoV-2 would likely render it much more susceptible to

IFN-I pretreatment as activated STAT1 has the capacity to enter the nucleus and induce ISGs

and the antiviral response. In these studies, we have found that following IFN-I pretreatment,

STAT1 phosphorylation is induced following SARS-CoV-2 infection. The increase in ISG

proteins (STAT1, IFITM1) suggests that SARS-CoV-2 ORF6 does not effectively block nuclear

transport as well as SARS-CoV ORF6. For SARS-CoV ORF3b, the viral protein has been

shown to disrupt phosphorylation of IRF3, a key transcriptional factor in the production of IFN-I

and the antiviral state (33). While its mechanism of action is not clear, the ORF3b absence in

SARS-CoV-2 infection likely impacts its ability to inhibit the IFN-I response and eventual STAT1

activation. Similarly, while NSP3 deubiquitinating domain remains intact, SARS-CoV-2 has a 24

AA insertion upstream of this deubiquitinating domain that could potentially alter that function

(32). While other antagonists are maintained with high levels of conservation (>90%), single

point mutations in key locations could modify function and contribute to increased IFN-I

sensitivity. Overall, the sequence analysis suggests that differences between SARS-CoV and

SARS-CoV-2 viral proteins may drive attenuation in the context of IFN-I pretreatment.

The increased sensitivity of SARS-CoV-2 suggests utility in treatment using IFN-I. While

IFN-I has been used in response to chronic viral infection (43), previous examination of SARS-

CoV cases found inconclusive effect for IFN-I treatment (44). However, the findings from the

SARS-CoV outbreak were complicated by combination therapy of IFN-I with other treatments

including ribavirin/steroids and lack of a regimented protocol. While IFN-I has been utilized to

treat MERS-CoV infected patients, no conclusive data yet exists to determine efficacy (45). Yet,

in vivo studies with MERS-CoV have found that early induction with IFN-I can be protective in

mice (46); importantly, the same study found that late IFN-I activation can be detrimental for

MERS-CoV disease (46). Similarly, early reports have described treatments using IFN-I in

combination for SARS-CoV-2 infection; yet the efficacy of these treatments and the parameters

of their use are not known (47). Overall, sensitivity data suggest that IFN-I treatment may have

utility for treating SARS-CoV-2 if the appropriate parameters can be determined. In addition, use

of type III IFN, which is predicted to have utility in the respiratory tract, could offer another

means for effective treatment for SARS-CoV-2.

In addition to treatment, the sensitivity to IFN-I may also have implications for animal

model development. For SARS-CoV, mouse models that recapitulate human disease were

developed through virus passage in immune competent mice (48). Similarly, mouse models for

MERS-CoV required adaptation in mice that had genetic modifications of their dipeptidyl-

peptidase 4 (DPP4), the receptor for MERS-CoV (49, 50). However, each of these MERS-CoV

mouse models still retained full immune capacity. In contrast, SARS-CoV-2 sensitivity to IFN-I

may signal the need to use an immune deficient model to develop relevant disease. While initial

work has suggested incompatibility to SARS-CoV-2 infection in mice based on receptor usage

(8), the IFN-I response may be a second major barrier that needs to be overcome. Similar to

the emergent Zika virus outbreak, the use of IFN-I receptor knockout mice or IFN-I receptor

blocking antibody may be necessary to develop a useful SARS-CoV-2 animal models for

therapeutic testing (51).

Overall, our results indicate that SARS-CoV-2 has a much higher sensitivity to type I IFN

than the previously emergent SARS-CoV. This augmented type I IFN sensitivity is likely due to

changes in viral proteins between the two epidemic CoV strains. Moving forward, these data

could provide important insights for both the treatment of SARS-CoV-2 as well as developing

novel animal models of disease. In this ongoing outbreak, the results also highlight a distinction

between the highly related viruses and suggest insights from SARS-CoV must be verified for

SARS-CoV-2 infection and disease.

Methods

Viruses and cells. SARS-CoV-2 USA-WA1/2020 was provided by the World Reference Center for Emerging Viruses and Arboviruses (WRCEVA) or BEI Resources and was originally obtained from the USA Centers of Disease Control as described (52). SARS-CoV-2 and mouse-adapted recombinant SARS-CoV (MA15) (48) were titrated and propagated on Vero E6 cells, grown in DMEM with 5% fetal bovine serum and 1% antibiotic/antimytotic (Gibco). Calu3 2B4 cells were grown in DMEM with 10% defined fetal bovine serum, 1% sodium pyruvate (Gibco), and 1% antibiotic/antimitotic (Gibco). Standard plaque assays were used for SARS-CoV and SARS-CoV-2 (26, 53). All experiments involving infectious virus were conducted at the University of Texas Medical Branch (Galveston, TX) or New York University School of Medicine (New York City, NY) in approved biosafety level 3 (BSL) laboratories with routine medical monitoring of staff.

Infection and type I IFN pre- and post-treatment. Viral replication studes in Vero E6 and Calu3 2B4 cells were performed as previously described (37, 54). Briefly, cells were washed two times with PBS and inoculated with SARS-CoV or SARS-CoV-2 at a multiplicity of infection (MOI) 0.01 for 60 minutes at 37 °C. Following inoculation, cells were washed 3 times, and fresh media was added to signify time 0. Three or more biological replicates were harvested at each described time. No blinding was used in any sample collections, nor were samples randomized. For type I IFN pretreatment, experiments were completed as previously described (37). Briefly, Vero E6 cells were incubated with 1000 U/mL of recombinant type I IFN-α (PBL Assay Sciences) 18 hours prior to infection (37). Cells were infected as described above and type I IFN was not added back after infection.

Generation of polarized human airway epithelial cultures (HAEC).

hTert-immortalized human normal airway tracheobronchial epithelial cells, BCi.NS1.1 (55) were

maintained in ExPlus growth media (StemCell). To generate HAEC, BCi.NS1.1 were plated

(7.5E4 cells/well) on rat-tail collagen type 1-coated permeable transwell membrane supports

(6.5mm; Corning Inc), immersed in ExPlus growth media in both the apical and basal chamber.

Upon reaching confluency, media in the apical chamber was removed (airlift), and media in the

basal chamber was changed to Pneumacult ALI maintenance media (StemCell). Pneumacult

ALI maintenance media was changed every two days for approximately 6 weeks to form

differentiated, polarized cultures that resemble in vivo pseudostratified mucociliary epithelium.

IFN-I treatment and infection of HAEC.

For interferon pretreatment of human airway epithelial cultures, 1000 U/mL of universal

recombinant interferon-alpha was added to the basolateral chamber 2 h prior to viral infections.

For viral infections, cultures were washed apically with 50 µl of prewarmed PBS three times for

30 min at 37°C. Virus inoculates (1.3E6 PFU for influenza A/California/07/2009 (H1N1) virus

and 1.35E5 PFU for SARS-CoV-2 USA/WA/1/2020 in 50 µl PBS Mg/Ca) were added apically for

two hours. Inoculates were then removed and cultures were washed three times with PBS, the

third wash was collected and then the cultures were incubated for the indicated times. Progeny

virus was collected in apical washes with 50 µl PBS Mg/Ca for 30 min at 8, 24 and 48 hpi

(endpoint). At endpoint, the membranes were collected and lysed in 300 µl LDS lysis buffer (Life

Technologies). Protein levels were measured by western blot. Briefly, 5 µl of cell lysates were

loaded into a SDS-PAGE gel and transferred to a nitrocellulose membrane. The membranes

were blocked for 1 h at room temperature with 5% skim milk and then incubated overnight at

4°C with appropriate primary antibody dilution: anti-STAT1 (1:1000, Cell Signaling), anti-

pSTAT1 (1:1000, Cell Signaling) or anti-beta actin (1:5000, Invitrogen). Then, the membranes

were incubated with the recommended dilution of HRP-conjugated secondary antibody (goat

anti-rabbit IgG HRP 1:10000, Invitrogen; goat anti-mouse IgG HRP 1:10000, Invitrogen) at room

temperature for 1 h. Super Signal West Dura Kit (Thermo Scientific) was used for signal

development according to manufacturer's recommendations and the chemiluminescence was

detected with a ChemiDoc Imager (BioRad).

Phylogenetic Tree and Sequence Identity Heat Map. Heat maps were constructed from a set

of representative group 2B coronaviruses by using alignment data paired with neighbor-joining

phylogenetic trees built in Geneious (v.9.1.5). Sequence identity was visualized using EvolView

(http://evolgenius.info/) and utilized SARS-CoV Urbani as the reference sequence. Tree shows

the degree of genetic similarity of SARS-CoV-2 and SARS-CoV across selected group 2B

coronaviruses.

Immunoblot Analysis and Antibodies:

Viral and host protein analysis were evaluated as previously described (52, 56). Briefly, cell

lysates were resolved on 7.5% Mini-PROTEAN TGX SDS-PAGE gels and then transferred to

polyvinylidene difluoride (PVDF) membranes using a Trans-Blot Turbo transfer system (Bio-

Rad). Membranes were blocked with 5% (w/v) non-fat dry milk in TBST (TBS with 0.1% (v/v)

Tween-20) for 1 hr, and then probed with the indicated primary antibody in 3% (w/v) BSA in

TBST at 4°C overnight. Following overnight incubation, membranes were probed with the

following secondary antibodies in 5% (w/v) non-fat dry milk in TBST for 1 hr at room

temperature: anti-rabbit or anti-mouse IgG-HRP conjugated antibody from sheep (both 1:10,000

GE Healthcare). Proteins were visualized using ECL or SuperSignal West Femto

chemiluminescence reagents (Pierce) and detected by autoradiography. The following primary

antibodies were used: anti-pSTAT1 (Y701) (1:1000 9171L Cell Signaling Technologies), anti-

STAT1 D1K9Y (1:1000 14994P Cell Signaling Technologies), anti-IFITM1 (1:1000 PA5-20989

Invitrogen), anti-SARS-CoV/CoV-2 Spike 1A9 (1:1000 GTX632604 GeneTex), and anti-β-Actin

(1:1000 ab8227 Abcam).

Statistical analysis. All statistical comparisons in this manuscript involved the comparison

between 2 groups, SARS-CoV or SARS-CoV-2 infected groups under equivalent conditions.

Thus, significant differences in viral titer were determined by the unpaired two-tailed student's T-

Test.

Acknowledgements. Research was supported by grants from NIA and NIAID of the NIH

(U19AI100625 and R00AG049092 to VDM; R24AI120942 to WRCEVA; R01AI134907 to RR;

1R01AI143639-01 and 1R21AI139374-01 to MD; Jan Vilcek/David Goldfarb Fellowship

Endowment Funds to AMVJ; and T32 AI007526 to AH). Research was also supported by

STARs Award provided by the University of Texas System to VDM and trainee funding provided

by the McLaughlin Fellowship Fund at UTMB.

References

1.	**Gralinski LE, Menachery VD.** 2020. Return of the Coronavirus: 2019-nCoV. Viruses **12**.

2.	**Gorbalenya AE, Baker SC, Baric RS, de Groot RJ, Drosten C, Gulyaeva AA, Haagmans BL, Lauber C, Leontovich AM, Neuman BW, Penzar D, Perlman S, Poon LLM, Samborskiy DV, Sidorov IA, Sola I, Ziebuhr J, Coronaviridae Study Group of the International Committee on Taxonomy of V.** 2020. The species Severe acute respiratory syndrome-related coronavirus: classifying 2019-nCoV and naming it SARS-CoV-2. Nature Microbiology doi:10.1038/s41564-020-0695-z.

3.	**Zhu N, Zhang D, Wang W, Li X, Yang B, Song J, Zhao X, Huang B, Shi W, Lu R, Niu P, Zhan F, Ma X, Wang D, Xu W, Wu G, Gao GF, Tan W, China Novel Coronavirus I, Research T.** 2020. A Novel Coronavirus from Patients with Pneumonia in China, 2019. N Engl J Med **382:**727-733.

4.	**Huang C, Wang Y, Li X, Ren L, Zhao J, Hu Y, Zhang L, Fan G, Xu J, Gu X, Cheng Z, Yu T, Xia J, Wei Y, Wu W, Xie X, Yin W, Li H, Liu M, Xiao Y, Gao H, Guo L, Xie J, Wang G, Jiang R, Gao Z, Jin Q, Wang J, Cao B.** 2020. Clinical features of patients infected with 2019 novel coronavirus in Wuhan, China. Lancet **395:**497-506.

5.	**Wu Z, McGoogan JM.** 2020. Characteristics of and Important Lessons From the Coronavirus Disease 2019 (COVID-19) Outbreak in China: Summary of a Report of 72314 Cases From the Chinese Center for Disease Control and Prevention. JAMA doi:10.1001/jama.2020.2648.

6.	**Xu X, Chen P, Wang J, Feng J, Zhou H, Li X, Zhong W, Hao P.** 2020. Evolution of the novel coronavirus from the ongoing Wuhan outbreak and modeling of its spike protein for risk of human transmission. Sci China Life Sci **63:**457-460.

7.	**Letko M, Marzi A, Munster V.** 2020. Functional assessment of cell entry and receptor usage for SARS-CoV-2 and other lineage B betacoronaviruses. Nat Microbiol doi:10.1038/s41564-020-0688-y.

8.	**Zhou P, Yang XL, Wang XG, Hu B, Zhang L, Zhang W, Si HR, Zhu Y, Li B, Huang CL, Chen HD, Chen J, Luo Y, Guo H, Jiang RD, Liu MQ, Chen Y, Shen XR, Wang X, Zheng XS, Zhao K, Chen QJ, Deng F, Liu LL, Yan B, Zhan FX, Wang YY, Xiao GF, Shi ZL.** 2020. A pneumonia outbreak associated with a new coronavirus of probable bat origin. Nature doi:10.1038/s41586-020-2012-7.

9.	**de Wit E, Feldmann F, Cronin J, Jordan R, Okumura A, Thomas T, Scott D, Cihlar T, Feldmann H.** 2020. Prophylactic and therapeutic remdesivir (GS-5734) treatment in the rhesus macaque model of MERS-CoV infection. Proc Natl Acad Sci U S A doi:10.1073/pnas.1922083117.

10.	**Wang M, Cao R, Zhang L, Yang X, Liu J, Xu M, Shi Z, Hu Z, Zhong W, Xiao G.** 2020. Remdesivir and chloroquine effectively inhibit the recently emerged novel coronavirus (2019-nCoV) in vitro. Cell Res doi:10.1038/s41422-020-0282-0.

11.	**Sheahan TP, Sims AC, Leist SR, Schafer A, Won J, Brown AJ, Montgomery SA, Hogg A, Babusis D, Clarke MO, Spahn JE, Bauer L, Sellers S, Porter D, Feng JY, Cihlar T, Jordan R, Denison MR, Baric RS.** 2020. Comparative therapeutic efficacy of remdesivir and combination lopinavir, ritonavir, and interferon beta against MERS-CoV. Nat Commun **11:**222.

12.	**Sheahan TP, Sims AC, Graham RL, Menachery VD, Gralinski LE, Case JB, Leist SR, Pyrc K, Feng JY, Trantcheva I, Bannister R, Park Y, Babusis D, Clarke MO, Mackman RL, Spahn JE, Palmiotti CA, Siegel D, Ray AS, Cihlar T, Jordan R, Denison MR, Baric RS.** 2017. Broad-spectrum antiviral GS-5734 inhibits both epidemic and zoonotic coronaviruses. Sci Transl Med **9**.

13.	**Ahmed SF, Quadeer AA, McKay MR.** 2020. Preliminary Identification of Potential Vaccine Targets for the COVID-19 Coronavirus (SARS-CoV-2) Based on SARS-CoV Immunological Studies. Viruses **12**.

14.   **Medzhitov R, Janeway CA, Jr.** 1997. Innate immunity: the virtues of a nonclonal
system of recognition. Cell **91:**295-298.
15.   **Meylan E, Tschopp J.** 2006. Toll-like receptors and RNA helicases: two parallel ways to
trigger antiviral responses. Mol Cell **22:**561-569.
16.   **Akira S.** 2006. TLR signaling. Curr Top Microbiol Immunol **311:**1-16.
17.   **Schoggins JW, Wilson SJ, Panis M, Murphy MY, Jones CT, Bieniasz P, Rice CM.**
2011. A diverse range of gene products are effectors of the type I interferon antiviral response.
Nature **472:**481-485.
18.   **Platanias LC.** 2005. Mechanisms of type-I- and type-II-interferon-mediated signalling.
Nat Rev Immunol **5:**375-386.
19.   **Rajsbaum R, Garcia-Sastre A.** 2013. Viral evasion mechanisms of early antiviral
responses involving regulation of ubiquitin pathways. Trends Microbiol **21:**421-429.
20.   **Lin FC, Young HA.** 2014. Interferons: Success in anti-viral immunotherapy. Cytokine
Growth Factor Rev **25:**369-376.
21.   **Zumla A, Hui DS, Perlman S.** 2015. Middle East respiratory syndrome. Lancet
386:995-1007.
22.   **Song Z, Xu Y, Bao L, Zhang L, Yu P, Qu Y, Zhu H, Zhao W, Han Y, Qin C.** 2019.
From SARS to MERS, Thrusting Coronaviruses into the Spotlight. Viruses **11**.
23.   **Diaz MO, Ziemin S, Le Beau MM, Pitha P, Smith SD, Chilcote RR, Rowley JD.** 1988.
Homozygous deletion of the alpha- and beta 1-interferon genes in human leukemia and derived
cell lines. Proc Natl Acad Sci U S A **85:**5259-5263.
24.   **Narayanan K, Ramirez SI, Lokugamage KG, Makino S.** 2015. Coronavirus
nonstructural protein 1: Common and distinct functions in the regulation of host and viral gene
expression. Virus Res **202:**89-100.
25.   **Menachery VD, Eisfeld AJ, Schafer A, Josset L, Sims AC, Proll S, Fan S, Li C,**
Neumann G, Tilton SC, Chang J, Gralinski LE, Long C, Green R, Williams CM, Weiss J,
Matzke MM, Webb-Robertson BJ, Schepmoes AA, Shukla AK, Metz TO, Smith RD, Waters
KM, Katze MG, Kawaoka Y, Baric RS. 2014. Pathogenic influenza viruses and coronaviruses
utilize similar and contrasting approaches to control interferon-stimulated gene responses. MBio
5:e01174-01114.
26.   **Sims AC, Tilton SC, Menachery VD, Gralinski LE, Schäfer A, Matzke MM, Webb-**
Robertson BJ, Chang J, Luna ML, Long CE, Shukla AK, Bankhead AR, Burkett SE,
Zornetzer G, Tseng CT, Metz TO, Pickles R, McWeeney S, Smith RD, Katze MG, Waters
KM, Baric RS. 2013. Release of severe acute respiratory syndrome coronavirus nuclear import
block enhances host transcription in human lung cells. J Virol **87:**3885-3902.
27.   **Österlund P, Pirhonen J, Ikonen N, Rönkkö E, Strengell M, Mäkelä SM, Broman M,**
Hamming OJ, Hartmann R, Ziegler T, Julkunen I. 2010. Pandemic H1N1 2009 Influenza A
Virus Induces Weak Cytokine Responses in Human Macrophages and Dendritic Cells and Is
Highly Sensitive to the Antiviral Actions of Interferons. Journal of Virology **84:**1414-1422.
28.   **Totura AL, Baric RS.** 2012. SARS coronavirus pathogenesis: host innate immune
responses and viral antagonism of interferon. Curr Opin Virol **2:**264-275.
29.   **Yang XL, Hu B, Wang B, Wang MN, Zhang Q, Zhang W, Wu LJ, Ge XY, Zhang YZ,**
Daszak P, Wang LF, Shi ZL. 2015. Isolation and Characterization of a Novel Bat Coronavirus
Closely Related to the Direct Progenitor of Severe Acute Respiratory Syndrome Coronavirus. J
Virol **90:**3253-3256.
30.   **Ge XY, Li JL, Yang XL, Chmura AA, Zhu G, Epstein JH, Mazet JK, Hu B, Zhang W,**
Peng C, Zhang YJ, Luo CM, Tan B, Wang N, Zhu Y, Crameri G, Zhang SY, Wang LF,
Daszak P, Shi ZL. 2013. Isolation and characterization of a bat SARS-like coronavirus that
uses the ACE2 receptor. Nature **503:**535-538.

31.   **Lau SK, Woo PC, Li KS, Huang Y, Tsoi HW, Wong BH, Wong SS, Leung SY, Chan**
KH, Yuen KY. 2005. Severe acute respiratory syndrome coronavirus-like virus in Chinese
horseshoe bats. Proc Natl Acad Sci U S A **102:**14040-14045.
32.   **Clementz MA, Chen Z, Banach BS, Wang Y, Sun L, Ratia K, Baez-Santos YM, Wang**
J, Takayama J, Ghosh AK, Li K, Mesecar AD, Baker SC. 2010. Deubiquitinating and
interferon antagonism activities of coronavirus papain-like proteases. J Virol **84:**4619-4629.
33.   **Kopecky-Bromberg SA, Martinez-Sobrido L, Frieman M, Baric RA, Palese P.** 2007.
Severe acute respiratory syndrome coronavirus open reading frame (ORF) 3b, ORF 6, and
nucleocapsid proteins function as interferon antagonists. J Virol **81:**548-557.
34.   **Frieman M, Yount B, Heise M, Kopecky-Bromberg SA, Palese P, Baric RS.** 2007.
Severe acute respiratory syndrome coronavirus ORF6 antagonizes STAT1 function by
sequestering nuclear import factors on the rough endoplasmic reticulum/Golgi membrane. J
Virol **81:**9812-9824.
35.   **Coutard B, Valle C, de Lamballerie X, Canard B, Seidah NG, Decroly E.** 2020. The
spike glycoprotein of the new coronavirus 2019-nCoV contains a furin-like cleavage site absent
in CoV of the same clade. Antiviral Res **176:**104742.
36.   **Tong ZD, Tang A, Li KF, Li P, Wang HL, Yi JP, Zhang YL, Yan JB.** 2020. Potential
Presymptomatic Transmission of SARS-CoV-2, Zhejiang Province, China, 2020. Emerg Infect
Dis **26**.
37.   **Menachery VD, Yount BL, Jr., Josset L, Gralinski LE, Scobey T, Agnihothram S,**
Katze MG, Baric RS. 2014. Attenuation and restoration of severe acute respiratory syndrome
coronavirus mutant lacking 2'-o-methyltransferase activity. J Virol **88:**4251-4264.
38.   **Thiel V, Weber F.** 2008. Interferon and cytokine responses to SARS-coronavirus
infection. Cytokine Growth Factor Rev **19:**121-132.
39.   **Frieman MB, Chen J, Morrison TE, Whitmore A, Funkhouser W, Ward JM,**
Lamirande EW, Roberts A, Heise M, Subbarao K, Baric RS. 2010. SARS-CoV pathogenesis
is regulated by a STAT1 dependent but a type I, II and III interferon receptor independent
mechanism. PLoS Pathog **6:**e1000849.
40.   **Channappanavar R, Fehr AR, Vijay R, Mack M, Zhao J, Meyerholz DK, Perlman S.**
2016. Dysregulated Type I Interferon and Inflammatory Monocyte-Macrophage Responses
Cause Lethal Pneumonia in SARS-CoV-Infected Mice. Cell Host Microbe **19:**181-193.
41.   **Menachery VD, Gralinski LE, Mitchell HD, Dinnon KH, 3rd, Leist SR, Yount BL, Jr.,**
Graham RL, McAnarney ET, Stratton KG, Cockrell AS, Debbink K, Sims AC, Waters KM,
Baric RS. 2017. Middle East Respiratory Syndrome Coronavirus Nonstructural Protein 16 Is
Necessary for Interferon Resistance and Viral Pathogenesis. mSphere **2**.
42.   **Falzarano D, de Wit E, Martellaro C, Callison J, Munster VJ, Feldmann H.** 2013.
Inhibition of novel beta coronavirus replication by a combination of interferon-alpha2b and
ribavirin. Sci Rep **3:**1686.
43.   **Finter NB, Chapman S, Dowd P, Johnston JM, Manna V, Sarantis N, Sheron N,**
Scott G, Phua S, Tatum PB. 1991. The use of interferon-alpha in virus infections. Drugs
42:749-765.
44.   **Stockman LJ, Bellamy R, Garner P.** 2006. SARS: systematic review of treatment
effects. PLoS Med **3:**e343.
45.   **de Wit E, van Doremalen N, Falzarano D, Munster VJ.** 2016. SARS and MERS:
recent insights into emerging coronaviruses. Nat Rev Microbiol **14:**523-534.
46.   **Channappanavar R, Fehr AR, Zheng J, Wohlford-Lenane C, Abrahante JE, Mack M,**
Sompallae R, McCray PB, Jr., Meyerholz DK, Perlman S. 2019. IFN-I response timing
relative to virus replication determines MERS coronavirus infection outcomes. J Clin Invest
130:3625-3639.
47.   **Pang J, Wang MX, Ang IYH, Tan SHX, Lewis RF, Chen JI, Gutierrez RA, Gwee**
SXW, Chua PEY, Yang Q, Ng XY, Yap RK, Tan HY, Teo YY, Tan CC, Cook AR, Yap JC, Hsu

LY. 2020. Potential Rapid Diagnostics, Vaccine and Therapeutics for 2019 Novel Coronavirus
(2019-nCoV): A Systematic Review. J Clin Med **9**.
48.   **Roberts A, Deming D, Paddock CD, Cheng A, Yount B, Vogel L, Herman BD,**
Sheahan T, Heise M, Genrich GL, Zaki SR, Baric R, Subbarao K. 2007. A mouse-adapted
SARS-coronavirus causes disease and mortality in BALB/c mice. PLoS Pathog **3**:e5.
49.   **Cockrell A YB, Scobey T, Jensen K, Douglas M, Beall A, Tang X-C, Marasco WA,**
Heise MT, Baric RS 2016. A Mouse Model for MERS Coronavirus Induced Acute Respiratory
Distress Syndrome. . Nature Microbiology **In Press**.
50.   **Li K, Wohlford-Lenane CL, Channappanavar R, Park JE, Earnest JT, Bair TB, Bates**
AM, Brogden KA, Flaherty HA, Gallagher T, Meyerholz DK, Perlman S, McCray PB, Jr.
2017. Mouse-adapted MERS coronavirus causes lethal lung disease in human DPP4 knockin
mice. Proc Natl Acad Sci U S A **114**:E3119-E3128.
51.   **Lazear HM, Govero J, Smith AM, Platt DJ, Fernandez E, Miner JJ, Diamond MS.**
2016. A Mouse Model of Zika Virus Pathogenesis. Cell Host Microbe **19**:720-730.
52.   **Harcourt J, Tamin A, Lu X, Kamili S, Sakthivel SK, Murray J, Queen K, Tao Y,**
Paden CR, Zhang J, Li Y, Uehara A, Wang H, Goldsmith C, Bullock HA, Wang L, Whitaker
B, Lynch B, Gautam R, Schindewolf C, Lokugamage KG, Scharton D, Plante JA,
Mirchandani D, Widen SG, Narayanan K, Makino S, Ksiazek TG, Plante KS, Weaver SC,
Lindstrom S, Tong S, Menachery VD, Thornburg NJ. 2020. Severe Acute Respiratory
Syndrome Coronavirus 2 from Patient with 2019 Novel Coronavirus Disease, United States.
Emerg Infect Dis **26**.
53.   **Josset L, Menachery VD, Gralinski LE, Agnihothram S, Sova P, Carter VS, Yount**
BL, Graham RL, Baric RS, Katze MG. 2013. Cell host response to infection with novel human
coronavirus EMC predicts potential antivirals and important differences with SARS coronavirus.
MBio **4**:e00165-00113.
54.   **Sheahan T, Rockx B, Donaldson E, Corti D, Baric R.** 2008. Pathways of cross-
species transmission of synthetically reconstructed zoonotic severe acute respiratory syndrome
coronavirus. J Virol **82**:8721-8732.
55.   **Walters MS, Gomi K, Ashbridge B, Moore MA, Arbelaez V, Heldrich J, Ding BS,**
Rafii S, Staudt MR, Crystal RG. 2013. Generation of a human airway epithelium derived basal
cell line with multipotent differentiation capacity. Respir Res **14**:135.
56.   **van Tol S, Atkins C, Bharaj P, Johnson KN, Hage A, Freiberg AN, Rajsbaum R.**
2020. VAMP8 Contributes to the TRIM6-Mediated Type I Interferon Antiviral Response during
West Nile Virus Infection. J Virol **94**.

Figure Legends

Figure 1. SARS-CoV-2 sensitive to type I IFN pretreatment. A) Vero E6 cells were treated

with 1000 U/mL recombinant type I (hashed line) IFN or mock (solid line) for 18 hours prior to

infection. Cells were subsequently infected with either SARS-CoV WT (black) or SARS-CoV-2

(blue) at an MOI of 0.01 as described above. Each point on the line graph represents the group

mean, N ≥ 3. All error bars represent SD. The two tailed students t-test was used to determine

P-values: *** $P < 0.001$. B) Vero cell protein lysates from IFN-I treated and untreated cells were

probed 48 hours post infection by Western blotting for phosphorylated STAT1 (Y701), STAT1,

IFITM1, SARS spike, and Actin.

Figure 2. SARS-CoV-2 attenuated and IFN-I sensitive in Calu3 respiratory cells. A) Calu3

2B4 cells were treated with 1000 U/mL recombinant type I (hashed line) IFN or mock (solid line)

for 18 hours prior to infection. Cells were subsequently infected with either SARS-CoV WT

(black) or SARS-CoV-2 (blue) at an MOI of 1. Each point on the line graph represents the

group mean, N ≥ 3. All error bars represent SD. The two tailed students t-test was used to

determine P-values: *** $P < 0.001$. B) Calu3 cell protein lysates from IFN-I treated and untreated

cells were probed 48 hours post infection by Western blotting for phosphorylated STAT1 (Y701),

STAT1, IFITM1, SARS spike, and Actin.

Figure 3. Differential IFN-I sensitivity and pSTAT1 phosphorylation following SARS-CoV-

2 or influenza A virus on polarized human airway epithelial cultures (HAEC). A) HAEC

were pretreated with 1000 U/mL IFN-α basolaterally for 2 hours prior to and during infection.

Cultures were then infected apically with influenza A/California/09 H1N1 virus or SARS-CoV-2.

Apical washes were collected at the indicated times and progeny titers determined by plaque

assay on MDCK cells (influenza virus) or Vero E6 cells (SARS-CoV-2). At the 48 h endpoint,

cultures were lysed for western blot analysis. B) Western for total STAT1 or phospho-STAT1 at

48 hpi, and actin as loading control. C) Influenza A virus titers by plaque assay on MDCK cells.

0 h, virus inoculate; 2 h, virus in third apical wash; 8, 24, 48 h, virus in apical washes at these

time points. D) SARS-CoV-2 titers by plaque assay on Vero E6 cells. 0 h, virus inoculate; 2 h,

virus in second apical wash; 8, 24, 48 h, virus in apical washes at these time points.

Figure 4, SARS-CoV-2 impacted by post IFN-I treatment. A-B) Vero E6 and Calu3 2B4 cells

were infected with either SARS-CoV WT (black) or SARS-CoV-2 (blue) at an MOI 0.01 (A, Vero

cells) or MOI 1 (B, Calu3 cells). Cells were subsequently treated with 1000 U/mL recombinant

type I IFN (hashed line) or mock (solid line) for 4 hours following infection. Each point on the line

graph represents the group mean, N ≥ 3. All error bars represent SD. The two tailed student's t-

test was used to determine P-values: **P <0.01 *** P < 0.001.

Figure 5, Conservation of SARS-CoV IFN antagonists. Viral protein sequences of the

indicated viruses were aligned according to the bounds of the SARS-CoV open reading frames

for each viral protein. Sequence identities were extracted from the alignments for each viral

protein, and a heat map of percent sequence identity was constructed using EvolView

(www.evolgenius.info/evolview) with SARS-CoV as the reference sequence. TR = truncated

protein.

Figure 1. SARS-CoV-2 sensitive to type I IFN pretreatment. A) Vero E6 cells were treated
with 1000 U/mL recombinant type I (hashed line) IFN or mock (solid line) for 18 hours prior to
infection. Cells were subsequently infected with either SARS-CoV WT (black) or SARS-CoV-2
(blue) at an MOI of 0.01 as described above. Each point on the line graph represents the group
mean, N ≥ 3. All error bars represent SD. The two tailed student's t-test was used to determine
P-values: *** P < 0.001. B) Vero cell protein lysates from IFN-I treated and untreated cells were
probed 48 hours post infection by Western blotting for phosphorylated STAT1 (Y701), STAT1,
IFITM1, SARS spike, and Actin.

655
656
Figure 2. SARS-CoV-2 attenuated and IFN-I sensitive in Calu3 respiratory cells. A) Calu3
2B4 cells were treated with 1000 U/mL recombinant type I (hashed line) IFN or mock (solid line)
for 18 hours prior to infection. Cells were subsequently infected with either SARS-CoV WT
(black) or SARS-CoV-2 (blue) at an MOI of 1. Each point on the line graph represents the
group mean, N ≥ 3. All error bars represent SD. The two tailed student's t-test was used to
determine P-values: *** P < 0.001. B) Calu3 cell protein lysates from IFN-I treated and untreated
cells were probed 48 hours post infection by Western blotting for phosphorylated STAT1 (Y701),
STAT1, IFITM1, SARS spike, and Actin.

666
667

Figure 3. Differential IFN-I sensitivity and pSTAT1 phosphorylation following SARS-CoV-2 or influenza A virus on polarized human airway epithelial cultures (HAEC). A) HAEC were pretreated with 1000 U/mL IFN-α basolaterally for 2 hours prior to and during infection. Cultures were then infected apically with influenza A/California/09 H1N1 virus or SARS-CoV-2. Apical washes were collected at the indicated times and progeny titers determined by plaque assay on MDCK cells (influenza virus) or Vero E6 cells (SARS-CoV-2). At the 48 h endpoint, cultures were lysed for western blot analysis. B) Western for total STAT1 or phospho-STAT1 at 48 hpi. C) Influenza A virus titers by plaque assay on MDCK cells. 0 h, virus inoculate; 2 h, virus in third apical wash; 8, 24, 48 h, virus in apical washes at these time points. D) SARS-CoV-2 titers by plaque assay on Vero E6 cells. 0 h, virus inoculate; 2 h, virus in second apical wash; 8, 24, 48 h, virus in apical washes at these time points.

680
Figure 4, SARS-CoV-2 impacted by post IFN-I treatment. A-B) Vero E6 and Calu3 2B4 cells
were infected with either SARS-CoV WT (black) or SARS-CoV-2 (blue) at an MOI 0.01 (A, Vero
cells) or MOI 1 (B, Calu3 cells). Cells were subsequently treated with 1000 U/mL recombinant
type I IFN (hashed line) or mock (solid line) for 4 hours following infection. Each point on the line
graph represents the group mean, N ≥ 3. All error bars represent SD. The two tailed student's t-
test was used to determine P-values: **P <0.01 *** P < 0.001.

Figure 5, Conservation of SARS-CoV IFN antagonists. Viral protein sequences of the
indicated viruses were aligned according to the bounds of the SARS-CoV open reading frames
for each viral protein. Sequence identities were extracted from the alignments for each viral
protein, and a heat map of percent sequence identity was constructed using EvolView
(www.evolgenius.info/evolview) with SARS-CoV as the reference sequence. TR = truncated
protein.