Universal nature of drug treatment responses in drug-tissue-wide model-animal experiments using tensor decomposition-based unsupervised feature extraction

Y-h. Taguchi [1,*]**, Turki Turki**[2]

[1]*Department of Physics, Chuo University, Tokyo 112-8551, Japan*
[2] *King Abdulaziz University, Department of Computer Science, Jeddah 21589, Saudi Arabia*

Correspondence*:
Corresponding Author
tag@granular.com

ABSTRACT

Gene expression profiles of tissues treated with drugs have recently been used to infer clinical outcomes. Although this method is often successful from the application point of view, gene expression altered by drugs is rarely analyzed in detail, because of the extremely large number of genes involved. Here, we applied tensor decomposition (TD)-based unsupervised feature extraction (FE) to the gene expression profiles of 24 mouse tissues treated with 15 drugs. TD-based unsupervised FE enabled identification of the common effects of 15 drugs including an interesting universal feature: these drugs affect genes in a gene-group-wide manner and were dependent on three tissue types (neuronal, muscular, and gastroenterological). For each tissue group, TD-based unsupervised FE enabled identification of a few tens to a few hundreds of genes affected by the drug treatment. These genes are distinctly expressed between drug treatments and controls as well as between tissues in individual tissue groups and other tissues. We also validated the assignment of genes to individual tissue groups using multiple enrichment analyses. We conclude that TD-based unsupervised FE is a promising method for integrated analysis of gene expression profiles from multiple tissues treated with multiple drugs in a completely unsupervised manner.

Keywords: tensor decomposition, unsupervised learning, gene selection, drug treatment

BACKGROUND

Drug design is a time-consuming and expensive process. Multiple coordinated experimental efforts, involving large-scale trial-and-error methods, are required to investigate new compounds. In general, this is due to the inherent difficulties in identifying novel therapeutic targets such as genes that cause disease. Even where potential target genes are identified robustly, it is difficult to find drug candidate compounds that successfully bind to the proteins they encode.

Computer-based methods have been introduced in an attempt to shorten the period of drug development and to reduce the expenses involved. The two major computer-aided drug design strategies are ligand-based drug design (LBDD) and structure-based drug design (SBDD). LBDD has various advantages including less required computational resources and better success rates for drug design. However, it also has the

disadvantage of limited ability to find drug candidate compounds with low structural similarity to known
drugs. To compensate for the weaknesses of LBDD, SBDD shows a greater ability to find drug candidate
compounds lacking in structural similarity with known drugs. This is because SBDD tries to screen drug
candidate compounds by investigating whether these can bind to target proteins. The weak point of SBDD
is that it requires massive computational resources, and this prevents its application to large-scale screening,
in which candidate drug compounds often number several million.
Considering the relatively low cost of obtaining gene expression profiles, a third computer-aided strategy
has been proposed: gene expression profile-based drug design. In this strategy, the gene expression profiles
of tissues/cell lines treated with candidate drug compounds are collected. The collected profiles are then
compared with those of tissues/cell lines treated with known drug compounds. If the candidate drug
compounds share a gene expression profile to some extent with known drug compounds, they are identified
as having therapeutic potential against target diseases/proteins.
Some databases have been established to assist gene expression profiling for drug design. For example,
chemical checker (10) includes gene expression in computer-aided drug design, whereas PharmacoDB (49)
is fully implemented to consider the dose dependence of drug-treated cell lines for drug design. Many
papers have been published on the use of gene expression profiles for computer-aided drug design (8, 3).
For instance, Huang et al (18) used combinatorial analysis of drug-induced gene expression for cancer
drugs, which were then experimentally confirmed *in vitro*. Lee et al. (31) proposed DeSigN, a robust and
useful method for identifying candidate drugs using an input gene signature obtained from gene expression
analysis. Kim et al. (25) performed computational drug repositioning for gastric cancer using reversal of
gene expression profiles, and Wolf et al. (9) analyzed high-throughput gene expression profiles to identify
similarities between drugs and to predict compound activity. Hodos et al. (17) tried to fill in missing gene
expression observations in cells treated with drugs by predicting cell-specific drug perturbation profiles
using available expression data from related conditions. Pabon et al. (40) predicted protein targets for
drug-like compounds using transcriptomics. In contrast, Liu et al. (33) performed comparative analysis of
genes that are frequently regulated by drugs based on connectivity to map transcriptome data.
In contrast to these successful applications of gene expression profile analysis to computer-aided drug
design, it is unclear how individual gene expression is affected by drug treatment. First, the number of
genes expressed in a dose dependent-manner is as large as the number of genes expressed. Thus, it is
not easy to invent a useful method to integrate and understand the dose dependent-genes pertaining to
individual gene expression profiles. For example, Luka et al. (35) employed principal component analysis
(PCA) to integrate the dose dependence of gene expression profiles upon combinatorial drug treatment.
They reported a convex (not monotonic) dependence on dose density and identified this as evidence of
the cooperative effects of dual drug treatments. Nevertheless, convex dependence on dose was reportedly
observed in a single drug treatment if tensor decomposition (TD) was employed to integrate multiple gene
expression profiles of cell lines treated with a single drug (13). Thus, it is primarily important to identify
an effective method that can integrate numerous gene expression profiles of tissues/cell lines treated with
drugs.
Recently, Kozawa et al. (28) used the gene expression profiles of mouse tissues treated with drugs to
predict human clinical outcomes. In this paper, we applied TD-based unsupervised feature extraction (FE)
to the gene expression profiles used in their study and attempted to identify the changes in gene expression
profiles of mouse tissues treated with individual drugs.

METHODS AND MATERIALS

71 Figure 1 shows the flow chart of analysis.

Gene expression profiles

The gene expression profiles used in this study were downloaded from the gene expression omnibus
(GEO) with GEO ID GSE142068. Twenty four profiles named "GSE142068_count_XXXXX.txt.gz" were
downloaded, where "XXXXX" indicates one of the 24 tissues, i.e., AdrenalG, Aorta, BM (Bone marrow),
Brain, Colon, Eye, Heart, Ileum, Jejunum, Kidney, Liver, Lung, Pancreas, ParotidG, PituitaryG, SkMuscle,
Skin, Skull, Spleen, Stomach, Spleen, Thymus, ThyroidG, and WAT (white adipose tissue), which were

treated with 15 drugs: Alendronate, Acetaminophen, Aripiprazole, Asenapine, Cisplatin, Clozapine,
Clozapine, Empagliflozin, Lenalidomide, Lurasidone, Olanzapine, Evolocumab, Risedronate, Sofosbuvir,
and Teriparatide, and Wild type (WT).

TD-based unsupervised FE

For applying TD-based unsupervised FE (14) to the downloaded gene expression profiles, they must be
formatted as a tensor. In this analysis, they were formatted as tensor, $x_{ijkm} \in \mathbb{R}^{N \times 24 \times 18 \times 2}$, for N genes,
24 tissues, 18 drug treatments, and two replicates. Then, the HOSVD (14) algorithm was applied to x_{ijkm}
and we obtained TD

$$x_{ijkm} = \sum_{\ell_1 \ell_2 \ell_3 \ell_4} G(\ell_1, \ell_2, \ell_3, \ell_4) u_{\ell_1 j} u_{\ell_2 k} u_{\ell_3 m} u_{\ell_4 i} \tag{1}$$

where $G \in \mathbb{R}^{N \times 24 \times 18 \times 2}$ is the core tensor, $u_{\ell_1 j} \in \mathbb{R}^{24 \times 24}$, $u_{\ell_2 k} \in \mathbb{R}^{18 \times 18}, u_{\ell_3 m} \in \mathbb{R}^{2 \times 2}$, and $u_{\ell_4 i} \in$
$\mathbb{R}^{N \times N}$, represents singular value matrices that are also orthogonal matrices. x_{ijkm} is considered to be
standardized as $\sum_i x_{ijkm} = 0$ and $\sum_i x_{ijkm}^2 = N$.
Mathematically, eq. (1) aims to decompose the dependence of x_{ijkm} upon i, j, k, m into a series of
products among $u_{\ell_1 j}, u_{\ell_2 k}, u_{\ell_3 m}$, and $u_{\ell_4 i}$, each of which is supposed to represent the dependence on
i, j, k, m. As it is unlikely that a single product of $u_{\ell_1 j}, u_{\ell_2 k}, u_{\ell_3 m}$, and $u_{\ell_4 i}$ can reproduce x_{ijkm}, we
need to consider various combinations of $u_{\ell_1 j}, u_{\ell_2 k}, u_{\ell_3 m}$, and $u_{\ell_4 i}$, where those associated with distinct
$\ell_1, \ell_2, \ell_3, \ell_4$ are supposed to be associated with distinct dependence on i, j, k, m. Then, the products of
$u_{\ell_1 j}, u_{\ell_2 k}, u_{\ell_3 m}$, and $u_{\ell_4 i}$, must be summed up with the weight of G to reproduce x_{ijkm}. Biologically, we
cannot expect that $u_{\ell_1 j}, u_{\ell_2 k}, u_{\ell_3 m}$, and $u_{\ell_4 i}$ can represent the biological aspect because eq. (1) is simply a
mathematical hypothesis; therefore, their association with a biological aspect after performing TD needs to
be validated.
To understand how gene expression profiles are altered by drug treatment in a tissue-group-wide manner,
we first need to investigate $u_{\ell_1 j}, u_{\ell_2 k}$, and $u_{\ell_3 m}$. After identifying which ℓ_1, ℓ_2 and ℓ_3 are biologically
interesting, we select ℓ_4 associated with $G(\ell_1, \ell_2, \ell_3, \ell_4)$ that have the largest absolute values with fixed
ℓ_1, ℓ_2 and ℓ_3, because $u_{\ell_4 i}$ associated with such ℓ_4 is supposed to represent the weight of gene i that is
expressed in association with j, k, m dependence represented by the selected $u_{\ell_1 j}, u_{\ell_2 k}, u_{\ell_3 m}$.
Using the identified $u_{\ell_4 i}$, the P-value, P_i, is attributed to gene i as

$$P_i = P_{\chi^2} \left[> \left(\frac{u_{\ell_4 i}}{\sigma_{\ell_4}} \right)^2 \right] \tag{2}$$

where $P_{\chi^2}[> x]$ is the cumulative probability of χ^2 distribution and σ_{ℓ_4} is the standard deviation. Here, we
assume that $u_{\ell_4 i}$ obeys a Gaussian distribution with zero mean because $\sum_i x_{ijkm} = 0$. P_i is corrected via
the BH criterion (6) and I, a set of genes i associated with adjusted P-values less than 0.01, is selected.
For a more detailed explanation of TD-based unsupervised FE, see the recently published monograph (14).

t **test and Wilcoxon test applied to sets of genes classified based on tissue groups and**
drugs groups

In order to determine whether the selected set of genes, I, are expressed distinctly between the two
assigned tissue groups, J, $\{x_{ijkm}|i \in I, j \in J\}$, and $\overline{J}$, $\{x_{ijkm}|i \in I, j \in \overline{J}\}$, we applied a two-way t
test and Wilcoxon test and computed the P-values. Similar analyses were done with two drug groups, K,
$\{x_{ijkm}|i \in I, k \in K\}$, and $\overline{K}$, $\{x_{ijkm}|i \in I, k \in \overline{K}\}$.

Enrichment analysis

The selected genes (gene symbols) were uploaded to Enrichr (30) and Metascape (56) in order to validate
the various biological functions of the selected genes.

RESULTS

Figure 2 summarizes the results obtained in this study.

Drug treatment specificity

After obtaining the TD, eq. (1), we first investigated $u_{\ell_2 k}$ attributed to the kth drug. Although the number
of drugs tested is as many as 15, the total number of drug treatments was considered to be 18 due to the
testing of three additional conditions. Usually, the first singular value vectors represent uniform values (i.e.,
components that are not distinct between samples) (14). In this case, u_{1k} does not represent any dependence
on drug treatment. This is reasonable because the expression of most genes is unlikely to be affected by
drug treatment. We thus considered the second and third singular value vectors, u_{2k} and u_{3k}, attributed
them to drug treatments (Fig. 3). In contrast to expectations, the drug treatments were quite universal.
Most of the drug treatments (other than (2), (9), (15), and (17)) were separated from the control treatments
((2), (9), (15), and (17)) along one direction (red arrow) whereas the diversity among drug treatments was
spread perpendicular (blue arrow) to that direction, only among drug treatments. This suggests that the
gene expression profiles are altered similarly, independently of the drug treatment.

Tissue-specificity

We further studied the relationship of universal drug treatments with individual tissues. For this, we next
investigated $u_{\ell_1 j}$ attributed to 24 tissues. We then found that several $u_{\ell_1 j}$ are expressed in a tissue-group
wide manner (Fig. 4) . The tissue-wide expression pattern identified by singular value vectors is described
as follows; As u_{1j} does not express any tissue specificities, it is unlikely to exhibit tissue specificity; as
u_{2j} has larger absolute values for the brain, eye, pituitary, and testis, it is likely to represent neuronal
tissue specificities; as u_{3j} has larger absolute values only for the parotid, we did not consider it further;
as u_{4j} exhibits larger absolute values for the heart and SkMuscle, we considered that it exhibits muscle
specificities; As u_{5j} and u_{6j} exhibit larger absolute values for the pancreas and stomach, we considered
that it exhibits gastric tissue specificities. It is thus obvious that the combination of tissue specificity is
quite reasonable biologically.
Aiming to specify singular value vectors attributed to genes, $u_{\ell_4 i}$, for gene selection, we then checked
which of $G(\ell_1, 2, 1, \ell_4)$ and $G(\ell_1, 3, 1, \ell_4)$ have larger absolute values, as u_{1m} always exhibits the same
values between two replicates (Table 1).
For $\ell_1 = 2$, which is supposed to be attributed to neuron-specific tissues (u_{2j}), Gs with $\ell_4 = 2$ have
larger absolute values. Thus, u_{2i} was employed for neuron-specific gene selection. For $\ell_1 = 4$, which is
supposed to be attributed to muscle-specific tissues (u_{4j}), Gs with $\ell_4 = 4$ have larger absolute values. Thus,
u_{4i} was employed for muscle-specific gene selection. For $\ell_1 = 5$, which is supposed to be attributed to
gastrointestinal-specific tissues (u_{5j}), Gs with $\ell_4 = 5$ have larger absolute values. Thus, u_{5i} was employed
for muscle-specific gene selection. For $\ell_1 = 6$, which is also supposed to be attributed to gastrointestinal-
specific tissues (u_{6j}), Gs with $\ell_4 = 6, 7$ have larger absolute values. Then, u_{6i} and u_{7i} were employed for
muscle-specific gene selection.
After computing the adjusted P-values, P_i, attributed to the genes (see methods), the genes associated
with adjusted P_i less than 0.01 were selected (Table 2. The lists of selected genes can be found in supporting
information (Additional File 1). Figure 5 shows a Venn diagram of the selected genes. As expected, two sets
of genes, Gas1 and Gas2, which are supposed to be gastrointestinal-specific, are quite common. Other than
these, the selected genes are quite distinct from one another. Thus, TD-based unsupervised FE successfully
identified the genes whose expression was affected by the drugs in a tissue group-specific manner.

Confirmation of differential expression

In order to check whether the selected genes are expressed distinctly between the specified tissues and
other tissues, as well as between drug treatments and controls, we first applied statistical tests to the
selected genes (Table 2). The data clearly showed that for all cases, gene expression was distinct between
the specified tissues and other tissues as well as between drug treatments and controls. Thus, TD-based
unsupervised FE allowed us to select the genes whose expression is coincident with $u_{\ell_2 k}$s in Fig. 3 and
$u_{\ell_1 j}$s in Fig. 4.

Biological evaluation

Next, we evaluated the selected genes biologically. For this purpose, we first uploaded the genes to
Metascape (Fig. 6). Initially, we noticed that Gas1 and Gas2 largely shared the enriched terms as expected,

even though these two gene sets were selected using distinct singular values (u_{5i} and u_{6i}, u_{7i}, respectively).
In particular, it is important to note that two KEGG terms, "mmu04971: Gastric acid secretion" and
"mmu04972: Pancreatic secretion" are shared by Gas1 and Gas2, which are supposed to be Pancreas- and
Stomach-specific. In contrast, various muscle-related terms are enriched in the Muscle gene set as expected,
whereas "GO:0002088: lens development in camera-type eye" is enriched in the neuronal gene set. All of
these results suggest that TD-based unsupervised FE selected the biologically reasonable genes.

Figure 7 shows the protein-protein interaction (PPI) network provided by Metascape. A high degree of
connectivity was obvious. Thus, TD-based unsupervised FE identified the sets of genes among which PPI
is enriched. Moreover, Gas1 and Gas2 largely share the PPI network, whereas the neuronal and muscular
gene sets form their own PPI network within which PPI is enriched. Thus, PPI analysis also indicated that
TD-based unsupervised FE identified biologically reasonable genes.

To eliminate the possibility that our results were specific to the Metascape data set, we uploaded the genes
selected by TD-based unsupervised FE to Enrichr (Table 3). With this data set, we observed clear detection
of at least one tissue-related disease for four sets of tissue-specific genes, validating the Metascape-based
results.

DISCUSSION

Although it is unclear why the 15 drugs affected the expression of many common genes, a detailed
investigation can allow further interpretation. Table 4 shows the drugs' effects on neuronal, muscular, and
pancreatic tissues. These data suggest that most drugs simultaneously affect these three groups of tissues.

Our results are in contrast to the study that inspired our work (28), in which the authors employed a fully
supervised approach requiring previous knowledge. Although Kosawa et al. (28) also aimed to infer the
therapeutic and side effects of drug treatments in humans based on gene expression in drug-treated tissues,
their analysis required previous knowledge that is not needed for TD-based unsupervised FE. In this sense,
our approach has distinct potential that the original study could not achieve.

In addition to the above-mentioned biological superiority of TD-based unsupervised FE, this approach
also has some methodological advantages as follows. First, although we classified 24 tissues into two
groups based on the observation of singular value vectors attributed to tissues, $u_{\ell_1 j}$ (Fig. 4) prior to
the identification of differentially expressed genes, it is computationally infeasible for other methods to
classify 24 tissues into two groups before starting to seek differentially expressed genes, as there are
no criteria on how to divide 24 tissues into two groups. It is thus practically impossible to analyze all
possible divisions, as they number in the millions. The same advantage is observed when grouping 18 drug
treatments into two. This may be much easier than classifying tissues, because some of the drug treatments
are obviously controls. Nevertheless, based upon the second and third singular value vectors attributed to
drug treatments, u_{2k} and u_{3k} (Fig. 3), acetaminophen (APAP) and sofosbuvir are grouped together with
two control treatments. Such a classification can never be proposed without TD. In this sense, there is no
computationally feasible method that can compete with our method.

The biological basis for the two groups of drugs seen in Fig. 3 may be questioned. To clarify this point,
we uploaded two groups of drugs to DrugEnrichr (29), which evaluates the coincidence of genes targeted
by the uploaded drugs (Additional file 2). Based on the "Geneshot Predicted from Co-expression" category
in DrugEnrichr, we found that there are at least as many as 164 genes targeted by two drugs (APAP and
Sofosbuvir) in group1 whereas 213 genes are targeted by at least two drugs among as many as thirteen
drugs included in group2 (Alendronate, Aripiprazole, Asenapine, Cisplatin, Clozapine, Dox, EMPA,
Lenalidomide, Lurasidone, Olanzapine, Repatha, Risedronate, Teriparatide). This suggests that these two
groups of drugs are quite distinct because there are no common targeted genes between these 164 and 213
genes. Thus, the groups of drugs identified by TD based unsupervised FE are likely based on the genes that
the drugs target.

In view of the two above-mentioned advantages, TD-based unsupervised FE might yield completely
distinct outcomes that other supervised methods cannot, and it therefore represents a worthwhile primary
or supplementary approach to gene-expression-based investigation of drug effects.

One might wonder if the results were confirmed only by single experiments. As the results shown in Table
3 indicate coincidence between the present result and other studies, the results derived in this study are not
dependent on a single study, but are coincident with numerous studies in the public domain database.
Moreover, TD-based unsupervised FE is a very useful strategy for repositioning known drugs. As shown
in Fig. 3, TD-based unsupervised FE can determine the effective tissue. Furthermore, as indicated in Table
3, the genes selected by TD-based unsupervised FE can indicate the diseases for which the drugs have
potential effectiveness. Therefore, applying TD-based unsupervised FE to gene expression profiles altered
by drug treatments can be a promising strategy to repurpose known drugs for new diseases.

CONCLUSIONS

224   In this paper, we applied TD-based unsupervised FE (14) to the gene expression profiles of 24 mouse tissues
225   treated with 15 drugs. Integrated analysis allowed us to identify the universal nature of drug treatments in a
226   tissue-group-wide manner, which is generally impossible to identify using any other supervised strategy
227   that requires prior information.

DATA AVAILABILITY

228 All datasets analyzed in this study were obtained from the GEO database.

AUTHOR CONTRIBUTIONS

229 YHT planned and performed the study. YHT and TT discussed the results and wrote the paper.

FUNDING

230 This study was supported by KAKENHI 19H05270, 20K12067, and 20H04848. This project was also
231 funded by the Deanship of Scientific Research (DSR) at King Abdulaziz University, Jeddah, under grant
232 no. KEP-8-611-38. The authors, therefore, acknowledge DSR with thanks for providing technical and
233 financial support.

CONFLICT OF INTEREST STATEMENT

234 The authors declare that the research was conducted in the absence of any commercial or financial
235 relationships that could be construed as a potential conflict of interest.

ACKNOWLEDGMENTS

236 The authors would like to thank the reviewers for very constructive comments and thoughtful suggestions.
237 This manuscript has been released as a pre-print at BioRxiv (Taguchi Y-h. et al, 2020 (50)).

REFERENCES

1 .Anwar, I. J., Miyata, K., and Zsombok, A. (2016). Brain stem as a target site for the metabolic side
effects of olanzapine. *Journal of Neurophysiology* 115, 1389–1398. doi:10.1152/jn.00387.2015. PMID:
26719086
2 .Badayan, I. and Cudkowicz, M. E. (2009). Profound muscle weakness and pain after one dose of
actonel. *Case Reports in Medicine* 2009, 1–3. doi:10.1155/2009/693014
3 .Bates, S. (2011). The role of gene expression profiling in drug discovery. *Current Opinion in*
Pharmacology 11, 549 – 556. doi:https://doi.org/10.1016/j.coph.2011.06.009. Anti-infectives/ New
technologies
4 .Bendeck, M. P., Conte, M., Zhang, M., Nili, N., Strauss, B. H., and Farwell, S. M. (2002). Doxycycline
modulates smooth muscle cell growth, migration, and matrix remodeling after arterial injury. *The*
American Journal of Pathology 160, 1089–1095. doi:10.1016/s0002-9440(10)64929-2
5 .Bergemann, N., Ehrig, C., Diebold, K., Mundt, C., and v. Einsiedel, R. (1999). Asymptomatic
pancreatitis associated with clozapine. *Pharmacopsychiatry* 32, 78–80. doi:10.1055/s-2007-979197
6 .Burgos, K., Malenica, I., Metpally, R., Courtright, A., Rakela, B., Beach, T., et al. (2014). Profiles of
extracellular mirna in cerebrospinal fluid and serum from patients with alzheimer's and parkinson's
diseases correlate with disease status and features of pathology. *PLOS ONE* 9, 1–20. doi:10.1371/
journal.pone.0094839
7 .Chen, S.-J., Lin, C.-S., Hsu, C.-W., Lin, C.-L., and Kao, C.-H. (2015). Acetaminophen poisoning and
risk of acute pancreatitis. *Medicine* 94, e1195. doi:10.1097/md.0000000000001195
8 .Chengalvala, M. V., , Chennathukuzhi, V. M., Johnston, D. S., Stevis, P. E., and Kopf, G. S. (2007).
Gene expression profiling and its practice in drug development. *Current Genomics* 8, 262–270.
doi:10.2174/138920207781386942
9 .De Wolf, H., Cougnaud, L., Van Hoorde, K., De Bondt, A., Wegner, J. K., Ceulemans, H., et al.
(2018). High-throughput gene expression profiles to define drug similarity and predict compound
activity. *ASSAY and Drug Development Technologies* 16, 162–176. doi:10.1089/adt.2018.845. PMID:
29658791
10 .Duran-Frigola, M., Pauls, E., Guitart-Pla, O., Bertoni, M., Alcalde, V., Amat, D., et al. (2020).
Extending the small molecule similarity principle to all levels of biology. *Nature Biotechnology*
doi:10.1101/745703. In press

.Elsworth, J. D., Groman, S. M., Jentsch, J. D., Valles, R., Shahid, M., Wong, E., et al. (2012). Asenapine effects on cognitive and monoamine dysfunction elicited by subchronic phencyclidine administration. *Neuropharmacology* 62, 1442 – 1452. doi:https://doi.org/10.1016/j.neuropharm.2011.08.026. Schizophrenia

.Ghanem, C. I., Pérez, M. J., Manautou, J., and Mottino, A. D. (2016). Acetaminophen from liver to brain: New insights into drug pharmacological action and toxicity. *Pharmacological Research* 109, 119 – 131. doi:https://doi.org/10.1016/j.phrs.2016.02.020. Country in focus: Pharmacology in Argentina

.h. Taguchi, Y. (2019). Drug candidate identification based on gene expression of treated cells using tensor decomposition-based unsupervised feature extraction for large-scale data. *BMC Bioinformatics* 19. doi:10.1186/s12859-018-2395-8

.h. Taguchi, Y. (2020). *Unsupervised Feature Extraction Applied to Bioinformatics: A PCA Based and TD Based Approach* (Switzerland: Springer International Publishing). doi:10.1007/978-3-030-22456-1

.Harada, A., Ito, S., Matsui, Y., Sakai, Y., Takemura, M., Tokuda, H., et al. (2015). Effect of alendronate on muscle mass: Investigation in patients with osteoporosis. *Osteoporosis and Sarcopenia* 1, 53 – 58. doi:https://doi.org/10.1016/j.afos.2015.07.005

.Hayden, M. R., Grant, D. G., Aroor, A. R., and DeMarco, V. G. (2019). Empagliflozin ameliorates type 2 diabetes-induced ultrastructural remodeling of the neurovascular unit and neuroglia in the female db/db mouse. *Brain Sciences* 9. doi:10.3390/brainsci9030057

.Hodos, R., Zhang, P., Lee, H.-C., Duan, Q., Wang, Z., Clark, N. R., et al. (2017). Cell-specific prediction and application of drug-induced gene expression profiles. In *Biocomputing 2018* (WORLD SCIENTIFIC). doi:10.1142/9789813235533_0004

.Huang, C.-T., Hsieh, C.-H., Chung, Y.-H., Oyang, Y.-J., Huang, H.-C., and Juan, H.-F. (2019). Perturbational gene-expression signatures for combinatorial drug discovery. *iScience* 15, 291 – 306. doi:https://doi.org/10.1016/j.isci.2019.04.039

.Hung, W. Y. (2014). Contemporary review of drug-induced pancreatitis: A different perspective. *World Journal of Gastrointestinal Pathophysiology* 5, 405. doi:10.4291/wjgp.v5.i4.405

.Huo, X., Reyes, T. M., Heijnen, C. J., and Kavelaars, A. (2018). Cisplatin treatment induces attention deficits and impairs synaptic integrity in the prefrontal cortex in mice. *Scientific Reports* 8. doi:10.1038/s41598-018-35919-x

.Jordan, J., Tank, J., Heusser, K., Heise, T., Wanner, C., Heer, M., et al. (2017). The effect of empagliflozin on muscle sympathetic nerve activity in patients with type ii diabetes mellitus. *Journal of the American Society of Hypertension* 11, 604 – 612. doi:https://doi.org/10.1016/j.jash.2017.07.005

.Kakaria, P. J., Nashel, D. J., and Nylen, E. S. (2005). Debilitating Muscle Cramps after Teriparatide Therapy. *Annals of Internal Medicine* 142, 310–310. doi:10.7326/0003-4819-142-4-200502150-00023

.Kerr, T. A., Jonnalagadda, S., Prakash, C., and Azar, R. (2007). Pancreatitis following olanzapine therapy: A report of three cases. *Case Reports in Gastroenterology* 1, 15–20. doi:10.1159/000104222

.Keyal, N., Shrestha, G., Pradhan, S., Maharjan, R., Acharya, S., and Marhatta, M. (2017). Olanzapine overdose presenting with acute muscle toxicity. *International Journal of Critical Illness and Injury Science* 7, 69–71. doi:10.4103/2229-5151.201962

.Kim, I.-W., Jang, H., Kim, J. H., Kim, M. G., Kim, S., and Oh, J. M. (2019). Computational drug repositioning for gastric cancer using reversal gene expression profiles. *Scientific Reports* 9. doi:10.1038/s41598-019-39228-9

.Kiraly, B. and Gunning, K. (2008). A case of pancreatitis associated with aripiprazole in the absence of hyperglycemia. *The Primary Care Companion to The Journal of Clinical Psychiatry* 10, 484–485. doi:10.4088/pcc.v10n0612e

.Kishimoto, M., Yamaoki, K., and Adachi, M. (2019). Combination therapy with empagliflozin and insulin results in successful glycemic control: A case report of uncontrolled diabetes caused by autoimmune pancreatitis and subsequent steroid treatment. *Case Reports in Endocrinology* 2019, 1–8. doi:10.1155/2019/9415347

.Kozawa, S., Sagawa, F., Endo, S., Almeida, G. M. D., Mitsuishi, Y., and Sato, T. N. (2020). Predicting human clinical outcomes using mouse multi-organ transcriptome. *iScience* , 100791doi:10.1016/j.isci.

2019.100791
29 .[Dataset] Kuleshov, M., Kropiwnicki, E., and Ma'ayan, A. (2020). Drugenrichr
30 .Kuleshov, M. V., Jones, M. R., Rouillard, A. D., Fernandez, N. F., Duan, Q., Wang, Z., et al. (2016).
Enrichr: a comprehensive gene set enrichment analysis web server 2016 update. *Nucleic Acids Research*
44, W90–W97. doi:10.1093/nar/gkw377
31 .Lee, B. K. B., Tiong, K. H., Chang, J. K., Liew, C. S., Rahman, Z. A. A., Tan, A. C., et al. (2017).
DeSigN: connecting gene expression with therapeutics for drug repurposing and development. *BMC*
Genomics 18. doi:10.1186/s12864-016-3260-7
32 .Li, C. H., Stratford, R. E., de Mendizabal, N. V., Cremers, T. I., Pollock, B. G., Mulsant, B. H.,
et al. (2014). Prediction of brain clozapine and norclozapine concentrations in humans from a scaled
pharmacokinetic model for rat brain and plasma pharmacokinetics. *Journal of Translational Medicine*
12. doi:10.1186/1479-5876-12-203
33 .Liu, X., Zeng, P., Cui, Q., and Zhou, Y. (2017). Comparative analysis of genes frequently regulated by
drugs based on connectivity map transcriptome data. *PLOS ONE* 12, 1–13. doi:10.1371/journal.pone.
0179037
34 .Lucchetti, J., Fracasso, C., Balducci, C., Passoni, A., Forloni, G., Salmona, M., et al. (2018). Plasma
and brain concentrations of doxycycline after single and repeated doses in wild-type and app23 mice.
Journal of Pharmacology and Experimental Therapeutics doi:10.1124/jpet.118.252064
35 .Lukačišin, M. and Bollenbach, T. (2019). Emergent gene expression responses to drug combinations
predict higher-order drug interactions. *Cell Systems* 9, 423–433.e3. doi:10.1016/j.cels.2019.10.004
36 .Margapuri, J. and Jubbal, S. (2019). 902: Acute pancreatitis following treatment with
ledipasvir/sofosbuvir for hepatitis virus infection. *Critical Care Medicine* 47, 430. doi:10.1097/
01.ccm.0000551651.45733.80
37 .Myrick, H., Li, X., Randall, P. K., Henderson, S., Voronin, K., and Anton, R. F. (2010). The effect of
aripiprazole on cue-induced brain activation and drinking parameters in alcoholics. *Journal of Clinical*
Psychopharmacology 30, 365–372. doi:10.1097/jcp.0b013e3181e75cff
38 .Nissen, S. E., Stroes, E., Dent-Acosta, R. E., and for the GAUSS-3 Investigators (2016). Efficacy
and Tolerability of Evolocumab vs Ezetimibe in Patients With Muscle-Related Statin Intolerance: The
GAUSS-3 Randomized Clinical Trial. *JAMA* 315, 1580–1590. doi:10.1001/jama.2016.3608
39 .Oliveira, J. R. M. and Oliveira, M. F. (2016). Primary brain calcification in patients undergoing
treatment with the biphosphanate alendronate. *Scientific Reports* 6. doi:10.1038/srep22961
40 .Pabon, N. A., Xia, Y., Estabrooks, S. K., Ye, Z., Herbrand, A. K., Evelyn, S., et al. (2018). Predicting
protein targets for drug-like compounds using transcriptomics. *PLOS Computational Biology* 14, 1–24.
doi:10.1371/journal.pcbi.1006651
41 .Patel, S., Trakroo, S., Sanaka, S., and Qureshi, K. (2015). Severe myositis with the use of
sofosbuvir/ledipasvir for hepatitis c infection: A case of unexpected interactions. *American Journal of*
Gastroenterology 110, S333. doi:10.14309/00000434-201510001-00761
42 .Raja, V., Sandanshiv, P., and Neugebauer, M. (2007). Risedronate induced transient ocular myasthenia.
Journal of Postgraduate Medicine 53, 274. doi:10.4103/0022-3859.37525
43 .Rawla, P. and Raj, J. P. (2017). Doxycycline-induced acute pancreatitis: A rare adverse event.
Gastroenterology Research 10
44 .Reece, D., Kouroukis, C. T., LeBlanc, R., Sebag, M., Song, K., and Ashkenas, J. (2012). Practical
approaches to the use of lenalidomide in multiple myeloma: A canadian consensus. *Advances in*
Hematology 2012, 1–14. doi:10.1155/2012/621958
45 .Reznik, I., Volchek, L., Mester, R., Kotler, M., Sarova-Pinhas, I., Spivak, B., et al. (2000). Myotoxicity
and neurotoxicity during clozapine treatment. *Clinical Neuropharmacology* 23, 276–280. doi:10.1097/
00002826-200009000-00007
46 .Rollin-Sillaire, A., Delbeuck, X., Pollet, M., Mackowiak, M.-A., Lenfant, P., Noel, M.-P., et al. (2013).
Memory loss during lenalidomide treatment: a report on two cases. *BMC Pharmacology and Toxicology*
14. doi:10.1186/2050-6511-14-41

.Sakai, H., Sagara, A., Arakawa, K., Sugiyama, R., Hirosaki, A., Takase, K., et al. (2014). Mechanisms of cisplatin-induced muscle atrophy. *Toxicology and Applied Pharmacology* 278, 190 – 199. doi:https://doi.org/10.1016/j.taap.2014.05.001

.Salman, A. G. (2016). Ocular surface changes with sofosbuvir in egyptian patients with hepatitis c virus infection. *Cornea* 35, 323–328. doi:10.1097/ico.0000000000000736

.Smirnov, P., Kofia, V., Maru, A., Freeman, M., Ho, C., El-Hachem, N., et al. (2017). PharmacoDB: an integrative database for mining in vitro anticancer drug screening studies. *Nucleic Acids Research* 46, D994–D1002. doi:10.1093/nar/gkx911

.Taguchi, Y.-h. and Turki, T. (2020). Universal nature of drug treatment responses of drug-tissue-wide model-animal experiments. *bioRxiv* doi:10.1101/2020.03.08.982405

.Trappe, T. A., Carroll, C. C., Dickinson, J. M., LeMoine, J. K., Haus, J. M., Sullivan, B. E., et al. (2011). Influence of acetaminophen and ibuprofen on skeletal muscle adaptations to resistance exercise in older adults. *American Journal of Physiology-Regulatory, Integrative and Comparative Physiology* 300, R655–R662. doi:10.1152/ajpregu.00611.2010. PMID: 21160058

.Ullenhag, G. J., Mozaffari, F., Broberg, M., Mellstedt, H., and Liljefors, M. (2017). Clinical and immune effects of lenalidomide in combination with gemcitabine in patients with advanced pancreatic cancer. *PLOS ONE* 12, 1–19. doi:10.1371/journal.pone.0169736

.Vinogradova, Y., Coupland, C., and Hippisley-Cox, J. (2013). Exposure to bisphosphonates and risk of common non-gastrointestinal cancers: series of nested case–control studies using two primary-care databases. *British Journal of Cancer* 109, 795–806. doi:10.1038/bjc.2013.383

.Yadav, Y. C. (2019). Effect of cisplatin on pancreas and testies in wistar rats: biochemical parameters and histology. *Heliyon* 5, e02247. doi:10.1016/j.heliyon.2019.e02247

.Yasui-Furukori, N. (2012). Update on the development of lurasidone as a treatment for patients with acute schizophrenia. *Drug Design, Development and Therapy* , 107doi:10.2147/dddt.s11180

.Zhou, Y., Zhou, B., Pache, L., Chang, M., Khodabakhshi, A. H., Tanaseichuk, O., et al. (2019). Metascape provides a biologist-oriented resource for the analysis of systems-level datasets. *Nature Communications* 10. doi:10.1038/s41467-019-09234-6

FIGURES

Figure 1. Schematic of the analysis performed in this study

Figure 2. Summary of the results obtained in this study

Figure 3. Scatter plot between u_{2k} and u_{3k} attributed to drug treatment. Red and blue arrows represent distinct controls and drug treatments, and diversity among drug treatments, respectively. (1) Alendronate, (2) APAP, (3) Aripiprazole, (4) Asenapine, (5) Cisplatin, (6) Clozapine, (7) Dox, (8) EMPA, (9) FivePercentSucrose, (10) Lenalidomide, (11) Lurasidone, (12) Olanzapine, (13) Repatha, (14) Risedronate, (15) Sofosbuvir, (16) Teriparatide, (17) WT.No.treated, (18) 5percentCMC0.25percentTween80.

Figure 4. Singular value vectors, $u_{\ell_1 j}$, attributed to tissues. u_{1j}: no tissue specificity. u_{2j}: Brain Eye, Pituitary, and Testis, thus mostly neuron-specific. u_{3j}: Parotid-specific, u_{4j}: Heart and SkMuscle, thus muscle-specific, u_{5j} and u_{6j}: stomach and pancreas, thus, gastrointestinal-specific.

Figure 5. Venn diagram of genes selected by TD-based unsupervised FE. Neuron: genes associated with u_{2j}, which is supposed to be neuron-specific. Muscle: genes associated with u_{4j}, which is supposed to be muscle-specific. Gas1 and Gas2: genes associated with u_{5j} and u_{6j} respectively, which is supposed to be gastrointestinal-specific.

Figure 6. Heatmap of enrichment analysis provided by Metascape.

Figure 7. PPI network provided by Metascape. Red: Gas1, Blue: Gas2, Green: Muscle, Purple: Neuron.

TABLES

Table 1. $G(\ell_1, 2, 1, \ell_4)$ and $G(\ell_1, 3, 1, \ell_4)$ for $\ell_1 = 2, 4, 5, 6$. Values in bold correspond to those of ℓ_4s used for gene selection with $u_{\ell_4 i}$.

ℓ_1	2		4	
ℓ_4	$G(2,2,1,\ell_4)$	$G(2,3,1,\ell_4)$	$G(4,2,1,\ell_4)$	$G(4,3,1,\ell_4)$
1	131.248442	19.7819438	-98.4349019	-13.498228
2	**-173.243689**	**-23.9915660**	-4.8528076	1.113899
3	-11.859736	-3.2551088	-0.1595594	-1.116396
4	13.669561	2.4373120	**-81.3734282**	**36.838277**
5	26.610843	-0.3136913	-22.2440356	9.820737
6	-1.275395	4.5339065	-1.3753621	-5.318282
7	-18.306263	15.9791077	21.3673134	-11.230437
8	20.891762	26.5918473	3.9733331	-7.152480
9	21.836494	16.1461476	9.2972447	2.232529
10	11.717415	-12.8960548	1.4137802	-7.748038

ℓ_1	5		6	
ℓ_4	$G(5,2,1,\ell_4)$	$G(5,3,1,\ell_4)$	$G(6,2,1,\ell_4)$	$G(6,3,1,\ell_4)$
1	97.897860	-42.9481806	72.181307	27.3218396
2	9.267391	4.5503920	3.780984	6.3881436
3	-3.744432	0.2003586	2.340165	-0.2130656
4	1.648558	3.4031386	-9.812308	2.8751439
5	**93.027741**	**-56.9322793**	6.435061	8.5776220
6	-57.463765	23.2247109	**-19.332916**	**34.1868710**
7	28.276681	-26.9479131	**30.604535**	**-18.8319412**
8	12.884351	-13.8270607	1.798188	-10.6484624
9	-5.865058	1.0216563	9.581512	0.3507831
10	15.683762	3.7893181	-14.429706	-4.7985105

Table 2. Statistical tests for distinct expression between the specified tissues and other tissues, and between drug treatments and controls.

				P values by statistical tests		
				Tissues	Drug treatment	
ℓ_1	tissue specificity	# of genes	Specified tissues	t test	Wilcoxon test	t test
2	Neuron	18	Brain, Eye, Pituitary, Testis	2.14×10^{-24}	9.65×10^{-49}	0.22
4	Muscle	51	Heart, SkMuscle	1.99×10^{-55}	2.67×10^{-77}	0.04
5	Gastrointestinal	97	Pancreas, Stomach	8.48×10^{-11}	2.73×10^{-40}	8.13×10^{-22}
6		128		6.67×10^{-8}	8.69×10^{-90}	8.69×10^{-90}

Table 3. Enrichment analysis for "Disease Perturbations from GEO up" and "Disease Perturbations from GEO down" by Enrichr. Diseases in bold correspond to those related to specific tissues. Up to the top 10 ranked terms are shown.

Disease Perturbations from GEO up			
Term	Overlap	P-value	Adjusted P-value
Neuron-specific genes			
amyotrophic lateral sclerosis DOID-332 mouse GSE3343 sample 685	5/138	1.16×10^{-7}	9.72×10^{-5}
Retinitis Pigmentosa C0035334 mouse GSE128 sample 33	5/338	9.57×10^{-6}	4.01×10^{-2}
Muscle specific			
Polycystic Ovary Syndrome C0032460 human GSE6798 sample 292	21/306	2.87×10^{-25}	2.41×10^{-22}
polycystic ovary syndrome DOID-11612 human GSE8157 sample 880	21/325	1.03×10^{-24}	4.32×10^{-22}
Insulin Resistance DOID-9352 human GSE36297 sample 581	20/290	4.52×10^{-24}	1.26×10^{-21}
Neurogenic Muscular Atrophy C0270948 rat GSE2566 sample 396	18/208	1.98×10^{-23}	4.15×10^{-21}
Nemaline myopathy DOID-3191 mouse GSE3384 sample 976	15/150	1.65×10^{-20}	2.77×10^{-18}
psoriasis DOID-8893 mouse GSE27628 sample 822	18/346	2.06×10^{-19}	2.88×10^{-17}
Nemaline Myopathy C0206157 mouse GSE3384 sample 160	16/276	5.20×10^{-18}	6.24×10^{-16}
Muscular Dystrophy C0026850 mouse GSE2507 sample 405	16/278	5.84×10^{-18}	6.13×10^{-16}
cystic fibrosis DOID-1485 mouse GSE3100 sample 1057	17/344	5.91×10^{-18}	5.51×10^{-16}
COPD - Chronic obstructive pulmonary disease C0024117 human GSE475 sample 343	16/289	1.09×10^{-17}	9.11×10^{-16}
Disease Perturbations from GEO down			
Term	Overlap	P-value	Adjusted P-value
Gas1 genes			
pancreatitis DOID-4989 mouse GSE3644 sample 513	36/238	9.21×10^{-45}	7.73×10^{-42}
Skin squamous cell carcinoma DOID-3151 human GSE2503 sample 627	37/373	5.24×10^{-39}	2.20×10^{-36}
Pancreatic ductal adenocarcinoma DOID-3498 mouse GSE53659 sample 699	26/101	1.24×10^{-38}	3.48×10^{-36}
Pancreatic invasive intraductal papillary-mucinous carcinoma DOID-8150 human GSE19650 sample 610	31/248	1.21×10^{-35}	2.54×10^{-33}
Cystic fibrosis DOID-1485 mouse GSE769 sample 1058	32/288	3.92×10^{-35}	6.58×10^{-33}
Acute pancreatitis C0001339 mouse GSE3644 sample 376	28/188	2.33×10^{-34}	3.26×10^{-32}
Cystic Fibrosis C0010674 mouse GSE769 sample 428	31/275	3.34×10^{-34}	4.00×10^{-32}
Chronic phase chronic myelogenous leukemia DOID-8552 human GSE5550 sample 456	30/270	7.05×10^{-33}	7.40×10^{-31}
Invasive ductal carcinoma DOID-3008 human GSE21422 sample 606	31/304	8.09×10^{-33}	7.54×10^{-31}
Eczema C0013595 human GSE6012 sample 268	26/163	1.06×10^{-32}	8.87×10^{-31}
Gas2 genes			
Skin squamous cell carcinoma DOID-3151 human GSE2503 sample 627	51/373	9.37×10^{-55}	7.86×10^{-52}
Pancreatitis DOID-4989 mouse GSE3644 sample 513	45/238	1.14×10^{-54}	4.77×10^{-52}
Systemic lupus erythematosus DOID-9074 human GSE10325 sample 691	43/210	9.36×10^{-54}	2.62×10^{-51}
Systemic lupus erythematosus (SLE) DOID-9074 human GSE36700 sample 512	47/294	1.50×10^{-53}	3.15×10^{-51}
Invasive ductal carcinoma DOID-3008 human GSE21422 sample 606	44/304	5.90×10^{-48}	9.90×10^{-46}
Eczema C0013595 human GSE6012 sample 268	37/163	7.47×10^{-48}	1.04×10^{-45}
Malignant Melanoma C0025202 human GSE3189 sample 117	41/250	6.75×10^{-47}	8.09×10^{-45}
Chronic phase chronic myelogenous leukemia DOID-8552 human GSE5550 sample 456	41/270	1.91×10^{-45}	2.00×10^{-43}
Sickle Cell Anemia C0002895 human GSE9877 sample 109	37/197	1.61×10^{-44}	1.50×10^{-42}
Actinic keratosis C0022602 human GSE2503 sample 350	46/429	4.69×10^{-44}	3.94×10^{-42}

Table 4. Previously reported drug effects on neuron (brain and eye), muscle and pancreas tissues. (*): Reported side effects.

Drugs	Tissue types		
	Neuron	Muscle	Pancreas or Stomach
Alendronate	Brain calcification (39)	Muscle mass (15)	Pancreatitis (19)
Acetaminophen (APAP)	Brain (12)	Skeletal muscle (51)	Pancreatitis (7)
Aripiprazole	Brain Activation (37)	Muscle spasms (*)	Pancreatitis (26)
Asenapine	Cognitive and monoamine dysfunction (11)	Muscle rigidity(*)	—
Cisplatin	Prefrontal cortex (20)	Muscle atrophy (47)	Pancreas (54)
Clozapine	Brain (32)	Myotoxicity (45)	Pancreatitis (5)
Doxycycline	Brain (34)	Smooth Muscle (4)	Acute pancreatitis (43)
Empagliflozin	Neurovascular unit and neuroglia (16)	Muscle sympathetic nerve activity (21)	Pancreatitis (27)
Lenalidomide	Memory loss (46)	Muscle cramp (44)	Pancreatic cancer (52)
Lurasidone	Acute schizophrenia (55)	Muscle(*)	—
Olanzapine	Brain stem (1)	Acute muscle toxicity (24)	Pancreatitis (23)
Repatha (Evolocumab)	—	Muscle-related statin Intolerance (38)	—
Risedronate (actonel)	Ocular myasthenia (42)	Muscle weakness (2)	Gastrointestinal cancer (53)
Sofosbuvir	Ocular surface (48)	Myositis (41)	Pancreatitis (36)
Teriparatide	—	Muscle Cramp (22)	Pancreatitis (*)

ADDITIONAL FILES

395 **Additional file 1 — List of genes selected by TD-based unsupervised FE**

396   List of genes shown in Table 2 (xlsx)

398 **Additional file 2 — List of genes predicted by DrugEnrichr, whose expression is likely**

399 **affected by the drugs investigated in this study**

400   List of genes (xlsx)