Tests of hybridisation in *Tetragonula* stingless bees using multiple
genetic markers.

Authors: James P. Hereward[1], Tobias J. Smith[1], Dean R. Brookes[1], Ros Gloag[2], Gimme H.
Walter[1].

Affiliations:

[1]School of Biological Sciences, The University of Queensland, Brisbane, Queensland 4072,

Australia.

[2]School of Life and Environmental Sciences, The University of Sydney, Sydney, New South

Wales 2006, Australia.

Corresponding author: James Hereward, School of Biological Sciences, The University of

Queensland, Brisbane, Queensland 4072, Australia

E-mail: j.hereward@uq.edu.au

Keywords: Hybridisation, gene flow, species, genetic marker, stingless bee

Abstract

Discrepancies in mitochondrial and nuclear genetic data are often interpreted as evidence of hybridisation. We re-examined reports of hybridisation in three cryptic stingless bee species in the genus *Tetragonula* in South East Queensland, Australia (*T. carbonaria*, *T. davenporti*, and *T. hockingsi*). Previous studies on this group using microsatellites proposed that rare hybrids do occur. In contrast, we find that allele frequencies at neutral regions of the nuclear genome (both microsatellites and random *SNPs*) reliably separated the three species and revealed no evidence of hybridisation. We found no inter-species variation in the nuclear gene *EF1alpha*, but low and moderate species-specific polymorphisms in the nuclear gene *Opsin* and the mitochondrial *16S* respectively, with no cases of mito-nuclear discord at these genes. We confirm that nuclear divergence between these species is low, based on 10-26kb of non-coding sequence flanking *EF1alpha* and *Opsin* (0.7-1% pairwise difference between species). However, we find mitogenomes to be far more diverged than nuclear genomes (21.6-23.6% pairwise difference between species). Based on these comprehensive analyses of multiple marker types, we conclude that there is no ongoing gene flow in the *Tetragonula* species of South East Queensland, despite their high morphological similarity and low nuclear divergence. The mitochondrial genomes indicate that divergence could be deep between these lineages but the low divergence in the nuclear genomes means that further investigation is required before divergence times can be estimated accurately. The mitogenomes, and draft nuclear genomes provided for these species will be a resource for further molecular studies on this group.

Introduction

Allopatry is a potent driver of speciation, but species boundaries may be tested when populations that have diverged in allopatry come back into secondary contact due to changes in habitat or climate, or as a result of dispersal by humans over geographic boundaries (Mallet *et al.*, 2011; Sánchez-Guillén *et al.*, 2013). The process of hybridisation is closely tied to species concepts and their designation in population genetic terms (Paterson 1978); by definition, hybridisation cannot occur within a species. When species are considered conceptually as recombining species gene pools (Ayala *et al.* 1994), hybridisation is a process of gene flow across two such species gene pools. High levels of 'hybridisation' in local areas of overlap indicate that the 'species' in question are subspecies, which, by definition, have independent geographical distributions but hybridise freely where and when they do overlap (Walter 2003, Ford 1974). Such situations are usually seen when closely-related but divergent lineages of the same species are brought back into sympatry (Feder *et al.* 2003), but sometimes this pattern is seen across more distantly related lineages (e.g. Shipman *et al.* 2019). The frequency of hybridisation, or gene flow, between gene pools is therefore key to understanding particular species and their limits, and is key to our understanding of speciation.

Understanding hybridisation relies not only on our concept of species generally and on definitions of particular species, but also on the methods that are available to detect it. Accurate assessment of hybridisation is the first step in shedding light on the mechanisms that might prevent gene flow between diverged populations. The more recently two populations have diverged, the more likely they have not yet become separate species, and thus the more likely that gene flow is re-established (Feder *et al.* 2003). Hybrid detection between close-relatives is expected to be especially sensitive to the molecular methodologies used (e.g. number and type

of markers), and more likely to be susceptible to sampling error because parent species are

similar or cryptic in appearance. A species complex of stingless bees (Meliponini) in the genus

Tetragonula from Australia illustrates this well.

The stingless bee (Meliponini) genus *Tetragonula* includes three cryptic sister species

(*T. carbonaria. T. hockingsi* and *T. davenporti*) native to eastern Australia that are part of a

monophyletic clade known as the "Carbonaria group" (Franck *et al*., 2004). These bees can be

readily kept in wooden hives, are popular pets, and are effective pollinators of some tropical

and subtropical crops (Dyer *et al.* 2016, Heard 1994), leading to many thousands of colonies

kept by beekeepers in Eastern Australia. Today, the distributions of all three species overlap in

parts of south-east Queensland (Fig 1). However, the historical distributions of the species are

poorly known due to their cryptic morphology. It is likely that beekeeping and other human

activities, such as land use change, have greatly increased the overlap of their distributions in

recent decades, particularly via the reported southward spread of *T. hockingsi*. Queens, males

and workers (females) cannot be reliably identified based on morphology alone (Dollin *et al.*

1997). Rather, species are typically identified by the structure of their brood comb, with *T.*

carbonaria producing a neat spiral of adjacent brood cells and *T. hockingsi* producing clusters

of cells in a form known as semi-comb (Fig. 2). *T. davenporti* shares the semi-comb brood

structure of *T. hockingsi*, and is known from only a handful of managed colonies in a sub-

coastal pocket of southern Queensland (Franck *et al*. 2004, Brito *et al.* 2014). No behavioural

barriers to reproduction between the species are known. Indeed, male mating congregations,

which form outside colonies with virgin queens, can contain a mix of both *T. carbonaria* and

T. hockingsi males (Cunningham *et al.* 2014). These two species are also known to engage in

interspecific nest usurpations, in which an invading colony kills a resident queen and installs

its own in the nest (Cunningham *et al.* 2014), a behaviour which may increase the chances that

males are attracted to interspecific queens.

Several previous efforts have been made to establish if, and how regularly, gene flow

occurs between these species, with equivocal results. First, microsatellite markers have been

used to look for evidence of hybrids. Frank *et al.* (2004) identified some Queensland colonies

of *Tetragonula* as being of hybrid origin based on microsatellite data and defined as a sample

having a posterior probability of assignment of less than 90% to a cluster in the *STRUCTURE*

(Pritchard *et al.* 2000) analysis. Franck *et al.* (2004) also attempted to characterise eight other

species of *Tetragonula* genetically, from Australia and Thailand, based on the *cytochrome b*

(*cytb*) mitochondrial gene, but the outcome was poor because they detected issues with

pseudogenes. The same three Australian *Tetragonula* species were re-investigated by Brito *et*

al. (2014), with microsatellite data and two mitochondrial genes. Three instances of

hybridisation in southeast Queensland were reported (within the distribution overlap zone of

T. carbonaria and *T. hockingsi*), using a cut-off of 95% posterior probability of belonging to a

cluster in a *STRUCTURE* analysis of the microsatellite data. Brito *et al*. (2014) also included

two hives reported to be *T. davenporti* in their study. One of the *T. davenporti* hives clustered

with *T. carbonaria* in a Principal Coordinate Analysis (PCoA) of genetic distance based on the

microsatellite data, and the other clustered slightly outside of the '*T. carbonaria*' cluster. In

interpreting these results, Brito *et al.* (2014) suggested that *T. hockingsi* represents two genetic

clusters (based on their microsatellite and mitochondrial data), one each in northern and

southern Queensland. Their *T. hockingsi x T. carbonaria* hybrids were from South East

Queensland and they suggested that this hybridisation was due to the anthropogenic movement

of northern *T. hockingsi* into the southern region.

Second, both nuclear and mitochondrial markers have been used to look for evidence

of introgression and thus past interbreeding, but nuclear copies of mitochondrial genes (numts)

have regularly hampered these attempts. Most recently, several numts of Cytochrome Oxidase

Subunit I (*COI*) were identified in *T. hockingsi* and *T. carbonaria* (Françoso *et al*. 2019). This

cast into question earlier work on hybrids that incorporated COI-numts, and even the

designation of *T. davenporti* as a true species. Surprisingly, the genuine COI sequences of these

species are highly diverged (16.5 %) relative to available nuclear genes (Cunningham 2014,

Francosco 2019). This pattern of high mitochondrial divergence but low nuclear divergence

might be consistent with recent or ongoing gene flow, and further shows that new analyses are

needed to understand the extent to which species in this complex are discrete, and the potential

impact of continuing movement of hives around Australia.

Here we aim to resolve whether there is evidence of ongoing gene flow between *T.*

carbonaria, T. hockingsi and *T. davenporti* kept by beekeepers in sympatry in South East

Queensland, using both microattelites and a *genotyping by sequencing* approach that assayed

2,049 *snp* markers. We also look for evidence of mitochondrial introgression, by considering

two nuclear gene sequences and one mitochondrial gene sequence of each species, and use

Illumina sequencing to reconstruct mitochondrial genomes and thus confirm that mitochondrial

sequences are not pseudogenes. Finally, we assembled draft nuclear genomes for these three

species as a resource for further genetic analysis of the species complex. This comprehensive

approach leads to new insights about the extent of hybridisation in these bees, and their species

status.

Materials and Methods

Sampling and DNA extraction

We sampled four workers per colony from hived colonies of each species between February 2017 and April 2019 in south-east Queensland (*T. carbonaria*: N=39, *T. hockingsi*, N=21, *T. davenporti*, N=3); (Fig 1). Most *T. carbonaria* and *T. hockingsi* colonies were sampled in the greater Brisbane region. We opened each hive and assigned it to species based on brood nest architecture (spiral comb or semi-comb). The three *T. davenporti* colonies were sampled from the border area of Queensland and New South Wales and were the same colonies used for past genetic work on this species (P. Davenport, pers. comm. 2019). A further two *T. carbonaria* colonies and one *T. hockingsi* colony were sampled from the same meliponary as the three *T. davenporti* colonies. We extracted DNA using a silica spin-column method (Ridley *et al* 2016) and normalized concentrations to 5ng/ul per sample following quantification (PicoGreen, Life Technologies, California, USA).

Neutral nuclear markers

Microsatellites: We first assessed species status and evidence of hybridisation using seven microsatellite loci (Green *et al.* 2001), as used in previous molecular studies of this group (Franck *et al.* 2004, Brito *et al.* 2014). We used PCR reactions of 2 µl DNA template, 0.5 U MyTaq polymerase (Bioline, Australia), 0.1 µM of forward primer, 0.2 µM of reverse primer, 0.2 µM M13 labelled primer with different fluorescent dyes (6-FAM, VIC, PET or NED) and 1x buffer, with reaction conditions of an initial denaturation at 95 °C for 3 min, followed by 10 cycles of 10 s at 95 °C, annealing at 45 °C for 25 s, and 1 min extension at 72 °C, then 40 cycles of 10 s at 95 °C, annealing at 55 or 52°C for 25 s, and 1 min extension at 72 °C, and the final extension was at 72 °C for 2 min (see Cunningham *et al.* (2014) for locus-specific annealing temperatures). Loci were then pooled according to dye, cleaned with 1U

Exonuclease I and one unit of Antarctic Phosphatase as above, and genotyped by Macrogen

Inc. (Seoul, Republic of Korea). Microsatellite peaks were analysed using the microsatellite

plugin in Geneious v. 2019.2.1 (http://www.geneious.com, Kearse *et al*. (2012)).

single nucleotide polymorphisms: We generated genome-wide *snp* data using a genotyping-by-

sequencing method. We adapted a protocol and adaptor regime based on the methods of Elshire

et al. (2011), Poland *et al.* (2012) and Petersen *et al*. (2012), with barcodes based on Caporaso

et al. (2012). Full details of the method are provided in supplementary data and can be found

online at http://www.jameshereward.org/GBS.html. We pooled 288 individuals per sequencing

lane, and sequenced the libraries with PE150 Illumina sequencing at Novogene (Beijing,

China).

The sequence data were demultiplexed, assembled, and *snps* called using STACKS

(Catchen *et al* 2013). We then filtered the vcf using vcftools, with the data first being filtered

to a minor allele count of three (one heterozygote and one homozygote), which allows the

conservative removal of singleton *snps* that are likely to be errors, without discarding rare

alleles (Linck & Battey 2019). We set a minimum depth of 5, and kept only biallelic *snps*. We

filtered the data for missing data in three steps. First, any marker missing more than 50% data

was discarded (i.e. 50% of individuals were not genotyped at that marker), to remove the

markers most affected by missing data. Second, any individual that had missing data at more

than 50% of the markers was discarded, to remove the individuals that had bad quality

genotyping. Finally, any marker missing more than 5% data was discarded to produce a final

dataset with relatively little missing data (~3%).

Analyses: For both our microsatellite and *snp* datasets, we performed principal component

analyses (PCA) using the adegenet package in R (Jombart, 2008) and *STRUCTURE* analyses

(Pritchard 2000). *STRUCTURE* is an individual-based clustering algorithm that assigns

individuals to each of *K* population clusters using Hardy-Weinberg equilibrium and linkage

information in genetic markers. To assess the effect of replication and missing data on

hybridisation inferences from the *STRUCTURE* output, we repeated the analyses on three

versions of each of our two datasets: (i) a dataset including one worker per colony, (ii) a dataset

including all four workers per colony, and (iii) the dataset with 20% randomly deleted. For

the microsatellite data, data was removed evenly across markers until 20% missing was

reached, for *snps* we changed the missing data threshold in the filtering step to 20%, and this

increased the number of markers to 13,409 loci. For each *STRUCTURE* run, we used the

admixture model and ran 500,000 iterations of burn-in followed by 1 million iterations. We

assumed both two or three populations (K=2 and K=3) and performed multiple runs that were

then permuted and plotted using Clumpak server (Kopelman *et al.* 2015).

Gene sequencing

To confirm whether common marker genes showed species-specific polymorphisms,

we sequenced two nuclear genes (*EF1alpha* and *Opsin*) and one mitochondrial gene (16S).

These genes were selected as they have previously been used for a phylogenetic study of

stingless bees (Rasmussen and Cameron, 2009). For *EF1alpha* we used primers F2-ForH

(GGRCAYAGAGATTTCATCAAGAAC) and F2-RevH2

(TTGCAAAGCTTCRKGATGCATTT) (Hines 2006). For *Opsin* we used LWRhF

(AATTGCTATTAYGARACNTGGGT) and LWRhR

(ATATGGAGTCCANGCCATRAACCA) (Mardulyn & Cameron 1999). For 16S we used

primers 874-16S1R (TATAGATAGAAACCAAYCTG) (Cameron 1992) and 16SWb

(CACCTGTTTATCAAAAACAT) (Dowtown & Austin 1992). We performed PCRs in 25µl

reactions containing 2 µl DNA template, 1U MyTaq polymerase (Bioline, Australia), 0.2 µM

of each PCR primer, and 1x buffer. For amplification of *EF1alpha* and 16S we performed PCR

with 35 cycles at 52ºC annealing, and for *Opsin* we used five cycles at 45ºC annealing followed

by 35 cycles at 50ºC. PCR products were cleaned using 1U of Exonuclease I and Antarctic

Phosphatase (New England Biolabs, Ipswich, Mass., USA) and incubating at 37 °C for 20 min

followed by 10 min enzyme denaturation at 80 °C. Sequencing was conducted using the same

forward and reverse primers used for PCR, by Macrogen Inc. (Seoul, Republic of Korea). We

edited sequence data with CodonCode aligner, and aligned them using MAFFT in Geneious

Prime 2019.2.1 (www.geneious.com, Kearse *et al*. 2012). We generated haplotype networks

using the TCS algorithm (Clement *et al.* 2000) in PopART version 1.7

(http://popart.otago.ac.nz/) (Leigh and Bryant 2015).

Mitogenome sequencing and assembly

The carbonaria species complex of Tetragonula are now known to have several

pseudogenes (or "numts", nuclear insertions of mitochondrial genes) (Françoso *et al.* 2019; see

also Franck *et al.* 20014, Brito *et al.* 2014). To ensure we were amplifying only the true 16S

gene, and to provide a reference genome for each species for future work, we conducted whole

genome sequencing on one individual of each of the three species to reconstruct the complete

mitochondrial sequence. Sequencing libraries were constructed using the NebNext UltraII

DNA kit for Illumina (New England Biolabs, Ipswich, Mass., USA), and PE150 sequencing

conducted by Novogene (Beijing, China).

Sequences were checked with FastQC (Andrews, 2019), then we removed adaptors,

and quality-trimmed the end of the sequences of all samples to q10 using bbduk from the

bbtools package (version 36, Bushnell, 2019), with a kmer of 8. We binned the sequences by

depth using a kmer-based approach in bbnorm (Bushnell, 2019). We put everything under 500x

coverage into one file (this includes the nuclear genome which was between 23x and 51x

coverage for each species, Table 1), everything between 500x and 3,000x into another file (this

is repetitive sequence and other artefacts below the depth of the mitochondria), everything over

3,000x coverage was placed in a third file, and this contained mitochondrial sequence and

anything else with a depth over 3,000. This process excluded nuclear-mitochondrial

pseudogenes and also reduced the amount of data going into subsequent steps. We then

reconstructed mitogenomes by mapping reads to the mitogenome of the stingless bee *Melipona*

bicolor (AF466146.2, Araujo & Arias 2019), followed by *de novo* assembly of the mapped

reads, mapping and extension to the longest contigs, and then joining of contigs and manual

verification of the final sequence by read-mapping (Geneious version 2019.2., Kearse *et al*.

2012). The final mitogenome assemblies were annotated using DOGMA (Wyman *et al*. 2004)

and MITOS (Bernt *et al.* 2013), with reference to the *M. bicolor* mitogenome and manual

editing in Geneious.

We aligned these three new bee mitogenomes with three more unpublished *Tetragonula*

mitogenomes and the *M. bicolor* mitogenome (AF466146.2) using MAFFT (Katoh and

Standley 2013). The most appropriate evolutionary model for each gene was estimated using

jModelTest 2.1.10 with the best model chosen based on Bayesian Information Criterion (BIC).

We then constructed a bayesian tree with MrBayes (Huelsenbeck & Ronquist 2001).

Nuclear genome assembly

To further investigate the low nuclear divergence suggested by our two nuclear genes

(*Opsin* and *EF1alpha*) we considered the remaining genome-wide data generated from our

Illumina sequencing. We took the lower genomic bin of the Illumina data (everything below

500x coverage based on kmer binning (Bushnell 2019) and assembled each species using

SPADES (Bankevich *et al.* 2012), which was run with mostly default settings, including the

in-built error correction module. We assessed the assemblies and generated basic statistics

using QUAST (Gurevich *et al.* 2013). We then took the 768bp *EF1alpha* PCR product and

searched it against custom-blast databases for each genome. We did the same for *Opsin*. These

larger sequences were aligned with MAFFT and a Bayesian tree was inferred from this

alignment with MrBayes.

Results

Tests of hybridization using microsatellites and snps

PCAs using both microsatellite and *snp* datasets separated samples into three non-

overlapping clusters consistent with their species assignment ((<5% missing data, Fig. 3A).

The microsatellite PCA placed *T. davenporti* closer to *T. carbonaria* than to *T. hockingsi* (Fig.

3A), but the *snp* PCA indicated an equal genetic distance between the three clusters (20% of

the variance explained by each of the first two components, Fig. 3B).

STRUCTURE analyses using both microsatellite and *snp* datasets (N= X colonies, 1

worker per colony) returned no evidence of hybrids, with all individuals showing close to 100%

assignment to a species' cluster (Fig. 4). Notably however, STRUCTURE results were

sensitive to duplicated and missing data. The same analyses using datasets comprising four

workers per colony (akin to resampling the same individual multiple times), assigned X-X

individuals at less than 90% posterior probability of belonging to their species' cluster, and

thus putative hybrids (Fig. 4). Likewise, subsampling to 20% missing microsattelite data

resulted in two *T. hockingsi* individuals being designated as hybrids with *T. davenporti* Missing

data had no impact on the *snp* dataset (i.e. every individual was still assigned 100% to its cluster

when lower threshold for missing data shifted marker number from 1,935 to 13,409).

Species-specificity of PCR-amplified gene regions

The mitochondrial gene *16S*, and the nuclear gene *Opsin*, each grouped samples into three

haplotypes consistent with species assignment (Fig. 2) although pairwise divergence between

species was greater for 16S (229 individuals representing all 63 colonies ; ~6% between *T.*

carbonaria and *T. hockingsi*) than *Opsin* (116 individuals representing all 63 colonies ; ~0.5%

difference between *T. carbonaria* and *T. hockingsi*). The percent difference in the *16S* gene

region amplified was lower than that of the complete mitochondria (see below), because one

half of the PCR region showed no variation. The nuclear gene *EF1alpha* was identical in

sequence for all samples (245 individuals representing all 63 colonies). No individuals showed

mitochondrial-nuclear discordance between *Opsin* and *16S*. That is, if an individual had a *16S*

sequence representative of *T. carbonaria* it also had an *Opsin* sequence representative of *T.*

carbonaria (Fig. 2). However, one colony that was labelled *T. carbonaria* during field

collections returned sequences (and nuclear markers) consistent with *T. hockinsgsi*, and this

was assumed to be the result of either labelling error, or a recent interspecific colony takeover

(1 of 69, 1.5% sample ID error rate). We also detected one instance of miss-labelling between

the sequence dataset and the microsatellite dataset because the same plate of individuals had

been extracted twice and two samples had switched positions in one of these plates.

Mitogenomes and genomes

We were unable to fully resolve the AT-rich control region of the mitochondria, but we did

recover all mitochondrial genes. Our assemblies of the mitogenomes for these three species

indicated 22.2% pairwise difference across *T. carbonaria* and *T. hockingsi*, 21.6% across *T.*

carbonaria and *T. davenporti,* and 23.6% across *T. davenporti* and *T. hockingsi* (Fig. 1). Using

these mitogenomes we were able to determine that the *16S* mitochondrial sequences were of

mitochondrial origin and not nuclear pseudogenes, the inclusion of which had hindered

interpretation in previous studies. An obvious gene inversion was detected, relative to *M.*

bicolor and all other bees (Supplementary Fig. 1) with *12S*, *16S*, *NAD1*, *CYTB* and *NAD6*

reversed and inserted in between *NAD3* and *NAD5*.

Nuclear genomes returned a mean *n50* of 14,954 across the three species (Table 1).

From this sequencing, we investigated further the lack of interspecific difference at *EF1alpha*.

We took the *EF1alpha* sequence and used it as a 'bait' to pull out the regions of the genome

surrounding the 768bp PCR product. We managed to select at least 10kb in each species that

aligned around the PCR product using MAFFT (Katoh and Standley 2013). Duplications were

not present as only one hit was found in each assembly. Variation was detected around the

EF1alpha locus and this separated the three species, based on the one individual sequenced of

each. We thus confirmed that the pairwise difference across these three bee species was indeed

extremely low in this region of the nuclear genome (0.7-1%, Fig. 5) relative to that detected in

the mitochondrial genome. Similarly, we recovered 26kb around the *Opsin* gene, which

returned the same low nucleotide difference between species (0.7-1%).

A Bayesian phylogenetic analysis based on either mitogenomes, or the flanking regions

of *EF1alpha*, resolved the evolutionary relationships between the *Tetragonula* species in the

same way, with *T. carbonaria* and *T. davenporti* diverging from each most recently, and *T.*

hockingsii as their sister group (Fig 5).

Discussion

Testing for hybridisation

Understanding hybridisation is crucial to understanding species and speciation, but detecting

hybridisation relies critically on the methods used to infer it, as illustrated here using a range

of techniques. Incorrect inference of hybridisation can lead to the species status of an organism

being questioned, as was the case with these stingless bees. Initial reports of hybridisation

(Franck *et al.* 2004 & Brito *et al.* 2014) would suggest these species were actually subspecies

that had recently come into secondary contact, especially given their close ecological

interaction in SE Queensland. We find, conversely, that these three species, *T. carbonaria*, *T.*

hockingsi and *T. davenporti*, are good species with no evidence for hybridisation in the areas

of sympatry that we sampled (including in colonies of all three species residing together at one

property for at least nine years). The mitogenome data indicate that the divergence between

these species is deep (21.6 to 23.6% difference, and at least 8 Mya for the closest relatives).

Although the nuclear genome does show much less variation (0.7-1%), this ratio of nuclear to

mitochondrial divergence (~22x) is within the known natural range (0.8-40x) described in a

recent comparative study across metazoans, and within the range reported specifically for

insects (Allio *et al.* 2017).

In this study we analysed multiple genetic markers to assess the extent of hybridisation

(if any) in these closely related sympatric species. Each marker set was found to have its own

set of issues that can lead to incorrect inferences about the extent of hybridisation. We discuss

issues highlighted by this study in the following sections.

Extremely low levels of nuclear divergence

We found extremely low levels of nuclear divergence in this study relative to the

divergence rates of the mitochondrial sequence. This pattern has often been considered

evidence for ongoing nuclear geneflow despite different mitochondrial backgrounds. A recent

study suggests that this ratio varies substantially across taxa (Allio *et al*. 2017). The nuclear-

mitochondrial ratio in this group of bees does seem to be extremely high relative to that reported

for other insects, but the pattern we observed when considering the larger fragments of the

nuclear genome was consistent and one cannot automatically infer hybridisation from this

pattern. Given the data we have generated it is not possible to determine whether nuclear gene

flow has occurred across divergent mitochondrial lineages for some period of time, but our

data indicates that no gene flow is occurring across the nuclear genome now.

We investigated the lack of variation in the *EF1alpha* PCR product further to see

whether it was restricted to the 700bp *pcr* product. We were able to recover more sequence

from the whole genome assemblies. These larger fragments did exhibit variation, and from

that, we recovered the same phylogeny (Fig. 5) as from the mitogenomes. The ratio of mito-

nuclear divergence is the same across the three species included in this analysis. Hymenoptera

are regarded as having a higher rate of mitochondrial divergence compared to other insects

(Kaltenpoth 2012). The extremely large discrepancy across the mitochondrial and nuclear rates

in these species requires further work to establish whether this derives from a fast

mitochondrial rate, a slow nuclear rate, or a combination of these two. This further investigation

is required before divergence times can be estimated more realistically in this group of bees.

Mito-nuclear pseudogenes

Two earlier studies on the three *Tetragonula* species we investigated amplified nuclear

pseudogenes instead of the true mitochondrial gene region. This is a common problem in

mitochondrial DNA based tests, but is more prevalent in some insect groups than others.

Recently, Françoso *et al.* (2019) successfully re-designed primers for COI that avoid picking

up pseudogenes in these *Tetragonula* species. Researchers are now much more aware of the

issue of pseudogenes and it is common to check for stop codons or frame-shifting indels, but

this scrutiny detects only a subset of pseudogenes. The ability to reconstruct whole

mitogenomes from short read data (and using a depth cut-off, as we did, to avoid nuclear

copies) provides a way to detect pseudogenes more comprehensively. There are likely many

reports of hybridisation based on undetected pseudogenes in the literature, inflating perceptions

of the extent of hybridisation between species, and contributing misinformation to the debate

on species concepts.

Mito-nuclear discordance

Hybridisation (and thus gene flow) between population samples is often inferred from

discordance across mitochondrial and nuclear gene sequences. This pattern may well be

representative of recent hybridisation, but may also be caused by rare events, as seen with

examples of mitochondrial capture (Good *et al.* 2015; Perea *et al.* 2016). Another issue with

this measure of hybridisation is that the pattern (a sample returning one mitochondrial sequence

and a different nuclear one) is the same as one gets when samples are mislabelled. A recent

study indicates that rates of mislabelling are around 4% in published genetic datasets (Zych *et*

al. 2017), so any cases of mito-nuclear discordance at these rates probably deserve additional

scrutiny. We detected one instance of mis-labelling across two of our datasets, by virtue of

having conducted multiple genetic tests on the same individuals. To detect this kind of

'accidental introgression' any samples showing mito-nuclear discordance should be re-

amplified and sequenced again.

Incorrect inference from STRUCTURE plots

The two earlier papers that report hybridisation among the three *Tetragonula* species

inferred it on the basis of individuals having less than 90% assignment to a cluster in a

STRUCTURE plot (Franck *et al.* 2004; Brito *et al.* 2014). However, *STRUCTURE* quite often

mis-assigns individuals and this can be influenced by many factors. Warnings have been

published about over-interpreting structure plots (Lawson *et al.* 2018), but it is still a prevalent

issue. With bees, living in colonies of closely-related individuals, the inclusion of four

individuals per hive (in our analysis), caused *STRUCTURE* to assign several individuals in a

manner consistent with hybridisation. Clearly, in this case, this is simply a consequence of a

biased sample of the population having been presented to the algorithm, because the analysis

on single hive representatives did not show any suggestions of hybridisation.

We investigated the effect of missing data on inferring hybridisation from

STRUCTURE plots. Our two original datasets had very low levels of missing data (<5%) and

every individual was assigned at close to 100% when only one bee per hive was used (Fig. 4).

When we subsampled the two datasets to 20% missing data, the microsatellites indicated

hybrids, but the *snp* dataset did not. This was probably because the '20% missing' *snp* dataset

had many more markers than the '5% missing' microsatellite dataset. The microsatellite results

illustrate how missing data can lead to incorrect inferences of hybridisation from population

assignment tests like STRUCTURE. Including more markers appears to compensate for the

effect of missing data, at least based on our *snp* results (Fig. 4).

Using multiple nuclear markers

In cases where the nuclear genome divergence is low, using multiple markers, and

including more rapidly evolving markers are two ways in which to assess hybridisation rates

more accurately. In this study we used 7 microsatellites (more variable than *snps*), and also

assayed 2,049 *snps* (using more markers). Both of these marker systems revealed no evidence

for hybridisation in the PCA plots, with no individuals intermediate between clusters (Fig. 3).

This kind of multi-marker nuclear genome assay is the most accurate way in which to infer

hybridisation (Twyford & Ennos 2011), with a parentage analysis providing the most

convincing evidence of recent hybridisation, where it is known to have occurred.

Our results indicated much tighter clustering of the *snp* data than the microsatellite data

in the PCA plots, and less ambiguous assignments in *STRUCTURE*. This observation indicates

that misidentification of hybridisation will be less common in *snp* datasets or around a couple

of thousand markers than might be the case when using relatively few microsatellites.

Tetragonula davenporti and the 'carbonaria complex'

Using multiple markers, we have comprehensively tested for hybridisation in three

Tetragonula bees that overlap in their geographic distribution. Although previous studies have

reported that these species hybridise in these areas, our approach found no evidence for it.

Despite swarms of these bees taking over each other's hives and having mixed species drone

aggregations, the three species are all good species, likely diverged from one another millions

of years ago, and do not hybridise even when in close sympatry. *Tetragonula davenporti* has

been confirmed from only one property in the Gold Coast hinterland. Its species status had

been questioned recently (Françoso *et al.* 2019), but we find strong evidence for the validity of

T. davenporti as a species. The natural range of this species is currently unknown, and the

difficulty in identifying adults (which are almost identical to those of *T. carbonaria*) means

that their distribution in managed colonies is also relatively unknown. South-east Queensland

is an area of high endemicity, so it may well be endemic in this area. This study investigated

managed colonies mostly from Brisbane and SE Queensland because this is where

hybridisation is most likely between these three species. Examining wild populations of this

genus may reveal additional species within the '*carbonaria* complex' (perhaps the 'northern'

T. hockingsi).

Conclusions

We have highlighted several ways in which hybridisation might be incorrectly inferred

when using certain marker types and certain types of inference. It is quite likely that many

published records of hybridisation might be affected by some of these issues. In particular,

hybridisation inferred from a low frequency (<5%) of mitochondrial-nuclear discordance. This

inference comes from the same pattern as miss-labelling, which is also usually around this

frequency. We suggest these considerations should inform the design of genetic studies that

investigate hybridisation and also inform the meta-analysis of genetic studies of hybridisation.

We have also shown how whole-genome data can provide additional resolution when the rate

of nuclear mutation is low, and this should allow further resolution of closely related lineages.

The genomic resources developed here will enable further molecular investigation of the

evolution of this group of social, stingless bees, and enable further analysis of the divergence

times and ratio of mitochondrial to nuclear divergence rates.

Acknowledgements

We would like to acknowledge all the bee-keepers and hive owners who donated bees

to this study, especially Tim Heard, Peter Davenport and Alan Lashmar. We thank the

University of Queensland for financial support.

Data Availability

Genome assemblies and mitogenomes have been deposited at genbank (pending

accession numbers).

Illumina data has been deposited at the Short Read Archive, (awaiting final SRA

numbers.)

Microsatellite data and *snp* vcf files are to be deposited at the University of Queensland

espace (doi to follow).

References

Andrews S. 2019. FastQC: a quality control tool for high throughput sequence data [Online]. Available: http://www.bioinformatics.babraham.ac.uk?/projects/fastqc/.

Araujo N de S and Arias MC (2019) Mitochondrial genome characterization of *Melipona bicolor*: Insights from the control region and gene expression data. Gene 705: 55–59.

Ayala, FJ, Escalante, A, O'Huigino, C, & J Klein. 1994. Molecular genetics of speciation and human origins. Proceedings of the National Academy of Science USA. 91: 6787-6794.

Bankevich, A, Nurk, S, Antipov, D, Gurevich, AA, Dvorkin, M, Kulikov, AS, Lesin, VM, *et al.* 2012. SPAdes: A new genome assembly algorithm and its applications to single-cell sequencing. Journal of Computational Biology: A Journal of Computational Molecular Cell Biology 19 (5): 455–77.

Barreto FS, Watson ET, Lima TG, *et al*. (2018) Genomic signatures of mitonuclear coevolution across populations of *Tigriopus californicus*. Nature Ecology & Evolution 2(8): 1250–1257.

Bernt M, Donath A, Jühling F, Externbrink F, Florentz C, Fritzsch G, Pütz J, Middendorf M, Stadler PF. MITOS: improved de novo metazoan mitochondrial genome annotation. Molecular Phylogenetics and Evolution. 69(2):313-9

Brito, RM, Francisco, FO, Ho, SYW, & BP Oldroyd. 2014. Genetic architecture of the Tetragonula carbonaria species complex of Australian stingless bees (Hymenoptera: Apidae: Meliponini). Biological Journal of the Linnean Society. 113(1) 149-161.

Bushnell B, 2019 Bbmap – a suite of bioinformatics tools. [Online]. Available: https://sourceforge.net/projects/bbmap/

Cameron, SA, Austin, AD, James Derr, & RA Wharton. 1992. The application of nucleotide sequence data to phylogeny of the Hymenoptera: a review. Journal of Hymenoptera Research. 1:63-79.

Caporaso, J, Gregory, CL, Lauber, WA, Walters, D, Berg-Lyons, J Huntley, N, Fierer, S & M Owens, *et al.* 2012. Ultra-High-Throughput Microbial Community Analysis on the Illumina HiSeq and MiSeq Platforms. The ISME Journal 6 (8): 1621–24.

Catchen, J, Hohenlohe, P, Bassham, S, Amores A, & W Cresko. 2013. Stacks: an analysis tool set for population genomics. Molecular Ecology, 22(11):3124-3140

Clement M, Posada D, & Crandall, KA. 2000. TCS: a computer program to estimate gene genealogies. Molecular Ecology. 9(10):1657-9.

Cunningham, JP, Hereward, JP, Heard, TA, De Barro, PJ, & SA West. 2014. Bees at war: interspecific battles and nest usurpation in stingless bees. American Naturalist 184 (6): 777–86.

Dollin AE, Dollin LJ & SF Sagakami. 1997. Australian Stingless Bees of the Genus *Trigona* (Hymenoptera:Apidae). Invertebrate Taxonomy. 11, 861-896.

Dowton, M & AD Austin. 1994. Molecular phylogeny of the insect order Hymenoptera: Apocritan relationships. Proc Natl Acad Sci USA. 91(21): 9911–9915.

Dyer AG, Streinzer M & J Garcia. 2016. Flower detection and acuity of the Australian native stingless bee Tetragonula carbonaria Sm. Journal of Comparative Physiology. 202: 629-639.

Elshire, RJ, Glaubitz, JC, Sun, Q, Poland, JA, Kawamoto, K, Buckler, ES & SE Mitchell. 2011. A Robust, Simple Genotyping-by-Sequencing (GBS) Approach for High Diversity Species. PloS One 6 (5): e19379.

Feder, JL *et al*. 2003. 'Allopatric genetic origins for sympatric host-plant shifts and race formation in Rhagoletis', Proceedings of the National Academy of Sciences of the United States of America. National Acad Sciences, 100(18), pp. 10314–10319.

Ford, J. 1974. Concepts of Subspecies and Hybrid Zones, and Their Application in Australian Ornithology. Emu - Austral Ornithology 74 (3): 113–23.

Françoso, E, Zuntini, AR, Ricardo, PC, Silva, JPN, Brito, R, Oldroyd, BP & MC Arias. 2019. Conserved Numts Mask a Highly Divergent Mitochondrial-COI Gene in a Species Complex of Australian Stingless Bees Tetragonula (Hymenoptera: Apidae). Mitochondrial DNA. Part A, DNA Mapping, Sequencing, and Analysis 30 (7): 806–17.

Franck P, Cameron E, Good G, Rasplus JY, & BP Oldroyd. 2004. Nest architecture and genetic differentiation in a species complex of Australian stingless bees. Molecular Ecology. 13(8) 2317-31.

Good, JM, Vanderpool, D, Keeble, S, & K Bi. 2015. Negligible Nuclear Introgression despite Complete Mitochondrial Capture between Two Species of Chipmunks. Evolution; International Journal of Organic Evolution 69 (8): 1961–72.

Green, CL, Franck, P & BP Oldroyd. 2001. Characterization of Microsatellite Loci for Trigona Carbonaria, a Stingless Bee Endemic to Australia. Molecular Ecology Notes 1 (1-2): 89–92.

Gurevich, A, Saveliev, V, Vyahhi, N, & G Tesler. 2013. QUAST: Quality Assessment Tool for Genome Assemblies. Bioinformatics. 29 (8): 1072–75.

Heard TA. 1994. Behaviour and pollinator efficiency of stingless bees and honey bees on macadamia flowers. Journal of Apicicultural Research 33(4):191–198.

Hines, HM, Cameron SA & PH Williams, 2006. Molecular phylogeny of the bumble bee subgenus Pyrobombus (Hymenoptera:Apidae:Bombus) with insights into gene utility for lower-level analysis. Invertebrate Systematics 20(3) 289-303 https://doi.org/10.1071/IS05028

Huelsenbeck, JP & F Ronquist. 2001. MRBAYES: Bayesian inference of phylogeny. Bioinformatics 17:754-755.

Jombart, T. (2008) adegenet: an R package for the multivariate analysis of genetic markers. Bioinformatics 24: 1403-1405.

Kaltenpoth M, Showers Corneli P, Dunn DM, *et al*. (2012) Accelerated evolution of mitochondrial but not nuclear genomes of Hymenoptera: new evidence from crabronid wasps. PloS one 7(3): e32826.

Katoh K, Standley DM. (2013). MAFFT Multiple Sequence Alignment Software Version 7: Improvements in Performance and Usability. Molecular Biology and Evolution. 30(4): 772-780.

Kearse, M, Moir, R, Wilson, A, Stones-Havas, S, Cheung, M, Sturrock, S, Buxton, S, Cooper,
A, Markowitz, S, Duran, C, Thierer, T, Ashton, B, Meintjes, P, & A Drummond. 2012.
Geneious Basic: An Integrated and Extendable Desktop Software Platform for the Organization
and Analysis of Sequence Data. Bioinformatics, 28, 1647-1649.

Kopelman, NM, Mayzel, J, Jakobsson, M, Rosenberg NA and I Mayrose. 2015. clumpak: a
program for identifying clustering modes and packaging population structure inferences across
K. Molecular Ecology Resources. 15(5): 1179–1191.

Lawson, DJ, van Dorp, L & D Falush. 2018. A Tutorial on How Not to over-Interpret
STRUCTURE and ADMIXTURE Bar Plots. Nature Communications 9 (1): 3258.

Leigh, JW, & D Bryant. 2015. Popart : Full-Feature Software for Haplotype Network
Construction. Edited by Shinichi Nakagawa. Methods in Ecology and Evolution / British
Ecological Society 6 (9): 1110–16.

Linck, E, and Battey, CJ. 2019. Minor Allele Frequency Thresholds Strongly Affect Population
Structure Inference with Genomic Data Sets. Molecular Ecology Resources 19 (3): 639–47.

Mallet J, Wynne IR, & CD Thomas. 2011. Hybridisation and climate change: brown argus
butterflies in Britain (*Polyommatus* subgenus *Aricia*). Insect Conservation and Diversity. 4 (3)
192-199.

Mardulyn, P, & SA Cameron. 1999. The Major Opsin in Bees (Insecta: Hymenoptera): A
Promising Nuclear Gene for Higher Level Phylogenetics. Molecular Phylogenetics and
Evolution. 12(2):168-76.

Oliveira DCSG, Raychoudhury R, Lavrov DV, *et al*. (2008) Rapidly evolving mitochondrial
genome and directional selection in mitochondrial genes in the parasitic wasp nasonia
(hymenoptera: pteromalidae). Molecular biology and evolution 25(10): 2167–2180.

Perea, S, Vukić, J, Šanda, R & I Doadrio. 2016. Ancient Mitochondrial Capture as Factor
Promoting Mitonuclear Discordance in Freshwater Fishes: A Case Study in the Genus Squalius
(Actinopterygii, Cyprinidae) in Greece. PloS One 11 (12): e0166292.

Peterson, BK, Weber, JN, Kay, EH, Fisher, HS & HE, Hoekstra. 2012. Double Digest RADseq:
An Inexpensive Method for de Novo snp Discovery and Genotyping in Model and Non-Model
Species. PloS One 7 (5): e37135.

Poland, JA, Brown, PJ, Sorrells, ME, & JL Jannink. 2012. Development of High-Density
Genetic Maps for Barley and Wheat Using a Novel Two-Enzyme Genotyping-by-Sequencing
Approach. PloS One 7 (2): e32253.

Pritchard, JK, Stephens, M, & P Donnelly. 2000. Inference of Population Structure Using
Multilocus Genotype Data. Genetics 155 (2): 945–59.

Ridley, AW, Hereward, JP, Daglish, GJ, Raghu, S, McCulloch, GA & GH Walter. 2016. Flight
of Rhyzopertha Dominica (Coleoptera: Bostrichidae)-a Spatio-Temporal Analysis With
Pheromone Trapping and Population Genetics. Journal of Economic Entomology 109 (6):
2561–71.

Sánchez-Guillén RA, Muñoz J, Rodríguez-Tapia G, Arroyo TPF, & A Córdoba-Aguilar. 2013.
Climate-Induced Range Shifts and Possible Hybridisation Consequences in Insects. PloS One.
8(11) e80531.

Schuelke, M. 2000. An Economic Method for the Fluorescent Labeling of PCR Fragments.
Nature Biotechnology 18 (2): 233–34.

Shipham, A, Joseph, L, Schmidt, DJ, Drew, A, Mason, I & JM Hughes. 2019. Dissection by
Genomic and Plumage Variation of a Geographically Complex Hybrid Zone between Two
Australian Non-Sister Parrot Species, *Platycercus adscitus* and *Platycercus eximius*. Heredity
122 (4): 402–16.

Toon, A, Daglish, GJ, Ridley, AW, Emery, RN, Holloway JC, & GH Walter. 2015. Random
Mating Between Two Widely Divergent Mitochondrial Lineages of *Cryptolestes ferrugineus*
(Coleoptera: Laemophloeidae): A Test of Species Limits in a Phosphine-Resistant Stored
Product Pest. Journal of Economic Entomology. 109(5):2221-8.

Twyford, AD & RA Ennos. 2011. Next-generation hybridization and introgression. Heredity
108: 179-189.

Walter GH (2003) Insect pest management and ecological research. Cambridge University
Press, Cambridge, p 162-165.

Wyman SK, Jansen RK, Boore JL. 2004. Automatic annotation of organellar genomes with
DOGMA. Bioinformatics. 20(17):3252-3255.

Xiao J-H, Jia J-G, Murphy RW, *et al*. (2011) Rapid evolution of the mitochondrial genome in
Chalcidoid wasps (Hymenoptera: Chalcidoidea) driven by parasitic lifestyles. PloS one 6(11):
e26645.

Zych, K, Basten, LS, Elvin, M, Rodriguez, M, Van der Velde, KJ, Arends, D, Westra, HJ *et al.*
2017. reGenotyper: Detecting Mislabeled Samples in Genetic Data. PloS One 12 (2):
e0171324.

Author Contributions

All authors conceived the project, TJS obtained samples, JPH conducted molecular lab work, JPH and DRB analysed the genetic data. All authors wrote the manuscript.

Supplementary data

Supplementary Figure 1. Mauve alignment of the mitogenomes showing the inversion between *Tetragonula* and other bees.

Figures and Tables

Table 1. Characteristics of the genomes and their draft assemblies for three species of stingless bees, including genome size, coverage, and % repeat, which were estimated with a kmer analysis in bbnorm. The mitogenome coverage was assessed by read mapping, and the assembly statistics (length, longest contig, n50, number of contigs) were calculated in Quast. The genome assemblies can be found at genbank under the provided bioproject and biosample IDs.

Species	Genome size estimate	Nuclear genome coverage	Mitogenome coverage	% repeat	Assembly length	Longest contig	N50	N contigs	BioProject	BioSample
T. carbonaria	457,237,826	23x	6,899	33%	284,886,424	187,309	14,476	156,012	PRJNA578948	SAMN13088838
T. hockingsi	376,762,141	35x	11,089	26%	288,263,186	200,717	11,462	225,627	PRJNA578994	SAMN13089661
T. davenporti	342,107,567	38x	26,907	17%	288,842,417	271,029	18,924	72,464	PRJNA578998	SAMN13089823

Figure 1. Map showing the known distributions of three species of *Tetragonula* in Australia, the study area is marked with a black rectangle.

Figure 2. Haplotype networks for the mitochondrial *16S* gene, and nuclear *Opsin* and *EF1alpha* (top). Brood of the three species we investigate is illustrated (middle), as are representative bees of each species (bottom).

Figure 3. PCA of nuclear genetic data, 7 microsatellites (A) and 2049 *snp*s (B).

One individual per hive

1935 snps - <5% missing data

13,409 snps - ~15% missing data

microsatellites - <5% missing data

microsatellites ~ 20% missing data

Multiple individuals per hive

microsatellites

1935 snps - <5% missing data

● *T. carbonaria*　　○ *T. hockingsii*　　● *T. davenporti*

Figure 4. Structure plots for 7 microsatellite loci (with < 5% missing data and 20% missing data) and the *snp* dataset (for the 1,935 *snps* with less than 5% missing data, and the 13,409 snps that were retained when the missing data limit was set to 20%). When the missing data limit for *snps* was raised to 20% the *snp* dataset had a final missing data rate of 15%. A. The analyses were run with multiple individuals per hive (mostly four but sometimes fewer). B. The same analyses were run with only one individual bee per hive.

Figure 5. Bottom. Alignment of 10kb surrounding the *EF1alpha* gene, blue rectangle represents the PCR product, (Left) Bayesian tree of this region. Right. A Bayesian tree was also constructed from the mitogenome alongside which the bees are illustrated.