Application of probiotic *Bacillus* spp. isolated from African nightcrawler (*Eudrilus*
***eugeniae*) on Nile Tilapia (*Oreochromis niloticus* L.)**

Samson J.[1 2]*, Quiazon K.M.[1 2], Choresca C.[3]
[1] College of Fisheries, Central Luzon State University, Science City of Muñoz 3120, Nueva
Ecija, Philippines

[2] Freshwater Aquaculture Center, Central Luzon State University, Science City of Muñoz 3120,
Nueva Ecija, Philippines

[3] National Fisheries Research and Development Institute - Fisheries Biotechnology Center,
Bureau of Fisheries and Aquatic Resources - National Freshwater Fisheries Technology Center,
Central Luzon State University Compound, Science City of Muñoz 3120, Nueva Ecija,
Philippines

Corresponding author's Email: samsonjaypee@clsu.edu.ph

Abstract

15     Due to the emergence of antibiotic-resistant pathogens, probiotics in aquaculture are used

for the prevention of infectious microbial diseases and substitute for antibiotics and

chemotherapeutics. In this study, we evaluated the effect of probiotic *Bacillus* spp. isolated from

African nightcrawler (*Eudrilus eugeniae*) on the growth, feed utilization, and disease resistance

of Nile tilapia (*Oreochromis niloticus*). Four probiotic strains of *Bacillus* spp. (ANSCI9, BFAR9,

RM3, and RM10) were individually incorporated in the commercial diet (control) at 10^8 CFU g^{-1}

of feed. The experimental fish were fed at 5% of their body weight for 30 days, and subjected to

a 14-day *Aeromonas hydrophila* challenge test afterward. The results showed the probiotic-

treated groups have higher ($P<0.05$) average body weight (ABW) (4.51 ± 0.34 g) than the

control (3.89 ± 0.17 g). The BFAR9 (2.73 ± 0.26 g) and RM10 (3.15 ± 0.30 g) showed higher

($P<0.05$) absolute growth (AG) than the control (2.20 ± 0.16 g). Furthermore, RM10 had higher

($P<0.05$) specific growth rate (SGR) (1.60 ± 0.10 % day^{-1}) and relative growth rate (RGR)

(181.39 ± 18.16 %) than the control (SGR=1.29 ± 0.07 % day^{-1}; RGR=129.84 ± 9.77 %).

Consequently, RM10 had significantly lower ($P<0.05$) feed conversion ratio (FCR) (1.99 ± 0.13)

than the control (2.60 ± 0.16). The challenge test revealed that the probiotic-treated groups have

higher ($P<0.05$) survival (81.25 ± 9.57 %) than the control (55.00 ± 19.15 %). These results

revealed that the probiotic *Bacillus* spp. isolated from *E. eugeniae* improved the growth, feed

utilization, and the disease resistance of Nile tilapia.

Keywords: microbiota, *Aeromonas hydrophila*, earthworms, aquaculture, immunomodulation,

disease resistance, feed utilization

Introduction

The need for sustainable aquaculture has encouraged exploration into the use of probiotics for growth promotion and health improvement of aquatic organisms. These beneficial microorganisms can be isolated from various sources, including in food (Demir and Başbülbül, 2017; Leite et al., 2015), aquatic organisms (Lamari et al., 2014; Lin et al., 2013; Sanchez-Ortiz et al., 2015), terrestrial animals (Brashears et al., 2003; Feng et al., 2017; Jena et al., 2013; Musikasang et al., 2009), and humans (Amenu, 2015; Reis et al., 2016). In this study, we evaluated the probiotic potential of the isolated *Bacillus* spp. from earthworms on Nile tilapia. Although some studies have used earthworms as a protein source for fish (Boaru et al., 2016; Mohanta et al., 2016), and explored the presence of microorganisms in them (Byzov et al., 2009; Horn et al., 2003; Kim et al., 2004; Singleton et al., 2003), to the best of our knowledge, no study has demonstrated the effect of these microorganisms on aquatic species.

The gut of earthworms provides an ideal and favorable environment for the growth and activity of bacteria (Govindarajan and Prabaharan, 2015). For several centuries, China and other parts of the Far East believed that earthworms promote overall health, and are sources of therapeutic drugs for various diseases (El-Kamali, 2000; Ismail, 2005). Furthermore, indigenous people all over the world, more particularly in Asia, including Vietnam, China, Korea, India, and Myanmar, extract and use the biologically active compounds from earthworms (Ranganathan, 2006). Unfortunately, there is no available information on the application of probiotic bacteria from earthworms, specifically on aquaculture. Hence, in this present study, we aimed to evaluate the effect of probiotic *Bacillus* spp. isolated from *E. eugeniae* on the growth, feed utilization, and disease resistance of Nile tilapia.

Materials and Methods

Source of Bacteria

Four probiotic strains of *Bacillus* spp. (ANSCI9, BFAR9, RM3, and RM10) isolated from *E. eugeniae* were used in this study. The 16s rRNA sequences of the probiotic isolates are available in the GenBank, with accession numbers: MH919310, MH919302, MH919306, and MH919308.

Experimental Diets and Design

A commercial diet (Tateh Aquafeeds, Philippines) served as the basal diet (control). The proximate composition was as follows: crude protein 31%, lipid 5%, crude fiber 8%, and crude

ash 12%. The probiotic-treated diets were prepared to contain a single strain of *Bacillus* spp.

(ANSCI9, BFAR9, RM3, and RM10) per group. The McFarland standard was used to estimate

the bacterial densities. Dilutions were done using phosphate buffer solution (PBS) to have a

similar bacterial density. The isolates were sprayed on commercial feeds (10^8 CFU·g^{-1}) and air-

dried for 2 hours before storing at 4°C.

Four hundred *O. niloticus* fingerlings were distributed into 20 glass aquaria ($52 \times 25 \times 30$

cm) divided into five experimental groups (four replicates per group). Each group consists of 80

fish (20 fish $\times$ 4 aquaria), which had an initial weight of 1.76 ± 0.07 g. The water quality

parameters during the study were the following: the temperature at 25.3 ± 0.9 ℃, pH at 8.0 ± 0.2,

and dissolved oxygen at 3.52 ± 1.16 ppm. The fish were fed their respective diets (at

approximately 5% of body weight day^{-1}) thrice daily at 8:00, 12:00, and 16:00 for 30 days.

Data Collection

At the end of the feeding trial, all fish were measured and weighed. Growth parameters

and survival rate were measured as follows:

$$\text{Specific growth rate (SGR)} = \frac{(\ln[\text{final weight}] - \ln[\text{initial weight}])}{(\text{time interval in days})} \times 100$$

$$\text{Relative growth rate (RGR)} = \frac{\text{Final weight of fish - Initial weight of fish}}{\text{Initial weight of fish}} \times 100$$

$$\text{AG (g)} = \text{Final weight} - \text{Initial weight}$$

$$\text{FCR} = \frac{\text{Amount of feed given (g)}}{\text{Weight gain (g)}}$$

$$\text{SR (\%)} = \frac{\text{No. of individuals at the end of the experiment}}{\text{No of individuals at the beginning of the study}} \times 100$$

Challenge Test

O. niloticus fingerlings (4.39 ± 0.40 g) were challenged with *A. hydrophila* (BIOTECH

10089) after the 30-day feeding trial. Fish were anesthetized in tricaine methanesulfonate

(MS222) solution and intramuscularly injected with 0.1 mL *A. hydrophila* suspension (10^7

CFU·mL^{-1}). The mortality was observed daily until the 14th day, and the relative level of

protection (RLP) among the challenged fish was determined using the equation: [100 –

(immunized mortality% / control mortality%)] × 100 (Ruangroupan et al., 1986)

Statistical Analysis

Results were subjected to a one-way analysis of variance (ANOVA), followed by

Duncan's Multiple Range Test (DMRT) at a significant level of *P*<0.05 when there was a

significant difference among the treatments.

Results

Growth Performance

The summary of the measurements and growth parameters of the experimental fish is

shown in Table 1. The treatments supplemented with probiotic *Bacillus* spp. showed higher

(*P*<0.05) ABW than the control. RM10 gained the highest ABW (4.88 ± 0.30 g) among the

treatments; whereas, the lowest was recorded from the control group (3.89 ± 0.17 g). Treatments

RM10 (6.91 ± 0.23 cm), BFAR9 (6.88 ± 0.07 cm), and RM3 (6.65 ± 0.19 cm) have significantly

higher (*P*<0.05) lengths than the control (6.16 ± 0.11 cm). Also, treatment RM10 had

significantly higher (*P*<0.05) SGR (1.60 ± 0.10 % day^{-1}) and RGR (181.39 ± 18.16 %) than the

control (SGR=1.29 ± 0.07 % day^{-1}; RGR=129.84 ± 9.77 %). In addition, treatments BFAR9

(2.73 ± 0.26 g) and RM10 (3.15 ± 0.30 g) showed significantly higher (*P*<0.05) AG than the

control (2.20 ± 0.16 g). Consequently, treatment RM10 had significantly lower (*P*<0.05) FCR

(1.99 ± 0.13) than the control (2.60 ± 0.16). While no significant difference was observed on the

survival of the treatments.

Table 1. Summary (Mean ± SD) of measurements and growth parameters of the experimental

fish (n=4).

	TREATMENTS				
	Control	ANSCI9	BFAR9	RM3	RM10
Initial ABW (g)	1.69 ± 0.04[a]	1.80 ± 0.07[a]	1.78 ± 0.09[a]	1.77 ± 0.08[a]	1.74 ± 0.02[a]
Initial Length (cm)	4.63 ± 0.21[a]	4.59 ± 0.12[a]	4.82 ± 0.24[a]	4.67 ± 0.10[a]	4.66 ± 0.08[a]
Final ABW (g)	3.89 ± 0.17[c]	4.31 ± 0.20[b]	4.51 ± 0.21[ab]	4.34 ± 0.35[b]	4.88 ± 0.30[a]
Final Length (cm)	6.16 ± 0.11[c]	6.18 ± 0.04[c]	6.88 ± 0.07[a]	6.65 ± 0.19[b]	6.91 ± 0.23[a]
AG (g)	2.20 ± 0.16[c]	2.51 ± 0.23[bc]	2.73 ± 0.26[b]	2.57 ± 0.35[bc]	3.15 ± 0.30[a]
SGR (% day^{-1})	1.29 ± 0.07[b]	1.35 ± 0.11[b]	1.44 ± 0.13[ab]	1.39 ± 0.14[b]	1.60 ± 0.1[a]
RGR (%)	129.84 ± 9.77[b]	139.71 ± 16.82[b]	153.16 ± 21.1[ab]	145.61 ± 22.44[b]	181.39 ± 18.16[a]

FCR	2.60 ± 0.16^{a}	2.47 ± 0.23^{a}	2.30 ± 0.21^{ab}	2.41 ± 0.31^{ab}	1.99 ± 0.13^{b}
SR (%)	95.00 ± 5.77^{a}	87.50 ± 25.00^{a}	97.50 ± 2.89^{a}	97.50 ± 5.00^{a}	91.25 ± 14.36^{a}

Note: Treatments with the same superscript are not significantly different at $P<0.05$.

Challenge Test

The cumulative mortality of the experimental groups was recorded daily until the 14[th] day (Fig 1). All treated groups showed significantly higher ($P<0.05$) survival rates rate (81.25 ± 9.57 %) than the control (55.00 ± 19.15 %). Treatment RM10 had the highest survival rate (85.0 ± 10.00 %) among the treatments, while the lowest was recorded on the control (55 ± 19.15%). The RLP of the treated groups is ≥50% with the highest value in RM10 (66.67%) while the lowest for RM3 (50.00%). The result of the challenge test is presented in Table 3.

Figure 1. Cumulative mortality of each treatment during the *A. hydrophila* challenge test

Table 3. Values (Mean ± SD) of SR RLP for each treatment after challenging with *A. hydrophila.*

	TREATMENTS				
	Control	ANSCI9	BFAR9	RM3	RM10
SR (%)	55.0 ± 19.15^{b}	82.5 ± 8.16^{a}	80.0 ± 10.00^{a}	77.5 ± 20.82^{a}	85.0 ± 10.00^{a}
RLP (%)	0.0	61.11	55.56	50.00	66.67

Note: Treatments with the same superscript are not significantly different at $P<0.05$.

Discussion

The beneficial effect of probiotic *Bacillus* spp. isolated from *E. eugeniae* was observed in
this study. Nile tilapia showed better growth when fed with probiotics. Previous studies have
reported similar results on the effect of *Bacillus* spp. on the growth and survival of Nile tilapia.
Soltan et al., (2016) reported that commercial probiotics containing allicin, high unit hydrolytic
enzyme, *Bacillus subtilis* spores, and ginseng extract improved the growth and feed utilization of
Nile tilapia. Similarly, *Bacillus pumilus* improved the growth performance of Nile tilapia due to
enhanced immune system, health status, and disease resistance (Aly et al., 2008). Furthermore,
the application of *Bacillus coagulans* and other *Bacillus* spp. in the water also enhanced the
growth and immune response of Nile tilapia (Sutthi et al., 2018; Zhou et al., 2009). According to
Buruian et al. (2014), the application of probiotic *Bacillus* spp. in the aquaculture system
improves the growth performance, immune response, disease resistance, and survival of the fish.
These probiotics produce extracellular enzymes (proteases, amylases, and lipases), and provide
vitamins, fatty acids, and amino acids that have a beneficial effect on the metabolism of aquatic
animals (Balcazar et al., 2006). The observed growth improvement of treated groups can be a
result of improved enzymatic activity on the fish gut, as the probiotics used in the experiment
produce extracellular enzymes. These extracellular enzymes aids in breaking down food
molecules, which makes the absorption of nutrients more efficient (EL-Haroun et al., 2006). In
contrast, Silva et al. (2015) reported that the addition of *Bacillus amyloliquefaciens* did not
improve the growth and proximal composition of Nile tilapia; however, they observed a
significant increase of villi height and number of goblet cells in the intestine. Their result
suggests that the digestion and nutrient absorption of fish had been improved. Similarly, Shelby
et al. (2006) observed no significant improvement in growth; but they recorded higher survival
rates after 39 to 63 days of probiotic application. As we noted in this study, only RM10 showed
significantly better SGR, RGR, and FCR; although the probiotics used are of the same genus
(i.e., *Bacillus*) and source (i.e., *E. eugeniae*). This result does not imply that the three probiotic
isolates (i.e., ANSCI9, BFAR9, RM3) are less efficient. In fact, all probiotic-treated groups
showed higher survival rates when challenged with *A. hydrophila*. This suggests that the effect of
the probiotics may not be observed on growth but rather to the immunity and intestinal
morphology of the fish. Thus, it is better to evaluate the effect of these probiotics in the
immunological parameters and intestinal morphology of the fish. Furthermore, this observation
also suggests that the effects of probiotics application differs from each bacterial species and

strain. Correspondingly, it is important to measure the effective rate of application for each
probiotic bacterium.

In this study, higher survival rates of probiotic-treated groups infected with *A. hydrophila*
was recorded. Similar results were reported by Selim and Reda (2015); they reported that dietary
supplementation of *B. amyloliquefaciens* improved the survival of Nile tilapia infected with
Yersinia ruckeri or *Clostridium perfringens* type D. Moreover, Gupta et al. (2014) also observed
an improvement in the survival of *Cyprinus carpio* fry fed with probiotics and challenged with *A.*
hydrophila. In our study, the improvement of survival rates of the probiotic-treated groups is
attributed to the ability of probiotics to enhance the immune system. It is a result of immunity
modulation or immunomodulation, which is one of the common health benefits credited to the
application of probiotics (Cross, 2002). It is well-known that probiotics may lessen the
occurrence of disease or decrease the danger of disease outbursts. One possible mechanism for
this is the prevention of pathogen colonization on the intestinal mucosa or other epithelial
surfaces; as probiotic strains colonize these surfaces (Balcazar et al., 2006). Furthermore, they
can also produce inhibitory substances against pathogenic organisms and compete for essential
nutrients (Ringø and Gatesoupe, 1998). The spore-forming ability of the *Bacillus* spp. used in
this study may perhaps aid in administering their effect. This ability increases their chance to
survive the harsh conditions in the intestine and the activities of the digestive enzymes. To
further explore the beneficial effects of these probiotic bacteria, we suggest assessing their
effects on the biochemical and hematological parameters of the fish. Additionally, the ability of
these probiotics in preventing the infection of other pathogenic bacteria should also be
considered.

Conclusion

This study is the first report on the application of probiotic *Bacillus* spp. isolated from *E.*
eugeniae in fish. The results of this study revealed the beneficial effect of *Bacillus* spp. isolated
from *E. eugeniae* on the growth performance, feed utilization and disease resistance of Nile
tilapia.

Acknowledgement

The authors acknowledge the financial support from the Department of Science and
Technology - Accelerated Science and Technology Human Resource Development Program

(DOST-ASTHRDP). Also, the authors acknowledge the assistance from the Fisheries

Biotechnology Center, College of Fisheries, and Freshwater Aquaculture Center, CLSU.

References

Aly, S.M., Mohamed, M.F., John, G., 2008. Effect of probiotics on the survival, growth and
challenge Infection in Tilapia nilotica (*Oreochromis niloticus*). Aquac. Res. 39, 647–656.
https://doi.org/10.1111/j.1365-2109.2008.01932.x

Amenu, D., 2015. Probiotic Properties of Lactic Acid Bacteria from Human Milk. J. Med.
Microbiol. Diagn. s3. https://doi.org/10.4172/2161-0703.S3-005

Balcazar, J., Blas, I., Ruizzarzuela, I., Cunningham, D., Vendrell, D., Muzquiz, J., 2006. The role
of probiotics in aquaculture. Vet. Microbiol. 114, 173–186.
https://doi.org/10.1016/j.vetmic.2006.01.009

Boaru, A., Struți, D., Dărăban, S.V., Georgescu, B., 2016. The effect of using earthworm meal
(*Eisenia foetida*) as protein supplement for the growth of *Xiphophorus hellerii* juveniles.
Poeciliid Res. 6, 4–9.

Brashears, M.M., Jaroni, D., Trimble, J., 2003. Isolation, selection, and characterization of lactic
acid bacteria for a competitive exclusion product to reduce shedding of *Escherichia coli*
O157:H7 in Cattle. J. Food Prot. 66, 355–363. https://doi.org/10.4315/0362-028X-
66.3.355

Buruian, C.-T., Profir, A.G., Vizreanu, C., 2014. Effects of probiotic *Bacillus* species in
aquaculture – an overview. Food Technol. 38, 9–17.

Byzov, B.A., Nechitaylo, T.Yu., Bumazhkin, B.K., Kurakov, A.V., Golyshin, P.N., Zvyagintsev,
D.G., 2009. Culturable microorganisms from the earthworm digestive tract. Microbiology
78, 360–368. https://doi.org/10.1134/S0026261709030151

Cross, M., 2002. Microbes versus microbes: immune signals generated by probiotic lactobacilli
and their role in protection against microbial pathogens. FEMS Immunol. Med.
Microbiol. 34, 245–253.

Demir, E., Başbülbül, G., 2017. Screening of bacteriocin production in lactic acid bacteria
isolated from fermented dairy products. Biotechnol. J. Int. 18, 1–9.
https://doi.org/10.9734/BJI/2017/33504

EL-Haroun, E.R., Goda, A.M.A.-S., Kabir Chowdhury, M.A., 2006. Effect of dietary probiotic
Biogen® supplementation as a growth promoter on growth performance and feed
utilization of Nile tilapia *Oreochromis niloticus* (L.). Aquac. Res. 37, 1473–1480.
https://doi.org/10.1111/j.1365-2109.2006.01584.x

El-Kamali, H.H., 2000. Folk medicinal use of some animal products in Central Sudan. J.
Ethnopharmacol. 72, 279–282.

Feng, Y., Qiao, L., Liu, R., Yao, H., Gao, C., 2017. Potential probiotic properties of lactic acid
bacteria isolated from the intestinal mucosa of healthy piglets. Ann. Microbiol. 67, 239–
253. https://doi.org/10.1007/s13213-017-1254-6

Govindarajan, B., Prabaharan, V., 2015. Gut bacterial load analysis of earthworms (*Eudrilus*
eugeniae)–a controlled laboratory study. Eur. J. Environ. Ecol. 2, 38–43.

Gupta, A., Gupta, P., Dhawan, A., 2014. Dietary supplementation of probiotics affects growth,
immune response and disease resistance of *Cyprinus carpio* fry. Fish Shellfish Immunol.
41, 113–119. https://doi.org/10.1016/j.fsi.2014.08.023

Horn, M.A., Schramm, A., Drake, H.L., 2003. The Earthworm Gut: an Ideal Habitat for Ingested
N2O-Producing Microorganisms. Appl. Environ. Microbiol. 69, 1662–1669.
https://doi.org/10.1128/AEM.69.3.1662-1669.2003

Ismail, S.A., 2005. The Earthworm Book. Other India Press, Mapusa, Goa.

Jena, P.K., Trivedi, D., Thakore, K., Chaudhary, H., Giri, S.S., Seshadri, S., 2013. Isolation and
characterization of probiotic properties of Lactobacilli isolated from rat fecal microbiota.
Microbiol. Immunol. 57, 407–416. https://doi.org/10.1111/1348-0421.12054
Kim, H.-J., Shin, K.-H., Cha, C.-J., Hur, H.-G., 2004. Analysis of aerobic and culturable
bacterial community structures in earthworm (Eisenia fetida) intestine. Agric. Chem.
Biotechnol. 47, 137–142.
Lamari, F., Sadok, K., Bakhrouf, A., Gatesoupe, F.-J., 2014. Selection of lactic acid bacteria as
candidate probiotics and in vivo test on *Artemia nauplii*. Aquac. Int. 22, 699–709.
https://doi.org/10.1007/s10499-013-9699-5
Leite, A.M.O., Miguel, M.A.L., Peixoto, R.S., Ruas-Madiedo, P., Paschoalin, V.M.F., Mayo, B.,
Delgado, S., 2015. Probiotic potential of selected lactic acid bacteria strains isolated from
Brazilian kefir grains. J. Dairy Sci. 98, 3622–3632. https://doi.org/10.3168/jds.2014-9265
Lin, Y.-H., Chen, Y.-S., Wu, H.-C., Pan, S.-F., Yu, B., Chiang, C.-M., Chiu, C.-M., Yanagida,
F., 2013. Screening and characterization of LAB-produced bacteriocin-like substances
from the intestine of grey mullet (*Mugil cephalus* L.) as potential biocontrol agents in
aquaculture. J. Appl. Microbiol. 114, 299–307. https://doi.org/10.1111/jam.12041
Mohanta, K.N., Subramanian, S., Korikanthimath, V.S., 2016. Potential of earthworm (*Eisenia*
foetida) as dietary protein source for rohu (*Labeo rohita*) advanced fry. Cogent Food
Agric. 2. https://doi.org/10.1080/23311932.2016.1138594
Musikasang, H., Tani, A., H-kittikun, A., Maneerat, S., 2009. Probiotic potential of lactic acid
bacteria isolated from chicken gastrointestinal digestive tract. World J. Microbiol.
Biotechnol. 25, 1337–1345. https://doi.org/10.1007/s11274-009-0020-8
Ranganathan, L.S., 2006. Vermibiotechnology: from soil health to human health. Agrobios
(India).
Reis, N.A., Saraiva, M.A.F., Duarte, E.A.A., de Carvalho, E.A., Vieira, B.B., Evangelista-
Barreto, N.S., 2016. Probiotic properties of lactic acid bacteria isolated from human milk.
J. Appl. Microbiol. 121, 811–820. https://doi.org/10.1111/jam.13173
Ringø, E., Gatesoupe, F., 1998. Lactic acid bacteria in fish. Aquaculture 160, 177–20.
Ruangroupan, L., Kitao, T., Yoshida, T., 1986. Protective efficacy of *Aeromonas hydrophila*
vaccines in Nile tilapia. Vet. Immunol. Immunopathol. 12, 345–350.
Sanchez-Ortiz, A.C., Luna-Gonzalez, A., Campa-Cordova, A.I., Escamilla-Montes, R., Flores-
Miranda, M. del C., Mazon-Suastegui, J.M., 2015. Isolation and characterization of
potential probiotic bacteria from pustulose ark (*Anadara tuberculosa*) suitable for shrimp
farming. Lat. Am. J. Aquat. Res. 43, 123–136. https://doi.org/10.3856/vol43-issue1-
fulltext-11
Selim, K.M., Reda, R.M., 2015. Improvement of immunity and disease resistance in the Nile
tilapia, *Oreochromis niloticus*, by dietary supplementation with *Bacillus*
amyloliquefaciens. Fish Shellfish Immunol. 44, 496–503.
https://doi.org/10.1016/j.fsi.2015.03.004
Shelby, R.A., Lim, C., Yildirim-Aksoy, M., Delaney, M.A., 2006. Effects of probiotic diet
supplements on disease resistance and immune response of young Nile tilapia,
Oreochromis niloticus. J. Appl. Aquac. 18, 23–34.
https://doi.org/10.1300/J028v18n02_02
Silva, T.F.A., Petrillo, T.R., Yunis-Aguinaga, J., Fernandes, P., 2015. Effects of the probiotic
Bacillus amyloliquefaciens on growth performance, hematology and intestinal

morphometry in cage-reared Nile tilapia. Lat. Am. J. Aquat. Res. 43, 963–971.
https://doi.org/10.3856/vol43-issue5-fulltext-16
Singleton, D.R., Hendrix, P.F., Coleman, D.C., Whitman, W.B., 2003. Identification of
uncultured bacteria tightly associated with the intestine of the earthworm Lumbricus
rubellus (Lumbricidae; Oligochaeta). Soil Biol. Biochem. 35, 1547–1555.
https://doi.org/10.1016/S0038-0717(03)00244-X
Soltan, M.A., Fouad, I.M., Elfeky, A., 2016. Growth and feed utilization of Nile tilapia,
Oreochromis niloticus fed diets containing probiotic. Glob. Vet. 17, 442–450.
https://doi.org/10.5829/idosi.gv.2016.442.450
Sutthi, N., Thaimuangphol, W., Rodmongkoldee, M., Leelapatra, W., Panase, P., 2018. Growth
performances, survival rate, and biochemical parameters of Nile tilapia (*Oreochromis*
niloticus) reared in water treated with probiotic. Comp. Clin. Pathol.
https://doi.org/10.1007/s00580-017-2633-x
Zhou, X., Tian, Z., Wang, Y., Li, W., 2009. Effect of treatment with probiotics as water additive
on tilapia (*Oreochromis niloticus*) growth performance and immune response. Fish
Physiol. Biochem. 36, 501–9. https://doi.org/10.1007/s10695-009-9320-z
296