Force-dependent recruitment of Piezo1 drives focal adhesion maturation and calcium entry in normal but not tumor cells

Mingxi Yao*[1,2,3], Ajay Tijore[3,4], Delfine Cheng[5], Jinyuan Vero Li[5], Anushya Hariharan[3], Boris Martinac[5], Guy

Tran Van Nhieu[6], Charles D Cox*[5] & Michael Sheetz*[3,7,8]

1. Department of Biomedical Engineering, Southern University of Science and Technology
2. Guangdong Provincial Key Laboratory of Advanced Biomaterials
3. Mechanobiology Institute, National University of Singapore Mechanobiology Institute, National University of Singapore
4. Centre for Biosystems Science and Engineering, Indian Institute of Science, Bangalore, 560012, India
5. Victor Chang Cardiac Research Institute
6. Ecole Normale Supérieure PARIS-SACLAY
7. Dept of Biological Sciences, National University of Singapore
8. Molecular MechanoMedicine Program, Dept. of Biochemistry and Molecular Biology, University of Texas Medical Branch

*Correspondence: yaomx@sustech.edu.cn, C.Cox@victorchang.edu.au or misheetz@utmb.edu

Significance Statement

Piezo1 localizes to focal adhesions under force in a cell-state dependent manner and faciliates calcium

entry and adhesion matruration/turnover.

Abstract:

Mechanosensing is an integral part of many physiological processes including stem cell differentiation,

fibrosis and cancer progression. Two major mechanosensing systems – focal adhesions and mechanosensitive

ion channels, can convert mechanical features of the microenvironment into biochemical signals. We report

here surprisingly that the mechanosensitive Ca^{2+}-permeable channel Piezo1, previously perceived to be

diffusive on plasma membranes, binds to matrix adhesions in a force-dependent manner, promoting adhesion

maturation and cell spreading in normal but not in tumor cells. In the absence of Piezo1, matrix adhesions are

smaller in normal cells mimicking transformed cells where adhesions do not change with or without Piezo1.

A novel adhesion-targeted calcium sensor shows robust Piezo1-dependent, calcium influx at adhesions in

30 normal cells; but not in transformed cells. A linker domain in Piezo1 is needed for binding to adhesions and

31 overexpression of the domain blocks Piezo1 binding to adhesions decreasing adhesion size and cell spread

32 area. Thus, we suggest that Piezo1 is a novel component of focal adhesions in non-transformed cells that

33 catalyzes adhesion maturation and growth through force-dependent calcium signaling, but this function is

34 absent in most cancer cells.

Introduction

Cellular mechanosensing, the cell's ability to detect and respond to its physical microenvironment, is

involved in important processes including development, tissue homeostasis and disease(*1*, *2*). For example,

sensing of shear flow in blood circulation is essential for the proper development of the vasculature and the

depletion of rigidity sensors enables cancer neoplastic transformation, i.e. substrate-rigidity-independent

growth(*3*, *4*). Mechanosensitive ion channels and integrin-mediated focal adhesions, are major

mechanosensors that are responsible for converting mechanical cues into biochemical signals in these

processes(*5*).

A mechanosensing function of focal adhesions is manifest by rigidity-dependent adhesion assembly and

turnover, which involves hundreds of different focal adhesion proteins that alter cell behavior during

spreading and migration(*1*). In addition, focal adhesion mechanosensing regulates pathways such as AKT,

MAPK and Hippo-signaling that influence cell fate decisions(*1*, *6*, *7*). Because adhesions affect cell behavior

and fate, the molecular components and dynamics of focal adhesions have been extensively studied. There

exists a robust process of adhesion protein recruitment and processing in the finely tuned steps of focal

adhesion formation, growth and turnover that correlates with cell state decision making based on cell-

microenvironment interactions(*8*). There is evidence for force-sensing proteins such as talin and vinculin

regulating the functions of focal adhesions as well(*2*, *8–13*). In contrast, although the ion conduction

properties of many mechanosensitive ion channels are well-characterized by structural and patch-clamp

studies, the mechanisms for converting matrix forces to channel gating are not well understood(*14*). This is

partly due to the complex nature of ion channel signaling and mechanically-induced calcium influx. For

example, mechanosensitive channels are linked to both cell proliferation and apoptosis depending on the

context, suggesting that both the distribution and dynamics of the mechanosensitive ion channels are

important for understanding the critical mechanosensing processes of the cell(*15*, *16*).

There is ample evidence of a crosstalk between mechanosensitive ion channels and focal adhesions. Local

calcium entry is critical for focal adhesion assembly and disassembly based upon the calpain dependence of

both processes (*17–21*). In adhesion assembly during cell spreading, rigidity sensing involves force-dependent

calpain cleavage of talin, which is needed for normal adhesions to form and for normal cells to grow(*20*). In

contrast, transformed cells do not sense matrix rigidity and adhesions are significantly smaller even though

tumor cells develop higher traction forces(*22*). This raises questions of whether or not transformed cells use

calcium in adhesion assembly like normal cells. In addition, several studies found that adhesion disassembly

depends upon force-dependent calpain cleavage(*18*, *19*). Thus, there are critical roles for calcium entry in

normal adhesion dynamics that are altered in transformed (tumor) cells and the analysis of the channels

involved can reveal important aspects of the life cycle of focal adhesions.

Several calcium-permeable channels have been implicated in focal adhesions dynamics (*23–27*) . One of those,

Piezo1, is particularly interesting, as early reports of Piezo1 document an interaction with integrins(*28*). Piezo1

has subsequently been identified as a bona-fide mechanosensitive ion channel in mammalian cells(*29*). Since

then, Piezo1 has been shown to have many roles in important biological processes and there is increasing

interest in how it functions *in situ*(*15*, *29–31*). Piezo1 is widely expressed in somatic cell types and is linked to

mechanosensing in stem cell lineage determination, blood pressure regulation, activation of innate immunity

and vascular development(*4*, *30*, *32*, *33*). Piezo1's activity has been shown to be related to the integrin-linked

actin cytoskeleton (*26*, *34*, *35*). When Neuro2A cells are mechanically agitated with micron-sized, collagen IV

or Matrigel coated beads, the tension threshold required to trigger Piezo-dependent Ca^{2+} entry is significantly

reduced(*36*). In addition, integrin signaling pathways are critical for proper function of Piezo1 in the context

of endothelial cell responses to flow(*37*). Consequently, we want to investigate the possible crosstalk between

Piezo1 and focal adhesions.

In this study, we show that in normal cell lines Piezo1 is needed for mature adhesion assembly and it

localizes to adhesions as the adhesions mature . In normal cell lines from many tissues, Piezo1 is enriched at

focal adhesions in a force-dependent manner and dissociates after the integrin adhesions disassemble. Its

localization correlates with local calcium entry and adhesion disassembly likely through a calpain-dependent

process[20]. Alternatively, in transformed cells, Piezo1 is diffusely distributed and does not affect either

adhesion assembly or disassembly and there is low adhesion calcium entry. These studies indicate that Piezo1

normally interacts with adhesions transiently during assembly and then its force-dependent recruitment to

mature adhesions catalyzes integrin-specific adhesion disassembly. These dynamic adhesion interactions are

lost in transformed cells and are likely a key neoplastic modification.

Results

Piezo1 stably localizes to mature adhesions in HFF cells

To investigate the role of Piezo1 in adhesion formation, we followed the distribution and dynamics of

Piezo1 localization during cell spreading in Human Foreskin Fibroblast (HFF) cells using total internal

reflection fluorescence microscopy (TIRFM). Transiently expressed Piezo1 tagged with GFP(*38*) or mRuby3

and paxillin-BFP were followed during the cell spreading in HFF cells (Figure 1A top panel, supplementary

movie 1). In the initial phase of spreading, Piezo1 did not colocalize with cell adhesions marked by paxillin

and it appeared concentrated around the nucleus. After 30-60 minutes when the cells developed mature

adhesions, Piezo1 dispersed to the entire spread area of the cell and was significantly enriched in maturing

adhesions.

Piezo1 localization to adhesions increased over time until the cell began to polarize. Upon cell

polarization, Piezo1 prominently localized to the retracting edges (Figure 1B, upper panel). After spreading

overnight, HFF cells reached a dynamic steady state and Piezo1 was enriched at mature focal adhesions

(Figure 1B, middle panel). In most cells, Piezo1 was recruited to a sub population of the focal adhesions,

mostly around the cell periphery at the ends of stress fibers. Quantification of the extent of Piezo1

recuitment to focal adhesions by the ratio of Piezo1 intensity in the adhesion to that in the plasma

membrane background showed that the recruitment of Piezo1 to adhesions increased over time as cells

spread further and started to polarize (Figure 1 C-D). Piezo1 was often recruited to a portion of individual

adhesions as well (Figure 1B lower panel, marked by arrow), implying that the regulation of Piezo1

recruitment happened locally. The localization of endogenous Piezo1 to adhesions in HFF cells was also

observed by Piezo1 immunostaining of fixed HFF cells (Figure S1) using two knockdown/knockout validated

antibodies(*15*). Thus, Piezo1 complexed with mature and retracting adhesions in polarized cells on

fibronectin.

Since Piezo1 was previously reported to be diffusive in plasma membranes(*26*), we confirmed that the

Piezo1 recruitment involved immobilization by checking the lifetime of Piezo1 in the focal adhesions of HFF

cells through Fluorescence Recovery After Photobleaching (FRAP). Surprisingly, as shown in Figure 1E, Piezo1

was stable at the adhesions after recruitment, with lifetimes of around 120 s that were considerably longer

compared to typical focal adhesion proteins such as paxillin ~25 s and even slightly longer than integrins

such as integrin $\beta 3$ ~90 s (Figure 1F). In addition, the mobile fraction of Piezo1 was considerably less than the

mobile fraction of paxillin and even integrin $\beta 3$ (Figure 1F). These data indicated that Piezo1 formed stable

interactions with certain focal adhesion components in HFF cells.

Piezo1 recruitment to adhesions requires myosin II contractility

Since focal adhesion maturation was linked to sustained traction forces and correlated with Piezo1

recruitment during spreading, we hypothesized that Piezo1-adhesion interactions were force dependent.

Treatment of Piezo1-GFP expressing HFF cells with 30 µM of Y-27632 to inhibit actomyosin contraction,

caused rapid loss of Piezo1 from adhesion sites and was followed by the disassembly of focal adhesions as

measured by paxillin (Figure 2A, top panel, supplementary movie2). Similar results were observed with 50

µM blebbistatin treatment (Figure S2A). When Y-27632 was washed out, Piezo1 returned rapidly to the

reforming adhesions (Figure 2A, top panel). Since Piezo1 recruitment during recovery was much more rapid

than the initial recruitment to adhesions during cell spreading (minutes versus hours), we postulated that

components remaining at adhesions in the absence of force catalyzed the rapid recovery of adhesion

proteins with the development of contractile forces to enable Piezo1 binding. Further, when the

contractility of the cell was enhanced by co-expressing the constitutively active version of RhoA along with

Piezo1 in HFF cells, the enrichment of Piezo1 at adhesions was strongly promoted (Figure 2A, bottom panel).

To check whether the reduction of traction forces alone could disrupt Piezo1's adhesion recruitment

without disruption of adhesion proteins such as talin and integrins, we tested the relationship between

Piezo1 localization and myosin II contractility in intact adhesions. Previous reports showed that inhibition of

the tyrosine phosphatase, PTP-PEST, as well as the expression of virulence bacterial protein IpaA, inhibited

adhesion disassembly upon addition of Y-27632 (*39*, *40*). In cells expressing GFP-labelled IpaA peptide,

addition of 30 µM Y-27632 caused cells to lose contractility while retaining adhesion structures (Figure 2B,

Supplemental movie 3). In this case, Piezo1-mRuby3 diffused out from adhesions within 10 min after Y-

27632 addition while the integrin β3-BFP remained intact. A similar effect was observed when 2.5 µM of

PTP-PEST inhibitor was added in combination with 30 µM Y-27632 (Figure S2B). Thus, the localization of

Piezo1 to focal adhesions depended on continued contraction forces possibly through the stretching of

adhesion proteins, and was independent of paxillin or integrin β3 localization to adhesions.

To further establish the role of myosin II-based contractility in Piezo1 recruitment, we checked the

distribution of Piezo1 in cells that had low traction forces such as COS-7 cells lacking myosin IIA

expression(*41*). In COS-7 cells transiently expressing Piezo1-GFP and paxillin-mapple, Piezo1 was diffusive in

the membrane with no apparent localization to adhesions (Figure 2C, top panel) and the cell adopted a

highly-spread disc morphology. When contractility was restored in COS-7 cells by overexpressing myosin IIA,

cells became polarized with myosin IIA forming stress-fiber-like cables (Figure 2C, bottom panel) and Piezo1

was recruited to the adhesions at the ends of the myosin cables. This further reinforced the hypothesis that

Piezo1 recruitment to adhesions required continued force on the adhesion complex.

Piezo1 recruitment triggers integrin β3 adhesion turnover

Since Piezo1 was recruited to only a fraction of focal adhesions, we proceeded to check whether different

integrin subtypes played a role in the regulation of Piezo1 localization. As shown in Figure 3A, we transiently

expressed Piezo1-1591-mRuby3 with integrin β3-mEmerald and analyzed the colocalization of Piezo1 with β3

adhesions during cycles of adhesion assembly and disassembly. Interestingly, the recruitment of Piezo1 to β3-

enriched adhesions correlated with the disassembly and turnover of the β3 adhesions. Figure 3A shows the

timelapse views of two such turnover events on fibronectin surfaces, one during spreading where the cell was

protruding and the other in migrating cells where the adhesion was retracting. In both cases, integrin β3-

based adhesions formed without apparent Piezo1 co-localization. Over time, Piezo1 was recruited to the

adhesions from the periphery of the adhesions and gradually occupied the whole adhesion. Piezo 1

recruitment appeared to initiate a decrease in integrin β3 fluorescence intensity within minutes (Figure 3A-B,

Figure S3A and supplementary movie 4-6). After integrin β3 reached background levels, the Piezo1

fluorescence often persisted in adhesions for a few minutes before dissociation (Figure 3A-B, marked by

arrows; and Figure 3 D for kymograph), suggesting that some adhesion components remained that supported

Piezo1 localization after integrin β3 dissociated from the adhesion.

In contrast, the anticorrelated behavior was not observed between Piezo1 and β1 integrins in adhesions.

Piezo1 recruitment at integrin β3 and β1 enriched adhesions was more stable, and they often co-localized for

tens of minutes ((Figure 3C, 3E for kymograph). In another interesting example (Figure S3B), Piezo1

recruitment correlated with the rapid dissociation of β3 integrins and enrichment of β1 integrins. Piezo1

recruitment persisted in adhesions even after dissociation of β3 integrins (Fig. S3B, arrows). These

observations indicated that Piezo1 recruitment was associated with adhesion turnover that was regulated

locally in an integrin β3 dependent manner. Considering Piezo1's function as a mechanosensitive Ca^{2+} channel,

adhesion turnover is likely caused by the calpain-dependent cleavage pathway. Based upon these findings we

hypothesized that Piezo1 recruitment was upstream of calpain in regulating adhesion turnover. In earlier

studies of cells on fibronectin, integrin β3 adhesions formed initially, and then the adhesions switched to

integrin β1. The greater stability of β1-containing adhesions with Piezo1 recruitment could potentially provide

a mechanistic explanation for such switching events(*42*).

Piezo1 localization to adhesions is lost in transformed cells

To check whether Piezo1 localization to adhesions was cell-type specific, we expressed Piezo1 in a variety

of normal and transformed cell lines (Figure 4A, Figure S4A). In all of the normal cell lines of fibroblastic (HFF,

MCF-10A), epithelial (HUMEC) or endothelial origins (HUVEC), Piezo1 localized to a significant subset of the

focal adhesions on fibronectin demarcated by paxillin (Figure 4A-B, Figure S4). In the case of 12 different

tumor cell lines from mesenchymal or epithelial tissues (fibrosarcoma, melanoma, breast and lung cancers,

Figure 4B-D), there was no significant colocalization of Piezo1 with paxillin containing adhesions. In the

HT1080 (fibrosarcoma) and MDA-MB-231 (breast cancer) cells, FRAP measurements of Piezo1 diffusion

showed that Piezo1 had a significantly higher mobile fraction than in HFF cells (Figure 4C). The difference in

Piezo1 immobilization was not due to absence of Piezo1 expression, since Western blots confirmed significant

Piezo1 expression in most transformed cell lines (Figure 5C, Figure S4B). Thus, the colocalization of Piezo1 with

adhesions was found only in normal cells. In all of the tumor cells tested, Piezo1 was not colocalizing with

adhesions and was rapidly diffusing.

Our previous studies showed that TPM 2.1 levels were low in many transformed cells(*22*). Knocking

down TPM 2.1 in HFF cells caused the cells to become transformed with smaller adhesions and the cells

were not able to differentiate between soft and rigid surfaces. In contrast, restoring TPM 2.1 expression in

MDA-MB-231 cells caused normal behavior with bigger adhesions and the cells died on soft agar(*22*). Based

on this, we checked if toggling between the non-transformed and transformed states in HFF and MDA-MB-

231 cells altered the localization of Piezo1 in these cell lines. When we knocked down TPM 2.1 in HFF cells,

the recruitment of Piezo1 to adhesions was significantly weakened compared with control cells or control

siRNA-treated cells (Figure S5A). In contrast, when we transiently expressed TPM 2.1-YFP in MDA-MB-231

cells, a fraction of cells formed stress fibers and Piezo1 was recruited to adhesions at the end of stress fibers

(Figure S5B). These data indicated that there was a correlation between normal cell behavior and Piezo1

adhesion localization and we recently showed that transformation resulted from the loss of rigidity sensing,

which was observed in the majority tumor cells(*22*). Thus, the change in cell state from normal to

transformed state diminished Piezo1 binding to adhesions.

Piezo1 is needed for adhesion maturation and polarity in normal but not in transformed cells

As the pattern of Piezo1 adhesion recruitment differs in normal and transformed cell lines, we checked if

Piezo1 regulated focal adhesion morphology differently in normal and transformed cell lines. To minimize

the possibility of cell-type variance, we used normal primary cells and transformed cancer cells of matching

origin (Human foreskin fibroblast with HT1080 human fibrosarcoma). As shown in Figure 5A, Piezo1 knock-

down HFF cells spread to smaller areas, had a lower aspect ratio and had smaller adhesions, while in the

transformed HT1080 cells, knocking down of Piezo1 caused a significant increase in cell spread area but had

no effect on aspect ratio or adhesion size (Figure 5 A-C).

This indicated that Piezo1 was important for adhesion maturation and cell polarization in normal cells

but had little effect on adhesion formation in transformed cells, while affecting spreading area. Previous

studies documented that transformed cell lines including cancer cells generally had smaller and less mature

adhesions compared with non-transformed cells(*22*, *43*). This was despite the fact that the majority of

adhesion components were present at similar levels in both normal and transformed cells(*44*, *45*). Thus,

Piezo1 recruitment to adhesions in normal cells appeared to be involved in important steps in the adhesion

maturation process whereas in transformed cells, Piezo1 depletion caused no change in adhesion

morphology(*46*).

Piezo1's adhesion recruitment leads to local calcium transients around adhesions in normal but not in transformed cells.

Next we checked the possible physiological significance of the Piezo1 adhesion recruitment. Since Piezo1 functions as a mechanosensitive ion channel, the recruitment of Piezo1 to adhesions may have altered the calcium entry pattern of the cells. Recent studies by Ellefsen et. al, showed that Piezo1-dependent calcium spikes were localized to the cell periphery near focal adhesions[26]. This implied that calcium spikes occurred near where there were concentrations of Piezo1 at adhesions. Because Piezo1 was not concentrated in adhesions of transformed cells such as MDA-MB-231 and HEK293T, there was a question of whether or not similar calcium spikes would occur near adhesions in transformed cells.

To determine if calcium entry near adhesions in fibroblasts also occurred in tumor cells, we loaded HFF and MDA-MB-231 cells with the cell-permeable calcium indicator Cal-520 and imaged transient calcium "puffs" on fibronectin surfaces. Similar to previous observations in HFF cells[26] , many local calcium entry events were detected in areas of high contractility at the cell edge and surrounding adhesions (Figure S6). Interestingly, although the transformed MDA-MB-231 cells generated higher local traction forces[22] and were killed by calcium-dependent apoptosis after stretch[16], they had very few calcium spikes on fibronectin surfaces (Figure S6). This indicated that the calcium influx in cancer cells was abnormal and was consistent with the lack of Piezo1 within adhesions after transformation.

Although the measurements with the Cal-520 calcium sensor provided interesting insights into the differences in spatial calcium signaling between normal and transformed cells, there remained several specific questions about whether these calcium fluctuations were localized to sites of adhesion. The diffusive nature of all Ca^{2+} dyes reduces the signal-to-noise ratio for calcium influx at sub-cellular locations, providing only limited spatio-temporal information on local Ca^{2+} changes. In addition, under imaging conditions sensitive to subcellular calcium events, Cal-520 had significant photo toxicity that caused rapid increases in intracellular calcium levels limiting the observation window. To better quantify and understand

the details of calcium entry at the adhesions, we generated a novel focal adhesion targeted calcium sensor,

in which one of the latest most sensitive recombinant calcium sensors, jGCaMP7s, was linked to a paxillin

mScarlet-I construct (Figure 6A)[68]. This improved the signal-to-noise ratio specifically at the adhesions, and

the ratio between the calcium sensor and the mscarlet-I fluorescence enabled normalization of the calcium

levels based on the local sensor concentration.

When the paxillin calcium sensor was transfected in HFF, HT1080 and MDA-MB-231 cells, it localized

robustly to apparent adhesions. After overnight spreading, the intensity ratio of jGCaMP7s and mScarlet-I in

the focal adhesions of HFF cells showed frequent calcium spikes (Figure 6B,supplementary movie 7). The

calcium fluctuations were greatly reduced in Piezo1 knock down cells, indicating that Piezo1 was required for

the calcium fluctuations around the adhesions (Figure 6B-C,supplementary movie 8). Interestingly, in the

transformed HT1080 and MDA-MB-231 cells the calcium spikes at the adhesions were significantly weaker,

often there was no detectable calcium signal during a 1-hour observation window(Figure 6B,supplementary

movie 9-10). In the transformed cells, matrix adhesions still formed and disassembled, suggesting that

calcium entry around adhesions was not required for adhesion turnover in the transformed cells. These

results indicated that Piezo1 recruitment to adhesions in HFF cells enabled calcium spikes that could in turn

induce adhesion turnover; but in transformed cells, adhesion dynamics and cellular calcium fluctuations

were de-coupled.

The central linker domain (1418-1656) of Piezo1 is required for Piezo1 adhesion

recruitment and alters cell morphology of non-transformed cells

We proceeded to investigate potential molecular mechanisms of Piezo1 adhesion recruitment. So far, no

binding partners of Piezo1 at focal adhesions have been reported in the literature. The long lifetime of Piezo1

at the adhesions and the dependence of Piezo1 recruitment on actomyosin contractility led us to hypothesize

that Piezo1 bound to adhesion-linked components of the actin cytoskeleton stretched by cellular contractility.

Upon examining the 3D architecture of Piezo1(*47*), we identified a long intracellular linker as a potential

interaction site between Piezo1 and the actin cytoskeleton, as it has the most exposed intracellular region and

is in close proximity to the CTD and beam region that is critical for its force dependent activation (Figure

7A)(*48*).

This long intracellular linker region was largely unresolved in the Cryo-EM structures and connected the

beam to the clasp and via another unresolved loop to Piezo1 repeat B(*47, 49*). Previous biophysical studies

showed that deletions at the proximal portion of the loop influenced the biophysical properties of the channel

via effects on the latch domain(*50*). However the distal end was not explored in those studies.

To check the potential function of the linker region of Piezo1, we first made a mNeongreen construct of the

larger region of human Piezo1 (residue 1418-1656;mNeonlinker) that contained the unstructured loop region,

the clasp and a second disordered loop previously shown to be able to house fluorescent tags(*38*). We next

checked for its localization and interaction with Piezo1. As shown in Figure 7B, the mNeonlinker construct

expressed in HFF cells appeared to be diffusely distributed with no apparent concentrations. However, Piezo1

recruitment to adhesions was reduced as was adhesion size with overexpression of the mNeonlinker (Figure

7B). This is consistent with the hypothesis that at high expression levels the mNeonlinker domain acted as a

dominant negative competitor and displaced Piezo1 from adhesions. When we checked the morphology of

the cells, HFF cells had significantly lower cell spread area and aspect ratio with the mNeonlinker over-

expression, which resembled the morphology change of HFF cells after Piezo1 knock-down (Figure 7B-C).

These results were consistent with the hypothesis that the recruitment of Piezo1 to focal adhesions in non-

transformed cells activated local calcium fluctuations in surrounding adhesions, and the calcium signal led to

adhesion maturation and cell polarization. In the case of transformed cells where Piezo1 did not localize to

adhesions, expression of mNeonlinker should have caused minimal effect on cell morphology. Indeed,

expression of mNeonlinker in HT1080 cells caused no change in morphology, cell spread area or aspect ratio

compared with controls (Figure 7B-C).

To further check if this linker region of Piezo1 indeed harboured focal adhesion binding sites, we generated

several Piezo1 mutants with the linker regions deleted (short: Piezo1 Δ1443-1556and long: Piezo1 Δ1443-

1473). The Δ1443-1473 deletion removed only residues from the oop region and the larger deletion Δ1443-

1556 removed much of the clasp domain. When we expressed these deletion mutants in FAK -/- MEF cells

that had prominent recruitment of WT Piezo1 to adhesions, we observed normal Piezo1 adhesion

recruitment of Piezo1 Δ1443-1473 mutant and diffusive Piezo1 localization of the Δ1443-1556 mutant (Figure

S7 and supplementary movie11-12). Electrophysiological characterization in Piezo1 KO HEK293T cells showed

that the Δ1443-1473 mutant had normal stretch induced channel activity while the Δ1443-1556 mutant was

not functional (Figure S8).

Taking these data together, it appeared that the linker domain (residues 1418-1656) and most likely the clasp

region of Piezo1 harbored a binding site for focal adhesion components that recruited Piezo1 to focal

adhesions in a force-dependent manner in non-transformed cells at the membrane-cytoplasm interface.

Discussion

In this study, we show that Piezo1 has important functions in focal adhesion dynamics in normal cells

but not in transformed cells. Normal fibroblasts, epithelial and endothelial cells form focal adhesions on

fibronectin where Piezo1 concentrates in a force-dependent process. The lifetime of Piezo1 in the adhesions

is similar to that of integrins and longer than typical adhesion proteins, which implies that it complexes

stably with adhesion bound proteins. Sustained contraction forces are needed for Piezo1 localization to

adhesions; and when contraction is inhibited, it could leave adhesions before the dissociation of adhesion

components such as integrin and talin. Piezo1 localization to focal adhesions leads to the dissociation of

integrin β3 containing adhesions but not integrin β1 adhesions on fibronectin. This localization is also lost in

transformed cells, including virus-transformed somatic cells (e.g. HEK293T) as well as tumor cell lines from

different tissues. When somatic cells are transformed such as in the case of TPM 2.1 knock down, the

recruitment of Piezo1 to adhesions is dramatically reduced similar to the transformed cancer lines tested.

Calcium levels in the vicinity of adhesions are higher and have many more local fluctuations in normal cells

compared with transformed cells. Binding to adhesions depends upon an unstructured cytoplasmic domain

of Piezo1 (aa 1418-1656). These findings indicate that Piezo1 has a number of complex interactions, which

depend upon contractility, the transformed state, as well as specific integrin-matrix interactions. Piezo1

appears to have multiple context-dependent functions in normal and transformed cells that need to be

analyzed temporally and spatially in a defined mechanical environment.

Piezo1 was first identified as an integrin-interacting protein which had a role in adhesion formation(*28*).

Our results reinforce those findings and further indicate that Piezo1 binds in a complex with integrins but

not with adhesion proteins such as talin or paxillin, which have much more rapid recovery rates after

photobleaching. Further, inhibition of PTPN12 activity or expression of the *Shigella* IpaA effector stabilizes

adhesion proteins even after inhibition of myosin but does not prevent Piezo1 dissociation. Piezo1's rapid

dissociation upon myosin inhibition is rapidly reversed upon washout of the myosin inhibitor. The rate of

return to the adhesions is much more rapid than during initial adhesion formation, indicating that the

recruitment of Piezo1 at adhesions may depend on a force-sensitive protein or activation of a force-

dependent enzyme that is retained at adhesion sites during myosin inhibition. Overexpression of a linker

region of human Piezo1 (aa 1418-1656) acts as a dominate negative, displacing Piezo1 from adhesions,

implying that this linker region of Piezo1 harbours the interacting site for adhesion binding. The lack of

adhesion maturation after depletion of Piezo1 strongly supports the hypothesis that Piezo1 has an important

role in the force-dependent maturation of adhesions on rigid surfaces, and this does not occur in

transformed cells.

Since the discovery of Piezo1's role as a mechanosensitive ion channel in eukaryotic cells, the regulation

mechanism of Piezo1's function in the various physiological activities of cells has been under heavy

investigation(*51*, *52*). The classical model of Piezo1's physiological function assumes that it is diffusive in the

plasma membrane and is activated locally through membrane bilayer tension or curvature changes that

occur through cytoskeletal-dependent mechanical events(*26*, *34*, *38*, *47*, *53*). Several recent studies have

clearly shown that the cystoskeletal environment influences Piezo1 function(*38*), including the observation

that Piezo1 localizes to the adhesions of some glioblastoma cell lines, which facilitates their growth and

invasion(*27*). The unusual nature of these glioblastoma lines indicates that they may be more like normal

cells than transformed cells (Yao et al unpublished results). In our study, we systematically examined

Piezo1's spatial distribution across many cell types from different tissues of origin and discovered that Piezo1

adhesion localization is much more prominent in non-transformed normal cells compared with transformed

cells such as most tumor cell lines. Binding to the adhesions in normal cells catalyzes adhesion maturation

and disassembly that correlates with cell growth.

Interestingly, in contrast to the reduced calcium transients that we observed in resting transformed

cells, we recently reported that mechanical perturbations, such as stretching or ultrasound treatment, cause

Piezo1-dependent increases in cytoplasmic Ca^{2+} levels and subsequent apoptosis of transformed but not

normal cells[16]. These observations are difficult to explain in terms of fluctuations in either mechanical

forces on adhesions or membrane tension, especially since fewer Ca^{2+} fluctuation events are observed in

transformed cells whereas Ca^{2+} flickering events near adhesion sites are common in normal cells. With

regard to membrane tension, typical measurements of membrane tension in cells report values that are 2-3

orders of magnitude smaller than those needed to activate Piezo1 in vitro[54, 55]. Certainly, the pattern of

Ca^{2+} activation in the HFF cells in the vicinity of adhesions is consistent with previous results[46, 56, 57], but

it is not consistent with the simplistic model of very high membrane tension for the whole cell. Rather, there

appear to be local activation events that involve elements from the adhesions that may alter local

membrane tension/curvature or the association of membrane/cytoskeletal proteins that can modulate the

mechanical gating of Piezo1 activity[53, 58, 59]. Although Piezo1 appears in many different mechanical

functions, the activation mechanism of Piezo1 is complex and both contributions from membrane tension

and interaction with cytoskeletal proteins may modulate its activity[35, 58].

We recently showed that transformed cancer cells are depleted of the rigidity sensing, local contraction

units[22, 43]. There is a conserved set of cytoskeletal proteins that are able to assemble into local rigidity

sensing contractile structures in normal cells. After the depletion of any one of several cytoskeletal proteins

critical for rigidity sensing and contractile pair assembly, the cell will be transformed and will no longer sense

substrate rigidity. Transformation also involves changes in adhesion morphology and composition[22]. The

differences in the adhesion components in normal and transformed cells that contribute to the adhesion

phenotype are still an open question but the mRNA levels of over 700 proteins are altered upon

transformation by tropomyosin 2.1 depletion[22]. An important function of the rigidity sensors is in the

maturation of focal adhesions that contain Piezo1 and without Piezo1 the mature adhesions do not form.

This further fits with the absence of any differences between transformed cell adhesions with and without

Piezo1. Thus, we suggest that normally both the maturation and the disassembly of focal adhesions is

catalyzed by Piezo1. In addition, whether the differences in Piezo1 recruitment in the transformed and

somatic cell lines are caused by changes in the molecular interactions or tension states of these cells remains

an open question and awaits further investigation.

Our model of the function of Piezo1 in Figure 8 emphasizes the catalytic role of Ca^{2+} entry in the life

cycle of the adhesions in normal cells. Initially, the nascent adhesions seed actin filament formation, myosin

bipolar filaments pull on the actin in the rigidity-sensing contractions, and there is a calcium-dependent

cleavage of talin1 that is needed for growth(*8*)[,20]. If the matrix is rigid, then adhesions mature through

enzymatic modifications that trigger the maturation process. Components in the adhesions are stretched by

traction forces and then can bind Piezo1 which permits Ca^{2+} entry in a positive feedback cycle for the further

maturation of the adhesion. However, when the concentration of Piezo1 reaches a threshold, there is

excessive Ca^{2+} entry and recruitment of calpain that now hydrolyzes components of the adhesion, causing

the disassembly of the adhesion. The maturation process requires the rigidity sensing complex and will not

occur in transformed (cancer) cells. This model indicates that covalent modification of adhesion

components during maturation as well as traction forces on those components are necessary for Piezo1

binding. Further understanding of Piezo1 functions requires dynamic studies of Piezo1 at the appropriate

time and location during cell spreading or migration.

There is growing evidence that covalent modifications in maturing adhesions alter adhesion proteins as

well as possibly membrane proteins in normal cells(*20, 60*). The release of Ca^{2+} near adhesions contributes to

protein processing by the adhesion signaling hub, which is needed for adhesion maturation and cell growth

on a rigid fibronectin matrix possibly through a localized calpain-dependent pathway. Our new paxillin-

associated calcium indicator shows that the Piezo1-dependent calcium transients are much more numerous

than observed with the cytoplasmic calcium indicator. Thus, the localization of Piezo1 to the adhesions

under force will provide a very localized activation of calpain and other calcium-dependent enzymes to

modify adhesions and membrane proteins in specific locations. Besides Piezo1, other membrane channels

such as TRPV4, Orai, L-type channels, TRPM4 and TRPM7 have been implicated in the regulation of adhesion

morphology, dynamics and cell migration(*20*, *23*, *61*, *62*). The non-interchangable role of these channels

suggests that there might exist coopertivity between these channels, such as calcium-dependent calcium

release (*63*).

In transformed cells, growth appears independent of local Ca^{2+} entry at rigid matrix sites and Piezo1 is

decoupled from traction forces and adhesion signaling. It is commonly believed that transformed cells are

related to wound healing where cell growth for repair is needed and localized signaling from matrices is not

needed(*64*) . Further, in many cancers, the elevated growth rates correlate with high levels of calpain that

could amplify small Ca^{2+} leaks to support the growth activity. This could help to explain how stretch could

cause general changes in the plasma membrane activating Ca^{2+} entry through Piezo1 by local membrane

tension or curvature changes that would cascade into an apoptotic process[16]. Most importantly, the general

activation of growth in cancer cells appears to occur without localized signaling at adhesions by Piezo1.

In the larger picture of adhesion maturation, there are many adhesion proteins that bind to adhesions in

a force-dependent manner(*65*, *66*). It is surprising, however, that the inhibition of the release of many

adhesion proteins by the inhibition of PTPN12 or expression of virulence bacterial protein IpaA did not block

the release of Piezo1 upon myosin inhibition. Further, Piezo1 is strongly bound to the adhesions and has a

lifetime similar to integrin β3 in adhesions that depends on contractility. The stronger retention of Piezo1 at

adhesions upon the transition from integrin β3 to β1 may aid in the stabilization of such adhesions for

development of higher forces on matrix contacts through continued local influx of calcium(*67*). The integrin

subtype dependence of Piezo1 dynamics could be of great importance for the physiological function of

Piezo1, e.g. in vascular development(*4*). Piezo1 knock-out mice die as embryos due to vascular defects, and

inactivation of β3 inhibits the angiogenesis process. Mechanical forces such as shear forces are very

important in these processes(*4*). The force dependent co-localization and turnover of Piezo1 at β3 integrin

adhesions could be synergistic with downstream events and provides a mechanism of tension-dependent

vascular growth that is critical in development. Thus, we suggest that the contractility dependent

recruitment of Piezo1 to focal adhesions is a novel crosstalk mechanism between the two major

mechanotrasduction hubs of the cell that play critical roles in adhesion function which are lost in most

tumor cells (Figure 8) .

Material and methods

Cell culture and constructs:

The mEmerald-integrin β3, mEmerald-integrin β1, paxillin-mapple, paxillin-BFP and talin-GFP were gifts

from Michael Davidson (Addgene plasmid # 56330, #54129, #54935). The piezo1-GFP-1591 and piezo1-

mruby3-1591 plasmids were generated as previously described(*38*). The RhoAV14 construct was from

Schwartz lab(*68*). TPM2.1-YFP construct was from Gunning Lab(*69*). The A431 ipa peptide plasmid was

generated as previously described(*70*). The paxillin calcium sensor was generated by PCR cloning of

jGCaMP7s segment from pGP-CMV-jGCaMP7s plasmid from addgene(*71*) and inserted into the N-terminal of

a paxillin-mScarleti construct derived from paxillin-mapple.

siRNA for tropomyosin 2.1 was synthesized as previously described by Dharmacon(*43*). siRNA for human

Piezo1 was ordered from Dharmacon (ON-TARGETplus human Piezo1 siRNA, SMARTpool, L-020870-03-0010)

and Merck Sigma-Aldrich (SASI_HS01_00208585) or Life Technologies(cat no #21816). RT-PCR analysis

showed that both siRNA caused >90% knock down at the mRNA level two days after transfection. The siRNA

were transfected using RNAiMAX reagent from ThermoFisher following manufacture's protocols.

Human Foreskin fibroblast (HFF), HUMEC, HUVEC, HT1080, A2058, A375P, and MCF-10A cells were

obtained from ATCC. MDA-MB-231, SKOV3, cells were from Ruby Huang's lab. HEK293T cells with piezo1 KO

were from Ardem Patapoutian's lab. HFF, MDA-MB-231, HEK293T Piezo1 KO and cells were cultured in 1X

DMEM supplemented with 5 mM sodium pyruvate and 10% of Hi-FBS. MCF-10A cells were cultured using

fomulated media as reported previously(*22*) and HUVEC cells were cultured in endothelial growth media

from Sigma.

449 Ca^{2+} *influx measurements*

For calcium level measurements using force sensors, cells were transfected with the GCAMP7S-paxillin-

mScarleti construct and seeded on fibronectin coated glass bottom coverslips overnight. Transfected cells

were imaged using TirfM microscrope using 488 and 561 laser lines. Specific band pass excitation/emission

filter cubes (GFP-HQ and mCherry-HQ from Nikon) were used to avoid fluorescence crosstalk between

channels. For comparison between different samples, all imaging parameters (laser power, tirf angle,

camera exposures) were kept constant. Only cells with comparable calcium sensor expression levels (judging

from fluorescence intensity from mScarlet channel) were picked for analysis to minimize potential bias due

to sensor concentration differences.

The local Ca^{2+} influx of the cells was measured following Ellefsen et. al(*26*). Briefly, the HFF cells were

loaded with 4 uM Ca^{2+} indicator Cal-520-AM for 2 hours. Then the cells were detached using enzyme free

dissociation media and resuspended in complete growth media. The cells were allowed to recover in

suspension for 20 min before seeding on glass bottom coverslips coated with fibronectin. After 2 hours the

buffer was switched to Ringer's balanced salt and proceeded for imaging on Nikon Tirf system with pulsed

laser lines to minimize photodamage of the sample(*72*). The calcium images of individuals cells were

acquired at 100Hz for 5000 frames. The deltaF/F images of the cells were generated using a custom matlab

script to reveal individual Ca^{2+} influx events. Such events were tracked using the Trackmate ImageJ

plugin(*73*).

Cell spreading assay

In the cell spreading assay, cells were transfected with respective plasmids 36-48 hours before experiments using Neon transfection system (Thermo Fisher) or JetOptimus chemical transfection reagent (PolyPlus) following manufacturer's recommendations. On the day of experiment, the cells were detached using an enzyme-free cell detaching buffer (Cell Dissociation Buffer, enzyme-free, PBS, Gibco), spun-down and resuspended in complete media (1X DMEM, 10% FBS). The suspended cells were allowed to recover by incubating in the cell culture hood for 20 min and then seeded onto glass bottom Petri dishes coated with ECM molecules of interest (10 ug/ml collagen I, 10 ug/ml fibronectin, 10 ug/ml VTN-N, and 10 ug/ml laminin for 1 hour). Imaging was done on a Nikon Tirf system with 405,488,561 and 640 laser lines. The cell spreading area and adhesion size were analyzed automatically using CellProfiler(*74*).

Immunostaining and antibodies

Cells of interest seeded on glass bottom Petri dishes were fixed with 4% paraformaldehyde (PFA) for 30 min at 37 degrees and washed with 1X PBS three times. The fixed cells were permeabilized/blocked by incubation with TBST (1X TBS, 0.1% Tween-20) containing 5% Bovine Serum Abumin (BSA) overnight at 4 degrees. Primary antibodies were diluted in TBST, 5% BSA solution according to manufactures' recommendations and added to the cells overnight at 4 degrees. The stained cells were washed with 1X PBS and stained with secondary antibody for 1 hour at room temperature.

Polyclonal Rabbit and monoclonal mouse anti-Piezo1 was from Novus Biologicals (NBP1-78446, NBP2-75617), both have been validated by Piezo1 knock down/knock off experiments(*15*, *75*). For all Piezo1 localization quantifications, the monoclonal NBP2-75617 antibody was used. Mouse anti-Paxillin antibody was from Merck Millipore (MAB 3060), Rabbit anti-paxillin antibody was from abcam(ab32115). Mouse anti-activated integrin $\beta 1$ antibody was from BD Biosciences (9EG7). Rat anti integrin $\alpha v\beta 3$ was from Millipore(LM609).

FRAP analysis

FRAP analyses were carried out on an ILAS2 TIRF image system similar to reported previously. Briefly,

HFF, HT1080, or MDA-MB-231 cells transiently co-transfected with Piezo1-GFP and Paxillin-mapple, or

Piezo1-mruby3 and mEmerald-integrin β3 were seeded on Fibronectin coated glass bottom dishes overnight.

2-5 square-shaped regions of interests (~3 um length) covering focal adhesions were selected for each cell

and bleached. The recovery of fluorescence in both fluorescent channels was observed for from 360 to 600

seconds after photobleaching. The recovery curves were normalized by dividing by average intensity of each

ROI before bleaching and the average and standard deviation of the recovery were calculated.

Electrophysiology of the Pieoz1 linker mutants

Piezo1 KO HEK293T cells were plated on 35 mm dishes and transfected with lipofectamine 3000 as per

the manufacturers instructions 48 h prior to patch-clamp analysis. The extracellular solution for cell-attached

patches contained high K^+ to zero the membrane potential; it consisted of 90 mM potassium aspartate, 50

mM KCl, 1 mM $MgCl_2$ and 10 mM HEPES (pH 7.2) adjusted with 5 M KOH. The pipette solution contained 140

mM NaCl, 3 mM KCl, 1 mM MgCl2, 1 mM CaCl2 with 10 mM HEPES (pH 7.2) adjusted with sodium hydroxide.

Negative pressure was applied to patch pipettes using a High-Speed Pressure Clamp-1 (ALA Scientific

Instruments, Farmingdale, NY, USA) and recorded in millimetres of mercury (mmHg) using a piezoelectric

pressure transducer (WPI, Sarasota, FL, USA). Borosilicate glass pipettes (Sigma-Aldrich) were pulled with a

vertical pipette puller (PP-83, Narashige, Tokyo, Japan) to produce electrodes with a resistance of ~2 MΩ.

The Piezo1 currents were amplified using an AxoPatch 200B amplifier (Axon Instruments, Union City, CA,

USA), and data were sampled at a rate of 10 kHz with 1 kHz filtration and analysed using pCLAMP10

software (Axon Instruments).

Acknoledgements

We would like to thank Professor Alexander Beshardsky and Dr Benjamin T. Goult for critical discussions and

the microscopy core and protein cloning expression core facility of Mechanobiology Institute for their

technical support.

Author Contributions:

M.Y, C.C and M.S designed research, M.Y, C. C, A.J, D.C, JV. L and A.H performed experiments and analyzed

the data. M.Y, C.C and G. TVN contributed to new reagents/methods. M.S and B.M supervised the research.

M.Y, C.C and M.S wrote the paper.

Conflict of Interest :

The authors declare no conflict of interest.

References

1. B. Geiger, J. P. Spatz, A. D. Bershadsky, Environmental sensing through focal adhesions. *Nat. Rev. Mol. Cell Biol.* **10**, 21–33 (2009).

2. V. Vogel, M. Sheetz, Local force and geometry sensing regulate cell functions. *Nat. Rev. Mol. Cell Biol.* **7**, 265–75 (2006).

3. R. K. Jain, J. D. Martin, T. Stylianopoulos, The Role of Mechanical Forces in Tumor Growth and Therapy. *Annu. Rev. Biomed. Eng.* **16**, 321–346 (2014).

4. J. Li, B. Hou, S. Tumova, K. Muraki, A. Bruns, M. J. Ludlow, A. Sedo, A. J. Hyman, L. McKeown, R. S. Young, N. Y. Yuldasheva, Y. Majeed, L. A. Wilson, B. Rode, M. A. Bailey, H. R. Kim, Z. Fu, D. A. L. L. Carter, J. Bilton, H. Imrie, P. Ajuh, T. N. Dear, R. M. Cubbon, M. T. Kearney, K. R. Prasad, P. C. Evans, J. F. X. X. Ainscough, D. J. Beech, Piezo1 integration of vascular architecture with physiological force. *Nature*. **515**, 279–282 (2014).

5. A. D. Bershadsky, N. Q. Balaban, B. Geiger, Adhesion-dependent cell mechanosensitivity. *Annu. Rev. Cell Dev. Biol.* **19**, 677–95 (2003).

6.   T. Panciera, L. Azzolin, M. Cordenonsi, S. Piccolo, Mechanobiology of YAP and TAZ in physiology and

disease. *Nat. Rev. Mol. Cell Biol.* (2017), doi:10.1038/nrm.2017.87.

7.   B. Trappmann, J. E. Gautrot, J. T. Connelly, D. G. T. Strange, Y. Li, M. L. Oyen, M. A. Cohen Stuart, H.

Boehm, B. Li, V. Vogel, J. P. Spatz, F. M. Watt, W. T. S. Huck, Extracellular-matrix tethering regulates

stem-cell fate. *Nat. Mater.* **11**, 642–649 (2012).

8.   H. Wolfenson, B. Yang, M. P. Sheetz, Steps in Mechanotransduction Pathways that Control Cell

Morphology. *Annu. Rev. Physiol.* **81**, 585–605 (2019).

9.   E. R. Horton, A. Byron, J. A. Askari, D. H. J. Ng, A. Millon-Frémillon, J. Robertson, E. J. Koper, N. R. Paul,

S. Warwood, D. Knight, J. D. Humphries, M. J. Humphries, Definition of a consensus integrin

adhesome and its dynamics during adhesion complex assembly and disassembly. *Nat. Cell Biol.* **17**,

1577–1587 (2015).

10.   A. del Rio, R. Perez-Jimenez, R. Liu, P. Roca-Cusachs, J. M. Fernandez, M. P. Sheetz, Stretching single

talin rod molecules activates vinculin binding. *Science (80-.).* **323**, 638–41 (2009).

11.   M. Yao, B. T. Goult, B. Klapholz, X. Hu, C. P. Toseland, Y. Guo, P. Cong, M. P. Sheetz, J. Yan, The

mechanical response of talin. *Nat. Commun.* **7**, 11966 (2016).

12.   B. Yang, Z. Z. Lieu, H. Wolfenson, F. M. Hameed, A. D. Bershadsky, M. P. Sheetz, *Nano Lett.*, in press,

doi:10.1021/acs.nanolett.6b02995.

13.   W. H. Goldmann, Role of vinculin in cellular mechanotransduction. *Cell Biol. Int.* **40**, 241–256 (2016).

14.   E. Bates, Ion Channels in Development and Cancer. *Annu. Rev. Cell Dev. Biol.* **31**, 231–247 (2015).

15.   S. A. Gudipaty, J. Lindblom, P. D. Loftus, M. J. Redd, K. Edes, C. F. Davey, V. Krishnegowda, J.

Rosenblatt, Mechanical stretch triggers rapid epithelial cell division through Piezo1. *Nature*. **543**,

118–121 (2017).

16. A. Tijore, M. Yao, Y. H. Wang, A. Hariharan, Y. Nematbakhsh, B. Lee Doss, C. T. Lim, M. Sheetz, B. Lee,

C. Teck, M. Sheetz, Biomaterials Selective killing of transformed cells by mechanical stretch.

Biomaterials. **275**, 120866 (2021).

17. A. Bhatt, I. Kaverina, C. Otey, A. Huttenlocher, Regulation of focal complex composition and

disassembly by the calcium-dependent protease calpain. *J. Cell Sci.* **115**, 3415–25 (2002).

18. S. J. Franco, M. a Rodgers, B. J. Perrin, J. Han, D. a Bennin, D. R. Critchley, A. Huttenlocher, Calpain-

mediated proteolysis of talin regulates adhesion dynamics. *Nat. Cell Biol.* **6**, 977–83 (2004).

19. S. J. Franco, Regulating cell migration: calpains make the cut. *J. Cell Sci.* **118**, 3829–3838 (2005).

20. M. Saxena, R. Changede, J. Hone, H. Wolfenson, M. P. Sheetz, Force-Induced Calpain Cleavage of Talin

Is Critical for Growth, Adhesion Development, and Rigidity Sensing. *Nano Lett.* **17**, 7242–7251 (2017).

21. T. Travers, H. Shao, B. a Joughin, D. a Lauffenburger, A. Wells, C. J. Camacho, Tandem

phosphorylation within an intrinsically disordered region regulates ACTN4 function. *Sci. Signal.* **8**,

ra51 (2015).

22. B. Yang, H. Wolfenson, V. Y. Chung, N. Nakazawa, S. Liu, J. Hu, R. Y. J. Huang, M. P. Sheetz, Stopping

transformed cancer cell growth by rigidity sensing. *Nat. Mater.* **19**, 239–250 (2020).

23. S. Yang, J. J. Zhang, X. Y. Huang, Orai1 and STIM1 Are Critical for Breast Tumor Cell Migration and

Metastasis. *Cancer Cell*. **15**, 124–134 (2009).

24. K. Clark, M. Langeslag, B. van Leeuwen, L. Ran, A. G. Ryazanov, C. G. Figdor, W. H. Moolenaar, K.

Jalink, F. N. van Leeuwen, TRPM7, a novel regulator of actomyosin contractility and cell adhesion.

EMBO J. **25**, 290–301 (2006).

25. C. Wei, X. Wang, M. Chen, K. Ouyang, L.-S. Song, H. Cheng, Calcium flickers steer cell migration.

Nature. **457**, 901–905 (2009).

26.   K. L. Ellefsen, J. R. Holt, A. C. Chang, J. L. Nourse, J. Arulmoli, A. H. Mekhdjian, H. Abuwarda, F.

580   Tombola, L. A. Flanagan, A. R. Dunn, I. Parker, M. M. Pathak, Myosin-II mediated traction forces

581   evoke localized Piezo1-dependent Ca2+ flickers. *Commun. Biol.* **2**, 298 (2019).

582   27.   X. Chen, S. Wanggou, A. Bodalia, M. Zhu, W. Dong, J. J. Fan, W. C. Yin, H. K. Min, M. Hu, D. Draghici,

583   W. Dou, F. Li, F. J. Coutinho, H. Whetstone, M. M. Kushida, P. B. Dirks, Y. Song, C. chung Hui, Y. Sun, L.

584   Y. Wang, X. Li, X. Huang, A Feedforward Mechanism Mediated by Mechanosensitive Ion Channel

585   PIEZO1 and Tissue Mechanics Promotes Glioma Aggression. *Neuron*. **100**, 799-815.e7 (2018).

586   28.   B. J. McHugh, R. Buttery, Y. Lad, S. Banks, C. Haslett, T. Sethi, Integrin activation by Fam38A uses a

587   novel mechanism of R-Ras targeting to the endoplasmic reticulum. *J. Cell Sci.* **123**, 51–61 (2010).

588   29.   B. Coste, J. Mathur, M. Schmidt, T. J. Earley, S. Ranade, M. J. Petrus, A. E. Dubin, A. Patapoutian,

589   Piezo1 and Piezo2 Are Essential Components of Distinct Mechanically Activated Cation Channels.

590   *Science (80-.).* **330**, 55–60 (2010).

591   30.   M. M. Pathak, J. L. Nourse, T. Tran, J. Hwe, J. Arulmoli, D. T. T. Le, E. Bernardis, L. A. Flanagan, F.

592   Tombola, Stretch-activated ion channel Piezo1 directs lineage choice in human neural stem cells.

593   *Proc. Natl. Acad. Sci.* **111**, 16148–16153 (2014).

594   31.   C. Li, S. Rezania, S. Kammerer, A. Sokolowski, T. Devaney, A. Gorischek, S. Jahn, H. Hackl, K.

595   Groschner, C. Windpassinger, E. Malle, T. Bauernhofer, W. Schreibmayer, Piezo1 forms

596   mechanosensitive ion channels in the human MCF-7 breast cancer cell line. *Sci. Rep.* **5**, 1–9 (2015).

597   32.   A. G. Solis, P. Bielecki, H. R. Steach, L. Sharma, C. C. D. Harman, S. Yun, M. R. de Zoete, J. N. Warnock,

598   S. D. F. To, A. G. York, M. Mack, M. A. Schwartz, C. S. Dela Cruz, N. W. Palm, R. Jackson, richard A.

599   Flavell, Mechanosensation of cyclical force by PIEZO1 is essential for innate immunity. *Nature*. **573**,

600   69–74 (2019).

33.   I. Borbiro, D. Badheka, T. Rohacs, Activation of TRPV1 channels inhibits mechanosensitive piezo

channel activity by depleting membrane phosphoinositides. *Sci. Signal.* **8** (2015),

doi:10.1126/scisignal.2005667.

34.   J. L. Nourse, M. M. Pathak, How cells channel their stress: Interplay between Piezo1 and the

cytoskeleton. *Semin. Cell Dev. Biol.* **71**, 3–12 (2017).

35.   N. Bavi, J. Richardson, C. Heu, B. Martinac, K. Poole, PIEZO1-Mediated Currents Are Modulated by

Substrate Mechanics. *ACS Nano*. **13**, 13545–13559 (2019).

36.   B. M. Gaub, D. J. Müller, Mechanical Stimulation of Piezo1 Receptors Depends on Extracellular Matrix

Proteins and Directionality of Force. *Nano Lett.* **17**, 2064–2072 (2017).

37.   J. Albarrán-Juárez, A. Iring, S. Wang, S. Joseph, M. Grimm, B. Strilic, N. Wettschureck, T. F. Althoff, S.

Offermanns, Piezo1 and G q /G 11 promote endothelial inflammation depending on flow pattern and

integrin activation. *J. Exp. Med.* **215**, 2655–2672 (2018).

38.   C. D. Cox, C. Bae, L. Ziegler, S. Hartley, V. Nikolova-Krstevski, P. R. Rohde, C. A. Ng, F. Sachs, P. A.

Gottlieb, B. Martinac, Removal of the mechanoprotective influence of the cytoskeleton reveals

PIEZO1 is gated by bilayer tension. *Nat. Commun.* **7**, 1–13 (2016).

39.   H. Park, C. Valencia-Gallardo, A. Sharff, G. Tran Van Nhieu, T. Izard, Novel vinculin binding site of the

IpaA invasin of Shigella. *J. Biol. Chem.* **286**, 23214–21 (2011).

40.   C. Valencia-Gallardo, D.-I. Aguilar-Salvador, H. Khakzad, C. Bou-Nader, C. Malosse, D. B. Lima, C. Bello,

B. Cocom-Chan, N. Quenech'Du, B. Mazhar, D. Javelaud, J. Fattaccioli, A. Mauviel, M. Fontecave, A.

Asnacios, J. Chamot-Rooke, L. Malmström, G. Tran Van Nhieu, Vinculin targeting by Shigella IpaA

promotes stable cell adhesion independent of mechanotransduction. *bioRxiv* (2019),

doi:10.1101/773283.

41.   A. N. Tullio, D. Accili, V. J. Ferrans, Z. X. Yu, K. Takeda, A. Grinberg, H. Westphal, Y. A. Preston, R. S.

624   Adelstein, Nonmuscle myosin II-B is required for normal development of the mouse heart. *Proc. Natl.*

625   *Acad. Sci. U. S. A.* **94**, 12407–12412 (1997).

626   42.   A. Elosegui-Artola, E. Bazellières, M. D. Allen, I. Andreu, R. Oria, R. Sunyer, J. J. Gomm, J. F. Marshall, J.

627   L. Jones, X. Trepat, P. Roca-Cusachs, Rigidity sensing and adaptation through regulation of integrin

628   types. *Nat. Mater.* **13**, 631–7 (2014).

629   43.   H. Wolfenson, G. Meacci, S. Liu, M. R. Stachowiak, T. Iskratsch, S. Ghassemi, P. Roca-Cusachs, B.

630   O'Shaughnessy, J. Hone, M. P. Sheetz, B. O'Shaughnessy, J. Hone, M. P. Sheetz, Tropomyosin controls

631   sarcomere-like contractions for rigidity sensing and suppressing growth on soft matrices. *Nat. Cell*

632   *Biol.* **18**, 33–42 (2015).

633   44.   J. Robertson, G. Jacquemet, A. Byron, M. C. Jones, S. Warwood, J. N. Selley, D. Knight, J. D.

634   Humphries, M. J. Humphries, Defining the phospho-adhesome through the phosphoproteomic

635   analysis of integrin signalling. *Nat. Commun.* **6**, 6265 (2015).

636   45.   A. Byron, J. A. Askari, J. D. Humphries, G. Jacquemet, E. J. Koper, S. Warwood, C. K. Choi, M. J. Stroud,

637   C. S. Chen, D. Knight, M. J. Humphries, A proteomic approach reveals integrin activation state-

638   dependent control of microtubule cortical targeting. *Nat. Commun.* **6**, 6135 (2015).

639   46.   G. Giannone, P. Rondé, M. Gaire, J. Beaudouin, J. Haiech, J. Ellenberg, K. Takeda, Calcium rises locally

640   trigger focal adhesion disassembly and enhance residency of focal adhesion kinase at focal adhesions.

641   *J. Biol. Chem.* **279**, 28715–28723 (2004).

642   47.   Y. R. Guo, R. MacKinnon, Structure-based membrane dome mechanism for piezo mechanosensitivity.

643   *Elife*. **6**, 1–19 (2017).

48.   Q. Zhao, H. Zhou, S. Chi, Y. Wang, J. Wang, J. Geng, K. Wu, W. Liu, T. Zhang, M. Q. Dong, J. Wang, X. Li,

B. Xiao, Structure and mechanogating mechanism of the Piezo1 channel. *Nature*. **554**, 487–492

(2018).

49.   K. Saotome, S. E. Murthy, J. M. Kefauver, T. Whitwam, A. Patapoutian, A. B. Ward, Structure of the

mechanically activated ion channel Piezo1. *Nature*. **554**, 481–486 (2018).

50.   J. Geng, W. Liu, H. Zhou, T. Zhang, L. Wang, M. Zhang, Y. Li, B. Shen, X. Li, B. Xiao, A Plug-and-Latch

Mechanism for Gating the Mechanosensitive Piezo Channel. *Neuron*. **106**, 438-451.e6 (2020).

51.   J. Wu, A. H. Lewis, J. Grandl, Touch, Tension, and Transduction – The Function and Regulation of Piezo

Ion Channels. *Trends Biochem. Sci.* **42**, 57–71 (2017).

52.   S. E. Murthy, A. E. Dubin, A. Patapoutian, Piezos thrive under pressure: Mechanically activated ion

channels in health and disease. *Nat. Rev. Mol. Cell Biol.* **18**, 771–783 (2017).

53.   Y.-C. Lin, Y. R. Guo, A. Miyagi, J. Levring, R. MacKinnon, S. Scheuring, Force-induced conformational

changes in PIEZO1. *Nature* (2019), doi:10.1038/s41586-019-1499-2.

54.   J. Dai, M. P. Sheetz, Membrane tether formation from blebbing cells. *Biophys. J.* **77**, 3363–3370

(1999).

55.   R. Syeda, M. N. Florendo, C. D. Cox, J. M. Kefauver, J. S. Santos, B. Martinac, A. Patapoutian, Piezo1

Channels Are Inherently Mechanosensitive. *Cell Rep.* **17**, 1739–1746 (2016).

56.   G. Giannone, P. Rondé, M. Gaire, J. Haiech, K. Takeda, Calcium oscillations trigger focal adhesion

disassembly in human U87 astrocytoma cells. *J. Biol. Chem.* **277**, 26364–26371 (2002).

57.   S. Sen, S. Kumar, Combining mechanical and optical approaches to dissect cellular mechanobiology. *J.*

Biomech. **43**, 45–54 (2010).

58.   C. D. Cox, N. Bavi, B. Martinac, Biophysical Principles of Ion-Channel-Mediated Mechanosensory

666   Transduction. *Cell Rep.* **29**, 1–12 (2019).

667   59.   Z. Shi, Z. T. Graber, T. Baumgart, H. A. Stone, A. E. Cohen, Cell Membranes Resist Flow. *Cell*. **175**,

668   1769-1779.e13 (2018).

669   60.   A. Schwartz[1], C. Hall[1], L. Barney[1], C. Babbitt[2], S. Peyton[1], Mechanosensing of Integrin α 6 and EGFR

670   Converges at Calpain 2 (2017), doi:10.1101/164525.

671   61.   C. L. Yankaskas, K. Bera, K. Stoletov, S. A. Serra, J. Carrillo-garcia, S. Tuntithavornwat, P. Mistriotis, J.

672   D. Lewis, M. A. Valverde, K. Konstantopoulos, The fluid shear stress sensor TRPM7 regulates tumor

673   cell intravasation (2021).

674   62.   A. K. Efremov, M. Yao, Y. Sun, Y. H. Tee, M. P. Sheetz, A. D. Bershadsky, B. Martinac, J. Yan,

675   Application of piconewton forces to individual filopodia reveals mechanosensory role of L-type Ca2+

676   channels. *Biomaterials*. **284**, 121477 (2022).

677   63.   C. L. Gilchrist, H. A. Leddy, L. Kaye, N. D. Case, K. E. Rothenberg, D. Little, W. Liedtke, B. D. Hoffman, F.

678   Guilak, TRPV4-mediated calcium signaling in mesenchymal stem cells regulates aligned collagen

679   matrix formation and vinculin tension. *Proc. Natl. Acad. Sci. U. S. A.* **116**, 1992–1997 (2019).

680   64.   M. Sheetz, *Annu. Rev. Cell Dev. Biol.*, in press, doi:10.1146/annurev-cellbio-100818-125227.

681   65.   S. W. Moore, P. Roca-Cusachs, M. P. Sheetz, Stretchy proteins on stretchy substrates: the important

682   elements of integrin-mediated rigidity sensing. *Dev. Cell*. **19**, 194–206 (2010).

683   66.   C. G. Galbraith, K. M. Yamada, M. P. Sheetz, The relationship between force and focal complex

684   development. *J. Cell Biol.* **159**, 695–705 (2002).

67.   P. Roca-Cusachs, A. del Rio, E. Puklin-Faucher, N. C. Gauthier, N. Biais, M. P. Sheetz, Integrin-

dependent force transmission to the extracellular matrix by -actinin triggers adhesion maturation.

Proc. Natl. Acad. Sci. **110**, E1361–E1370 (2013).

68.   X. D. Ren, W. B. Kiosses, M. A. Schwartz, Regulation of the small GTP-binding protein Rho by cell

adhesion and the cytoskeleton. *EMBO J.* **18**, 578–585 (1999).

69.   G. Gateva, E. Kremneva, T. Reindl, T. Kotila, K. Kogan, L. Gressin, P. W. Gunning, D. J. Manstein, A.

Michelot, P. Lappalainen, Tropomyosin Isoforms Specify Functionally Distinct Actin Filament

Populations In Vitro. *Curr. Biol.* **27**, 705–713 (2017).

70.   C. Valencia-Gallardo, C. Bou-Nader, D. Aguilar, N. Carayol, N. Quenech'Du, L. Pecqueur, H. Park, M.

Fontecave, T. Izard, G. T. Van Nhieu, Cell adhesion promoted by a unique Shigella IpaA vinculin- and

talin-binding site. *bioRxiv*, 329136 (2018).

71.   H. Dana, Y. Sun, B. Mohar, B. K. Hulse, A. M. Kerlin, J. P. Hasseman, G. Tsegaye, A. Tsang, A. Wong, R.

Patel, J. J. Macklin, Y. Chen, A. Konnerth, V. Jayaraman, L. L. Looger, E. R. Schreiter, K. Svoboda, D. S.

Kim, High-performance calcium sensors for imaging activity in neuronal populations and

microcompartments. *Nat. Methods*. **16**, 649–657 (2019).

72.   G. Donnert, C. Eggeling, S. W. Hell, Major signal increase in fluorescence microscopy through dark-

state relaxation. *Nat. Methods*. **4**, 81–86 (2007).

73.   J. Y. Tinevez, N. Perry, J. Schindelin, G. M. Hoopes, G. D. Reynolds, E. Laplantine, S. Y. Bednarek, S. L.

Shorte, K. W. Eliceiri, TrackMate: An open and extensible platform for single-particle tracking.

Methods. **115**, 80–90 (2017).

74.   C. McQuin, A. Goodman, V. Chernyshev, L. Kamentsky, B. A. Cimini, K. W. Karhohs, M. Doan, L. Ding,

S. M. Rafelski, D. Thirstrup, W. Wiegraebe, S. Singh, T. Becker, J. C. Caicedo, A. E. Carpenter,

CellProfiler 3.0: Next-generation image processing for biology. *PLoS Biol.* **16**, e2005970 (2018).

75.   J. V. Li, C.-A. Ng, D. Cheng, Z. Zhou, M. Yao, Y. Guo, Z.-Y. Yu, Y. Ramaswamy, L. A. Ju, P. W. Kuchel, M.

709   P. Feneley, D. Fatkin, C. D. Cox, Modified N-linked glycosylation status predicts trafficking defective

710   human Piezo1 channel mutations. *Commun. Biol.* **4**, 1038 (2021).

Figures:

Figure 1

Figure 1. Piezo1 is stably recruited to maturing adhesions on fibronectin surfaces. (A)Time lapse of cell

spreading of HFF cells seeded on fibronectin coated glass surface. The cells are transiently expressing Piezo1-

mruby3 and paxillin-BFP. Scale Bars denote 10 um. (B) Localization of Piezo1 spreading on fibronectin

surface during polarization and in stable condition. Scale Bars denote 10 um in the main figure and 5um in

the zoom. (C) Adhesion enrichment of Piezo1 at 1h, 2h and 4h after cell seeding. The adhesion enrichment

was defined as the average intesity ratio of Piezo1 in the adhesions and background. (D) Cell spreading area

and aspect ratio 1h, 2h and 4h after cell seeding. The significance were calculated using two sample t-test.

(E) Left panel: Live TIRF images of Human Foreskin Fibroblast (HFF) cells transiently transfected with Piezo1-

GFP and paxillin-mapple on fibronectin coated surfaces. The adhesion region marked by white square was

bleached and snapshots of fluorescent recovery for Piezo1 and talin are shown in the zoom in figure on

bottom. Scale Bars denote 5 um. Right panel: The FRAP recovery curve of the bleached region (mean $\pm$ std).

(F) The fitted mobile fraction and recovery rates from the FRAP measurements. More than five independent

cells and twenty adhesions were picked for each condition. The bracket after the fitting value denote 95%

confidence interval of the fitting parameters. In the Frap experiment, Piezo1 is either co-expressed with

Integrin β3 or Paxillin.

Figure 2

Figure 2. Piezo1 localization in HFF cells is controlled by contractility. (A) Upper panel: Distribution of

Piezo1-mRuby3 in HFF cells 15 min after treatment with 30 µM Y-compound and 30 min after washout.

Experiments were repeated on > 5 cells with similar results. Lower panel: representative Piezo1 localization

and Piezo1 adhesion enrichment quantification in control HFF cells and HFF cells expressing constitutive-

active RhoA V14 mutant. Scale bars denote 10 µm.   (B) Time-lapse image of HFF cells transfected with

Piezo1-mruby3, integrin β3-Emerald and IPA-GFP with the addition of 30 µM Y-27632. Experiments were

repeated on > 3 cells with similar results. (C) Fluorescence image of Piezo1-GFP in COS7 cells with/without

overexpressing myosin IIA-mCherry.

Figure 3

Figure 3. Piezo1 recruitment of adhesions precede integrin β3 disassembly. (A-C) Timelapse of HFF cells

co-transfected with Piezo1-mruby3 and mEmerald-Integrin β3 spreading on fibronectin surfaces.A: A

spreading cell one hour after seeding. B: A polarized cell after overnight seeding. (C) Timelaspe of HFF cells

transfected with Piezo1-mruby3 and mEmerald-Integrin β1 spreading on fibronectin surfaces overnight. The

normalized intensity profile of adhesions highlighted by the red line were shown below the figure. (D)

Kymograph of Piezo1 and Integrin β3 dynamics of the focal adhesion marked by red line in panel B. (E)

Kymograph of Piezo1 and Integrin β1 dynamics of the focal adhesion marked in red in panel C.

Figure 4

Figure 4. Piezo1 localizes to the adhesions of non-transformed cells but is diffusive in transformed

cells. (A) Immunostaining of Piezo1 localization in paired normal and transformed cell lines with similar

orgin. (B) Quantification of Piezo1 adhesion enrichment for cells in Figure (A). p-values were calculated using

two-sample t-test. (C) FRAP recovery curve of Piezo1 in HFF cells and transformed HT1080 and MDA-MB-231

cells. The error bar denotes 95% confidence interval of the mean. (D) Quantification of adhesion

entrichment of Piezo1 across various normal and transformed cancer cell lines. Error bars denote standard

error of the mean. There were statistically significant differences (p-value < 0.001) between HFF/HUMEC and

all the cancer cells.

Figure 5

Figure 5. Piezo1 localization affects adhesion maturation and correlates with transformation states of

cells. (A) TIRF image of fixed cell staining of non-transformed HFF cell line and transformed fibrosarcoma

HT1080 with/without Piezo1 knock down after overnight spreading on fibronectin surfaces. Scale Bars

denote 20 um. (B) Box plots of the cell spreading areas, cell aspect ratio and mean adhesion sizes of the cells

in figure A. Box and error bar denotes the s.e.m and s.t.d of the measurements. The significance value were

calculated using two sample t-tests. (C) Western blots of Piezo1 level in control siRNA and piezo1 knock-

down cells.

Figure 6

Figure 6. Normal and transformed cells have distinct adhesion calcium signals . (A) Domain Illustration of

the novel adhesion targeting calcium sensor based on jGCaMP7s from Janelia. (B) Left: TIRF image montages

of representative HFF cells (with/without Piezo1 knock down), HT1080 and MDA-MB-231 cells transfected

with paxillin calcium sensor. The cells were seed on fibronectin surfaces overnight before imaging.

Normalized ratio represents the ratio image between jGCaMP7s and mScarlet-I, which is an measure of local

Ca^{2+} concentrations. Right: calcium level fluctuations over time in the tracked adhesions of the cell presented

in the left panel, each line in the image denotes a single adhesion. The interval of measurement is 30 s. (C)

Quantification of average calcium ratio in tracked adhesions over 60 min for different cell types.

Figure 7

Figure 7, The linker domain of Piezo1 is linked to adhesion recruitment and alters cell morphology in HFF

cells. (A) Domain map of Piezo1 and the long-linker construct. The residue number is based on mouse Piezo1

structures . (B) Immunostaining of HFF and HT1080 cells transfected with mNeongreen control vector or

mNeongreen labeled Piezo1 linker construct. The cells were seeded on fibronectin surface overnight and

stained with paxillin and Piezo1. (C) Box plot of cell spreading area and cell aspect ratio of positively

transfected cells. Box and error bar present s.e.m and s.t.d respectively. Statistical significance were

calculated using two sample t-test.

Figure 8

Figure 8, Model of Piezo1's regulation in normal and transformed cells. (Top Panel) In normal cells, Piezo1

is recruited to maturing focal adhesions in a force dependent manner, where it regulates both the growth

and turnover of focal adhesions. (Bottom Panel) In transformed cells Piezo1's localization is diffusive and it

no-longer regulate adhesion morphology, which might contribute to the mis-regulated calcium signaling

observed in cancer cells including stretch dependent apoptosis(*16*).

  Supplementary Figures

Figure S1

Figure S1, Immunostaining of Piezo1 localization of HFF cells. (A) HFF cells were seeded on fibronectin

surfaces overnight, fixed with paraformaldehyde and stained for Piezo1 with two validated Piezo1 antibody

and paxillin. (B) Sequence information of Piezo1 siRNA used in the study (SASI_HS01_00208585 from Merck

Sigma-Aldrich).

Figure S2

A

B

Figure S2. (A) HFF cells transiently transfected with Piezo1-mruby3 were treated with 50 μM nitro-
Blebbistatin at time 0. After 30 min incubation, the localization of Piezo1 at focal adhesions were lost. Scale
bars denote 10 μm. (B) HFF cells transiently transfected with mEmerald-integrin β3 and Piezo1-mruby3 were
pre-treated with 2.5 μM PTP-PEST inhibitor that inhibits adhesion turnover by phosphatases before addition
of 20 μM Y-27632. In this condition, Piezo1 dissociated from adhesions before integrin β3 upon tension release.

Figure S3

Figure S3. Dynamics of Piezo1 localization in relation to integrins. (A) Adhesion time lapse HFF cells transiently transfected with mEmerald-integrin β3 and Piezo1-mruby3. The right panel showed normalized intensity profiles of the adhesions marked by red line over time. (B) Adhesion time lapse of HFF cells transfected with BFP-integrin β3, mEmerald-integrin β1 and Piezo1-mruby3. The right panel showed normalized intensity profiles of the adhesions marked by red line over time. The normalizations were done by dividing the intensity profiles by the maximum intensity value observed for each protein in the whole time series. Scale Bars denote 10 µm.

Figure S4

A

B

Figure S4 (A) TIRF live images of Piezo1-GFP localization in more normal cell lines tested. (B) Western blot images of Piezo1 expression in various cancer cell lines compared against HFF cells.

Figure S5

Figure S5. Piezo1 localization in cells with altered transformation state. (A) Representative HFF cells with

Control siRNA and TPM 2.1 siRNA knock down were transfected with Piezo1-mruby3, Integrin β3-BFP and

talin-GFP. The cells were seeded on fibronectin surfaces overnight and imaged by TIRFM. Right Panel:

quantification of adhesion enrichment with Ctrlsi and TPM 2.1 si (C) Transient expression of TPM 2.1 -YFP in

MDA-MB-231 cells led to the recruitment of Piezo1 to the end of stress fibres decorated by TPM 2.1 (TOP

panel). The images present representative cells out of > 20 imaged. Scale Bars denote 10 um.

Figure S6

A

B

Figure S6 **HFF and MDA-MB-231 cells showed distinct local calcium entry pattern**. (A) Locations of local calcium entry events (marked by red dots) in HFF cells and MDA-MB-231 cells measured at 100 frames/s for 50s, overlaid with focal adhesions marked by paxillin. Significantly more local calcium signal is observed in HFF cells. (B). Box plot of the rate of local calcium events per cell for HFF and MDA-MB-231 cells. Each point denotes an independent cell.

Figure S7

Figure S7 Linker domain mutants of Piezo1. Representative images of FAK-/- MEF cells transfected with
paxillin-mapple and wild type/linker deletion mutant of Piezo1. Similar behavior were observed in 3 biological
repeats with > 10 cells for each condition.

Figure S8

Figure S8. Determination of stretch activation of Piezo1 deletion mutants. (A) Representative patch-
clamp traces from human Piezo1 c-terminal EGFP fusion (hP1-cGFP) in cell-attached patches at -60 mV holding
potential expressed in Piezo1 KO HEK293T cells compared to four deletion mutants. Current traces are shown
in black and square wave negative pressure pulses increasing from 0 mmHg in increments of -10 mmHg are
shown in red with the peak negative pressure documented for clarity. (B) Box and whiskers plots indicating
the peak current elicited per patch from negative pressure application with all mutant Piezo1 channels. *
represents statistical significance from hP1-cGFP using one-way ANOVA with Dunnett's post hoc test.

864