Mutagenic effectiveness and efficiency of gamma rays and sodium azide in

M$_2$ generation of Cowpea [*Vigna unguiculata* (L.) Walp.]

Aamir Raina[1,2]*, Samiullah Khan[1]

[1]Mutation Breeding Laboratory, Department of Botany, Aligarh Muslim University, Aligarh,

India

[2]Botany Section, Women's College, Aligarh Muslim University, Aligarh, India

Corresponding author: *aamir854@gmail.com

These authors contributed equally to this work

Abstract

Legumes play a pivotal role in combating the chronic hunger and malnutrition in the developing nations and are also ideal crops to achieve global food and nutrition security. In the era of climate change, erratic rainfalls, depleting arable land and water resource, feeding the rapidly growing population is a challenging task. Among breeding programs for crop improvement, induced mutagenesis has proven to be the viable, effective, robust and coherent tool to facilitate the process of creating varieties with improved yield. Like most other pulses, cowpea is a self pollinated, nutritious, versatile and widely adapted grain legume, but harbour a little accessible genetic variability. Lack of sufficient genetic variability and small size of flowers, traditional plant breeding methods are not enough to meet the demands of improvement of cowpea. Hence, induced mutagenesis was employed to induce significant genetic variability across a range of agro-economical traits in two cowpea varieties Gomati VU-89 and Pusa-578 from M_1 to M_4 generations. The success of induced mutagenesis largely depends on the selection of appropriate mutagen, its dose, effectiveness and efficiency. Hence present study was conduct to assess the effectiveness and efficiency of single and combined doses of sodium azide and gamma rays to set an appropriate protocol for induced mutagenesis experimentation in cowpea.

1. Introduction

Pulses are the edible grains of members of the Leguminosae family, and have been defined as "Leguminosae crops harvested wholly for their grain, including beans, lentils and cowpeas" by the Food and Agricultural Organization of the United Nations [1]. Pulses are considered as vital constituents in the diets of people and hence are most suitable crops to achieve the global food and nutrition security. The production of pulses in India is much

lower as compared to global pulse production and it may be due to the lack of high yielding varieties of pulses (Table 1). Additional there is a progressive decline in the per capita pulses consumption from 1951 (60.7 g/day) to 2017 (52.9 g/day) [2]. The low yielding potential of crops is one of the primary obstacles in achieving the desired goals of production. The common asynchronous and longer maturity period, more flower drop and less seed set, natural and anthropogenic pressures are other obstacles in achieving the higher production of pulses.

Table 1. Average yield of pulses in India and world compared during 2007-2016.

YEAR	PULSES YEILD (Q/ha)	
	INDIA	WORLD
2007	5.970	8.154
2008	6.042	8.607
2009	6.706	9.285
2010	6.496	9.043
2011	6.163	8.769
2012	6.460	9.381
2013	7.162	9.745
2014	7.056	9.436
2015	6.465	9.505
2016	6.644	9.929

Source: FAOSTAT | © FAO Statistics Division, 2017.

Among the ten primary pulse crops as recognised by FAO, Cowpea is an important member on the basis of its use, nutritive value and other desired qualities. It belongs to fabaceae family and is one among the most ancient pulse crops used by man [3]. It is nutritious, versatile and widely adapted grain legume to many areas of the humid tropics and temperate zones. The dry and immature pods serve as a rich source of dietary protein for millions of people in developing countries including India and are also used as fodder for livestock [4]. Cowpea has substantial tolerance to high temperatures and severe drought stress and could withstand both acidic and alkaline soils but is intolerant of frost [5]. It is suitable for cultivation in subtropical areas of the world and usually grown alone but can be intercropped with a range of other crops including sorghum, millet, maize, cassava and cotton [6]. Cowpea is considered as a vital constituent of traditional cropping system because of its capability to reinstate soil fertility for succeeding cereal crops grown in rotation with it and enhances soil fertility mostly in smallholder farming systems where very less or no fertilizer is used. Cowpea growing countries in the Asian region are India, Sri Lanka, Bangladesh, Myanmar, China, Korea, Thailand, Indonesia, Nepal, Pakistan, Philippines and Malaysia. According to FAO cowpea is cultivated annually on 12.3 million ha and the estimated production worldwide is about 7.69 million tonnes [7]. Globally Africa contributes about 95% of total world cowpea production (Fig 1). Nigeria is the top most producer of cowpea with an estimated 45% of the world cowpea production (Table 2). The objective of this study was to study effect of mutagens on physiological parameters and their mutagenic effectiveness and efficiency.

Fig 1. Production share of Cowpeas by region (Average 2005 – 2016).

Table 2: World's top 10 producers of Cowpea (Average 2016).

RANK	COUNTRY	PRODUCTION (mt)
1	Nigeria	3.02
2	Niger	1.98
3	Burkina Faso	0.60
4	Cameroon	0.19
5	United Republic of Tanzania	0.18
6	Sudan	0.16
7	Mali	0.14
8	Kenya	0.14
9	Myanmar	0.11
10	Sri Lanka	0.01

Source: FAOSTAT | © FAO Statistics Division, 2017.

2. Materials and methods

2.1. Experimental materials and Seed Irradiation

In the present study, M_0 seeds of two cowpea varieties viz., Gomati VU-89 and Pusa-578 were procured from NBPGR, New Delhi (S Table 1). Initially, LD50 was calculated, the doses of gamma rays and sodium azide were optimised by calculating values based on germination rate and survival percentage [6]. A set of 300 dry and healthy M_0 seeds were pre-soaked in the double distilled water for 9 hours and then treated with 0.01%, 0.02%, 0.03% and 0.04% concentrations of sodium azide (SA) solution for 6 hrs at 25 C at Mutation Breeding Laboratory, Department of Botany, Aligarh Muslim University, Aligarh, India. Another set of 600 M_0 seeds per dose were irradiated with 100, 200, 300 and 400 Gy at a dose rate of 11.58 Gy/min of gamma radiation using a Cobalt 60 (^{60}Co) gamma source under standard conditions at NBRI, Lucknow, India. Each set of irradiated 300 M_0 seeds were also treated with respective SA concentration to produce M_1 seeds. Each set of seeds were

98 thoroughly washed under tap water twice to get rid of any excess mutagen adhered to the

99 seed surface before sowing in the field.

2.2. Experimental site and crop cultivation

The experimentation were performed at Agriculture field, Aligarh Muslim University,

Aligarh (27° 29′ to $28^\circ 11'$ N' 77° 29′ to 78° 38′ E). The net field size was 23.5×40 meter,

with 13 blocks (each 3×1.8 m) with a 0.6 m (row to row) and 0.3 m (seed to seed) distance

(S Fig 1). Cowpea M_1 seeds were sown in 10 replications of 30 seeds in a randomized

complete block design (RCBD) to raise the M_1 generation during mid-april 2014-October

2014 along with respective controls. Field was irrigated at regular intervals after seed sowing

and; weeding was carried out at seedling and flowering stage. All the M_1 plants were

harvested separately at pod stage to collect M_2 seeds. During the mid-april 2015-October

2015, 10 healthy M_2 seeds collected from each M_1 plant were sown treatment wise for raising

M_2 generation. In M_3 generation seven and four high yielding stable mutant lines of Gomati

VU-89 and Pusa-578, respectively were identified and advanced to M_4 generation during the

mid-april 2017-October 2017 [8]. All the recommended cultivation practices were taken care

of; such as 15 - 20 N, 50 - 60 P_2O_5 and 50 - 60 K_2O kg/ha of fertilizers were added during

seedbed preparation.

S1 Fig. Field plots layout of M_1 generations in a randomized complete block design (RCBD)

for each cultivar (Source: Raina et al., 2020. http://creativecommons.org/licenses/by/4.0/).

2.3. Field Analysis

2.3.1. Seed germination

Seed germination was recorded after 15 days of seed sowing and the number of seeds

germinated in each treated and untreated population was counted. After recording

germination counts, the percentage of seed germination was calculated on the basis of total

number of seeds sown in the field using the following formulla.

$$\text{Germination (\%)} = \frac{\text{No. of seeds germinated}}{\text{No. of seeds sown}} \times 100$$

2.3.2. Chlorophyll mutations

Chlorophyll mutants were observed in M_2 generation after 10-20 days of seed sowing

in the field. The identification and categorisation proposed by Gustafsson [9] and Khan [10]

was followed using the following formula. The influence of combination doses of gamma

rays and sodium azide on the frequency of chlorophyll mutations was analyzed by calculating

the Coefficient of interaction (k) proposed by Sharma [11].

$$\text{Mutation frequency (\%)} = \frac{\text{Number of mutant seedlings}}{\text{Total number of } M_2 \text{ seedlings}} \times 100$$

$$\text{Coefficient of interaction (k)} = \frac{(a + b)}{(a) + (b)}$$

Where,

(a + b) = The frequency of mutation due to combined mutagen doses.

(a) + (b) = The frequency of mutation due to individual mutagen doses

k = Hypothetical interaction coefficient

If k = 1 additive interaction, and if it is less or more than 1 that indicates synergistic

interaction.

2.4. Pollen fertility

The determination of pollen fertility was necessary to assess the pollen sterility based mutagenic effectiveness and efficiency. At the time of flowering, pollens were collected from 30 random plants per treatment and control in M_1 generation. Pollens were then placed on slides and stained with 1% acetocarmine solution. The uniformly stained pollens were counted as fertile, while as unstained pollens were counted as sterile.

2.5. Mutagenic effectiveness and efficiency

The mutation frequency induced by unit mutagen dose refers to mutagenic effectiveness whereas percentage of mutations in relation to biological damage refers to mutagenic efficiency. Mutagenic effectiveness and efficiency was calculated by following formulae given by Konzak *et al*. [12].

$$\text{Mutagenic effectiveness (Gamma rays)} = \frac{\text{Rate of mutation (Mp)}}{\text{Dose in Gray (Gy)}}$$

$$\text{Mutagenic effectiveness (SA)} = \frac{\text{Rate of mutation (Mp)}}{\text{Concentration} \times \text{duration of treatment}}$$

$$\text{Mutagenic effectiveness(Gamma rays + SA)} = \frac{\text{Rate of mutation (Mp)}}{\substack{\text{Dose of physical mutagen (Gy)} \times \\ \text{concentration of chemical mutagen} \times \\ \text{duration of treatment}}}$$

$$\text{Mutagenic efficiency} = \frac{\text{Rate of mutation (Mp)}}{\text{* Biological damage in } M_1 \text{ generation}}$$

159 *Biological damage: For measuring the biological damage, three different criteria were used;

160 Injury, Sterility and Meiotic abnormalities refers to %age decrease in seedling height (Mp/I),

161 pollen fertility (Mp/S) and meiotic abnormalities (Mp/Me), respectively.

2.6. Morphological mutations

Data on twenty three qualitative and quantitative traits were recorded in M_2 generation of

both the varieties. These traits include tall, dwarf, semi dwarf, compact/bushy, prostrate, semi

dwarf spreading, one sided branching, axillary branching, broad leaf, narrow leaf, altered leaf

architecture, elongated rachis, multiple flower, flower color, open flower, non flowering, late

flowering, early maturity, small/narrow pods, bold seeded pods, coat color, coat pattern,

shape and surface traits which were examined and modified on the basis of available cowpea

descriptors from the International Union for the Protection of New Varieties of Plants [13]

and International Board for Plant Genetic Resources [14]. The characteristics of each mutant

is presented in results section. In the field, morphological mutant features for plant height

variations, growth habit, leaf, flower, pod and seed were observed, recorded and analysed.

The frequency (F) of the morphological mutant was evaluated using the following formula.

174
$$F\ (\%) = \frac{\text{No. of mutants}}{\text{Total number of plants studied}} \times 100$$

3. Results

3.1. Seed germination

In M_2 generation, compared to the control, the gamma irradiated and SA treated

population reflected reduced seed germination (Table 3). In both the varieties, germination

decreased with the increase in mutagen dose. In the var. Gomati VU-89, control showed a

germination percentage of 93.33%, while it reduced from 88.33 to 80.67% in 100 - 400Gy

gamma rays and 87.33 to 81.33% in 0.01-0.04% SA and 87.00 to 77.67% in 100Gy + 0.01%

SA - 400Gy+0.04% SA treatments. Variety Pusa-578 showed much reduced seed

germination, the germination percentage for control was 92.00%, while it decreased from

85.00 to 80.33% in 100 - 400Gy gamma rays and 88.33 to 76.33% in 0.01 - 0.04% SA and

83.67 to 75.33% in 100Gy + 0.01% SA - 400Gy + 0.04% SA in gamma rays+SA treatments.

As compared to M_1 generation, the %age of seed germination was improved in M_2 generation

in both the varieties.

Table 3. Effects of gamma rays, SA and their combinations on seed germination and pollen fertility in M_2 generation two varieties of cowpea

Treatment	Gomati VU-89				Pusa-578			
	Seed germination	% inhibition	Pollen fertility (%)	% reduction	Seed germination	% inhibition	Pollen fertility (%)	% reduction
Control	**93.33**	**---**	**96.53**	**---**	**92.00**	**---**	**97.77**	**---**
100Gy γ rays	88.33	5.36	90.35	6.41	85.00	7.61	88.12	9.87
200Gy γ rays	86.00	7.86	89.11	7.69	83.33	9.42	86.88	11.14
300Gy γ rays	82.67	11.43	85.40	11.54	81.67	11.23	81.19	16.96
400Gy γ rays	80.67	13.57	81.19	15.90	80.33	12.68	77.23	21.01
Mean	**84.42**	**9.55**	**86.51**	**10.38**	**82.58**	**10.24**	**83.35**	**14.75**
0.01% SA	87.33	6.43	90.84	5.90	88.33	3.99	89.11	8.86
0.02% SA	86.00	7.86	87.13	9.74	84.00	8.70	84.65	13.42
0.03% SA	83.33	10.71	81.68	15.38	79.67	13.41	77.72	20.51
0.04% SA	81.33	12.86	77.23	20.00	76.33	16.67	75.00	23.29
Mean	**84.50**	**9.46**	**84.22**	**12.76**	**82.17**	**10.69**	**81.62**	**16.52**
100Gy γ rays +0.01% SA	87.00	6.79	91.58	5.13	83.67	9.06	87.13	10.89
200Gy γ rays +0.02% SA	83.67	10.36	86.14	10.77	80.67	12.32	84.41	13.67
300Gy γ rays +0.03% SA	80.33	13.93	83.66	13.33	77.33	15.94	76.49	21.77
400Gy γ rays +0.04% SA	77.67	16.79	74.50	22.82	75.33	18.12	72.03	26.33
Mean	**82.17**	**11.96**	**83.97**	**13.01**	**79.25**	**13.86**	**80.01**	**18.16**

3.2. Pollen fertility

Variable degree of pollen fertility was recorded in different doses of mutagens employed individually and in combination (Table 3). In the var. Gomati VU-89, per cent pollen fertility reduction ranged from 6.41% in 100Gy to 15.90% in 400Gy in gamma rays treatments, while as it ranged from 5.90% in 0.01% SA to 20.00% in 0.04% SA treatments and in case of combination treatments it ranged from 5.13% in 100Gy+0.01% SA to 22.82% in 400Gy+0.04% SA. In the var. Pusa-578, per cent pollen fertility reduction ranged from 9.87% in 100Gy to 21.01% in 400Gy in gamma rays treatments, while as it ranged from 8.86% to 23.29 in 0.01% SA-0.04% SA in SA treatments and in case of combination treatments it ranged from 10.89% in 100Gy + 0.01% SA to 26.33% in 400Gy + 0.04% SA. The individual mutagen treatments caused more reduction in pollen fertility than combination treatments in both the varieties.

3.3. Chlorophyll mutations: Frequency and Spectrum

Chlorophyll mutations are frequently employed for the determination of mutagenic potency in inducing genetic variability as they are vital indicator in the assessment of induced genetic changes of mutagenized population. In both the varieties six chlorophyll mutants viz., albina, chlorina, xantha, tigrina, viridis and xanthaviridis were observed in M_2 generation. A concise account of chlorophyll mutants are presented in Table 4 and Fig 2. The spectrum and frequency of the chlorophyll mutants are presented in Table 5. The chlorophyll mutant frequencies were estimated on M_2 seedling basis. In both the varieties, except highest dose in each treatment of gamma rays, SA and gamma rays+SA, mutation frequency increased linearly with the increase in mutagen dose. The combination treatment induced higher

mutation frequency in comparison to individual treatments. SA treatments were more

effective in inducing higher chlorophyll mutation frequency than gamma rays treatments in

Pusa-578. In the var. Gomati VU-89, the chlorophyll mutation frequency ranged from 0.72 to

1.06%, 0.88 to 0.92% and 0.53 to 1.68% in gamma rays, SA and gamma rays+SA treatments,

respectively. Var. Pusa-578 responded with the lesser mutation frequency compared to the

var. Gomati VU-89. The chlorophyll mutation frequency ranged from 0.58 to 0.77% in

gamma rays, 0.89 to 1.02% in SA treatment and the range of chlorophyll deficient mutants

induced by the combined mutagens was 0.48 to 1.78% in the var. Pusa-578.

Table 4. Characteristic features of chlorophyll mutants and percentage of mutated plant progenies induced by SA, gamma rays and their combinations in
M_2 generation of two varieties of cowpea.

Mutant types and their characteristics	Treatment	No. of M₁ plant progenies	No. of plant progenies segregating in M₂	% mutated plant progenies (Mp)
		Gomati VU-89		
1.Albina	Control	270.00	0.00	0.00
	100Gy γ rays	245.00	2.00	0.82
Lethal mutation with complete absence of pigments,	200Gy γ rays	223.00	11.00	4.93
white leaves; seedlings survive for 10-12 days after germination	300Gy γ rays	208.00	7.00	3.37
	400Gy γ rays	195.00	9.00	4.62
	Total	871.00	29.00	3.33
2. Chlorina	0.01% SA	235.00	2.00	0.85
Leaves are light green in colour; mostly seedlings	0.02% SA	218.00	5.00	2.29
survive for 20-25 days but few may survive till maturity	0.03% SA	203.00	5.00	2.46
	0.04% SA	183.00	6.00	3.28
3. Xantha	Total	839.00	18.00	2.15
Yellow colour leaves due to absence of chlorophyll;	100Gy γ rays + 0.01% SA	230.00	7.00	3.04
seedlings survive for 15-20 days	200Gy γ rays + 0.02% SA	210.00	8.00	3.81
4.Tigrina	300Gy γ rays + 0.03% SA	192.00	8.00	4.17
Leaves with yellow and green patches;	400Gy γ rays + 0.04% SA	178.00	10.00	5.62
seedlings survive for 15-25 days	Total		33.00	4.07
		Pusa-578		
	Control	260.00	0.00	0.00
	100Gy γ rays	235.00	1.00	0.43
5. Viridis	200Gy γ rays	213.00	5.00	2.35
Leaves possess viridine green colour;	300Gy γ rays	200.00	6.00	3.00
reduced seedling height survive till maturity	400Gy γ rays	185.00	6.00	3.24
	Total	833.00	18.00	2.16
	0.01% SA	225.00	1.00	0.44
	0.02% SA	208.00	5.00	2.40
	0.03% SA	193.00	4.00	2.07
	0.04% SA	173.00	6.00	3.47
6.Xanthavirdis	Total	799.00	16.00	2.00
Leaves possess viridine green colour.	100Gy γ rays + 0.01% SA	220.00	6.00	2.73
	200Gy γ rays + 0.02% SA	200.00	6.00	3.00
	300Gy γ rays + 0.03% SA	182.00	7.00	3.85
	400Gy γ rays + 0.04% SA	168.00	6.00	3.57
	Total	770.00	25.00	3.25

226 **Table 5.** Frequency and spectrum of chlorophyll mutants induced by gamma rays, SA and their combinations in M_2 generation of two varieties of cowpea.

Treatment	N	Chlorophyll mutant types						CMS	F (%)	K
		Albina	Chlorina	Xantha	Tigrina	Viridis	Xanthaviridis			
Gomati VU-89										
Control	3255.00	-	-	-	-	-	-	-	-	-
100Gy γ rays	3205.00	12.00	8.00	5.00	1.00	3.00	-	29.00	0.90	-
200Gy γ rays	3146.00	13.00	7.00	5.00	-	4.00	-	29.00	0.92	-
300Gy γ rays	3102.00	10.00	8.00	4.00	2.00	4.00	5.00	33.00	1.06	-
400Gy γ rays	3050.00	9.00	8.00	2.00	-	1.00	2.00	22.00	0.72	-
0.01%SA	3185.00	13.00	10.00	1.00	3.00	1.00	-	28.00	0.88	-
0.02%SA	3102.00	14.00	8.00	1.00	-	3.00	2.00	28.00	0.90	-
0.03%SA	3052.00	10.00	7.00	5.00	4.00	-	2.00	28.00	0.92	-
0.04%SA	2925.00	9.00	10.00	4.00	-	2.00	1.00	26.00	0.89	-
100Gy γ rays+0.01% SA	3025.00	11.00	-	-	1.00	-	4.00	16.00	0.53	0.30
200Gy γ rays+0.02% SA	2956.00	13.00	12.00	-	1.00	-	3.00	29.00	0.98	0.54
300Gy γ rays+0.03% SA	2854.00	15.00	15.00	6.00	4.00	6.00	2.00	48.00	1.68	0.85
400Gy γ rays+0.04% SA	2705.00	15.00	13.00	7.00	1.00	7.00	1.00	44.00	1.63	1.02
Pusa-578										
Control	3147.00	-	-	-	-	-	-	-	-	-
100Gy γ rays	3097.00	10.00	6.00	0.00	1.00	1.00	-	18.00	0.58	-
200Gy γ rays	3038.00	11.00	5.00	3.00	-	2.00	-	21.00	0.69	-
300Gy γ rays	2994.00	9.00	5.00	3.00	2.00	2.00	2.00	23.00	0.77	-
400Gy γ rays	2942.00	7.00	7.00	0.00	-	2.00	1.00	17.00	0.58	-
0.01%SA	3077.00	11.00	10.00	1.00	4.00	3.00	-	29.00	0.94	-
0.02%SA	2994.00	13.00	10.00	2.00	-	2.00	2.00	29.00	0.97	-
0.03%SA	2944.00	13.00	7.00	3.00	3.00	2.00	2.00	30.00	1.02	-
0.04%SA	2817.00	11.00	9.00	2.00	-	2.00	1.00	25.00	0.89	-
100Gy γ rays+0.01% SA	2917.00	10.00	-	-	1.00	-	3.00	14.00	0.48	0.32
200Gy γ rays+0.02% SA	2848.00	14.00	9.00	-	2.00	-	2.00	27.00	0.95	0.57
300Gy γ rays+0.03% SA	2746.00	16.00	14.00	6.00	5.00	6.00	2.00	49.00	1.78	1.02
400Gy γ rays+0.04% SA	2597.00	14.00	14.00	6.00	3.00	5.00	2.00	44.00	1.69	1.15

227 N= Number of M_2 seedlings, CMS= Chlorophyll mutated seedlings, F= Frequency, K= Coefficient of interaction based on % M_2 chlorophyll mutation

228 frequency.

Fig 2. Chlorophyll mutants.

a Gomati VU-89 seedling (Control) showing green color of leaf.

b Albina mutant lacking both chlorophyll and carotenoid contents.

c Xantha mutant lacking chlorophyll content.

d Tigrina mutant with yellow and green patches.

e Pusa-578 Seedling (Control) showing green color of leaf.

f Chlorina mutant with light green leaves.

g Xanthaviridis mutant with yellow leaf apices.

h Viridis mutant with viridine green colour leaves.

The spectrum of chlorophyll mutations was albina> chlorina> xantha> viridis> xanthaviridis> tigrina in the var. Gomati VU-89, whereas it was albina> chlorina> viridis> xantha> tigrina> xanthaviridis in the var. Pusa-578 (Table 6). Among the mutants the occurrence of albina type of chlorophyll mutants was the highest in varieties Gomati VU-89 and Pusa-578. Among the doses the frequency of albina mutants was highest in individual and combination doses of gamma rays+SA, whereas SA treatments induced higher frequency of chlorina mutants in varieties Gomati VU-89 and Pusa-578 (Fig 3). The estimation of coefficient of interaction (k) showed synergistic effects of gamma rays and SA.

Fig 3. Comparative frequency and spectrum of chlorophyll mutations in cowpea based on pooled values of two varieties Gomati VU-89 and Pusa-578.

252 **Table 6.** Mutagen induced chlorophyll mutation frequency and spectrum in two varieties of cowpea.

Mutagen/ Variety	Comparative frequency of chlorophyll mutation spectrum						Total Frequency
	Albina	Chlorina	Xantha	Tigrina	Viridis	Xanthaviridis	
Gomati VU-89							
Gamma radiations	0.35	0.25	0.13	0.02	0.10	0.06	0.90
SA	0.38	0.29	0.09	0.06	0.05	0.04	0.90
Gamma radiations + SA	0.47	0.35	0.11	0.06	0.11	0.09	1.19
Average Frequency	0.40	0.30	0.11	0.04	0.09	0.06	1.00
Pusa-578							
Gamma radiations	0.31	0.19	0.05	0.03	0.06	0.02	0.65
SA	0.41	0.30	0.07	0.06	0.08	0.04	0.96
Gamma radiations + SA	0.49	0.33	0.11	0.10	0.10	0.08	1.21
Average Frequency	0.40	0.27	0.07	0.06	0.09	0.04	0.94

253 *Data based on pooled values of treatments

3.4. Mutagenic effectiveness and efficiency

The mutagen effectiveness and efficiency in a induced mutagenesis determines its usefulness and applicability for crop improvement. Its assessment has been evaluated on the basis of chlorophyll mutation frequency (Tables 7 and 8). The effectiveness and efficiency were highest at the intermediate treatments of mutagens, however lower doses of combined mutagen treatments were most effective and efficient in both the varieties. SA was effective than gamma rays. In both the varieties, effectiveness of single and combined mutagens treatments decreased progressively with the increase in dose in both the varieties. In the gamma irradiated population, the effectiveness ranged from 0.02 to 0.06 in the var. Gomati VU-89 and 0.01 to 0.03 in the var. Pusa-578. The effectiveness of SA treatments ranged from 25.00 to 41.66 and 16.66 to 41.66 in the varieties Gomati VU-89 and Pusa-578, respectively. The effectiveness in combination treatments fluctuated from 0.10 to 1.16 in the var. Gomati VU-89 and 0.06 to 1.00 in the var. Pusa-578. The effectiveness of mutagens was SA> gamma rays + SA> gamma rays.

271 **Table 7.** Effectiveness and efficiency of gamma rays, SA and their combinations in cowpea var. Gomati VU-89.

Treatment	(%) Seedling injury	(%) Pollen sterlity	% meiotic abnormalities	% Mutated plant progenies	Mutagenic effectiveness	Mutagenic efficiency		
	(I)	(S)	(Me)	(Mp)		Mp/I	Mp/S	Mp/Me
100Gy γ rays	10.91	6.41	3.19	2.00	0.02	0.18	0.31	0.63
200Gy γ rays	16.37	7.69	5.14	11.00	0.06	0.67	1.43	2.14
300Gy γ rays	18.88	11.54	5.47	7.00	0.02	0.37	0.61	1.28
400Gy γ rays	25.61	15.90	8.51	9.00	0.02	0.35	0.57	1.06
0.01%SA	17.40	5.90	1.63	2.00	33.33	0.11	0.34	1.23
0.02%SA	19.64	9.74	3.50	5.00	41.66	0.25	0.51	1.43
0.03%SA	22.31	15.38	6.02	5.00	27.77	0.22	0.33	0.83
0.04%SA	29.13	20.00	7.63	6.00	25.00	0.21	0.30	0.79
100Gy γ rays+ 0.01% SA	16.89	5.13	5.44	7.00	1.16	0.41	1.37	1.29
200Gy γ rays +0.02% SA	20.35	10.77	6.43	8.00	0.33	0.39	0.74	1.25
300Gy γ rays+0.03% SA	25.05	13.33	9.05	8.00	0.14	0.32	0.60	0.88
400Gy γ rays+0.04% SA	31.79	22.82	10.31	10.00	0.10	0.31	0.44	0.97

Mutagenic efficiency was evaluated on the basis of mutation frequency in relation to Mp/I, Mp/S and Mp/Me. Mp/I and Mp/S based maximum efficiency was noted at 200 Gy of gamma rays followed by 100 Gy+0.01% SA and 0.02% SA in both the varieties of cowpea. Mp/Me based efficiency was maximum at 200Gy gamma rays followed by 0.02% SA and 100Gy+0.01% SA in the var. Gomati VU-89. However, the trend was slightly altered in var. Pusa-578 as the maximum efficiency was achieved in 0.02% SA followed by 100Gy+ 0.01% SA and 100 Gy gamma rays. In both the varieties injury and sterility based mutagenic efficiency was found in the decreasing order i.e., gamma rays> gamma rays+ SA> SA. In var. Gomati VU-89 and Pusa-578, meiotic abnormalities based efficiency was higher in gamma rays + SA treatments gamma rays treatments and SA. Meiotic abnormalities based mutagenic efficiency was higher than injury and sterility based efficiency.

 Table 8. Effectiveness and efficiency of gamma rays, SA and their combinations in cowpea var. Pusa-578.

Treatments	(%) Seedling injury (I)	(%) Pollen injury (S)	% meiotic abnormalities (Me)	% Mutated plant progenies (Mp)	Mutagenic effectiveness	Mutagenic efficiency		
						Mp/I	Mp/S	Mp/Me
100Gy γ rays	12.28	9.87	3.19	1.00	0.01	0.08	0.10	0.31
200Gy γ rays	17.17	11.14	5.77	6.00	0.03	0.35	0.54	1.04
300Gy γ rays	21.51	16.96	5.83	6.00	0.02	0.28	0.35	1.03
400Gy γ rays	23.47	21.01	8.48	6.00	0.02	0.26	0.29	0.71
0.01%SA	11.99	8.86	1.05	1.00	16.66	0.08	0.11	0.95
0.02%SA	14.84	13.42	3.56	5.00	41.66	0.34	0.37	1.41
0.03%SA	19.18	20.51	5.00	4.00	22.22	0.21	0.20	0.80
0.04%SA	20.12	23.29	6.62	6.00	25.00	0.30	0.26	0.91
100Gy γ rays+ 0.01% SA	13.11	10.89	4.05	5.00	1.00	0.38	0.46	1.23
200Gy γ rays+0.02% SA	17.72	13.67	4.90	6.00	0.25	0.34	0.44	1.22
300Gy γ rays+0.03% SA	21.12	21.77	5.63	7.00	0.13	0.33	0.32	1.24
400Gy γ rays+0.04% SA	26.88	26.33	8.43	8.00	0.06	0.30	0.30	0.95

3.5. Morphological mutations

A diverse range of mutagen induced morphological mutants were obtained in the mutagenized cowpea population in M_2 generation, many of which may be useful from yield improvement prospectives. These mutants revealed differences in the traits such as plant height, growth habit, leaf, flower pod and seed. The morphological mutations, detected at different growth stages of M_2 plants, were categorically inspected throughout the growing season. The maximum frequency was recorded in gamma rays+SA treatments (5.86 and 6.56%) and the minimum frequency with gamma rays treatments (4.49 and 5.05%), while the SA treatments induced intermediate frequency of morphological mutants (5.38 and 6.05%) in the varieties Gomati VU-89 and Pusa-578, respectively (Tables 9 and 10). The overall frequencies of morphological mutants were more in Pusa-578 than Gomati VU-89. The spectrum of morphological mutations was relatively wider in combined mutagen treatment than separate treatments (Fig 4). The most prominent morphological mutations were found to be associated with the seed and flower followed by growth habit, plant height and pod length (Tables 11 and 12). In general, the occurrence of tall, bushy, broad leaves, elongated rachis, multiple and open flower, bold podded and seeded mutants were found in more numbers at the lower doses of gamma rays and SA. The gamma rays+SA treatments produced significant effects on morphological mutations, however no additive effects were observed in both the varieties. A concise account of the morphological mutants isolated is discussed below.

Fig 4. Comparative percentage and spectrum of induced morphological mutations in cowpea based on pooled values of two varieties Gomati VU-89 and Pusa-578

310 **Table 9.** Frequency and spectrum of morphological mutants induced by gamma rays, SA and their combinations in M_2 generation of cowpea var. Gomati
311 VU-89 at different stages of growth.

Treatment	N	Morphological mutant types						TMP	% Mutated plants	k
		Plant height	Growth habit	Leaf	Flower	Pod	Seed			
Control	3050.00	-	-	-	-	-	-	-	-	-
100Gy γ rays	3000.00	8.00	-	6.00	2.00	9.00	4.00	29.00	0.97	-
200Gy γ rays	2941.00	4.00	10.00	13.00	5.00	-	5.00	37.00	1.26	-
300Gy γ rays	2897.00	-	16.00	9.00	7.00	-	6.00	38.00	1.31	-
400Gy γ rays	2845.00	8.00	8.00	-	2.00	6.00	3.00	27.00	0.95	-
Total	**11683.00**	**20.00**	**34.00**	**28.00**	**16.00**	**15.00**	**18.00**	**131.00**	**4.49**	**0.00**
0.01%SA	2980.00	9.00	-	11.00	12.00	-	3.00	35.00	1.17	-
0.02%SA	2897.00	5.00	-	6.00	5.00	11.00	6.00	33.00	1.14	-
0.03%SA	2847.00	-	12.00	7.00	10.00	-	7.00	36.00	1.26	-
0.04%SA	2720.00	10.00	7.00	13.00	8.00	7.00	4.00	49.00	1.80	-
Total	**11444.00**	**24.00**	**19.00**	**37.00**	**35.00**	**18.00**	**20.00**	**153.00**	**5.38**	**0.00**
100Gy γ rays +0.01% SA	2820.00	6.00	-	-	15.00	-	9.00	30.00	1.06	0.44
200Gy γ rays+0.02% SA	2751.00	6.00	-	19.00	13.00	12.00	10.00	60.00	2.18	0.80
300Gy γ rays+0.03% SA	2649.00	-	16.00	9.00	-	-	4.00	29.00	1.09	0.37
400Gy γ rays+0.04% SA	2500.00	-	17.00	7.00	-	14.00	0.00	38.00	1.52	0.52
Total	**10720.00**	**12.00**	**33.00**	**35.00**	**28.00**	**26.00**	**23.00**	**157.00**	**5.86**	**2.13**

312 N = Number of M_2 plants, TMP = Total mutated plants, k = coefficient of interaction based on % M_2 mutated plants.

Table 10. Frequency and spectrum of morphological mutants induced by gamma rays, SA and their combinations in M_2 generation of cowpea var. Pusa-578 at different stages of growth

Treatments	N	Morphological mutant types						TMP	% Mutated plants	k
		Plant height	Growth habit	Leaf	Flower	Pod	Seed			
Control	2960.00	-	-	-	-	-	-	-	-	-
100Gy γ rays	2910.00	10.00	-	8.00	2.00	-	9.00	29.00	1.00	-
200Gy γ rays	2851.00	2.00	9.00	14.00	4.00	11.00	10.00	50.00	1.75	-
300Gy γ rays	2807.00	-	14.00	8.00	5.00	-	5.00	32.00	1.14	-
400Gy γ rays	2755.00	7.00	10.00	-	4.00	8.00	3.00	32.00	1.16	-
TOTAL	**11323.00**	**19.00**	**33.00**	**30.00**	**15.00**	**19.00**	**27.00**	**143.00**	**5.05**	**-**
0.01%SA	2890.00	10.00	-	11.00	12.00	-	4.00	37.00	1.28	-
0.02%SA	2807.00	7.00	13.00	6.00	5.00	9.00	8.00	48.00	1.71	-
0.03%SA	2757.00	-	-	8.00	10.00	9.00	9.00	36.00	1.31	-
0.04%SA	2630.00	11.00	8.00	10.00	10.00	-	7.00	46.00	1.75	-
TOTAL	**11084.00**	**28.00**	**21.00**	**35.00**	**37.00**	**18.00**	**28.00**	**167.00**	**6.05**	**-**
100Gy γ rays +0.01% SA	2730.00	8.00	-	-	14.00	-	10.00	32.00	1.17	0.46
200Gy γ rays+0.02% SA	2661.00	8.00	-	17.00	15.00	13.00	11.00	64.00	2.41	0.60
300Gy γ rays+0.03% SA	2559.00	3.00	15.00	11.00	-	2.00	5.00	36.00	1.41	0.48
400Gy γ rays+0.04% SA	2410.00	-	14.00	9.00	-	15.00	0.00	38.00	1.58	0.50
TOTAL	**10360.00**	**19.00**	**29.00**	**37.00**	**29.00**	**30.00**	**26.00**	**170.00**	**6.56**	**2.04**

N = Number of M_2 plants, TMP = Total mutated plants, k = coefficient of interaction based on % M_2 mutated plants.

320 **Table 11.** Frequency and spectrum of morphological mutants induced by various mutagens in M$_2$ generation of cowpea var. Gomati VU-89.

Mutant Type		Gamma rays		SA		Gamma rays + SA		Total		Grand Total	
		N	F %	N	F%	N	F%	N	F%	N	F%
Plant height	Tall	13.00	0.11	15.00	0.13	9.00	0.08	37.00	0.11	**56.00**	**0.17**
	Dwarf	5.00	0.04	8.00	0.07	2.00	0.02	15.00	0.04		
	Semi dwarf	2.00	0.02	1.00	0.01	1.00	0.01	4.00	0.01		
Growth habit	Compact/Bushy	8.00	0.07	7.00	0.06	12.00	0.11	27.00	0.08	**63.00**	**0.19**
	Prostrate	1.00	0.01	2.00	0.02	5.00	0.05	8.00	0.02		
	Semi dwarf Spreading	3.00	0.03	4.00	0.03	3.00	0.03	10.00	0.03		
	One sided branching	7.00	0.06	1.00	0.01	1.00	0.01	9.00	0.03		
	Axillary branching	3.00	0.03	2.00	0.02	4.00	0.04	9.00	0.03		
Leaf	Broad leaf	8.00	0.07	7.00	0.06	6.00	0.06	21.00	0.06	**59.00**	**0.17**
	Narrow leaf	5.00	0.04	5.00	0.04	7.00	0.07	17.00	0.05		
	Altered leaf architecture	2.00	0.02	2.00	0.02	5.00	0.05	9.00	0.03		
	Elongated rachis	4.00	0.03	6.00	0.05	2.00	0.02	12.00	0.04		
Flower	Multiple flower	4.00	0.04	3.00	0.03	3.00	0.03	10.00	0.03	**66.00**	**0.19**
	Colour	2.00	0.02	5.00	0.04	4.00	0.04	11.00	0.03		
	Open flower	7.00	0.06	7.00	0.06	6.00	0.06	20.00	0.06		
	Non flowering	1.00	0.01	1.00	0.01	2.00	0.02	4.00	0.01		
	Late flowering	6.00	0.05	6.00	0.05	2.00	0.02	14.00	0.04		
	Early maturity	4.00	0.03	2.00	0.02	1.00	0.01	7.00	0.02		
Pod	Small/Narrow pods	7.00	0.06	8.00	0.07	10.00	0.09	25.00	0.07		
	Bold pods	5.00	0.04	6.00	0.05	9.00	0.08	20.00	0.06	**45.00**	**0.13**
Seed	Coat colour	7.00	0.06	8.00	0.07	7.00	0.07	22.00	0.06		
	Coat pattern	6.00	0.05	9.00	0.08	8.00	0.07	23.00	0.07	**75.00**	**0.22**
	Shape and surface	10.00	0.09	11.00	0.10	9.00	0.08	30.00	0.09		
	Grand Total	**120.00**	**1.03**	**126.00**	**1.10**	**118.00**	**1.10**	**364.00**	**1.08**		

321 N= Number of mutants, F %= Mutant frequency of M$_2$ plants.

323 **Table 12.** Frequency and spectrum of morphological mutants induced by various mutagens in M_2 generation of cowpea var. Pusa-578.

	Mutant Type	Gamma rays		SA		Gamma rays + SA		Total		Grand Total	
		N	F %	N	F%	N	F%	N	F%	N	F%
Plant height	Tall	10.00	0.09	11.00	0.10	8.00	0.08	29.00	0.09	**44.00**	**0.13**
	Dwarf	4.00	0.04	7.00	0.06	1.00	0.01	12.00	0.04		
	Semi dwarf	1.00	0.01	1.00	0.01	1.00	0.01	3.00	0.01		
Growth habit	Compact/Bushy	7.00	0.06	6.00	0.05	11.00	0.11	24.00	0.07	**57.00**	**0.17**
	Prostrate	4.00	0.04	1.00	0.01	6.00	0.06	11.00	0.03		
	Semi dwarf Spreading	2.00	0.02	3.00	0.03	2.00	0.02	7.00	0.02		
	One sided branching	5.00	0.04	2.00	0.02	1.00	0.01	8.00	0.02		
	Axillary branching	2.00	0.02	3.00	0.03	2.00	0.02	7.00	0.02		
Leaf	Broad leaf	7.00	0.06	6.00	0.05	5.00	0.05	18.00	0.05	**50.00**	**0.15**
	Narrow leaf	4.00	0.04	4.00	0.04	5.00	0.05	13.00	0.04		
	Altered leaf architecture	1.00	0.01	3.00	0.03	4.00	0.04	8.00	0.02		
	Elongated rachis	5.00	0.04	5.00	0.05	1.00	0.01	11.00	0.03		
Flower	Multiple flower	3.00	0.03	2.00	0.02	4.00	0.04	9.00	0.03	**64.00**	**0.20**
	Color	3.00	0.03	4.00	0.04	5.00	0.05	12.00	0.04		
	Open flower	6.00	0.05	8.00	0.07	5.00	0.05	19.00	0.06		
	Non flowering	2.00	0.02	2.00	0.02	1.00	0.01	5.00	0.02		
	Late flowering	5.00	0.04	5.00	0.05	1.00	0.01	11.00	0.03		
	Early maturity	3.00	0.03	4.00	0.04	1.00	0.01	8.00	0.02		
Pod	Small/Narrow pods	5.00	0.04	7.00	0.06	9.00	0.09	21.00	0.06	**45.00**	**0.14**
	Bold seeded pods	4.00	0.04	5.00	0.05	8.00	0.08	17.00	0.05		
Seed	Coat color	6.00	0.05	7.00	0.06	6.00	0.06	19.00	0.06	**75.00**	**0.23**
	Coat pattern	5.00	0.04	8.00	0.07	7.00	0.07	20.00	0.06		
	Shape and surface	9.00	0.08	10.00	0.09	8.00	0.08	27.00	0.08		
	Grand Total	**103.00**	**0.91**	**114.00**	**1.03**	**102.00**	**0.98**	**319.00**	**0.97**		

324 N= Number of mutants, F%= Mutant frequency of M_2 Plants

I. Plant height mutants (Fig 5)

(i) Tall mutants: Such mutants were induced with frequency of 0.11 and 0.09% of the total morphological mutations in the varieties Gomati VU-89 and Pusa-578, respectively at lower doses of gamma rays and SA in both the cowpea varieties. The mutants were much taller with increased length of internodes, reduced number of branches and pods than control plants. Plants attained a height 180-185 cm while, the height of control plants was 178-183 cm.

(ii) Dwarf mutants: Such mutants were mostly induced with a frequency of 0.04% of the total morphological mutations. These mutants were induced at higher doses of all the treatments except the combination treatments in the var. Gomati VU-89. These mutants possess shorter internodes, smaller and lesser leaves, reduced pod and seed size and poor yield. The plant height was severely reduced and it ranged from 110-115 cm.

(iii) Semi dwarf mutants: Such mutants were induced by the moderate doses of the mutagens at a frequency of 0.01% of the total morphological mutations in var. Gomati VU-89 and Pusa-578. These mutants were semi dwarf attaining height of 140-150 cm, harbour shorter internodes, reduced branching, less pods and lower yield in comparison to the untreated plants. Pod length was increased, however, the seeds were smaller in size.

II. Growth habit mutants (Fig 5)

(i) Compact/Bushy mutants: Such mutants are characterised with 150.55 cm average height, robust growth, increased branches, condensed internodes that imparted it a bushy appearance. The yield per plant was slightly higher and matured five days earlier compared to control plants. Such mutants were mostly induced by the moderate and higher treatments. The bushy mutants appeared at 0.08% of the total morphological mutations in the var. Gomati VU-89 and 0.07% in var. Pusa-578.

(ii) Prostrate mutants: Such mutants were flat on ground, branches spread several meters,

possess long internodes, and reflected trailing tendency at the soil surface. They were low

yielding with small sized pods, less number of pods containing 6-7 shriveled seeds with

rough and hard seed coat. In both the varieties of cowpea prostrate mutants were commonly

induced with combination treatments at a frequency of about 0.02% and 0.03% in the

varieties Gomati VU-89 and Pusa-578, respectively.

(iii) Spreading mutants: Such mutants were observed in 200Gy gamma rays treatment

(Gomati VU-89) and 0.03% SA treatment (Pusa-578). These mutants possess long internodes

and large canopy. The leaves were mostly broad and branches appeared spreading with wide

branch angle. Seed yield per plant was significantly reduced. Their frequency was 0.03% and

0.02% in varieties Gomati VU-89 and Pusa-578, respectively.

(iv) One sided branching mutants: Mutant plants had thick stems bearing branches on one

side and were late in flowering and maturing, low yielded with few pods and shriveled seeds.

These mutants were isolated in SA doses in both the cowpea varieties and in combination

treatment of 200Gy gamma rays + 0.02% SA in the var. Pusa-578. Their frequency was

recorded as 0.03% and 0.02% in the var. Gomati VU-89 and Pusa-578, respectively.

(v) Axillary branched mutants: Mutant plants with profuse axillary branching, condensed

internodes, small seeded pods and low seed set were mostly isolated from the moderate and

higher mutagen treatments. Their frequency was recorded as 0.03% and 0.02% in var.

Gomati VU-89 and var. Pusa-578, respectively.

Fig 5. Morphological mutations (plants with altered height and growth habit).

a Plant showing normal height and growth habit (Control).

b Tall mutant with less branches and long internodes.

c Dwarf mutant with stunted growth, shorter internodes and profuse branching.

d Semi dwarf mutant with sparse branching and short internodes.

e Spreading mutant with herbaceous branching and long vegetative phase.

f Bushy mutant with vigorous growth, profuse branching and condensed internodes.

g Axillary branched mutant.

h Prostrate mutant with horizontal growth habit bearing slender, soft and light green
378   branches.

III. Leaf mutants (Fig 6-7)

(i) Broad/Giant leaves mutants: Such mutants were characterised by broad sized leaves,

elongated ranchis, tall and vigorous growth. The yield of these mutant plants was found to be

higher with the increase in mean number of pods than control plants. Moderate doses of

gamma rays and SA treatments induced higher frequency of broad leaf mutants. Their

frequency was 0.06% and 0.05% in the var. Gomati VU-89 and var. Pusa-578, respectively.

(ii) Narrow leaves mutants: Such mutants possess narrow leaves with pointed leaf tips.

Compared to control, the yield was lower, small pods with few seeds per pod. Higher doses

of mutagens induce higher frequency of mutants in both the cowpea varieties. Their

frequency was recorded as 0.05% and 0.04% in the var. Gomati VU-89 and var. Pusa-578,

respectively.

(iii) Altered leaf architecture mutants: This type of morphological mutations in plants showed

wide variations in the leaf margins, apices leaf size and leaflet umber. Plants showed reduced

height (about 95 cm) at maturity. Such mutants were induced by all the mutagen treatments

with the frequency of 0.03% and 0.02% in the var. Gomati VU-89 and var. Pusa-578

respectively.

(iv) Elongated rachis: Mutants possessed increased length of rachis with narrow leaflets.

They appeared mostly at the lower mutagen doses with a frequency of 0.04% and 0.03% in

the var. Gomati VU-89 and var. Pusa-578, respectively.

Fig 6. Morphological mutations (plants with altered leaf morphology).

a Trifoliate leaf of Control (Ovate).

b Broad leaf mutant (Obovate).

c Narrow leaf mutant (hastate).

d Elongated rachis mutant (hastate).

e Trifoliate mutant (Orbicular).

f Altered leaf architecture.

Fig 7. Morphological mutations (plants with altered leaflet number)

g Mutant showing incision of leaf.

h Tetrafoliate with fused terminal leaflets.

i Pentafoliate with variations in leaflet size and shape.

j Tetrafoliate with free leaflets.

k Broad tetrafoliate leaflets.

l Bifoliate leaf.

IV. Flower mutants (Fig 8)

(i) Multiple flower mutants: Such mutants were characterized by the peduncle bearing

three/four flowers against two in control plants. Pods and seeds were small and shrivelled.

Mutants were late in maturity compared to the control. These mutants appeared at frequency

of about 0.03% in both the varieties.

(ii) Flower colour mutants: Such mutants were characterized by white or blue coloured

flowers compared to yellow coloured flowers in control. The 300 Gy gamma rays showed

higher frequency of white or blue flower colour mutants. Blue colour flower mutants were

observed at higher frequency than white colour flower mutants in the var. Gomati VU-89,

whereas in the var. Pusa-578, white colour flower mutants were at higher frequency than the

blue colour mutants. Their frequency was recorded as 0.03% in the var. Gomati VU-89 and

0.04% in the var. Pusa-578.

(iii) Open flower mutants: Such mutants harboured bell type flowers with wide open keel

and wings and were induced at a frequency 0.06% of total mutation at moderate and higher

doses of SA in var. Gomati VU-89 and Pusa-578. The anthers and stigmas were exposed and

the flowers withered without forming pods.

(iv) Non-flowering/Vegetative mutants: These mutants were devoid of flowers throughout the

life cycle, induced at a frequency of 0.01% and 0.02% in the var. Gomati VU-89 and var.

Pusa-578, respectively. They possessed dark green leaves, profuse branching, petite

internodes and big leaf. These mutants were noticed in SA and gamma rays+SA treatments.

(v) Late flowering mutants: In these mutant plants, the flowering as well as maturity period

was delayed by 8-10 days in comparison to control plants. Higher doses of mutagens induced

large number of late flowering mutants. Their frequency was 0.04% and 0.03% in the var.

Gomati VU-89 and var. Pusa-578 respectively.

(vi) Early maturity mutants: Such mutants matured 10-15 days earlier than control and were

induced at the lower mutagens doses used singly or in combination. The frequency of such

mutants was 0.02% in both the varieties.

Fig 8. Morphological mutations (altered flower colour and number per peduncle).

441   **a** Yellow colour flowers (Control).

442   **b** Blue and yellow colour petals.

443   **c** Single peduncle bearing three and four flowers.

444   **d** Open flower mutant.

445   **e** Light blue colour flower mutant.

446   **f** White flowers mutant.

V. Pod mutants (Fig 9)

(i) Small/Narrow pods: Such mutants possessed small and /or narrow pods compared to long

pods in control. The seed yield per plant was also reduced as compared to control plants.

Higher doses of individual mutagens induced a broad spectrum of these mutants and no such

mutants were observed in combined mutagen treatments. Their frequency was recorded as

0.07% and 0.06% in the var. Gomati VU-89 and var. Pusa-578, respectively.

(ii) Bold seeded pod mutants: The mutant plants are characterised by robust growth, normal

height, long pods containing 14-15 seeds and were induced at a frequency of 0.06% and

0.05% in the var. Gomati VU-89 and Pusa-578 respectively. The bold size of pods led to

significant increase in yield per plant as compared to control. Lower and moderate doses of

individual and combined mutagens induced broad spectrum of these mutants.

Fig 9. Morphological mutations (altered pod size).

459   **a** Normal pod (control).

b Long and narrow pods.

c Broad pod with bold seeds.

d Broad pod with small seeds.

e Variations in pod width.

f Variations in pod length.

VI. Seed mutants (Fig 10)

(i) Seed coat colour mutants: Such mutants are characterized by the different seed coat color

observed after harvest. Seeds were brown and white colour in control plants of the var.

Gomati VU-89 and var. Pusa-578, respectively. The colour of seeds changed to dark red,

light red, and black in the treated population of both the varieties. These seed mutants

appeared at frequency of 0.06% in both the varieties.

(ii) Seed coat pattern mutants: The mutants exhibited marbled, dotted and spotted seed coat

patterns and such mutants occurred at frequency of 0.07% and 0.06% in the var. Gomati VU-

89 and var. Pusa-578, respectively.

(iii) Seed shape and surface mutants: Seeds of control plants of both the varieties were of

kidney shaped with rough seed surface and were induced at a frequency of 0.09% and 0.08%

in the var. Gomati VU-89 and Pusa-578, respectively. Mutants with globular shape and

smooth seed surface were isolated in the mutagen treated population.

Fig 10. Morphological mutations (altered seed pattern, colour and shape)

a Variations in the seed coat colour.

b Bold and small seed.

c Light and dark red seeds with smooth surface.

d Wrinkled seeds with irregular surface.

4. Discussion

A considerable decline in seed germination, seedling height, pollen fertility and plant survival of treated population was observed in mutagen treatments. However, the extent of diminution in the bio-physiological parameters differed among mutagen doses. The diminution in plant survival was recorded in all the mutagen doses, however, no direct correlation between the mutagen doses and the bio-physiological traits studied was observed which confirmed the fact that the mutagens interact independently with the genes governing a particular trait and that mutations occur in a random manner. These results are in line with earlier reports on several grain legumes [15], [16], [17], [18], [19], [20], [21]. However, the promoting effects of mutagen doses on bio-physiological parameters were also reported in several other crops [22], [23], [24], [25], [26].

5.1. Seed Germination

The assessment of seed germination gives an idea about the mutagenic potency, the dose at which germination is below 50 per cent is treated as lethal or undesirable. Since, the purpose of this present study was to develop large mutagenized population to facilitate effective screening of the high yielding mutants, therefore, the lethal doses of mutagens were discarded to ensure high frequency of desirable mutation. Decreased germination due to mutagen treatments may be the attributed to mutagen induced germination inhibition. The inhibition in seed germination was found to be more or less comparable in gamma rays and SA treatments, while the combined treatments induced higher seed germination inhibition in both the cowpea varieties, which may be attributed to the synergistic effect of the two mutagens. Chauhan and Singh [27] reported that gamma rays interact and cause disruption of

tunica zone that leads to the decrease in seed germination. Interference of the mutagens with

the auxin synthesizing enzymes may be another reason for seed germination inhibition [28].

In the present study, the decrease in pollen fertility has been noticed in the

mutagenized population. Pollen sterility was maximum in combined mutagen doses than

individual mutagen doses. This may be attributed to meiotic anomalies induced by gamma

rays, SA and their combination treatments leading to the formation of aberrant pollen grains.

Structure and physiology of the pollen grains is under genetic control and any irregular

meiotic cycle may induce significant variations in the pollen properties. Reduction in pollen

fertility after mutagen treatments have also been reported in lentil [29], [30], cowpea [31],

fenugreek [32], faba bean [33]. The pollen sterility percentage was found remarkably less in

M_2 than M_1 generation that depict the operation of recovery mechanism between the two

generations.

The survival at maturity was declined with increasing mutagen dose in both the

varieties of cowpea. These findings support the earlier reports in *Lathyrus sativus* [34] in

Capsicum annum, [35]. The reduction of plant survival at maturity may be attributed to the

disturbed physiological processes [36], [37] or cytological damages Manjaya [38]. The

observations recorded on several bio-physiological and cytological parameters revealed that

cowpea var. Pusa-578 was comparatively more sensitive towards the individual and

combination doses than Gomati VU-89. The differential response of varieties from the same

species towards the mutagens have also been reported earlier in *Cajanas cajan* [39], *Vigna*

spp [40], [41], [42], *Phaseolus vulgaris* [43], [44], *Jatropha curcus* [45] and *Lens culinaris*

[46]. Sparrow *et al.* [47] suggested that differential genotypic sensitivity towards mutagens

may be due to the gene loci mutated by one mutagen in one variety were not necessarily

mutated by the second mutagen in another variety. The two cowpea varieties viz., Gomati

VU-89 and Pusa-578, used in the present study, were from different races, therefore, it is

obvious that the two varieties have substantial variation in their genome and thereby the

heritable genetic identity of the two varieties is immensely different. Although the spectrum

of variations was more or less similar in both the varieties but the mutagen doses to achieve

that were not identical in various parameters studied. It can be concluded from the present

bio-physiological that mutagenesis induced a comparable spectrum of apparent variation

through random mutation in both the varieties of diverse origin with the variable treatments

of gamma rays and SA.

5.2. Chlorophyll mutations

In mutation breeding, the increase in mutation frequency is necessary for crop

improvement. In order to acquire desired mutations at highest rates, the picking of optimum

mutagen and appropriate dose is important. Both the varieties of cowpea showed a

differential response towards gamma rays, SA and gamma rays+SA treatments. The different

chlorophyll deficient mutants such as albina, chlorina, xantha, viridis, tigrina and

xanthaviridis were observed in M_2 generation were identified as per the categorization of

Gustafsson [9]. Gamma rays+SA doses induced higher frequency of chlorophyll mutations

than the gamma rays and SA. Intermediate doses showed the maximum frequency of

mutations compared to the lower and higher doses. Similar trends of higher chlorophyll

mutation frequency with intermediate mutagen doses were also reported in *Cicer arietinum*

[48], *Oryza sativa* [49], *Vigna mungo* [50] *Zingiber officinale* [51], *Lablab purpureus* [52].

The reduced chlorophyll mutation frequency at the higher mutagen doses may be due to

saturation of mutations which lead to the exclusion of the mutant cells during plant growth

and development. On contrary Pavadai *et al*. [53] in soybean, Barshile *et al*. [54] in chickpea,

Kumar *et al*. [55] in mungbean reported the linear increase in chlorophyll mutation frequency

with increase in doses of different mutagen treatments. In the present study, albina and

chlorina mutants were found more frequently in both the cowpea varieties. SA was found to

be more potent in the production of albina and chlorina mutants than gamma rays. The

gamma rays+SA treatments have induced maximum frequency of albina and chlorina

mutants than gamma rays and SA treatments employed individually. The synergistic effects

of gamma rays and SA in the combination treatments are very much evident from the higher

frequency of chlorophyll mutations. The possible reason for synergism might be that the

mutagen first applied may expose the accessible protected mutable sites to the second

mutagen, the repair enzymes may be rendered non functional by the second mutagen which

indirectly facilitates the fixation of mutations induced by first mutagen [11]. The synergism

between physical and chemical mutagens in combined mutagen treatments were earlier

documented in *Cicer arietinum* [56], *Phaseolus vulgaris* [57], *Vigna spp.*, [58], *Pisum*

sativum [59], *Linum usitatissimum* [60].

5.3. Mutagenic effectiveness and efficiency

Induced mutagenesis has been known as a vital tool for the improvement of wide

range of crops. The success of mutation breeding depends on enhancing the efficiency of

desired mutation. Further the mutagen choice is imperative to achieve the desired mutation

rate. Assessment of mutagenic effectiveness and efficiency provides an idea about the utility

of mutagens in inducing the desired mutation. Mutagenic effectiveness refers to mutation

frequency induced by the unit mutagen dose [12], [61]. Whereas mutagenic efficiency on the

contrary indicate the mutation induction (genetic damage) in M_2 generation in relation to the

biological damage viz., injury, sterility and meiotic aberrations induced in M_1 generation.

Henceforth, mutagenic effectiveness reveal genotypic sensitivity while as mutagenic

efficiency reveals mutagenic potency.

The SA doses were superior among mutagen doses employed in present study. On the

basis of frequency of chlorophyll mutants, the order of mutagenic effectiveness was SA>

gamma rays + SA> gamma rays. Similar results have been reported by Balai and Krishna

[62] while studying the effectiveness of ethylmethane sulphonate, sodium azide and

hydroxylamine on mungbean. Earlier workers such as Kaul and Bhan [63] in rice, Laskar and

Khan [64] in lentil, Wani [65] and Shah *et al*. [66] in chickpea, Khan and Tyagi [67] in

soybean and Dhulgande *et al*. [68] in pea have also reported the more effectiveness of

chemical mutagens over physical mutagens. In both the varieties Gomati VU-89 and Pusa-

578, intermediate doses of gamma rays and SA and the lower doses of the combined mutagen

treatments were found to be more effective. The results were in close agreement with the

results of Girija and Dhanavel [69] while studying the mutation breeding in cowpea. As per

them, in order to attain higher effectiveness and efficiency, the mutation effect was more

prominent in the cell that led to meiotic aberrations, physiological changes, which decreased

cell survival and eliminated the mutation. The reduced effectiveness at the higher mutagen

doses revealed that the mutations and their frequency are random events irrespective of the

mutagen doses employed. Our results reflected a differential effectiveness in two varieties of

cowpea which depict the genetic divergence and mutagenic sensitivity towards different

mutagen doses.

The efficiency of the gamma rays, SA and gamma rays+SA treatments were assessed

on the basis of three criteria such as Mp/I, Mp/S and Mp/Me. Mutagenic efficiency shows

wide variation in the values depending on the criteria used for its evaluation. Each criterion

revealed wide variation in the values of efficiencies for same mutagen dose, thereby

reflecting the flexibility of using one or all at a time depending on the objectives of

assessment. Lower doses showed enhanced mutagenic efficiency as compared to higher

doses. Mp/Me based mutagenic efficiency was higher in comparison to Mp/I and Mp/S basis

in both the varieties. This may be due to low frequency of meiotic abnormalities than

seedling injury and pollen sterility in mutagen treatments. In the present study, the gamma

rays+SA treatments were more efficient than individual gamma rays and SA treatments and

this may be attributed to the higher frequency of chlorophyll mutants induced at combination

treatments. On contrary Zeerak [70] in brinjal reported that gamma rays+ EMS doses were

less efficient than the individual doses of gamma rays and EMS. In the present study, lower

and medium doses of gamma rays and SA and gamma rays+SA treatments were found to be

more efficient in both the varieties of cowpea. The results were in line with the earlier results

in *Lens culinaris* [64] *Glycine max* [67], *Vigna mungo* [71]. However, contrary results of

linear increase in mutagenic efficiency with the increase in the mutagen concentrations have

also been reported in black gram [72] and millet [73]. Makeen *et al*. [74] also reported higher

mutagenic efficiency at lower and medium doses in urdbean. The higher efficiencies of the

lower and medium doses may be due to the less biological damages as compared to higher

mutagen doses.

5.4. Morphological mutations

In the present study, several morphological mutants showing variations in characters

like altered plant height, growth habit, leaf architecture, flower colour and number, pod and

seed shape were isolated in M_2 generation of cowpea. Even though the majority of the

induced morphological mutants were uneconomical, however, these mutants can serve as a

source of valuable genes in hybridisation programmes. Toker [75] reported that the

morphological mutants are useful in gene mapping and phylogentic studies of crops. Mutagen

induced chromosomal anomalies or the pleiotropic effects of mutated genes could be the

probable reason for the induction of morphological mutants [76], [77]. The present study

morphological mutants revealed varying frequency and broader spectrum both between and

within the varieties at different mutagen doses reflecting the differential response of varieties

towards mutagen doses. Different mutagen doses and duration of treatment may be attributed

to the relative proportion of different mutation types [78]. In both the varieties Gomati VU-89

and Pusa-578, combined mutagen doses induced higher frequency of morphological

mutations followed by individual SA and gamma rays doses. Khursheed *et al*. [79] reported

highest frequency of morphological mutants in gamma rays and lowest with combined

treatments while investigating the frequency and spectrum of M_2 morphological faba bean

mutants. Previous reports of a high frequency and boarder spectrum of morphological

mutations in crops such as *Vigna mungo* [41], *Cicer arietinum* [80], [81], *Lens culinaris* [82],

Phaseolus vulgaris [83] *Glycine max* [67] also confirm that mutagens are effective in

inducing morphological mutations in wide range of crops that may serve as a source of

valuable genes for the crop improvement programs.

Some morphological mutants (early maturing, dwarf, bushy, bold seeded pods and

flower colour), screened in M_2 generation in the present study, can serve as source of

valuable genes or used as a parents in cross breeding programmes. Dwarf mutants, as

recorded in the present study, have been reported earlier in chickpea [21] barley [84],

mungbean spp. [77], lentil [85]. The decreased length of internodes and/or number of

internode could be attributed to the induction of dwarf and bushy mutants [86]. In our

experimentation dwarf and bushy mutants showed substantial decrease in internode length

but number of internodes were similar to that of control plants. The mutated genes that

control the altered plant height and growth habits are monogenic recessives [87]. Whereas

polygenes govern the semi dwarf trait in mutants in wheat [12] and triticale [88]. However,

Qin *et al*. [89] reported a dominant mutation in a single gene in dwarf mutants in rice. Both

dwarfness and bushy habit of a genotype ensure more plants per unit area, thereby,

significantly contribute to the production and productivity. The bold seeded pod mutants

induced are associated with high yield which may be valuable in increasing seed size that

lead to enhanced cowpea yield. Singh [90] reported that gene mutations govern the bold

seeded trait in *Vigna mungo*. Few early maturing mutants with seed yield comparable to the

controls were induced in the mutagenized population of both the cowpea varieties. The

mutants with modified phenotypes may be due to mutagen induced DNA breaks and altered

auxin, mineral and amino acids metabolism [91], [92], [41]. The morphological mutants

induced in present study were stable, non segregating and true breeding. Similarly, several

workers have also reported that the progenies of tall, dwarf, prostrate, bushy and bold seeded

bred true for the altered traits in subsequent generations [93], [94], [64]. The flower colour

trait is governed by multigene family, henceforth, flower colour mutants obtained in the

present study, can serve as genetic markers in mutation breeding experiments [95], [96], [97].

Conclusions

Increase in mutagen doses lead to a progressive decline in the values of all bio-

physiological parameters. Combination treatments showed an enhanced effect on seed

germination, seedling height and pollen fertility compared to individual mutagen treatments.

A dose dependant increase in the frequency of chlorophyll mutations was observed in both

the varieties. The estimation of coefficient of interaction (k) did not show any additive effects

of mutagens. The spectrum of chlorophyll mutations in the var. Gomati VU-89 was albina>

chlorina> xantha> viridis> xanthaviridis> tigrina, whereas it was albina> chlorina> viridis>

xantha> tigrina> xanthaviridis in the var. Pusa-578. In general, mutagenic effectiveness and

efficiency was highest at the moderate doses of gamma rays and SA, however, lower doses of

gamma rays+SA were most effective and efficient in both the varieties. The effectiveness of

mutagens was found as SA> gamma rays + SA> gamma rays. Mp/I and Mp/S based

mutagenic efficiency was gamma rays> gamma rays+ SA> SA whereas Mp/Me based order

was gamma rays + SA> gamma rays> SA. A diverse range of mutagen induced

morphological mutants were obtained in the population of two varieties of cowpea in M_2

generation, many of which are useful from yield improvement prospective. Of all the mutant

types, seed and flower mutants followed by growth habit, plant height and pod length were of

maximum occurrence in the two varieties. The spectrum of morphological mutations induced

by combined mutagens was relatively boarder than gamma rays and SA treatments in both

the varieties. The highest frequency was noted in combination treatments of gamma rays+SA

followed by SA and gamma rays treatments in both the varieties.

References

1. FAO. Definition and Classification of Commodities: Pulses and Derived Products. 1994.

2. DES. Agriculture statistics at a glance. Directorate of Economics and Statistics, Department of Agriculture and Economics, Ministry of Agriculture, Government of India 2017.

3. Summerfield RJ, Huxley PA, Dart PJ, Hughes AP. Some effects of environmental stress on seed yield of cowpea (*Vigna unguiculata* (L.) Walp.) cv. Prima. Plant and Soil. 1976; 44: 527–546.

4. Sprent IJ, Odee WD, Dakora FD. African Legume: a vital but under-utilized resource. J Exp. Bot. 2009; 61: 1257–1265.

5. Fery RL. The cowpea: production, utilization, and research in the United States. Hort. Reviews. 1990; 12: 197–222.

6. Raina A, Khursheed S, Khan S. Optimisation of mutagen doses for gamma rays and sodium azide in cowpea genotypes. Trends. Biosci. 2018; 11: 2386–2389

7. FAO. Production Statistics. Food and Agriculture Organization, Rome, Italy 2017.

8. Raina A, Laskar RA, Tantray YR, Khursheed S, Wani MR, Khan S. Characterization of Induced High Yielding Cowpea Mutant Lines Using Physiological, Biochemical and Molecular Markers. Sci. Rep. 2020; 10: 1–22.

9. Gustafsson A. The mutation system of the chlorophyll apparatus. Lund Univ. 1940

10. Khan S. Studies on Chemical Mutagenesis in Mungbean (*Vigna radiata* (L.) Wilczek). Ph.D. Thesis. Aligarh Muslim University, Aligarh. 1990.

11. Sharma RP. Increased mutation frequency and wider mutation spectrum in barley induced by combining gamma rays with ethyl methane sulphonate. Indian J. Genet. 1970; 30: 180–186.

12. Konzak CF, Nilan RA, Wagner J, Foster RJ. Efficient chemical mutagenesis. Rad. Bot. 1965; 5: 49–70.

13. UPOV. Cowpea (Vigna unguiculata L. Walp. subsp. *sesquipedalis* (L.) Verdc.). International Union for The Protection of New Varieties of Plants Nairobi, Kenya, 2007; 1–20

14. IBPGR. Cowpea descriptors. IBPGR Secretariat, Rome, Italy, 1983; 1–30.

15. Toker C, Cagirgan MI. Spectrum and frequency of induced mutations in chickpea. Intl. Chickpea Pigeonpea Newslet. 2004; 11: 8–10.

16. Khan S, Wani MR. Comparison on the effect of chemical mutagens on mungbean. Advanced Plant Sci. 2005; 18: 533–535.

17. Lal GM, Toms B, Lal SS (2009) Mutagenic sensitivity in early generation in black gram. Asian J. Agric. Sci. 2009; 1: 9–11.

18. Bhat M, Khan S, Kozgar MI. Studies on induced mutations in chickpea (*Cicer arietinum* L.) I. Responses of the mutagenic treatments in M1 biological parameters. Electronic J. Plant Breed. 2011; 2: 422–424.

19. Khan H, Laskar RA, Khan S. Comparative estimation of induced cytotoxicity and mutagenic potency of EMS and SA using *Vicia faba* L. as a biological system, In: Krishnan S., Rodrigues BF., editors. Advances in Plant Sciences and Biotechnology. Goa University; 2015. pp. 172–186.

20. Laskar RA, Laskar AA, Raina A, Khan S, Younus H. Induced mutation analysis with biochemical and molecular characterization of high yielding lentil mutant lines. Intl. J.Biol. Macromolecules. 2018; 109: 167–179.

21. Raina A, Laskar RA, Khursheed S, Khan S, Parveen K, Amin R, Khan S. Induce
physical and chemical mutagenesis for improvement of yield attributing traits and
their correlation analysis in chickpea. Intl. Let. Nat. Sci. 2017; 61: 14–22.

22. Venkateshwarlu S, Singh RM, Singh RB, Singh BD. Radiosensitivity and frequency
of chlorophyll mutations in pigeonpea. Indian J. Genet. 1978; 38: 90–94.

23. Vandana, Dubey DK. Effect of ethylmethane sulphonate (EMS) and diethyl sulphate
(DES) in faba bean. FABIS Newslet. 1988; 32: 18–22.

24. Mensah JK, Obadoni B. Effects of sodium azide on yield parameters of groundnut
(*Arachis hypogaea* L.). Afr. J. Biotechnol. 2007; 6: 668–671.

25. Boureima S, Diouf M, Silme RS, Diop T, Van Damme P, Çağirgan Mİ.
Radiosensitivity of African sesame cultivars to gamma rays. Turk. J. Field Crops.
2009; 14: 181–190.

26. Verma AK, Banerji BK, Chakrabarty, Datta SK. Studies on Makhana (*Euryale ferox*
Salisbury). Current Sci. 2010; 99: 795–800.

27. Chauhan MP, Singh IS. Inheritance of protein content in lentil (*Lens culinaris*
Medik.). Legume Res. 1995; 18: 5–8.

28. Usuf KK, Nair PM. Effect of gamma irradiation on the indole acetic acid synthesizing
system and its significance in sprout inhibition of potatoes. Rad. Bot. 1974; 14: 251–
256.

29. Reddy VRK, Annadurai M. Cytological effects of different mutagens in lentil (*Lens*
culinaris Medik). Cytologia. 1992; 57: 213–216

30. Taziun T, Laskar RA, Amin R, Parveen SK. Effects of dosage and durations of
different mutagenic treatment in lentil (*Lens culinaris* Medik.) cultivars Pant L 406
and DPL 62. Legume Res. 2017; 41: 500–509.

31. Singh VV, Krishna RK, Arya RK. Induced chemical mutagenesis in cowpea (*Vigna*
unguiculata (L) Walp. Indian J. Genet. 2006; 66: 312-315.

32. Choudhary S, Ansari MYK, Khan Z, Gupta H. Cytotoxic action of lead nitrate on
cytomorphology of *Trigonella foenum-graecum* L. Turk. J. Biol. 2012; 36: 267–273.

33. Bhat TA, Wani AA. Mutagenic Effects on Meiosis in Legumes and a Practical Case
Study of *Vicia faba* L. In: Bhat TA., Wani AA. editors. Chromosome Structure and
Aberrations. Springer; 2007. pp. 219–244.

34. Kumar S, Dubey DK. Effect of separate and simultaneous application of gamma rays
and EMS on germination, growth, fertility and yield in cultivars Nirmal and LSD-3 of
khesari (*Lathyrus sativus* L.) Var. P-505. J. Phytol. Res. 1998; 11: 165–70.

35. Dhamayanthi KPM, Reddy VRK. Cytogenetic effects of gamma rays and ethyl methane sulphonate in chilli pepper (*Capsicum annuum* L.). Cytologia. 2000; 65: 129–133.

36. Sato M, Gaul H. Effect of ethyl methane sulphonate on the fertility of barley. Rad. Bot. 1967; 7: 7–15.

37. Meherchandani M. Effects of gamma radiation on the dormant seeds of *Avena futua* L. Rad. Bot. 1975;15: 439–443.

38. Manjaya JG. Genetic improvement of soybean variety VLS-2 through induced mutations. In: Shu QY. editors. Induced Plant Mutation in the Genomic Era. Food and Agriculture Organization of the United Nations, Rome; 2009. pp. 106–110.

39. Pandey N, Ojha CB, Jha VB, Singh NB. Effect of chemical mutagens on the rate of germination, seedling mortality, and induced sterility in pigeonpea. Int. Chickpea and Pigeonpea Newslet. 1996; 3: 65–67.

40. Khan S, Siddiqui BA. Mutation genetic studies in mungbean. I. Variability components and genetic parameters. Thailand J. Agric. Sci. 1995; 28: 113–124.

41. Goyal S, Khan S. Differential response of single and combined treatment in moist seeds of urdbean. Indian J. Bot. Res. 2010; 6: 183–188.

42. Kozgar MI, Khan S, Goyal S. EMS induced mutational variability in *Vigna radiata* and *Vigna mungo*. Res. J. Bot. 2011; 6: 31–37.

43. Al-Rubeai MAF, Godward MBE. Genetic control of radiosensitivity in *Phaseolus vulgaris* L. Env. Exp. Bot. 1981; 21: 211–216.

44. Svetleva DL. Crino P. Effect of ethyl methanesulfonate (EMS) and n- nitrosen -ethyl urea (ENU) on callus growth of common bean. J. Central European Agric. 2006; 6: 59–64.

45. Dhakshanamoorthy D, Selvaraj R, Chidambaram A. Physical and chemical mutagenesis in *Jatropha curcas* L. to induce variability in seed germination, growth and yield traits. Rom. J. Biol. 2010; 55: 113–125.

46. Gaikwad NB, Kothekar VS. Mutagenic effectiveness and effi ciency of ethylmethane sulphonate and sodium azide in lentil (*Lens culinaris* Medik.). Indian J. Genet. 2004; 64: 73–74.

47. Sparrow AH, Sparrow RC, Thompson KH, Schairer DA. The use of nuclear and chromosome variables in determining and predicting radiosensitivity. In: Use of Induced Mutation in Plant Breeding. Rep/IAEA, Rome; 1965; 102–132.

799 48. Khursheed S, Raina A, Laskar RA, Khan S. Effect of gamma radiation and EMS on
800   mutation rate: their effectiveness and efficiency in faba bean (*Vicia faba* L.).
801   Caryologia. 2018; 71: 397–404.

802 49. Yamaguchi H, Hase Y, Tanaka A, Shikazono N, Degi K, Shimizo A, Morishita T.
803   Mutagenic effects of ion beam irradiation on rice. Breed. Sci. 2009; 59: 169–177.

804 50. Thilagavathi C, Mullainathan L. Isolation of macro mutants and mutagenic
805   effectiveness, efficiency in Black Gram (*Vigna mungo* (L.) Hepper). Global J. Mol.
806   Sci. 2009; 4: 76–79

807 51. Pawar N, Pai S, Nimbalkar M, Kolar F, Dixit G. Induction of chlorophyll mutants in
808   Zingiber officinale Roscoe by gamma rays and EMS. Emirates J Food Agric. 2010;
809   22: 406–411.

810 52. Srinivas RT, Veerabadhiran P. Efficiency and effectiveness of physical and chemical
811   mutagens and their combination in inducing chlorophyll mutations in M_2 generation
812   of Lablab [*Lablab purpureus* (l.) Sweet var. Typicus.]. Electronic J. Plant Breed.
813   2010; 1: 752–757.

814 53. Pavadai P, Girija M, Dhanavel D. Effectiveness, efficiency and biochemical content
815   of physical and chemical mutagens in soybean (*Gylcine max* (L.) Merr.). J. Phytol.
816   2009; 1: 444–447.

817 54. Barshile JD, Auti SG, Dalve SC, Apparao BJ. Mutagenic sensitivity studies in
818   chickpea employing SA, EMS and gamma rays. Indian J. Pulses Res. 2006; 19: 43–
819   46.

820 55. Kumar A, Parmhansh P, Prasad R. Induced chlorophyll and morphological mutations
821   in mungbean (*Vigna radiata* (L.) Wilczek). Legume Res. 2009; 32: 41–45.

822 56. Kozgar MI. Induced Mutagenesis: Biophysiological Damages and Cytological
823   Abberrations. In: Mutation Breeding in Chickpea. 2014. pp. 13–30.

824 57. Agarkova SN, Yakovlev AG. Effects of gamma rays and EMS on mutation frequency
825   in *Phaseolus vulgaris*. Referationyl Zhurnal. 1977; 3: 55–56.

826 58. Singh AK. Mutagenic effectiveness and efficiency of gamma rays and ethylmethane
827   sulphonate in mungbean. Madras Agric. J. 2007; 94: 7–13.

828 59. Mehandjiev A, Kosturkova G, Mihov M. Enrichment of Pisum sativum gene
829   resources through combined use of physical and chemical mutagens. Israel J. Plant
830   Sci. 2001; 49(4): 280–284.

60. Bhat IA, Pandit UJ, Sheikh IS, Hassan ZU. Physical and chemical mutagenesis in
Linum usitatissimum L. to induce variability in seed germination, survival and growth
rate traits. Current Bot. 2016; 7: 28–32.

61. Bhosale SS, Kothekar VS. Mutagenic efficiency and effectiveness in Cluster Bean
(*Cyamopsis tetragonoloba* (L.) Taub.). J. Phytol. 2010; 2: 21–27.

62. Balai OP, Ram K. Efficiency and effectiveness of chemical mutagens in mungbean. J.
Food Legumes. 2009; 22: 105–108.

63. Kaul MLH, Bhan AK. Mutagenic effectiveness and efficiency of EMS, DES and
gamma rays in rice. Theor. Appl. Genet. 1977; 50: 241–246.

64. Laskar RA, Khan S. Mutagenic Effectiveness and Efficiency of Gamma Rays and HZ
with Phenotyping of Induced Mutations in Lentil Cultivars. Intl. Let. Nat. Sci. 2017;
64: 17–31.

65. Wani AA. Mutagenic effectiveness and efficiency of gamma rays, ethyl methane
sulphonate and their combination treatments in chickpea (*Cicer arietinum* L.). Asian
J. Plant Sci. 2009; 8: 318.

66. Shah TM, Mirza JI, Haq MA, Atta BM. Induced genetic variability in chickpea
(*Cicer arietinum* L.) II. Comparative mutagenic effectiveness and efficiency of
physical and chemical mutagens. Pak. J. Bot. 2008; 40(2): 605–613.

67. Khan MH, Tyagi SD. Studies on effectiveness and efficiency of gamma rays, EMS
and their combination in soybean (*Glycine max* (L.) Merrill). J. Plant Breed. Crop Sci.
2010; 2(3): 55–58.

68. Dhulgande GS, Dhale DA, Pachkore GL, Satpute RA. Mutagenic effectiveness and
efficiency of gamma rays and Ethyl methanesulphonate in Pea (*Pisum sativum* L.). J.
Exp. Sci. 2011; 2(3): 7–8.

69. Girija M, Dhanavel D. Mutagenic effectiveness and efficiency of gamma rays, ethyl
methane sulphonate and their combination treatment in Cowpea (*Vigna unguiculata*
L. Walp). Global J. Mol. Sci. 2009; 4: 68–75.

70. Zeerak NA. Mutagenic effectiveness and efficiency of gamma rays and ethyl methane
sulphonate in brinjal. J. Nuc. Agric. Biol. 1992; 21: 84–87.

71. Sharma A, Plaha P, Rathour R, Katoch V, Singh Y, Khalsa GS. Induced mutagenesis
for improvement of garden pea. Intl. J. Vegetable Sci. 2010; 16: 60–72.

72. Gautam AS, Sood KC, Richaria AK. Mutagenic effectiveness and efficiency of
gamma rays, ethylmethane sulphonate and their synergistic effects in black gram
(*Vigna mungo* L.). Cytologia. 1992; 57: 85–89.

73. Ganapathy S, Nirmalakumari A, Senthil N, Souframanien J, Raveendran TS. Isolation of macromutations and mutagenic effectiveness and efficiency in little millet varieties. World J. Agric. Sci. 2008; 4: 483–486.

74. Makeen K, Babu GS. Mutagenic effectiveness and efficiency of gamma rays, sodium azide and their synergistic effects in urd bean (*Vigna mungo* L.). World J. Agric. Sci.2010; 6 (2): 234–237.

75. Toker C. A note on the evolution of kabuli chickpeas as shown by induced mutations in (*Cicer reticulatum* Ladizinsky.) Genet. Resources Crop Evol. 2009; 56: 7–12.

76. Gottschalk W. The genetic basis of variation. In: Improving Vegetatively Propagated Crops, Academic Press Limited, London, UK; 1987. pp. 317–334.

77. Wani MR, Khan S, Kozgar MI, Goyal S. Induction of morphological mutants in mungbean (*Vigna radiata* (L.) Wilczek) through chemical mutagens. Nucleus. 2011; 48: 243–247

78. Nilan RA. Nature of induced mutations in higher plants. Induced mutations and their utilization. In: Proc Symp Erawim Baur Gedachtnis Orle Singen IV 1966. Acverlag, Berlin; 1967. pp 5–20

79. Khursheed S, Raina A, Parveen K, Khan S. Induced phenotypic diversity in the mutagenized populations of faba bean using physical and chemical mutagenesis. J. Saudi Society Agric. Sci. 2019; 18: 113–119.

80. Khan MR, Qurshi AS, Hussain SA, Ibrahim M. Spectrum of morphological mutations induced by separate and simultaneous application of gamma rays with GA3 in chickpea (*Cicer arietinum* L.). Pak. J. Biol. Sci. 2000; 3:1431–5.

81. Laskar RA, Khan S, Khursheed S, Raina A, Amin R. Quantitative analysis of induced phenotypic diversity in chickpea using physical and chemical mutagenesis. J. Agron. 2015; 14 (3): 102–111.

82. Solanki IS, Sharma B. Induction and isolation of morphological mutations in different damage groups in lentil (*Lens culinaris* Medik). Indian J. Genet. 1999; 59(4): 479–485.

83. Yankulov MT, Isasi EM, Abreu F. Some aspects of the sensitivity and mutability of two French bean varieties under the influence of gamma radiation from 60Co and ethyl methanesulphonate (EMS). Ciencias de la Agricultura. 1980; 7: 59–64.

84. Sethi GS. Long-penduncled mutant: a new mutant type induced in barley. Euphytica, 1974; 23: 237–239.

85. Shahwar D, Ansari MY, Choudhary S. Induction of phenotypic diversity in mutagenized population of lentil (Lens culinaris Medik) by using heavy metal. Heliyon. 2019; 5(5):e01722.

86. Sjodin J. Induced morphological variations in *Vicia faba* L. Hereditas. 1971; 67: 155–180.

87. Reddy VRK, Gupta PK. Induced mutations in hexaploid triticale. Frequency and spectrum of morphological mutations. Genet. Agric. 1988; 42: 241–254.

88. Shakoor A, Sadiq MS, Hasan MU, Saleem M. Selection for useful semi dwarf mutants through induced mutation in bread wheat. Proceed. In: 5th Wheat Genetics Symposium, New Delhi; 1978. pp. 540–546.

89. Qin R, Qiu Y, Cheng Z, Shan X, Guo X, Zhai H, Wan J. Genetic analysis of a novel dominant rice dwarf mutant 986083D. Euphytica. 2008; 160: 379–387.

90. Singh RK. Gamma ray induced bold seeded mutant in *Vigna mungo* (L.) Hepper. Indian J. Genet. 1996; 56: 104–108.

91. Gun-kel JE, Sparrow AH. Ionizing radiations: biochemical, physiological and morphological aspects of their effect on plants. In: Ruhland W. editor. Encyclopedia of Plant Physiology. (ed.), Springer, Berlin; 1961. pp. 555–611.

92. Sharma SK, Sharma B. Pattern of induced mutability in different genotypes of lentil (*Lens culinaris* Medik.). Zeitscrift fur Pflanzenzuchttung. 1979; 83: 315–320

93. Solanki IS, Sharma B. Morphological mutations induced by gamma rays, ethylene amine and N-Niroso-N-Ethyl urea in Lentil. J. Genet. 2003; 53(2): 168–173.

94. Dhole VJ, Maheswari JJ, Patil S. Studies on mutations induced by EMS in Soybean (*Glycine max* L. Merrill.). Agric. Sci. Digest. 2003; 23(3): 226–228.

95. Davies KM, Schwinn KE. Flower colour. In: Geneve RL., Preece JE., Merkle SA. editors. Biotechnology of ornamental plants. CAB International; 1997. pp. 259–294

96. Mol J, Grotewold E, Koes R. How genes paint flowers and seeds. Trends Plant Sci. 1998; 3: 212–217.

97. Kolar FR, Ghatge SR, Nimbalkar NS, Dixit GB. Mutational changes in *Delphinium malabaricum* (Huth.) Munz.: A potential ornamental. Plant J. Hort. Res. 2015; 23: 5–15.

Acknowledgements

The authors are thankful to the Department of Botany, Aligarh Muslim University, Aligarh,
India for providing research facilities for providing research facilities and seeds of two
varieties of cowpea. We are also thankful to National Botanical Research Institute, Lucknow,
India for providing the gamma irradiation facility. The authors received no specific funds for
this work.

Author Contributions

Conceptualization: Aamir Raina, Samiullah Khan.

Data curation: Aamir Raina.

Formal analysis: Aamir Raina.

Investigation: Aamir Raina.

Methodology: Aamir Raina.

Project administration: Aamir Raina.

Resources: Aamir Raina.

Software: Aamir Raina.

Supervision: Samiullah Khan.

Validation: Aamir Raina.

Visualization: Aamir Raina.

Writing - original draft: Aamir Raina.

Writing - review & editing: Aamir Raina.

Competing interests

The authors declare no competing interests.

Figure

Figure

Gamma rays

SA

Gamma rays + SA

Figure

Gomati VU-89

Pusa-578

■ Plant Size ■ Growth habit ■ Leaf ■ Flower ■ Pod ■ Seed

Figure

Figure

Figure

Figure

Figure

Figure

Figure