Dynamic dot displays reveal material motion network in the human brain

Authors:

Alexandra C. Schmid [1.3]

Huseyin Boyaci [1,2]

Katja Doerschner [1,2]

[1] Department of Psychology, Justus Liebig University Giessen, Giessen 35394, Germany

[2] Department of Psychology, A.S. Brain Research Center, and National Magnetic Resonance

Research Center, Bilkent University, Ankara 06800, Turkey

[3] Present address: Laboratory of Brain and Cognition, National Institute of Mental Health,

National Institutes of Health, Bethesda, MD 20892, USA

Correspondence:

Alexandra.Schmid@psychol.uni-giessen.de (A.C.S.);

hboyaci@bilkent.edu.tr (H.B.);

Katja.Doerschner@psychol.uni-giessen.de (K.D.)

ABSTRACT

There is growing research interest in the neural mechanisms underlying the recognition of

material categories and properties. This research field, however, is relatively more recent and

limited compared to investigations of the neural mechanisms underlying object and scene

category recognition. Motion is particularly important for the perception of non-rigid materials,

but the neural basis of non-rigid material motion remains unexplored. Using fMRI, we

investigated which brain regions respond preferentially to material motion versus other types

of motion. We introduce a new database of stimuli – dynamic dot materials – that are

animations of moving dots that induce vivid percepts of various materials in motion, e.g.

flapping cloth, liquid waves, wobbling jelly. Control stimuli were scrambled versions of these

same animations and rigid three-dimensional rotating dots. Results showed that isolating

material motion properties with dynamic dots (in contrast with other kinds of motion) activates

a network of cortical regions in both ventral and dorsal visual pathways, including areas

normally associated with the processing of surface properties and shape, and extending to

somatosensory and premotor cortices. We suggest that such a widespread preference for

material motion is due to strong associations between stimulus properties. For example

viewing dots moving in a specific pattern not only elicits percepts of material motion; one

perceives a flexible, non-rigid shape, identifies the object as a cloth flapping in the wind, infers

the object's weight under gravity, and anticipates how it would feel to reach out and touch the

material. These results are a first important step in mapping out the cortical architecture and

dynamics in material-related motion processing.

Keywords: material perception, motion, fMRI, point-light motion, dynamic dot, structure from

motion

1. INTRODUCTION

Recognizing and estimating the material qualities of objects is an essential part of our visual

experience. Perceiving material qualities quickly and correctly is critical for guiding decisions

and actions, whether we are deciding what fruit to eat, if a blanket is soft enough, or how we

should grip a porcelain cup. Despite the importance of recognizing and estimating the

properties of materials, it is still not well understood how the brain accomplishes these tasks

and, compared to object and scene perception, neuroscientific studies investigating material

perception are relatively recent (see Komatsu & Goda, 2018 for a review).

Literature investigating the neural mechanisms of visual material perception has so far heavily

focused on how the brain discriminates between different surface optical appearances; that is,

the cortical areas involved in the visual processing of the surface properties of materials

(Komatsu & Goda, 2018). These properties, such as a surface's micro- and meso-structure, or its

reflective-, transmissive- and refractive properties, give a material its characteristic visual

appearance (e.g. glossy, plastic-y, metallic, etc.). Nearly all these studies have used static

images as stimuli (but see Okazawa et al. 2012; Kam et al. 2015; see Sun et al. 2016a for

binocular stimuli).

The use of static images when investigating the visual processing of materials is somewhat

justified because, although material perception is a multisensory (Baumgartner et al. 2013),

dynamic experience (Schmid & Doerschner, 2019), many material qualities can be conveyed

through images alone (Paulun et al. 2017; Schmid & Doerschner, 2018; Schmidt et al. 2017; van

Assen & Fleming 2016, Baumgartner et al. 2013, Xiao et al. 2016). But not all material

properties can be faithfully conveyed through static images, and most mechanical material

properties become much more vividly apparent with motion: watching a rubber band

stretching, a jelly jittering, and hair bending elicits strong impressions of elasticity, wobbliness,

and softness, respectively. In fact, image motion has been shown to provide information about

material qualities over and above the information available from 2D images (Doerschner et al.

2011, Schmid & Doerschner 2018; Schmidt et al. 2017). Furthermore, recognizing materials in

natural environments likely entails integrating spatiotemporally segregated information into

coherent percepts, similar to detecting animate objects (think of detecting a tiger through long,

swaying grass), making it a special kind of event perception (Johansson, 1974; Jannson,

Bergström & Epstein, 1994; Radvansky & Zacks, 2011; Bingham, Schmidt & Rosenblum, 1995).

Given that many brain regions are sensitive to certain kinds of motion structure (for reviews see

Erlikhman et al., 2018; Kourtzi et al., 2008; Nishida et al., 2018), it is surprising that, to date, no

studies have examined the cortical processing of non-rigid material motion. One possible

reason for this is that it is difficult to find adequate control conditions for complex, dynamic

stimuli. Recent improvements in computing power now puts us in a position to be able to

simulate complex material motion stimuli in a controlled way and generate adequate control

conditions (described below).

The first step to investigate the neural correlates of non-rigid material motion is to find whether

cortical areas exist that show a preference for non-rigid material motion over other types of

motion. In this study, we developed a new class of stimuli that utilizes the fact that mechanical

and tactile qualities of materials can be convincingly conveyed through image motion alone

(Schmid & Doerschner, 2018; Bi et al., 2019). Previously these have been called point-light

stimuli by Schmid & Doerschner (2018), analogous to "point light walkers" that have been used

extensively in biological motion research (Johansson, 1973). Similar stimuli have also been

called "dynamic dot stimuli" (Bi et al., 2019). Here, we name our new movie database "**dynamic

dot materials**", where specific non-rigid material types are solely depicted through the motion

of black dots on a gray background. Dynamic dot materials not only depict mechanical material

properties convincingly; they also isolate dynamic properties from optical (surface) cues and

can provide more stimulus control compared to "full cue" videos that contain surface

information. This is because the behavior of trajectories and speed of individual dots can be

manipulated.

Using fMRI, we investigated which brain regions respond preferentially to dynamic dot

materials versus motion-scrambled and structure-from-motion control stimuli. Comparing

material motion to these two control conditions allowed us to identify regions that responded

to material motion over and above just any motion (scrambled motion condition), and over and

above general structure from motion, motion coherence, and bounded "object-ness"

(structure-from-motion control stimuli). Anticipating our results, we find widespread

preferential activation for non-rigid material motion versus the control conditions. This

suggests that dynamic dot materials are very suitable for mapping the cortical network involved

in the perception of non-rigid materials, and we recommend that our stimuli can be used in

future studies – for example as a localizer – to further explore the specific function of these

different regions in this "material motion network".

2. MATERIALS & METHODS

2.1. Participants

Ten volunteers participated in the experiment (age range: 21-42, 2 males, mean age: 27.3, 1 left

handed). All participants had normal or corrected to normal vision and had no known

neurological disorders. Participants gave their written consent prior to the MR imaging session.

Protocols and procedures were approved by the local ethics committee of Justus Liebig

University Giessen and in accordance with the Code of Ethics of the World Medical Association

(Deklaration of Helsinki).

2.2. Stimuli

2.2.1. Dynamic dot materials. Stimuli were non-rigid materials, generated with Blender

(version 2.7) and Matlab (release 2012a; MathWorks, Natick, MA) and presented using Matlab

and the Psychophysics Toolbox (Brainard, 1997). Each dynamic dot material was a 2s animation

(48 frames) that depicted the deformation of a specific material under force. Object

deformations were simulated using the Particles System physics engine in Blender, with either

the fluid dynamics or Molecular addon (for technical specifications we refer the reader to

Schmid & Doerschner, 2018). For variety, we simulated non-rigid materials with different

mechanical properties under various forces (details shown in Table 1), including cloth flapping

in the wind, liquids rippling and waving, breaking materials of high, medium, and low elasticity,

as well as non-breaking elastic materials. Upon creation of these different material types we

exported the 3D coordinates of each particle at each frame. Using Matlab, we calculated the 2D

projection for each particle from a specific viewing angle. For each material only 200 random

particles were selected for display, as a medium density yielded best perceptual impressions in

previous work (Schmid & Doerschner, 2018), with the remaining particles set to invisible. The

particles were sampled from throughout the volume of the substance with the following

exception: for liquids the particles were sampled from the surface because sampling from the

volume in these cases did not convey the material qualities as convincingly. In total we

rendered ten animations (Table 1). Frames from example stimuli can be seen in Figure 1a, and

the trajectory of the dots over time can be seen in Figure 2 (left column) and Supplementary

Figure 1. We also informally confirmed with participants that they perceived the dynamic dot

materials as intended. We formally recorded four participants' responses, which are shown in

Supplementary Table 1.

Material motion name	Particle physics	Blender simulation details	Part of simulation rendered
cloth	Linked, unbreaking particles	A sheet of linked particles was attached at the top of the scene and blown by a wind force field.	Cloth blowing in the wind.
Cloth_rot	Linked, unbreaking particles	Same as cloth.	Cloth movie rotated 90 degrees.
Ripple	Fluid particles	Fluid particles were dropped into a small round container, causing it to ripple.	Liquid rippling.
Waves	Fluid particles	Fluid particles were dropped into a large square container and stirred by an invisible rod causing larger waves.	Liquid waves.
pokeWobble	Linked, unbreaking particles	Elastic cube made of linked particles was attached to the ground and poked with an invisible rod.	Cube wobbling.

pokeWobble_rot	Linked, unbreaking particles	Same as pokeWobble.	Cube wobbling, rendered from a different camera angle to pokeWobble.
stretchBounce	Linked, unbreaking particles	Elastic cube made of linked particles was attached to invisible solid walls that moved horizontally apart. Elastic cube stretched and bounced.	Cube stretching and bouncing.
stretchWobble	Linked, breaking particles	Same as stretchBounce but elastic cube ripped and wobbled like hard jelly.	Cube stretching, ripping, and wobbling.
stretchHighWobble	Linked, breaking particles	Same as stretchBounce but elastic cube ripped and wobbled substantially.	Cube stretching, ripping, and wobbling.
stretchDough_rot	Linked, breaking particles	Same as stretchBounce but the cube was made of low-elastic material that ripped apart softly with no wobble motion.	Cube stretching and ripping. Movie was rotated 90 degrees to provide variation to the other three "stretch" movies.

Table 1: Simulation and rendering details for each material motion animation. *Animations*

were 2s clips of materials being deformed in various ways. The first column shows the name of

the movie provided in the Supplementary Material. The second column shows whether each

material was simulated as linked particles (breaking or unbreaking) or fluid particles (see main

text). The third column shows details of how the simulation was set up in Blender. The fourth

column shows what part of the simulation was rendered in the final animation.

2.2.2. Scrambled control stimuli. We wanted to find cortical regions that are sensitive to

material motion over and above other kinds of motion. That is, we wanted to exclude regions

that are sensitive to motion generally but do not show a preference for material motion.

Therefore, we created "scrambled" control stimuli that were matched in motion energy to the

dynamic dot materials (measured as velocity magnitude, see Supplementary Figures 2-4). We

created a scrambled version of each material stimulus by shuffling the position of each particle

on the first frame. All dots in the scrambled motion stimuli had the same velocity magnitude as

the material motion; however, the direction of trajectory was rotated by a random (uniform

distribution) amount between 0 and 2π every 2nd frame. Without this rotation the scrambled

stimuli would still look like non-rigid materials. The trajectory of a given dot was forced to

remain inside the spatial range of the dynamic dot materials (in the 1st frame) by forcing it to

change its direction if it is outside the boundary. This was accomplished by rotating the dot

trajectory to a different direction. As a consequence, acceleration between material and control

stimuli were not matched (See Supplementary Figures 5-7). Example scrambled control stimuli

are shown in Figure 1C, Figure 2 (middle column), and Supplementary Figure 1.

2.2.3. Structure from Motion (SfM) control stimuli. We also wanted to make sure any

preference for material motion within a brain region was not just a preference for global

coherent 3D motion, or "object-ness". Therefore, we created 3D structure-from-motion (SfM)

control stimuli. To generate these control stimuli, we selected one frame of each material

motion movie and then rotated the camera back and forth around the center of the scene. This

gave the object an appearance of rotating in depth around the horizontal (or vertical axis). SfM

control stimuli are shown in Figure 1C, Figure 2 (right column), and Supplementary Figure 1.

2.3. Stimulus display

Visual stimuli were presented on an MR-safe LCD screen placed near the rear end of the

scanner bore (Cambridge Research Systems Ltd, Rochester, UK; resolution: 1920 by 1080;

refresh rate 120 Hz). Participants viewed the screen through an angled mirror attached to the

head-coil while lying supine inside the scanner bore. Total eye-to-screen optical distance was

140 cm, and the screen subtended a visual angle of 28 degrees horizontally at this distance.

Stimuli were presented at the center of the screen and approximately subtended 15.75 by

15.75 degrees visual angle.

2.4. Fixation task

To ensure fixation during the scanning session, aid maintaining vigilance and wakefulness, and

limit attentional effects, we asked participants to perform a demanding fixation task

throughout the functional runs. In this task, participants were required to report a brief

reduction in the size of the fixation cross via a button press. The cross shrank by a small amount

every 3 seconds with a random (+/- up to 1.5 seconds) time jitter to make it unpredictable

when the shrinking would occur.

2.5. MR Image Acquisition

Magnetic resonance images were collected on a 3 Tesla MRI scanner (Magnetom Prisma,

Siemens AG, Erlangen, Germany) equipped with a 64-channel head coil in BION imaging center

of JLU Giessen. MR sessions contained a structural run and 4-8 functional runs. Structural

images were acquired using a T1-weighted 3-D anatomical sequence (sagittal MP-RAGE, Spatial

resolution: 1 mm^3 isotropic; number of slices: 176). Functional images were acquired while

participants viewed the visual stimuli and were acquired with a T2*-weighted gradient-recalled

echo-planar imaging (EPI) sequence (TR: 2000 ms; TE: 30 ms; spatial resolution: 3x3x3 mm^3;

number of slices: 36; slice orientation: parallel to calcarine sulcus). Each participant took part in

one scanning session that lasted about an hour and a half. During functional runs participant

responses were collected using an MR-safe button box.

2.6. Experimental Design

During the functional runs, different types of motion stimuli (material, scrambled control, SfM

control) were presented in alternating blocks. The order of blocks was randomized and

counterbalanced. Each block lasted 22 seconds and contained ten short clips (2 s) of animation

separated by 200 ms interstimulus interval (ISI). Figure 1B depicts the experimental protocol.

There were 3 repeats of each stimulus type (material, scrambled control, SfM control) per run,

4-8 runs in a session, thus 12-24 repeats of each stimulus per participant. One run lasted 286

seconds.

Figure 1. Example stimuli and block design. *During the functional scans participants were presented with alternating blocks of dynamic dot materials, motion scrambled control stimuli, and structure from motion (SfM) control stimuli. On the right side of panel **A** we show selected frames of 4 of the 10 possible dynamic dot material animations that were shown in random order in a block. On the left side of the same panel the timing of presentation during a dynamic dot material block is shown. Panel **B** depicts an example run, and panel **C** shows selected frames*

of 1 of the 10 possible random motion control animations (left) and the SfM control stimuli

(right). The timing during a block of these control conditions was identical to that of dynamic

dot material blocks.

Figure 2. Dot Trajectories. *Shown are part of the trajectories for half of the dots for three*

sample dynamic dot stimuli across all three conditions: material motion, velocity-matched

scrambled motion control, and structure from motion (SfM) control. Each line shows the path a

given dot traveled between frames 7 and 19. This also corresponds to the number of frames

between reversals in the SfM condition. Trajectories for the remaining seven stimuli are shown

in Supplementary Figure 1.

2.7. MR Data Analysis

All MR image preprocessing and further analyses were performed using BrainVoyager QX,

except an initial inhomogeneity correction step on T1-weighted images, which was conducted

using Freesurfer version 4 software (http://surfer.nmr.mgh.harvard.edu/). After the initial

inhomogeneity correction with Freesurfer4, anatomical images were imported into

BrainVoyager for further preprocessing. Preprocessing for the anatomical images included the

following steps: another inhomogeneity correction using BrainVoyager (for 8 out 10

participants this led to a better white-gray matter segmentation), aligning the images in AC-PC

plane and converting to Talairach space, white-gray matter segmentation. After these steps a

3D cortical mesh was created for each subject and the resulting individual meshes were

morphed and aligned using cortical surface information (sulci and gyri). Finally, an average

mesh was created and inflated. Preprocessing steps on functional images included slice

acquisition time correction, motion correction, linear trend removal and high pass filter

(temporal). The resulting functional images were coregistered with the anatomical images per

participant. Functional data were spatially transformed and projected on the inflated 3D

average mesh for further analyses. These analyses included whole brain random-effects

surface-based group analyses using the general linear model (GLM), and region of interest (ROI)

analyses.

For the whole brain analyses, we performed several separate GLM analyses (material vs.

scrambled, material vs. SfM, SfM vs. scrambled). Active voxels were identified at $p < 0.05$ level

(using BrainVoyager's cluster correction for multiple comparisons). For the ROI analyses, we

used half the data (odd runs) to define ROIs, and half the data (even runs) to extract volume

time courses for each participant. ROIs were identified using fixed-effects surface-based group

GLM analyses (material vs. scrambled, material vs. SfM; active voxels were identified at $p < 0.05$

level). For each participant and ROI, average percent signal change (from fixation baseline,

using BOLD signal and middle 5 TRs from each block) was calculated for each condition

(material, SfM, scrambled), and these data were subjected to a 3 (condition: material, SfM,

scrambled) X 2 (hemisphere: RH, LH) repeated-measures ANOVA. Significant ANOVA results

were followed up with paired t-tests using Sidak-corrected p-values for multiple comparisons

(single pooled variance). We also performed sensitivity and effect size analyses for each ROI

and hemisphere, which are reported in Supplementary Table 2. Sensitivity analyses were

performed using G*Power (version 3.1.9.7; Faul et al., 2007), where we computed required

effect size given α (0.05), Power (1-β err prob = 0.95), and sample size (n=10) for a repeated

measures within-subjects ANOVA. Effect sizes were calculated as Cohen's f (Cohen, 1988).

2.8. Data/code availability statement

The stimuli used in this study come from the "Dynamic dot materials database". We provide

access to this database, which contains each stimulus in the form of movies, individual movie

frames, and the coordinates of each point for each frame. We also provide MR volume time

course data separately for each subject, brain region, and measurement. These can be accessed

through the following link:

https://www.dropbox.com/sh/nakqpn022lpaptn/AACQkbsgFuVvUIZlRXnoWXrya?dl=0).

3. RESULTS

3.1 Whole Brain GLM Results

Figure 3 shows the results of the whole-brain GLM analyses. The activation map in Figure 3A

shows that, overall, there are many cortical areas that have a larger BOLD response to dynamic

dot materials than to scrambled motion stimuli. This widespread network of brain areas (hot

colours in Figure 3A) includes dorsal and ventral visual regions in addition to multisensory,

somatosensory, and premotor areas. Only early visual areas responded more strongly to the

scrambled motion (cool colours in Figure 3A). This is in line with our motion energy

measurements of the stimuli (Supplementary Figures 2-7) and suggests that higher activation

for material versus control motion in other regions is not due to low-level differences in motion

energy between the conditions (See Discussion, section 4.5: Motion energy and motion

coherence).

A black outline of the activation map from Figure 3A is superimposed over the activation maps

in Figure 3C and D. Figure 3C shows that cortical areas with a material "preference" (black

outlines) appear to constitute a superset of those that respond to 3D rigid motion (SfM). The

overlap in activation for these two contrasts (material vs. scrambled and SfM vs. scrambled) is

interesting and suggests that 3D rigid motion could simply be a special type of material motion

and thus activity to this type of stimuli should be contained within the general material motion

network. If rigid motion is a subclass of all possible material motions then this might also

explain why cortical responses to dynamic dot materials are almost always stronger than those

to 3D rigid motion (Figure 3D): seeing just one type of material motion is likely to cause a

relatively weaker cortical response than the rich set of motions that occur in the dynamic dot

condition.

Broadly, the stronger response for material motion versus scrambled and 3D rigid motion (SfM)

overlap with the results of our ROI analysis (next section). This "material motion network"

encompasses ventral visual regions (light green regions labelled 3 in Figure 3B); occipito-

temporal and -parietal regions (yellow regions labelled 4), dorsal visual and

somatosensory/multisensory regions (though not primary somatosensory cortex; orange

regions labelled 5); pre-motor regions (light purple regions labelled 6); and superior temporal

regions (blue region labelled 7).

The average accuracy in the fixation task was 83% across observers, suggesting that participants

followed our instructions and fixated at the center of the screen during presentation of the

stimuli.

Figure 3. GLM group analyses. For all analyses, active voxels were identified at p < 0.05 level

(cluster corrected; see section 2.7: MR Data Analyses). ***A.*** *Results of the GLM contrasting*

average BOLD responses to dynamic dot materials with those to scrambled motion stimuli. Only

early visual areas respond more weakly to material motion (cool colours). A large network of

areas is more strongly active under the material motion condition (hot colours). The results of

this contrast (material vs. scrambled) are overlaid as a black outline in (C) and (D). **B.** *ROIs*

identified for subsequent analyses (see section 3.2: ROI Results). The ROIs largely overlap with

activation maps in (A), (C), and (D). Broadly the "material motion network" encompasses ventral

visual regions (3. Light green); occipito-temporal and -parietal regions (4. Yellow); dorsal visual

and somatosensory/multisensory regions (5. Orange); pre-motor regions (6. Light purple); and

superior temporal regions (7. Blue). **C.** *Results of a GLM analysis that contrasts average BOLD*

responses to 3D rigid motion (SfM) with those to scrambled stimuli. The resulting activity maps

suggest that cortical areas that respond strongly to material motion (black outline) are a

superset of those that respond to 3D rigid motion (hot-coloured activation maps). **D.** *The results*

of the same dynamic dot materials vs. scrambled motion contrast (black outline) together with a

GLM contrast of dynamic dot materials versus 3D rigid motion (SfM; hot-coloured activation

maps). Here we see that responses to dynamic dot materials were almost always stronger than

those to 3D rigid motion (SfM) stimuli. See section 3.1: Whole Brain GLM Results for further

details. Overall, lower visual areas tended to respond stronger to the scrambled motion stimuli.

This is consistent with the literature (e.g. Murray et al. 2002).

3.2 ROI Results

To evaluate the reliability of the results found with the whole-brain analyses, we also

conducted a region of interest (ROI) analysis. Six ROIs (early visual cortex (EVC), cingulate,

ventral, occipital, somatosensory, and premotor) were defined with the contrast material minus

scrambled, and one ROI (superior temporal sulcus; STS) was defined with the contrast material

minus SfM (active voxels were identified at $p < 0.05$ level). Data used to define ROIs (shown in

Figure 3B) were separate to the data extracted from the ROIs for analysis (see section 2.7: MR

Data Analyses). Figure 4 plots the mean percent signal change from baseline (fixation blocks)

for each ROI, condition, and hemisphere. A 3 (condition: material, SfM, scrambled) X 2

(hemisphere: RH, LH) repeated-measures ANOVA revealed a main effect of hemisphere for only

for the premotor ROI, $F(1, 9) = 5.317$, $p=0.0466$, where on average percent signal change was

larger for the left versus the right hemisphere. There was a main effect of condition for six of

the ROIs: cingulate, $F_{(2, 18)} = 6.914$, p=0.0059; ventral, $F_{(2, 18)} = 19.03$, p<0.0001; occipital, $F_{(2, 18)} = 59.44$, p<0.0001; somatosensory, $F_{(2, 18)} = 11.43$, p=0.0006; premotor, $F_{(2, 18)} = 6.886$,

p=0.0060; and STS[1], $F_{(2, 18)} = 8.467$, p=0.0026 (starred ROIs in Figure 4). There was a significant

interaction between condition and hemisphere for two ROIs: occipital, $F_{(2, 18)} = 6.035$,

p=0.0099, and somatosensory $F_{(2, 18)} = 5.530$, p=0.0134 (double starred ROIs in Figure 4).

For these two ROIs (occipital and somatosensory), follow-up tests showed significant

differences between all conditions ($p < 0.0001$ for all tests, Sidak-adjusted p-values). That is, for

both hemispheres, there was a greater BOLD response to material motion versus both control

motions (SfM and scrambled), and a greater response for SfM versus scrambled motion (see

Figure 4, occipital and somatosensory ROIs).

For the other ROIs where there was a main effect of condition (cingulate, ventral, premotor,

and STS; single starred ROIs in Figure 4), follow-up tests revealed a larger response for material

motion versus scrambled motion ($p < 0.01$ for all tests, Sidak-adjusted). Ventral and STS ROIs

additionally showed a larger response for material motion versus SfM ($p < 0.01$ for all tests,

Sidak-adjusted). However, the cingulate and premotor ROIs did not show a difference between

these two conditions (p>0.05, Sidak-adjusted). The cingulate ROI, however, did show a greater

response to SfM versus scrambled motion, suggesting that this region does not prefer material

motion per se, but rather coherent motion structure.

For EVC there were no significant main effects of condition, $F_{(2, 18)} = 1.090$, p=0.3574, or

hemisphere, $F_{(1, 9)} = 0.3470$, p=0.5703, nor a significant interaction between condition and

hemisphere, $F_{(2, 18)} = 1.110$, p=0.3511.

---

[1]   Since the STS ROI was only present in the right hemisphere, a one-way ANOVA was performed.

More details about the follow-up tests and effect sizes for each contrast (in the form of

confidence intervals) are presented in Supplementary Figure 8. We also show that the effects

reported here are extremely consistent across participants in Supplementary Figures 9 and 10.

Figure 4. ROI analysis. *Each graph shows percent signal change for each hemisphere (left*

hemisphere, LH; right hemisphere; RH), and each condition (Mat = material motion, SfM = shape

from motion control; Scram = scrambled motion control). Starred ROI labels indicate a

significant main effect of condition. Double starred labels indicate a significant interaction

between condition and hemisphere. Starred contrasts indicate a significant difference between

those conditions from a follow-up test (p < 0.05 level, Sidak-adjusted). See section 3.2: ROI

Results for details. Error bars are one standard error of the mean and show variability between

observers (see also Supplementary Figure 8 for confidence intervals associated with each

contrast, and Supplementary Figures 9 and 10 for plots of each subject's data separately).

4. DISCUSSION

Although motion compellingly conveys the properties of non-rigid materials such as stiffness and elasticity (e.g. Schmid & Doerschner 2018; Schmidt et al. 2017), until now the neural basis of non-rigid material motion remained unexplored. Here we investigated whether specialized dynamic dot animations portraying non-rigid material motion elicit meaningful differences in brain responses when compared to other kinds of motion using functional magnetic resonance imaging (fMRI). We found robust and widespread increased activation across the human brain in response to dynamic dot materials when compared to activation in response to other types of motion stimuli (Figures 3 and 4).

4.1 General activation maps in response to dynamic dot materials

Surprisingly, regions preferring dynamic dot materials included several areas in occipito-temporal and, -parietal cortices, secondary somatosensory cortex, and premotor regions. Ventral visual areas (labelled 3 in Figure 3b; light green) have previously been implicated in the processing of surfaces and textures in static scenes (Cant & Goodale, 2007; Cavina-Pratesi et al., 2010a, 2010b; Hiramatsu et al., 2011). Occipito-temporal and -parietal areas (labelled 4; yellow) include motion-, object-, face-, and place-selective areas (Grill-Spector & Mallach, 2004). Posterior parietal areas are sensitive to aspects of 3D shape, structure-from-motion, optic flow, multisensory information, and visuomotor control (Culham, et al., 2006; Erlikhman et al., 2018; Uesaki & Ashida, 2015). Recently it has been found that activity patterns in secondary somatosensory cortex (labelled 5; orange) could reliably discriminate visual properties, such as surface gloss and roughness (Sun et al., 2016b). Greater responses to biological versus scrambled point-light stimuli have been found in superior temporal areas (labelled 7; blue) and premotor areas (labelled 6; light purple) (Saygin et al., 2004). Note that the higher responses for material versus the two control motions does *not* include primary somatosensory nor primary motor cortices, but does encompass secondary somatosensory, multisensory, and premotor areas, in addition to nearly all extrastriate regions that are responsive to visual stimuli.

Discovering a network of preferentially more active areas during dynamic dot material viewing

does not mean that all these areas must be involved specifically in the recognition and

differentiation of materials: to find such finer-tuned responses would require further studies,

and dynamic dot stimuli provide a convenient way to investigate this, as discussed next.

4.2 Neural mechanisms of material perception

Traditionally, material perception has been portrayed as a hierarchical processes that proceeds

from the processing of image features to the perception of stimulus properties to the

recognition of material identity or category (e.g. Komatsu & Goda, 2018). For example,

neuroimaging studies typically investigate how particular image features (e.g. form, motion, or

texture cues) might encode particular stimulus properties (e.g. 3D shape or gloss; Georgieva et

al., 2008; Kam et al., 2015). A consequence of this is that stimulus properties related to material

recognition are often investigated separately to one another. This separate focus on different

visual properties (3D shape, texture, gloss, etc.) reflects our perceptual experience of layered

visual representations and is consistent with some neuroimaging data. When we look at a

scene, we interpret retinal image structure as distinct perceptual dimensions like 3D shape,

surface material (gloss, transparency), colour, shading, motion, etc. (Anderson, 2011). In line

with this, fMRI studies have demonstrated that different brain regions show a preference (in

terms of stronger BOLD response) when attention is directed to one visual dimension (material,

shape, colour, motion) over others (e.g. Cant & Goodale, 2007; Cant & Xu, 2012, 2015; 2016;

Cant, Arnott, & Goodale, 2009; Caviana-Pratesi et al., 2010; Peuskens et al. 2004). This is

interpreted as preferential processing of different stimulus properties in different brain regions

and mirrors our perceptual experience of layered representations. If different stimulus

properties are processed separately before being later integrated in a hierarchical manner

(Komatsu & Goda, 2018), then to understand our multifaceted visual experience it would make

sense to first study the component perceptual dimensions.

Our results challenge this view, however, as we find a preference for materials defined purely

by motion properties in many areas implicated in the processing of surface material properties.

For example, the ventral activity we find overlaps with posterior fusiform gyrus (pFs), fusiform

gyrus (FG), collateral sulcus (CoS), lingual gyrus (LG), which have been implicated in the

processing of gloss (pFs), texture (CoS, LG, FG), and material categories (FG, CoS; see Komatsu &

Goda, 2018 for a review). This might suggest that these regions are not specialized for

processing the image properties that depict surface properties per se, or, alternatively, that

these regions are activated due to associations between material motion and surface

properties (Schmid & Doerschner, 2019). We expand on this idea in section 4.3: Encoding

directly perceived versus associated properties.

Our results are in line with recent neuroimaging studies that have found that regions previously

associated with shape processing are also sensitive to changes in material properties (e.g.

V3b/KO; Kam et al. (2015), Sun et al. (2015, 2016a); LOC, Hiramatsu et al. (2011); V4, Kim et al.

(2019), as these regions overlap with occipital activation in the present study. Furthermore, our

results are consistent with neuroimaging data that show crossmodal associations between

visual and tactile properties. For example, visual discrimination of rough and smooth surfaces

can be decoded in somatosensory cortex (Sun et al., 2016b).

The widespread activation that we find in response to quite sparse stimuli suggests that

material perception is an inherently distributed process (see Schmid & Doerschner, 2019).

Consistent with this idea, there is a growing body of literature suggesting that object category

representations are grounded in distributed networks (e.g. Kravitz et al., 2011; 2013; Martin,

2016). Under this framework, a widespread cortical preference for dynamic dot materials is

perhaps expected, given the properties of these stimuli: the stimuli are objects (LOC, e.g. Grill-

Spector & Malach, 2004), they are non-rigidly moving structures (e.g. hMT+, MST, PPC, e.g. see

review by Erlikhman, et al., 2018; STS, e.g. Saygin et al., 2004), they elicit a distinct tactile

experience (e.g. Schmid & Doerschner, 2018; Bi et al., 2019), and such tactile experiences of

material qualities are often associated with certain optical material qualities (CoS, e.g. Arnott et

al., 2008; Podrebarac et al., 2014; Sun et al., 2016b; and see Komatsu & Goda, 2018 for a

review).

Future work using dynamic dot materials could help reveal the underlying computations carried

out by each region. For example, they could be used to investigate whether areas previously

associated with materials (e.g. CoS, e.g. Cant & Goodale 2007; Cant & Xu, 2012, 2015, 2016;

Gallivan et al. 2014; Eck et al., 2016; Kitada et al. 2014) are indeed specialized for processing

visual (optical) properties or whether they represent material properties more generally. Our

results so far suggest the latter, but further investigation, for example using a multivariate

design and analysis approach, would allow one to better test this (Schmid & Doerschner, 2019).

We expand on this below.

4.3 Encoding directly perceived versus associated properties

Materials are inherently multidimensional in that they have multiple stimulus properties that

are portrayed through multiple perceptual dimensions (e.g. motion or surface optical cues), and

they are inherently multimodal in that their properties can be inferred through multiple

modalities (e.g. vision or touch). Over time, specific visual (or auditory, proprioceptive, or

olfactory) information becomes associated with specific tactile information and vice versa. It is

possible that mechanical and tactile material qualities like softness, viscosity, and roughness

can be conveyed through the surface properties, 3D structure, and motion of visual objects

(e.g. Baumgartner et al. 2013; Fleming, 2014, 2017; Fleming et al. 2013; Giesel & Zaidi, 2013; Ho

et al. 2006; van Assen & Fleming 2016; Xiao et al. 2016). For example, a velvet cloth looks soft

(optical properties), moves (visual mechanical motion) and folds (visual 3D shape) in a way that

suggests that it is soft, but it also feels soft to the touch (tactile). This multidimensionality and

multimodality make materials the ideal candidate to develop experimental designs that can

help to understand computational architecture of the cortical representations involved in

recognition. As an example, if a cortical region represents material/object category A based on

visual property X but not the same category based on the visual property Y then this suggests

that this region encodes the visual property X but not the category. Conversely, if this cortical

region has a shared representational structure, i.e. category A is encoded through both visual

properties X and Y, then it likely encodes the concept. Thus, by investigating neural responses

to stimuli like the dynamic dot materials proposed here, in conjunction with stimuli defined

along other visual dimensions, we may be able to tease apart the direct encoding of visual

properties from the indirect activation of associated properties (Schmid & Doerschner, 2019).

Note that we are referring to a generic kind of association between properties; we did not find

a preference for material motion in cortical regions involved in memory or contextual

associations, such as the angular gyrus, medial parietal cortex, or anterior parahippocampal

cortex (Kravitz et al., 2013; Bar et al., 2008).

4.4 Non-rigid motion and event perception

Dynamic dot materials convey the non-optical properties of material qualities purely based on

2D image motion patterns. Here we introduced various types of non-rigid materials, yet also

rigid materials, like breakable substances, can be rendered convincingly by the means of this

technique (e.g. see Schmid & Doerschner, 2018). In their creation, these stimuli are

conceptually closely related to "point light walkers" (Johannson, 1973), where the motion of

small light sources affixed to different parts of limbs of biological species can elicit a very vivid

impression of animacy. Bingham et al. (1995) generated point-light-like displays with

retroflecting materials to capture the motion structure of animate and inanimate events,

including the splashing of water or blowing of leaves in the wind. Similarly, we "attached" small

dots to an otherwise invisible substance and recorded the motion of these dots while the

material reacted to a force. How individual materials change their shape in response to a force

strongly depends on their mechanical properties, and it is this idiosyncratic change of shape

information over time that appears to convey the mechanical qualities of a material. It would

be very interesting to pin-point the specific motion characteristics that elicit a particular

material quality (Schmid & Doerschner, 2018; also see Bi et al. 2019), just as it has been done in

the field of biological motion where researchers have tried to understand what it is that makes

point-light walkers look "biological" (Chang & Troje, 2009; or Troje, 2013 for a review). The fact

that the location, direction, and speed of individual dots in our stimuli can be manipulated

renders investigations of research questions like these more feasible.

538 Such an interest intersects with general event perception, which aims to understand how we

539 parse the continuous flow of visuo-spatial information into segments. Gunnar Johansson was a

540 pioneer of this research and proposed a general framework of vector analysis, which relies on

541 the decomposition of the spatiotemporal information into common and relative motion

542 components (see Jannson et al., 1994 for an overview of Johanssons work). It is not

543 immediately obvious what might constitute common and relative motion components in our

544 stimuli since our shapes are rather compact, unlike a moving human or animal shape with

545 limbs. More research is needed to understand what type of motion events observers are

546 sensitive to (Todd, 1981). And, in pursuing this line of work, our stimuli could help to discover

547 the perceptual boundary between non-rigid animate and inanimate objects, as well as the

548 corresponding neural maps and mechanisms (Long, Yu, Konkle, 2018; Grill-Spector & Weiner,

549 2014). Moreover, since our activation maps overlap with areas also implicated in biological

550 motion perception (e.g. STS, premotor; Saygin et al., 2004; Chang et al., 2018), our stimuli could

551 also be suitable for teasing apart the contribution of non-rigid motion to neural activity

552 observed while watching biological motion stimuli.

554 Just as all biological motion is also non-rigid motion, rigid motion is also a special case of the

555 general event type of non-rigid motion. Our stimuli seem to 'trigger' this wide-spread non-rigid

556 object motion network, with areas responding to rigid motion constituting a subset of this

557 general network. These results are consistent with results by Bingham, et al. (1995) who found

558 that observers can identify non-rigid types of motions as well as rigid motion events and point

559 out, that "rigid motion is but one of a variety of types of motion" (Bingham, et al., 1995, also

560 see Todd 1984).

562 Most of the recent work on event perception involves human agents or rather coarse physical

563 interactions of objects (see Radvansky & Zacks 2011, for a review). For these kinds of scenarios

564 it is quite intuitive to break down events into their subcomponents, like large- and small-scale

565 events (Zacks et al. 2001). However, it is less clear what constitutes a small-scale or large-scale

566 event in the waving motion of a cloth, or the wobbling of a jelly. This would be a very

interesting topic to pursue in future work and could potentially reveal general mechanisms of

event perception.

4.5 Motion energy and motion coherence

Our results showed that higher visual areas in the ventral and dorsal streams responded more

strongly to material motion and structure from motion (SfM) over scrambled motion, and more

strongly to SfM than scrambled motion, while only early visual areas (EVC) "preferred"

scrambled stimuli over the other two conditions. We tested whether the preference of higher

cortical areas to materials could be explained by differences in velocity of the dots in our

stimuli. If we order our stimulus conditions according to the mean velocity it would yield

material=scrambled=SfM; if we ordered according to the median velocity it would yield

SfM>material=scrambled; and if we ordered them according to velocity range it would yield

material=scrambled>SfM (Supplementary Figure 4). We can therefore rule out that differences

in velocities per se drove the cortical pattern of responses we observed. In Supplementary

Figure 7 we examine weather acceleration patterns can account for the measured cortical

activity. Ordering the stimuli according to their acceleration profiles would yield

scrambled>material>SfM. Comparing this to our fMRI measurements, we conclude that this

statistic does not correspond to the observed cortical activity patterns in higher visual regions,

though it might account for the preference seen for scrambled motion over SfM stimuli in early

visual areas. Finding good quantitative descriptors of the motion profiles for stimuli like ours is

an exciting area of research (see work in this spirit, for example in biological motion Chang &

Treue 2009).

Another interesting difference between our stimulus conditions is the perception of motion

coherence. Could differences in coherence explain the differences in activation we find

between the conditions? This is not a trivial question, as it is not clear how we should define

motion coherence: the dynamic dot materials are clearly a coherent appearance – the dots are

easily grouped as belonging to the same non-rigid object – yet this coherence seems

qualitatively quite different than that of rotating rigid objects (SfM stimuli). Traditional

definitions involving the percentage of dots moving coherently, as for example in Shadlen et al.

(1996), does not apply here. We tested several possibilities to capture coherence of our stimuli

(e.g. entropy in motion direction, or divergence in optic flow), but the one that corresponds to

an intuitive ordering of our stimulus classes (SfM>material>scrambled), seems to be the shape

consistency across two consecutive frames. For this we resorted to a Procrustes analysis across

2 consecutive frames. The rationale was that if dots stay in their configuration (shape) across

two consecutive frames it should be possible to superimpose them, i.e. match the shape of

their configuration. Procrustes analysis does precisely this: it computes the linear

transformation (rotation, translation scaling) between two shapes, providing an index of the

error (sum of squared errors) that remains after applying this transform. We reasoned that this

error should be much larger for scrambled dot stimuli, compared to the other two types of

stimuli, since the configuration changes maximally between any 2 frames, and that the SfM

stimuli should yield the smallest error. Supplementary Figure 11 shows the results of this

analysis. Indeed, we can see that the coherence (as we defined it) in scrambled stimuli was

quite low for all stimuli, and coherence of the SfM stimuli tended to be high, yet not always

higher than that of the material motion stimuli. Note, that this overall pattern remains if we

increase the frame distance to 4 for computing the Procrustes analysis. Overall, Supplementary

Figure 11 shows that the difference in coherence between the material motion and SfM stimuli

is rather small. It is possible, of course, that other measures capture a different definition of

coherence (e.g. overall collinearity of motion trajectories), which we are currently investigating

in a separate project.

Interestingly, Murray et al. (2002) showed that perceiving visual elements as a coherent shape

increases activity in the lateral occipital complex, while at the same time reducing activity in

lower visual areas, with an opposite pattern found for randomly arranged visual elements.

Similarly, it is likely that the visual grouping of individually moving dots into a coherent shape is

much stronger for SfM stimuli compared to randomly moving dots, potentially contributing to

the stronger activation in early visual areas for scrambled stimuli. However, whether this

grouping differs between material motion and SfM stimuli is not obvious, and another potential

avenue to investigate in future work. Dynamic dot materials could provide a way to investigate

whether areas associated with coherent motion preference (e.g. hMT+/V5 and posterior

parietal cortex, e.g. Orban et al. 1999; Peuskens, 2004); show a preference for specific types of

coherent motion (non-rigid vs. rigid).

5. CONCLUSION

Dynamic dot materials are a novel class of stimuli that convey the non-optical properties of

materials purely based on 2D image motion patterns. Our results act as a proof of principle that

such stimuli can be used for mapping the cortical network involved in the perception of

material qualities and can act as a localizer for future studies. From a broader perspective,

dynamic dot materials constitute a novel class of stimuli that holds promise for understanding

general event perception, and, owing to their inherently multidimensional and multimodal

nature, materials in general are a unique type of stimulus that can help neuroimaging research

to advance our understanding of the computational architecture of the cortical representations

involved in recognition.

6. ACKNOWLEDGEMENTS

This work was supported by a Sofja Kovalevskaja Award endowed by the German Federal

Ministry of Education; a Marie Sklodowska-Curie Action – Innovative Training Network (MSCA-

ITN/ETN) Grant, DyViTo: Dynamics in Vision and Touch – the look and feel of stuff. We thank

Chris Baker for helpful comments on earlier versions of this manuscript.

7. REFERENCES

Arnott SR, Cant JS, Dutton GN, Goodale MA (2008) Crinkling and crumpling: an auditory fMRI study of material properties. *NeuroImage* 43:368-378.

Bingham, Geoffrey P., Richard C. Schmidt, and Lawrence D. Rosenblum. "Dynamics and the Orientation of Kinematic Forms in Visual Event Recognition." *Journal of Experimental Psychology: Human Perception and Performance* 21, no. 6 (1995): 1473–93. https://doi.org/10.1037//0096-1523.21.6.1473.

Bar M, Aminoff E, Schacter, D (2008) Scenes unseen: The parahippocampal cortex intrinsically subserves contextual associations, not scenes of places per se. The Journal of Neuroscience 28(34): 8539-8544. DOI:10.1523/JNEUROSCI.0987-08.2008

Baumgartner E, Wiebel CB, Gegenfurtner KR (2013) Visual and Haptic Representations of Material Properties. *Multisensory Research* 26: 429–55. https://doi.org/10.1163/22134808-00002429.

Bi, W, Jin, P, Nienborg, H, Xiao, B (2020) Manipulating patterns of dynamic deformation elicits the impression of cloth with varying stiffness. *Journal of Vision, 19*(5):18: 1-18. https://doi.org/10.1167/19.5.18

Cant JS, and Goodale MA (2007) Attention to Form or Surface Properties Modulates Different Regions of Human Occipitotemporal Cortex. *Cerebral Cortex* 17, no. 3: 713–31. https://doi.org/10.1093/cercor/bhk022.

Cant JS, Xu Y (2012) Object ensemble processing in human anterior- medial ventral visual cortex. *Journal of Neuroscience* 32:7685–7700.

Cant JS, Xu Y (2015) The impact of density and ratio on object- ensemble representation in human anterior-medial ventral visual cortex. *Cerebral Cortex* 25:4226–4239.

Cant JS, Xu Y (2016) The contribution of object shape and surface properties to object

ensemble representation in anterior-medial ventral visual cortex. *Journal of Cognitive*

Neuroscience 29:398–412.

Cavina-Pratesi, C Kentridge, RW, Heywood, CA, Milner, AD (2010a) Separate processing of

texture and form in the ventral stream: evidence from fMRI and visual agnosia.

Cerebral Cortex, 20: 433-446. doi:10.1093/cercor/bhp111

Cavina-Pratesi, C, Kentridge, RW, Heywood, CA, Milner, AD (2010b) Separate channels for

processing form, texture, and colour: evidence from fMRI adaptation and visual

object agnosia. *Cerebral Cortex, 20:* 2319-2332. Doi:10.1093/cercor/bhp298

Chang, DHF, Troje, N (2009) Acceleration Carries the Local Inversion Effect in Biological

Motion Perception. *Journal of Vision,* 9(1), 19–19. https://doi.org/10.1167/9.1.19.

Chang, DHF, Ban, H, Ikegaya, Y, Fujita, I, Troje, N (2018) Cortical and subcortical responses

to biological motion. *NeuroImage, 174,* 87-96.

Cohen, J. E. (1988). Statistical Power Analysis for the Behavioral Sciences. Hillsdale, NJ:

Lawrence Erlbaum Associates, Inc.

Culham, JC, Cavina-Pratesi, C, Singhal, A (2006) The role of parietal cortex in visuomotor

control: What have we learned from neuroimaging? Neuropsychologia 44, 2688-2684.

doi:10.1016/j.neuropsychologia.2005.11.003

Doerschner K, Fleming RW, Yilmaz O, Schrater PR, Hartung B, Kersten DJ (2011) Visual

Motion and the Perception of Surface Material." *Current Biology* 21, no. 23: 2010–16.

https://doi.org/10.1016/j.cub.2011.10.036

Eck J, Kaas AL, Mulders JL, Hausfeld L, Kourtzi Z, and Goebel R (2016) The Effect of Task

Instruction on Haptic Texture Processing: The Neural Underpinning of Roughness and

Spatial Density Perception. *Cerebral Cortex* 26, no. 1: 384–401.

https://doi.org/10.1093/cercor/bhu294

Erlikhman G, Gurariy G, Mruczek REB, Caplovitz GP (2016) The Neural Representation of
Objects Formed through the Spatiotemporal Integration of Visual Transients.
NeuroImage 142 (2016): 67–78. https://doi.org/10.1016/j.neuroimage.2016.03.044

Faul, F, Erdfelder, E, Lang, AG, & Buchner, A (2007). G*Power 3: A flexible statistical power
analysis program for the social, behavioral, and biomedical sciences. *Behavior*
Research Methods, 39, 175-191.

Fleming RW, Wiebel CB, Gegenfurtner KR (2013) Perceptual Qualities and Material Classes.
Journal of Vision 13, no. 8: 1–20. https://doi.org/10.1167/13.8.9

Fleming RW (2014) Visual Perception of Materials and Their Properties. *Vision Research*
94: 62–75. https://doi.org/10.1016/j.visres.2013.11.004

Fleming RW (2017) Material Perception. *Annual Review of Vision Science, 3:* 365-388.
https://doi.org/10.1146/annurev-vision-102016- 061429

Gallivan JP, Cant JS, Goodale MA, Flanagan JR (2014) Representation of Object Weight in
Human Ventral Visual Cortex. *Current Biology* 24, no. 16: 1866–73.
https://doi.org/10.1016/j.cub.2014.06.046.

Giesel M, Zaidi Q (2013) Frequency-Based Heuristics for Material Perception." *Journal of*
Vision 13, no. 14: 7–7. https://doi.org/10.1167/13.14.7.

Grill-Spector K, Weiner KS (2014) The Functional Architecture of the Ventral Temporal
Cortex and Its Role in Categorization. *Nature Reviews Neuroscience* 15, no. 8: 536–48.
https://doi.org/10.1038/nrn3747.

Grill-Spector K, Malach R (2004) The human visual cortex. *Annual Review of Neuroscience*
27: 649–77. https://doi.org/10.1146/annurev.neuro.27.070203.144220.

Ho YX, Landy MS, Maloney LT (2006) How Direction of Illumination Affects Visually

Perceived Surface Roughness. *Journal of Vision* 6, no. 5: 8.

https://doi.org/10.1167/6.5.8.

Hiramatsu, C, Goda, N, Komatsu, H (2011) Transformation from image-based to perceptual

representation of materials along the human ventral visual pathway. *NeuroImage, 57:*

482-494. Doi:10.1016/j.neuroimage.2011.04.056

Jansson, G., Bergström, S. S., & Epstein, W. (Eds.). (1994). *Resources for ecological*

psychology.Perceiving events and objects. Lawrence Erlbaum Associates, Inc.

Johansson G (1973) Visual perception of biological motion and a model for its analysis.

Perception & Psychophysics 14, 201–211. https://doi.org/10.3758/BF03212378

Johansson, G. (1974). Vector analysis in visual perception of rolling motion. *Psychological*

Research, 36(4), 311-319.

Kam TE, Mannion DJ, Lee SW, Doerschner K, Kersten DJ (2015) Human Visual Cortical

Responses to Specular and Matte Motion Flows. *Frontiers in Human Neuroscience* 9:

579. https://doi.org/10.3389/fnhum.2015.00579.

Kim T, Bair W, Pasupathy A (2019). Neural coding for shape and texture in macaque area

V4. *Journal of Neuroscience 39*, 4760-4774.

Kitada R, Sasaki AT, Okamoto Y, Kochiyama T, Sadato N (2014). Role of the precuneus in

the detection of incongruency between tactile and visual texture information: a

functional MRI study. *Neuropsychologia* 64:252-262.

Komatsu H, Goda N (2018) Neural Mechanisms of Material Perception: Quest on

Shitsukan. *Neuroscience*. https://doi.org/10.1016/J.NEUROSCIENCE.2018.09.001

Kourtzi, Z, Krekelberg, B, van Wezel, JA (2008). Linking form and motion in the primate

brain. *Trends in Cognitive Sciences, 12*(6): 230-236.

Kravitz, DJ, Saleem, KS, Baker, CI, Mishkin, M (2011) A new neural framework for

visuospatuial processing. *Nature Reviews Neuroscience, 12:* 217-230.

doi:10.1038/nrn3008

Kravitz, DJ, Saleem, KS, Baker, CI, Ungerleider, LG, Mishkin, M (2013). *Trends in Cognitive*

Sciences, 17(1): 26-49. http://dx.doi.org/10.1016/j.tics.2012.10.011

Long B, Yu CP, Konkle T (2018) Mid-Level Visual Features Underlie the High-Level

Categorical Organization of the Ventral Stream. *Proceedings of the National Academy*

of Sciences 115, no. 38: E9015–24. https://doi.org/10.1073/pnas.1719616115.

Martin, A. (2016) GRAPES—Grounding Representations in Action, Perception, and Emotion

Systems: How Object Properties and Categories Are Represented in the Human

Brain." *Psychonomic Bulletin and Review* 23, no. 4: 979–90.

https://doi.org/10.3758/s13423-015-0842-3.

Murray SO, Kersten D, Olshausen BA, Schrater P, Woods DL (2002) Shape perception

reduces activity in human primary visual cortex. *PNAS*, 99, 15164-15169.

Nishida, S, Kawabe, T, Sawayama, M, Fukiage, T (2018). Motion perception: from detection

to interpretation. *Annual Review of Vision Science, 4(20):* 1-23.

https://doi.org/10.1146/annurev-vision-091517-

Okazawa G, Goda N, Komatsu H (2012) Selective Responses to Specular Surfaces in the

Macaque Visual Cortex Revealed by FMRI." *NeuroImage* 63, no. 3 (2012): 1321–33.

https://doi.org/10.1016/j.neuroimage.2012.07.052.

Orban GA, Sunaert S, Todd JT, Van Hecke P, Marchal G (1999) Human cortical regions

involved in extracting depth from motion. *Neuron*. Dec;24(4):929-940. DOI:

10.1016/s0896-6273(00)81040-5

Paulun VC, Schmidt F, van Assen JJR, Fleming RW (2017) Shape, Motion, and Optical Cues
to Stiffness of Elastic Objects. *Journal of Vision* 17, no. 1: 20.
https://doi.org/10.1167/17.1.20

Peuskens H, Claeys KG, Todd JT, Norman JF, Van HP, Orban GA (2004) Attention to 3-D
Shape, 3-D Motion, and Texture in 3-D Structure from Motion Displays. *J.Cogn*
Neurosci. 16, no. 4: 665–82. https://doi.org/10.1162/089892904323057371

Podrebarac SK, Goodale MA, Snow JC (2014) Are visual texture- selective areas recruited
during haptic texture discrimination? *NeuroImage* 94: 129-137.

Radvansky, Gabriel A., and Jeffrey M. Zacks. "Event Perception." *Wiley Interdisciplinary*
Reviews: Cognitive Science, November 2011. https://doi.org/10.1002/wcs.133.

Saygin, AP, Wilson, SM, Hagler, DJJ, Bates, E, Sereno, MI (2004). Point-light biological
motion perception activates human premotor cortex. The Journal of Neuroscience,
24(27): 6181-6188. DOI:10.1523/JNEUROSCI.0504-04.2004

Schmid AC, Doerschner K (2018) Shatter and Splatter: The Contribution of Mechanical and
Optical Properties to the Perception of Soft and Hard Breaking Materials. *Journal of*
Vision 18, no. 1: 14. https://doi.org/10.1167/18.1.14

Schmid AC, Doerschner K (2019) Representing stuff in the human brain. *Current Opinion in*
Behavioral Sciences, 30, 178–185. https://doi.org/10.1016/j.cobeha.2019.10.007

Schmidt F, Paulun VC, van Assen JJR, Fleming RW (2017) Inferring the stiffness of
unfamiliar objects from optical, shape, and motion cues. *Journal of Vision*, 17(3):18.
doi: 10.1167/17.3.18

Servos P, Osu R, Santi A, Kawato M (2002) The Neural Substrates of Biological Motion
Perception: an fMRI Study, *Cerebral Cortex*, Volume 12, Issue 7, Pages 772–782,
https://doi.org/10.1093/cercor/12.7.772

Shadlen MN, Britten KH, Newsome WT, Movshon JA. A computational analysis of the
relationship between neuronal and behavioral responses to visual motion. *Journal of*
Neuroscience. 1996;16:1486–1510.

Sun HC, Ban H, Di Luca M, & Welchman AE (2015). fMRI evidence for areas that process
surface gloss in the human visual cortex. *Vision Research, 109*, 149-157.

Sun HC, Di Luca M, Ban H, Muryy A, Fleming RW, Welchman AE (2016a). Differential
Processing of Binocular and Monocular Gloss Cues in Human Visual Cortex. *Journal of*
Neurophysiology, http://jn.physiology.org/content/jn/115/6/2779.full.pdf

Sun, HC, Welchman, AE, Chang, DHF, Di Luca, M. (2016b). Look but don't touch: visual cues
to surface structure drive somatosensory cortex. *Neuroimage, 128:* 353-361.
http://dx.doi.org/10.1016/j.neuroimage.2015.12.054

Todd, J. T. (1981). Visual information about moving objects. *Journal of Experimental*
Psychology: Human Perception and Performance, 7(4), 795–
810. https://doi.org/10.1037/0096-1523.7.4.795

Troje N (2013) What Is Biological Motion? Definition, Stimuli, and Paradigms. In:
Rutherford, M. D. and Kuhlmeier, V. A. (eds.) Social Perception: Detection and
Interpretation of Animacy, Agency, and Intention. MIT Press. 13 - 36.
10.7551/mitpress/9780262019279.003.0002.

Uesaki, M, Ashida, H (2015) Optic-flow selective cortical sensory regions associated with
self-reported states of vection. *Frontiers in Psychology. 6*(775), 1-9. doi:
10.3389/fpsyg.2015.00775

van Assen J JR, Fleming R W (2016) Influence of optical material properties on the
perception of liquids. *Journal of Vision*, *16*(15), 12. https://doi.org/10.1167/16.15.12

Xiao B, Bi W, Jia X, Wei H, Adelson EH (2016) Can You See What You Feel? Color and

Folding Properties Affect Visual–Tactile Material Discrimination of Fabrics. *Journal of*

Vision 16, no. 3 (February 23, 2016): 34. https://doi.org/10.1167/16.3.34

Zacks, Jeffrey M., Todd S. Braver, Margaret A. Sheridan, David I. Donaldson, Abraham Z.

Snyder, John M. Ollinger, Randy L. Buckner, and Marcus E. Raichle. "Human Brain

Activity Time-Locked to Perceptual Event Boundaries." *Nature Neuroscience* 4, no. 6

(2001): 651–55. https://doi.org/10.1038/88486.

817