Article

UNRAVELLING THE PHYLOGENOMIC RELATIONSHIPS OF THE MOST DIVERSE AFRICAN PALM GENUS *Raphia* (CALAMOIDEAE, ARECACEAE)

Andrew J. Helmstetter [1,†], **Suzanne Mogue Kamga [2]**, **Kevin Bethune [3]**, **Thea Lautenschläger [4]**, **Alexander Zizka [5]**, **Christine D. Bacon [6]**, **Jan J. Wieringa [7]**, **Fred Stauffer [8]**, **Alexandre Antonelli [9]**, **Bonaventure Sonké [10]** and **Thomas L.P. Couvreur [11,*]**

1. IRD, DIADE, Univ Montpellier, Montpellier, France; andrew.j.helmstetter@gmail.com
2. Université de Yaoundé I, Ecole Normale Supérieure, Département des Sciences Biologiques, Laboratoire de Botanique systématique et d'Ecologie, B.P. 047, Yaoundé Cameroon ; mogueblue@yahoo.com
3. IRD, DIADE, Univ Montpellier, Montpellier, France; kevin.bethune@hotmail.fr
4. Department of Biology, Institute of Botany, Faculty of Science, Technische Universität Dresden, 01062 Dresden, Germany ; thea.lautenschlaeger@tu-dresden.de
5. German Center for Integrative Biodiversity Research (iDiv) Halle -Leipzig-Jena, Leipzig, Germany
6. Department of Biological and Environmental Sciences and Gothenburg Global Biodiversity Centre, University of Gothenburg, Gothenburg, Sweden; christine.bacon@bioenv.gu.se
7. Naturalis Biodiversity Center, Darwinweg 2, 2333 CR, Leiden, the Netherlands; jan.wieringa@naturalis.nl
8. Department of Botany and Plant Biology, Conservatory and Botanical Garden of the City of Geneva, University of Geneva, Geneva, Switzerland; Fred.Stauffer@ville-ge.ch
9. Department of Biological and Environmental Sciences and Gothenburg Global Biodiversity Centre, University of Gothenburg, Gothenburg, Sweden; and Royal Botanic Gardens, Kew, Richmond, UK; a.antonelli@kew.org
10. Université de Yaoundé I, Ecole Normale Supérieure, Département des Sciences Biologiques, Laboratoire de Botanique systématique et d'Ecologie, B.P. 047, Yaoundé Cameroon; bsonke_1999@yahoo.com
11. IRD, DIADE, Univ Montpellier, Montpellier, France; Thomas.Couvreur@ird.fr
* Correspondence: Thomas.Couvreur@ird.fr
† Current address: FRB-Cesab, Institut Bouisson Bertrand, 5, rue de l'Ecole de médecine - 34000 Montpellier, France

Abstract: Palms are conspicuous floristic elements across the tropics. In continental Africa, even
though there are less than 70 documented species, they are omnipresent across the tropical landscape.
The genus *Raphia* has 20 accepted species in Africa and one species endemic to the Neotropics. It is
the most economically important genus of African palms with most of its species producing food
and construction material. *Raphia* is divided into five sections based on inflorescence morphology.
Nevertheless, the taxonomy of *Raphia* is problematic with no intra-generic phylogenetic study
available. We present a phylogenetic study of the genus using a targeted exon capture approach
sequencing of 56 individuals representing 18 out of the 21 species. Our results recovered five well
supported clades within the genus. Three sections correspond to those based on inflorescence
morphology. *R. regalis* is strongly supported as sister to all other *Raphia* species and is placed
into a newly described section: Erectae. Overall, morphological based identifications agreed
well with our phylogenetic analyses, with 12 species recovered as monophyletic based on our
sampling. Species delimitation analyses recovered 17 or 23 species depending on the confidence
level used. Species delimitation is especially problematic in the Raphiate and Temulentae sections.
In addition, our clustering analysis using SNP data suggested that individual clusters matched
geographic distribution. The Neotropical species *R. taedigera* is supported as a distinct species,
rejecting the hypothesis of a recent introduction into South America. Our analyses support the

hypothesis that the *Raphia* individuals from Madagascar are potentially a distinct species different from the widely distributed *R. farinifera*. In conclusion, our results support the infra generic classification of *Raphia* based on inflorescence morphology, which is shown to be phylogenetically useful. Classification and species delimitation within sections remains problematic even with our phylogenomic approach. Certain widely distributed species could potentially contain cryptic species. More in-depth studies should be undertaken using morphometrics, increased sampling and more variable markers. Our study provides a robust phylogenomic framework that enables further investigation on the biogeographic history, morphological evolution and other eco-evolutionary aspects of this charismatic, socially and economically important palm genus.

Keywords: Africa, exons, Madagascar, rain forests, phylogenomics, *Raphia*, sequence capture

1. Introduction

Palms are iconic floristic elements across the tropics both in terms of diversity and the natural resources they provide, playing important roles for the welfare of rural and urban people at equatorial latitudes. Worldwide, there are an estimated 2500 palm species [1], mainly occurring in tropical rain forests. Africa, however, harbours less than 70 species (excluding Madagascar) [2,3], a pattern that contrasts strongly with the Neotropics or South East Asia, which contain 800 and 1200 species respectively [1,4,5]. Despite this low diversity, palms are omnipresent across the African landscape, particularly in the tropical rain forests of the continent [2,6].

Among African palms, the genus *Raphia* (subfamily Calamoideae, tribe Raphiaeae) is the most species rich, with 21 species described to date [2,7]. Of these, one, *R. taedigera*, is endemic to the Neotropics, with a disjunct distribution in Brazil and central America. The presence of this species in the Neotropics was suggested as either pre-Colombian and natural (biogeographic long distance dispersal/vicariance [8,9]) or as recently naturalized by Africans during the slave trade some 400 years ago [6,10,11]. *Raphia* species mainly occur in tropical rain forests, most often in swampy or periodically inundated areas where they can dominate the vegetation, producing dense monospecific stands (known as "Raphiales" in French). A few species, however, have adapted to drier conditions restricted to river systems in the Sahel or southern Africa.

Raphia is one of the most economically important genus of African palms across tropical African communities. One recent study documented over 100 different uses across the genus, with the most important ones being extraction of palm wine, grubs and construction material [12,13]. Exploitation of its species in the wild also represents an important source of income for populations across tropical Africa, especially for low-income households [12,14,15]. In addition, *Raphia* species play vital ecological roles in wet land ecosystems [16] where they dominate the landscape, such as in peatlands of the Congo Basin where they are highly abundant [17]. *Raphia* dominated swamps are also important ecosystems for the protection for critically endangered animals, such as lowland gorillas, because such areas are hard to access or bring into cultivation (e.g. [18]).

Raphia species are massive palms with very long pinnate leaves. One species holds the record for the longest measured leaf in angiosperms, reaching up to 26 meters (*R. regalis*). The stipe is generally above-ground and is solitary or clustered, while three species have very short (*R. palma-pinus* and *R. vinifera*) or subterranean (acaulescent) (*R. regalis*) stipes. When present, the stipe can be covered by old leaf sheath remains or a dense network of fibres (decomposed leaf sheaths), which can be curly or straight, an important character to identify species (e.g. [19,20]). *Raphia* species are monoecious, with male and female flowers on the same individual and are hapaxanthic, meaning that individual stipes die after a single flowering event [1]. The inflorescences structure is relatively simple and branched to two orders [1]. The first and second order branches, or rachillae, are referred to as the "partial inflorescence" [21]. The shape and overall morphology of these partial inflorescences are one of the

most important taxonomic characters for species identification and to define the different sections of the genus [20,21].

Despite its importance, *Raphia* remains one of the least understood palm genera in terms of taxonomy and phylogenetic relationships [1,20]. This is mainly due to their massive size, making them difficult to collect for non-specialists, which leads to few informative herbarium specimens or specimens that are incomplete or fragmentary and hence uninformative for taxonomy. Several attempts have been undertaken to tackle the taxonomy of the genus, beginning in the early 1900s with the first complete monograph of the genus [22]. This was followed by more regional attempts through the last century [23,24]. The last major revision of the genus was undertaken by Otedoh [21], who placed species into five different sections based on the structure of the partial inflorescence: Moniliformes (including the subsection Erectae), Temulentae, Raphiate, Flabellatae and Obclavatae.

The six species within the Moniliformes section are characterized by thin and easily breakable rachillae when fresh (Figure 1 B). Otedoh [21] also created a subsection, Erectae, where he placed two species in which the inflorescences are defined as erect (Figure *R. australis, R. regalis*) (Figure 1 G, O). The Temulentae section has robust and tightly appressed rachillae. The partial inflorescences are racquet-shaped with the apical second order rachillae shorter than the basal ones (Figure 1 E). This section contains three (possibly four) species, including one of the most widespread and important species *R. hookeri*. With seven species, the Raphiate section is the most complex group of the genus. Several species in this section are only known from a few collections or just the type. This section is characterized by species having second order rachillae that are robust (thick) but loosely disposed between them (Figure 1 D). The inflorescence within this section can be semi-erect or drooping (Figure 1 I). The Flabellatae section contains two species with very characteristic partial inflorescence structures. The second order rachillae are tightly packed in a single plane being racket-shaped in appearance (Figure 1 F). The inflorescence also has very conspicuous bracts that cover completely or partially the partial inflorescences (Figure 1 O). Finally, the Obclavatae section contains one species (*R. sudanica*) with distinct club-shaped and compact partial inflorescences with large bracts covering too (Figure 1 C).

To date, no in depth morphological or molecular phylogenetic study of *Raphia* has been undertaken. The most recent phylogenetic analysis of the Calamoideae subfamily included several individuals of a single species, namely *R. farinifera* [25]. The main objective of this study is to generate a densely sampled phylogenetic tree of the genus and test the validity of the taxonomic sections of Otedoh [21]. In particular, we test if the partial inflorescence structure has a phylogenetic signal and is useful for *Raphia* species classification. In addition, by sampling several individuals per morphologically identified species, we also tested species delimitation and monophyly. In order to achieve these objectives we sequenced more than 150 palm specific nuclear markers across 56 *Raphia* accessions. We used a species delimitation approach to define species limits and generated SNP data to study at fine-scale genetic relationships in identified species complexes.

2. Results

2.1. DNA sequencing

We sequenced 56 individuals representing 18 species or 87.5% of the species diversity within the genus. A total of 15.4 million reads were generated and mapped to the reference exons belonging to 176 genes of the Heyduk et al. [26] baiting kit. Across all *Raphia* and outgroup individuals the average coverage depth was 139.6x. We identified 102 genes for which 75% of the exon length was recovered in at least 25% of individuals. 20 loci were flagged by Hybpiper as paralogs because multiple assembled contigs matched a single reference locus. Of these 20, those that occurred in the 75/25 set were removed, resulting in a final dataset of 85 supercontigs equalling 162kb of sequence data. Our SNP calling approach applied filters on mapping quality (>40%) depth (>25), quality by depth (>2), minimum depth across individuals (>10) minor allele frequency (>0.01) and we excluded

112 monomorphic site. This ultimately yielded 915 and 1,627 high-quality, biallelic SNPs for the *R. hookeri*
113 and *R. zamiana* species complexes, respectively (see below).

2.2. Evolutionary history of Raphia

We generated two phylogenetic hypotheses for *Raphia* using two distinct methods. The first
analysis was conducted based on a gene-tree coalescent approach using ASTRAL while the second
inferred phylogenetic relationships based on a concatenated approach using IQ-TREE.

Support varied throughout the *Raphia* ASTRAL tree - about 50% of branches had a local posterior
probability (LPP) above 75% (Figure 1 A). Major clades were well supported (LPP > 80%) while
relationships towards the tips of the tree generally had lower support. The final normalized quartet
score, the proportion of quartet trees that agree with the species tree, was 65%, indicating that there is
gene tree conflict in the genus.

The IQ-TREE concatenated approach (see Figure A1 in the supplementary materials) had increased
bootstrap support compared to ASTRAL. More than 88% of branches had bootstrap support greater
than 75%. The best partitioning scheme put the 85 loci into 20 different partitions. Major clades were
again well-supported in this tree (bootstrap > 80%).

Our phylogenetic analyses recovered five well supported clades. Three clades matched the
sections as defined by Otedoh [21]. *Raphia regalis* was always inferred with strong support as sister to
the rest of the genus independent of the inference method (Figures 1, A1). When comparing the two
phylogenetic approaches we identified a topological difference in the phylogenetic placement of the
section Temulentae, the species *R. matombe* and the Moniliformes and Flabellatae sections (Figure 2).
In the IQ-TREE analyses we recovered weak support for the Temulentae as sister to all *Raphia* (except
R. regalis) (Figure 2a) while the ASTRAL analysis indicates with higher support that Temulentae is
sister to a clade containing *R. matombe*, Moniliformes & Flabellatae (Figure 2b).

The relationships between species in the Raphiate section are weakly to moderately supported
in both analyses (Figures 1, A1). Nevertheless, we do recover monophyletic groups in some species
consistent with prior morphological identifications. This is the case for individuals of *R. laurentii*
and *R. monbuttorum*, which despite low support are monophyletic. Furthermore, both these species
are recovered as sister, with moderate to high support. However, our species delimitation analysis
suggested that individuals identified under both species are conspecific (Figure 1 A). Support is
generally higher in the ASTRAL tree, even when taking into account different gene histories, so we
suggest that the ASTRAL tree represents a more accurate reconstruction of the phylogeny of *Raphia*
(Fig. 2b) so we will principally refer to the relationships in this tree from now on.

Figure 1. A: Cladogram of the genus *Raphia* inferred using 85 gene trees and ASTRAL. Values of local posterior probabilities (LPP) are shown above the branches. Branch lengths are represented in Figure A2. Individuals are colour coded based on the hypothesis of species delimitation inferred using SODA with α = 0.01. For a single clade, the Temulentae section marked with a black star and referred to as the "hookeri" complex in the main text, varied between the two values of α ued here. The orange boxes represent the species limits using SODA with a more stringent value of α = 0.005. The black square shows the "zamiana" complex clade. Tip names contain the species name as well as the sequencing ID. B: *R. regalis* partial inflorescence representing the Erectae section (described here, see discussion). C: *R. sudanica* inflorescence, representing the Obclavatae section. D: *R. palma-pinus* inflorescence, representing the Raphiate section. E: *R. hookeri* inflorescence, representing the Temulentae section. F: *R. farinifera* inflorescence, representing the Moniliformes (and ex Flabellatae) section. G: *R. regalis*, note the acaulescent habitat and inflorescences subtended by the leaves (Couvreur 398, Cameroon). H: *R. zamiana* (Mogue Kamga 17, Gabon). I: *R. monbuttorum* (Couvreur 1212, Cameroon). J: detail of *R. monbuttorum* rachillae (Couvreur 1212, Cameroon). K: detail of *R. laurentii* rachillae (Mogue Kamga 39, Democratic Republic of Congo). L: *R. hookeri* (no voucher, Cameroon). M: *R. gabonica* (Mogue Kamga 22, Gabon). N: *R. australis* (no voucher, South Africa, Kirstenbosch Botanic Garden). O: Inflorescence of *R. vinifera* (Couvreur 638, Cameroon). B-F: Line drawings by Mary Grierson reproduced with permission from [23]; Photos G-J, L-O: T.L.P. Couvreur; Photo K: S. Mogue Kamga.

Figure 2. Major incongruences between the (**A**) concatenation (IQ-TREE) and (**B**) gene tree (ASTRAL) phylogenetic approaches. Both trees have been modified to show the relationships among major *Raphia* clades. Support values are indicated on the nodes as either (**a**) bootstrap or (**b**) local posterior probability.

2.3. Species delimitation

Our species delimitation approach yielded between 17 (α = 0.005) and 23 (α = 0.01) species (Figure 1). Higher values of α split a clade of closely related individuals (marked with a star in Figure 1), predominantly belonging to *R. hookeri*, into seven different species. Generally, our species delimitation results corresponded well with our field identifications and using available floras (e.g. [19,20]). In some cases we found that SODA split individuals belonging *a priori* to a single species into multiple species, for example *R. farinifera* and *R. sudanica* (Fig. 1). Conversely, individuals assigned to different species such as *R. laurentii* and *R. monbuttorum* were classified as the same species according to SODA delimitation independent of α values. In general, the support among different species as delimited by SODA was high (Figure 1).

2.4. Fine scale structure in two species-complexes

To further explore genetic structure among our two main species complex, namely the "zamiana" and "hookeri complexes (marked with a black square and a black star in Figure 1, respectively), we used SNPs extracted from the sequence data to look at the variation among individuals. The "hookeri" complex showed little evidence of clustering, with most individuals evenly spread out on the first two principal component (PC) axes (Figure 3a). We observed two major groups of >8 individuals in the "zamiana" complex along PC1 (Figure 3b), separating all of the *R. laurentii* and *R. monbuttorum* from the rest of the individuals. The first two PCs in both analyses explained 7-10% of the variance in the dataset. In general, our SNP data supports SODA species delimitation as the assigned species grouped together along one or both of the first two PCs in most cases (Figure 3a, b). Finally, our SNP data revealed that individuals within the "hookeri" complex clustered into four major groups (Figure 3a, c): the single individual from Togo; individuals from western Cameroon; individuals from East Cameroon and individuals from Gabon.

The plotting of these complexes on maps of the sampling region revealed that the delimited species clustered geographically (Figure 3c, d). In the "hookeri" complex, the *R. sese* individuals were sampled at a great distance from each other and *R. gabonica* falls in the middle of the *R. hookeri* distribution range. Likewise, in the "zamiana" complex *R. laurentii* and *R. monbuttorum* are widespread, overlapping with other taxa. Many of the delimited species co-occur or are adjacent to one another in Cameroon.

Figure 3. (**a**) Scatterplot of *R. hookeri* complex based on 915 SNPs. (**b**) Scatterplot of *R. zamiana* complex based on 1627 SNPs. Clades representing the (**c**) *R. hookeri* and (**d**) *R. zamiana* complexes were extracted from the ASTRAL (Figure 1) tree and linked to their locations on a map of central Africa. Individuals are coloured by the colours corresponding to SODA species delimitation for α = 0.01 in (**a**) & (**c**) and α = 0.005 in (**b**) & (**d**). An individual belonging to *R. taedigera* (RA_TA) is not shown in panels (**a**) & (**c**) due to missing data.

3. Discussion

3.1. Synthesizing morphology and molecules: the sections of Otedoh reevaluated.

Our phylogenomic analyses of *Raphia* provide a novel and overall well supported phylogenetic framework for this important African genus (Figure 1 A). Although some of the morphology based sections of Otedoh [21] were recovered, we also recovered some topological differences (Figure 1 A).

The Moniliformes and Flabellatae are not recovered as monophyletic. The Moniliformes are split into two clades, while the two Flabellatae species (*Raphia regalis*, *R. australis*) are not recovered as sister and are nested within the Moniliformes section (Figure 1). In addition, the phylogenetic placement of the Moniliformes species *R. matombe* from the Democratic Republic of the Congo and Angola (including Cabinda) was different between the two types of analyses (Figure 2).

In all analyses, the acaulescent central African species *Raphia regalis* is recovered with strong support as sister to the rest of the genus (Figures 1 A, A1). This species, together with *R. australis*, were placed within the subsection "erectae" within the Moniliformes section [20,21] because the inflorescences were suggested to be "erect", in contrast to the rest of the *Raphia* species whose inflorescences are hanging or semi-erect (except for *R. palma-pinus* which also has an erect inflorescence). Our results do not support this classification, as *R. australis* is recovered as sister to *R. farinifera* (of the Flabelattae section, Figure 1) and phylogenetically divergent from *R. regalis*. A closer observation in the field showed that only the inflorescences of *R. australis* are truly erect (Figure 1 N). In contrast, the inflorescences of *R. regalis* appear erect but are in fact "supported" by the large leaves and not truly erect (Figure 1 G).

The close relationships between the Moniliformes and Flabellatae section is not surprising. The inflorescences, although different in some aspects such as the clearly racket-shaped partial inflorescences of the Flabellatae section, show certain similarities not encountered in other *Raphia* species. Both have thin rachillae and the partial inflorescences are subtended by large showy bracts at least in the younger stages of development. These morphological similarities thus support the close phylogenetic relationships recovered here between these two sections.

The Obclavate section, composed of the sole species *R. sudanica*, is recovered with strong or moderate support as sister to the Raphiate section. This species presents a unique inflorescence structure within the genus that is reduced and compressed into a cylindrical shape (Figure 1 C), with large bracts covering the partial inflorescences almost completely [20,21,27]. In addition, and in contrast to most species, *R. sudanica* thrives within the drier regions of the Sahel along small stream courses. These distinctive characters and its phylogenetic position support it being placed in a section of its own, confirming the classification of Otedoh [21].

Finally, the two remaining sections, Raphiate and Temulentae, are recovered as monophyletic, although with varying levels of support from strong to moderate (Figures 1 A,A1). This also confirms the classification of Otedoh [21] and the usefulness of partial inflorescence shapes in the classification of *Raphia* species.

Our results, however, suggest that certain sections erected by Otedoh [21] are not monophyletic and need to be re-evaluated. Differences in phylogenetic relationships between the concatenated and coalescent approaches have been increasingly reported in this genomic era [28–30]. Our results were similar to those in Couvreur et al. [19] where higher bootstrap support were obtained when using the concatenation approach, despite the coalescent approach highlighting considerable gene tree conflict. Here, we favour the phylogenetic hypothesis recovered when using the coalescent approach (Figure 1) because these methods allow gene history to be taken into account [31] and provide an arguably more realistic reconstruction of phylogenetic relationships when using a large number of independently evolving nuclear markers as used here. Our analyses suggest that we can retain five sections, only slightly different than those initially defined by Otedoh [21]. Three sections have been reconstructed in the phylogeny: Obclavatae (with its only species *R. sudanica*), Raphiate and Temulentae. The latter two sections are internally complicated, and more discussion about the phylogenetic relationship within

sections is provided below. The main problem thus comes from the Moniliformes and Flabellatae sections, which are not monophyletic. *Raphia regalis* should be placed in a section of its own, linked to its unique morphology being an acaulescent species with large inflorescences subtended by large leaves (Figure 1 G). Finally, the last section should regroup all the other species from both the Moniliformes and Flabellatae.

We thus recognize five main sections within *Raphia* based on the phylogenetic results presented here. We then discuss these results in more detail below.

1. Section Erectae (Otedoh) Couvreur, Mogue Kamga & Sonké **sect. nov.**. Type species: *R. regalis* Becc..

Diagnose: Acaulescent palms with less than ten leaves. Inflorescences erect amongst leaves, rachillae thin and brittle. Contains to date one species: *R. regalis*.

2. Section Moniliformes Otedoh. Type species: *R. textilis* Welw. This section also includes the species formally placed in section Flabellatae by Otedoh [21]. Because the name Moniliformes was published before Flabellatae (page 148 versus 163), we retained the former name here. A section with seven recognized species: *R. australis, R. farinifera, R. gentiliana, R. matombe, R. ruwenzorica, R. textilis* and *R. vinifera* (sensus Kamga Mogue et al. [32]).

3. Section Temulentae Otedoh. Type species: *R. hookeri* Mann & Wendl. This section remains the same as defined by Otedoh [21]. We also include the newly described species *R. gabonica* [7] in this section.

A section with four recognized species: *R. gabonica, R. hookeri, R. sese* and *R. rostrata*.

4. Section Raphiate Otedoh. Type species: *R. palma-pinus* (Gaertn.) Hutch.. Otedoh [21] had an erroneous vision of *R. vinifera* (the type of this section) which does not belong to the Raphiate section (see [32] for details). We thus choose a new type species being the second oldest name within this section.

A section with eight recognized species: *R. africana, R. laurentii, R. longiflora, R. mannii, R. monbuttorum, R. palma-pinus, R. taedigera* and *R. zamiana*.

5. Section Obclavatae Otedoh. Type species: *R. sudanica* Chev. One single single species is recognised in this section: *R. sudanica*

3.2. Species delimitation and species complexes

Phylogenetic relationships between species are well to weakly resolved depending on the section, as discussed below.

3.2.1. The Moniliformes/Flabellatae section

Within the Moniliformes/Flabellatae section species relationships are generally strongly supported (Figures 1, A1) and several species are recovered as monophyletic (*R. australis, R. farinifera, R. matombe*) while species limits in others are less clear (*R. textilis, R. vinifera*).

Once again, there is a conflict between the concatenated and coalescent analyses. *Raphia textilis* is split into two well-supported clades in the former analysis (Figure A1), while it is recovered as monophyletic with strong support in the latter (Figure 1). Nevertheless, there is little doubt that these samples represent the same species as they are morphologically similar. This is also confirmed by the species delimitation analysis at both levels of α (Figure 1, A2).

Another result recovered is the close relationship of the two montane species of *Raphia*: *R. ruwenzorica* been included within *R. vinifera*. Both species occupy a similar ecological and altitudinal range, despite being geographically separated by ca. 2,500 km. *Raphia vinifera*, of which most samples were in the past identified as *R. mambillensis* and now a synonym of *R. vinifera*, occurs mainly in Cameroon and Nigeria, where it grows between 1,200 and 2,000m in grassland or open vegetation and is very abundant along streams and rivers [10,32]. *Raphia ruwenzorica* occurs between 800 and 1,500 m in the Albertine rift region in eastern Democratic Republic of the Congo and Burundi and has been suggested to grow in "savanna country" along valleys [20,21,33,34]. In addition, both species present

similar partial inflorescences that are flat and racket shaped. However, both species differ markedly in their port with *R. ruwenzorica* reported to have a distinct tall stipe reaching up to 15 m [10,21,33] whereas as *R. vinifera* is acaulescent or with a short stipe (less than 1 m; [32]). This, in addition to the 2,500+ km separating these species, suggests that they could be recognized as distinct, despite our results. Interestingly, a within species CVL / Albertine rift disjunction has been documented in different taxa such as *Isolona congolana* (Annonaceae, [35]) and *Prunus africana* [36].

Raphia farinifera is the most widespread species of *Raphia*, occurring from West Africa to East Africa and Madagascar, and has also been reported from the Republic of Congo and Angola [23,34,37–42]. Our limited (3) but widespread sampling (West Africa and Madagascar) of individuals clustered together with maximum support (Figures 1, A1). However, our species delimitation analysis suggests that the Malagasy individual (R41_T15) is a different species (Figure 1). Interestingly, *Raphia* individuals from Madagascar were initially described as a different species (*R. ruffia* and also once described as *R. tamatavensis* Sadeb.) [22,34] and the name subsequently synonymized with *R. farinifera* [34,40]. In Madagascar, *Raphia* is widely used (one of the most useful palms) and, today at least, not found in natural forests across the island [43]. This has led to the hypothesis that *Raphia* was introduced 1,500 years ago during the first wave of human colonization of the island [43]. However, several authors suggested that this species use to occur naturally and abundantly in some places of the Island [22,44]. Our sampling is not extensive enough to answer this question conclusively, but our results suggest that Malagasy individuals might indeed belong to a different species (*R. ruffia*) as concluded by Beccari [22] (p. 53) thus not supporting the recent introduction hypothesis. Finally, *R. farinifera* is recovered as sister to *R. australis*, a relationship already suggested based on morphology [34].

3.2.2. The Raphiate section

This is one of the complex and least understood sections. Some of these species are poorly known and rarely collected, sometimes only known from a single poor quality specimen (*R. africana*; *R. longiflora*; *R. mannii*). In our study we were not able to sample *R. mannii* and *R. longiflora*, thus our results for this section are still incomplete. Overall, the relationships between species in the Raphiate section are weakly to moderately supported in both analyses (Figures 1, A1). Nevertheless, we do recover monophyletic groups in some species consistent with prior morphological identifications. This is the case for individuals of *R. laurentii* and *R. monbuttorum*, which despite low support are monophyletic. Furthermore, both these species are recovered as sister, with moderate to high support. However, our species delimitation analysis suggests that individuals identified under both species are conspecific (Figure 1 A). Indeed, these two species are morphologically similar [19], having clustering stipes covered with straight fibres in addition to having semi-erect inflorescences when young (Figure 1 I), an unusual character within the genus. Nevertheless, it is hard morphologically to consider these two species as conspecific. Indeed, the shape of the rachillae is quite different between these species (Figure 1 J,K). *Raphia laurentii* is characterized by rather thick rachillae covered by numerous tightly packed rachis bracts leading to an overall digitate aspect of the rachillae (Figure 1 J). In *R. monbuttorum* the rachillae are thinner and the rachis' bracts are less tightly packed around the rachillae (Figure 1 K). These differences appear to be consistent and provide useful identification characters [19].

Raphia zamiana was recently described [7]. Our broad sampling of this species, however, recovers *R. zamiana* as polyphyletic, with individuals grouping into two main clades, flagged as two different species by our species delimitation analysis (Figures 1, A2). Interestingly, these two species are geographically distinct, with one clade sampled across Gabon and one across Cameroon (Figure 3b, d), the latter containing the type of *R. zamiana*. The Gabon cluster is particularly well supported in both analyses. At this point, however, it is hard to pin point clear morphological characters differentiating these two clusters, as extensive field observations have yet to distinguish them properly.

We sampled two individuals of the Neotropical species *R. taedigera*, both from Brazil. As expected from the morphology of the partial inflorescence [21], this species grouped within the Raphiate section (Figure 1). Both individuals clustered together with strong support, and, in turn, were recovered as

sister to either *R. africana* (Figure 1) or *R. palma-pinus* (Figure A1, in both cases with weak support
values). Otedoh [21], following certain authors [23,45] suggested that *R. taedigera* was very close
morphologically to a species he called *R. vinifera*. However, early on, the taxonomic concept of *R.*
vinifera has been confusing, erroneously mistaking this species for a Raphiate type species [23,45].
Mogue Kamga et al. [32] clarified the situation showing that the name *R. vinifera* refers to a Flabelattae
species mainly occurring in the CVL. To date, it remains unclear to what species Otedoh and others
[21,23,45] were referring to when invoking *R. vinifera*.

Despite these taxonomic confusions, our results provide some results as to the origin of *R. taedigera*
in the Neotropcis. It has been hypothesised that this species originated as a result of vicariance during
the breakup of Gondwana [4]. The deeply nested position of *R. taedigera* within the genus does not
support this hypothesis. Instead, our results lend some support to the conclusion of Otedoh [10]
who suggested that *R. taedigera* did not show any "primitive" characters within the genus. Otedoh
went further to suggest that *R. taedigera* was the result of a recent introduction in South and Central
America during the slave trade some 400 years ago [10,11]. Our species delimitation results suggest,
however, that *R. taedigera* is a valid species (Figure 1), at least based on the individuals sampled
from Brazil. Finally, Otedoh also suggested the presence of *R. taedigera* in coastal west-central Africa
[10,11]. However, to date we have not been able to locate this species in African collections and this
hypothesis remains doubtful [1]. Our phylogenetic analyses suggest that *R. taedigera* is genetically
quite different from other *Raphia* species (Figure A2) supporting the hypothesis that it must have
dispersed to the Neotropics more than 400 years ago. This would fit with paleoecological data from
Nicaragua documenting *R. taedigera* pollen over the last 2,500 years [9]. A more detailed sampling of
R. taedigera from the Neotropics together with a dated molecular phylogeny approach will provide a
better understanding of the biogeographic history of this interesting trans-Atlantic disjunction.

3.2.3. The Temulentae section

This section contains the species referred to as the "wine" palms [20,21] with three species
previously included in this section (*R. hookeri*, *R. rostrata*, *R. sese*), all of which are sampled here.
In addition, our results show that the newly described species, *R. gabonica* [7], is also part of the
Temulentae section. This was not clear at the time of the publication as the partial inflorescence
suggested a possible relationship with the Moniliformes section [7]. Overall, species identified based
on morphology clustered together (e.g. *R. gabonica*, *R. sese*) with strong or low support. Nevertheless,
all four species show a very close phylogenetic proximity, suggesting that this section could be regarded
as a species complex. Indeed, depending on the level of stringency, our species delimitation analysis
recovered either seven distinct species or one single species (Figures 1, A2). It is important to note
that changing levels of α did not impact species delimitation in the other sections. Morphologically,
however, these species are different and can easily be identified in the field, which is partly supported
by our phylogenetic analysis. For example, *R. gabonica* resembles *R. hookeri* in the clearly visible single
stipe covered with characteristic curly fibers, but differs markedly by being a *terra firma* low-density
species with thin (Moniliformes-like) and densely packed rachillae. In contrast, *R. hookeri* is a swampy
species, growing in large, monodominant stands with robust and more evenly-spaced rachillae [7,19].
In the same way, *R. rostrata* is characterized by a small but clustering stipe with curly mixed with
straight hanging fibers and occurs along rivers with strong currents [19].

Raphia hookeri is recovered here as polyphyletic, possibly including four different cryptic species.
This is one of the most important, abundant and widespread *Raphia* species and its overall morphology
is rather constant across its range. However, individuals appear to be geographically structured like in
R. zamiana (see above). Interestingly, this mirrors patterns of genetic structure recovered across a wide
range of central African plant species [46,47], including *R. zamiana*.

4. Conclusions

Our results provide a new step forward in understanding the phylogenetic relationships and taxonomy within this major African palm genus. We show that the morphological sections based on partial inflorescence shape defined by Otedoh [21] are relatively robust overall. Sections Obclavatae, Temulentae and Raphiate are recovered as monophyletic with good support, while sections Moniliformes and Flabellatae are not. We thus redefined these later two sections into sections Erectae, composed of *R. regalis*, and Moniliformes, including all species previously included in Flabellatae. Our results also uncover important species delimitation problems defined here as species complexes (*R. hookeri*, *R. zamiana*) that must be solved if we are to have a thorough understanding of *Raphia* systematics. Given the economic and ecological importance of *R. hookeri*, clarifying its species delimitation will be important in the future. Different approaches could rely on more in-depth population level studies using more variable markers (e.g. microsatellites) combined with detailed morphometric measurements as has been done in other African tree species, e.g. [48–50]. A better comprehension of the taxonomy and the phylogenetic relationships in *Raphia* represents a fundamental tool towards the proposal of conservation strategies aiming to characterize genetic and morphologic diversity in this ecological and economically important genus. Finally, we show here that the Heyduk et al. baiting kit [26] is useful for understanding relationships within the *Raphia* genus and between species as was shown in other groups e.g. [51], although it appears to be limited for untangling species complexes. Resolving relationships within *Raphia* will thus rely on more data, including increased infra-species sampling, detailed morphological studies in certain species and larger baiting kits, e.g. [52].

5. Materials and Methods

5.1. Species sampling, library preparation and DNA sequencing

We sampled a total of 56 individuals (see A1 for details) representing 18 out of the 21 species accepted to date [7,19] and representing all sections described by Otedoh [21]. In order to collect proper material for sequencing, several field trips were undertaken across several African countries including Ivory Coast, Ghana, Gabon, Cameroon, Angola and the Demographic Republic of the Congo between 2012 and 2017. We were not able to access material from three accepted species: *R. gentiliana*, *R. mannii* and *R. longiflora*. We sampled two to seven individuals per species in order to test for monophyly. However, only a single specimen was available for *R. ruwenzorica*. Finally, we sampled four species within Calamoideae as outgroups: *Eremospatha cabrae*, *Eremospatha quinquecostulata*, *Laccosperma cristalensis* and *Mauritiella armata* following [25,53]. We extracted DNA from leaves dried in silicagel, except for one individual of *R. taedigera* and the only individual of *R. ruwenzorica* for which DNA was extracted from herbarium material.

Methods for DNA extraction, preparation of sequencing libraries, hybridization, Illumina MiSeq DNA sequencing and read cleaning followed [19]. In brief, barcoded Illumina libraries were constructed based on a modified protocol of Rohland and Reich [54]. We hybridized DNA to defined exons using the palm-specific nuclear baiting kit of Heyduk et al. [26]. This kit allows to sequence exons from 176 nuclear genes across the palm family.

5.2. Contig assembly and multi-sequence alignment

We used HybPiper (v1.2) [55] to process our cleaned reads (following [19]) to obtain sequences corresponding to the target exons plus associated intronic sequence data (referred to as supercontigs). We aligned each set of supercontigs using MAFFT (v7.305) [56] with the –auto option and cleaned these alignments with GBLOCKS (v0.91b) [57] using the default parameters and all allowed gap positions.

To identify a suitable set of loci for phylogenetic inference we selected only those supercontigs that had 75% of their exon length reconstructed in at least 25% of individuals (referred to as 75/25).

411 We used only those loci in which at least 75% of the exon length was recovered because the use of
412 fragmented sequences is known to increase gene tree error, whereas the number of individuals has
413 little effect as long as the gene tree is accurate [58].

5.2.1. Paralog identification

HybPiper flags potential paralogs when multiple contigs are discovered mapping well to a
single reference sequence. We ran hybpiper on the 837 exons that made up the baiting kit [26],
identified flagged loci and constructed exon trees using RAxML (v8.2.9) [59]. We examined each tree to
determine whether putative paralogs formed a species clade. When sequences concerning more than
three individuals were flagged for a locus, we examined whether the 'main' and alternative sequences
formed separate clades. If so this locus was classified as a paralog and discarded from the dataset. For
each gene, we then calculated at the proportion of exons that we confirmed as paralogs after inspection.
If this proportion was <%50 we removed the entire gene from our analyses.

5.3. Coalescent phylogenetic inference

424 Individual gene trees were constructed with 100 bootstraps and the GTRGAMMA model using
425 RAxML (v8.2.9) [59] (option "-f a"). If after inference, branches had bootsrap support values >10 they
426 were collapsed using the program nw_ed [60] because this approach has been shown to improve the
427 accuracy of ASTRAL [31]. We used the selected 75/25 gene trees as our input to run ASTRAL-III
428 (v5.5.11) [31] using the default options.

5.4. Species delimitation

After constructing our ASTRAL tree we used the associated approach SODA [61]. Simulations
using this approach have shown it to be of similar accuracy or more accurate [61] than other popular
species delimitation methods such as BPP [62] at a fraction of the computational cost. SODA uses
frequencies of quartet topologies to determine if each branch in a guide tree inferred from gene trees
(i.e. the ASTRAL tree from above) is likely to have a positive length. This identifies where in the tree
coalescence is random, and where it is non-random. It then uses the results to infer a new, extended
species tree that defines boundaries among species. We used two cut-off values of α (confidence level):
0.01 and 0.005.

5.5. Maximum-likelihood phylogenetic inference

After suitable loci were identified we filled any missing individuals in each alignment with
an empty sequence. We then concatenated all aligned loci using the *pxcat* function in the program
phyx [63]. We used IQ-TREE (v1.6.8; [64])to infer a maximum likelihood tree of all individuals. We
partitioned our dataset so that each supercontig had a separate substitution model and used the
following options when running the program: "-m MFP+MERGE -rcluster 10 -bb 1000 -alrt 1000". We
selected the optimal partitioning scheme using ModelFinder [65], choosing the best model based on
Bayesian Information Criterion (BIC) score and merging genes until model fit stopped increasing. We
also used rcluster [66] to decrease computational load. We made use of the ultrafast bootstrapping
([67]; 1000 replicates) and the SH-like approximate likelihood ratio test ([68]; 1000 replicates) to assess
branch support in the tree.

5.6. SNP calling

450 To call SNPs we first used SeCaPr (v1.1.4; [69]) to build a psuedoreference. After filtering out
451 low coverage and paralogous loci, consensus sequences are built and combined to form a reference
452 file that is closer to the study group than the original, and will recover more data. We mapped our
453 cleaned, paired reads to this new, dataset-specific reference using BWA (v0.7.12; [70]). Duplicates were
454 removed and we called SNPs using the program HaplotypeCaller in GATK (v4.0; [71]). We applied

thresholds to mapping quality (>40%) depth (>25), quality by depth (>2), minimum quality across all individuals (>10) and minor allele frequency (>0.01) to filter SNPs using bcftools (v1.8; [72]). We kept only biallelic SNPs and excluded monomorphic sites.

5.7. Genetic clustering

We performed Discriminant Analysis of Principal Components (DAPC) [73] to identify genetic clusters in two species complexes of *Raphia*. We used the function *find.clusters* in the R package 'adegenet' [74] to infer the number of clusters using successive K-means with 100,000 iterations per value of k up to k = 20. We used BIC to identify the best-fitting number of clusters. We then used the function *dapc* [73] to define the diversity among the clusters identified. We chose the optimum number of axes to use with the function *optim.a.score*.

Author Contributions: Conceptualization, TLPC and BS; methodology KB and AJH; validation AJH, FS, KB and MKS; formal analysis AJH; resources LT, FS, CDB, JJW, AZ, AA, MKS, TLPC; data curation KB, AJH and MKS; writing–original draft preparation AJH and TLPC; writing–review and editing, all authors; supervision, TLPC and BS; project administration, TLPC; funding acquisition, TLPC. All authors have read and agreed to the published version of the manuscript.

Funding: This study was supported by Agropolis Fondation (RAPHIA project) under the reference ID 1403-026 through the «Investissements d'avenir» program (ANR-10-LABX-0001-01; ISITE MUSE: ANR-16-IDEX-0006) and Agence Nationale de la Recherche (ANR-15- CE02-0002-01, AFRODYN project), both to T.L.P.C.. This work was also part of the PALM-A-GOV (project number: 1702-007) project to T.L.P.C., publicly funded through the ANR (the French National Research Agency) under the "Investissements d'avenir" program with the reference ANR-10-LABX001-01 Labex Agro and coordinated by Agropolis Fondation within the I-SITE MUSE (ANR-16-IDEX-0006). A.Z. is funded by iDiv via the German Research Foundation (DFG FZT 118), specifically through sDiv, the Synthesis Centre of iDiv. C.D.B. is funded by the Swedish Research Council (2017-04980). A.A. is funded by the Swedish Research Council, the Swedish Foundation for Strategic Research, the Knut and Alice Wallenberg Foundation and the Royal Botanic Gardens, Kew. J.J.W. received funding for travel from the Alberta Mennega foundation.

Acknowledgments: The authors acknowledge the IRD itrop HPC (South Green Platform) at IRD Montpellier for providing HPC resources that have contributed to the research results reported within this paper. We are grateful to the Centre National de la Recherche Scientifique et Technique (CENAREST) in Gabon and the Agence National des Parques Nationaux (ANPN) for delivering the necessary research permits for our field work in Gabon in 2015 and 2016 (permits AR0020/16; AR0036/15, AR0035/15 (CENAREST) and AE16014; AE15027 (ANPN)). Prof. Paul Henri Bourobou Bourobou is thanked for his assistance in obtaining these research permits, while Raoul Niangadouma and Narcisse Kamdem are thanked for there help in the field. Fieldwork in Cameroon was undertaken under the "accord cadre de cooperation" between the IRD and Ministère de la Recherche Scientifique et Technique (MINRESI). TU Dresden maintains a Memorandum of Understanding with the Instituto Nacional da Biodiversidade e Áreas de Conservação (INBAC) of the Ministério do Ambiente da República de Angola that includes collection and export permits.

Conflicts of Interest: The authors declare no conflict of interest.

Abbreviations

The following abbreviations are used in this manuscript:

CVL	Cameroon Volcanic Line
SNP	Single Nucleotide Polymorphism
PC	Principal Components Analysis

Appendix A Supplementary Tables

Genus	Species epithet	ID	Collector (herbarium)	Country	Latitude	Longitude	Run	Tag	No. mapped	Mean depth	Stdev depth
Eremospatha	*cabrae*	R227	Couvreur 1165 (WAG)	Cameroon	3.128984	9.973292	RUN67	TAG-28	441790	315.48	469.205
Eremospatha	*quinquecostulata*	R162	Couvreur 1079 (WAG)	Gabon	-1.45585	12.5863	RUN41	TAG39	91982	46.1243	68.317
Laccosperma	*cristensis*	R164	Couvreur 1142 (WAG)	Gabon	0.6059	10.4118	RUN41	TAG41	105250	54.4592	86.7057
Mauritiella	*armata*	R135	Couvreur 257 (NY)	Bolivia	-13.496289	-68.019923	RUN37	TAG6	406142	213.159	1244.61
Raphia	*africana (cf)*	R072	Couvreur 971 (WAG)	Cameroon	4.87036	9.26579	RUN41	TAG10	121972	63.533	69.0331
Raphia	*africana (cf)*	R077	No voucher, close to Couvreur 971 (WAG)	Cameroon	4.87064	9.26582	RUN33	TAG43	315915	159.931	188.084
Raphia	*africana (cf)*	R174	No voucher, close to Couvreur 971 (WAG)	Cameroon	4.12977	9.21399	RUN41	TAG47	50272	26.0156	28.656
Raphia	*australis*	R092	MBC 99 874D (SANBI)	Cultivated, Kirstenbosch National Botanical Garden, South Africa	N/A	N/A	RUN33	TAG53	255329	129.65	156.525
Raphia	*australis*	R130	MBC 99 874D (SANBI)	Cultivated, Kirstenbosch National Botanical Garden, South Africa	N/A	N/A	RUN37	TAG4	913167	473.13	627.945
Raphia	*farinifera*	R128	Baker 1410 (K)	Burkina Faso	5.39625	-1.38278	RUN37	TAG1	476130	245.7	292.787
Raphia	*farinifera*	R127	Baker 1357 (K)	Cultivated, Royal Botanic Gardens, Kew, U.K.	NA	NA	RUN37	TAG105	486723	250.42	309.47
Raphia	*farinifera*	R132	Dransfield 7516 (K)	Madagascar	-1.84988	13.85903	RUN41	TAG15	133441	68.7643	85.4757
Raphia	*gabonica*	R034	Kamga Mogue 22 (WAG)	Gabon	-0.07916	11.00836	RUN33	TAG23	284243	143.822	174.031
Raphia	*gabonica*	R036	Kamga Mogue 23 (WAG)	Gabon	-1.03044	10.51881	RUN33	TAG24	212852	108.979	136.141
Raphia	*hookeri*	R063	Kamga Mogue 1 (WAG)	Cameroon	3.07485	13.3663	RUN33	TAG35	319096	162.68	210.004
Raphia	*hookeri*	R069	Kamga Mogue 12 (WAG)	Cameroon	3.97037	13.2367	RUN33	TAG39	197508	100.49	122.815
Raphia	*hookeri*	R071	Kamga Mogue 26 (WAG)	Cameroon	4.11224	9.56915	RUN33	TAG41	292201	149.509	181.385
Raphia	*hookeri*	R124	Couvreur 984 (WAG)	Cameroon	3.52108	11.74376	RUN41	TAG14	78554	40.8646	44.4911
Raphia	*hookeri*	R037	Kamga Mogue 25 (WAG)	Gabon	-0.82955	10.52294	RUN33	TAG25	278678	142.219	175.028
Raphia	*hookeri*	R089	Michon 01 (G)	Togo	6.39183	2.67703	RUN33	TAG51	201880	103.103	124.193
Raphia	*laurentii*	R208	Lautenschläger 806 (JACQ)	Angola	-6.14997	15.40333	RUN46	TAG18	64379	34.3542	37.7478
Raphia	*laurentii*	R040	Ayole 01 (YA)	Cameroon	2.15329	15.7367	RUN33	TAG26	272509	138.397	168.066
Raphia	*laurentii*	R186	Kamga Mogue 39 (WAG)	Democratic Republic of Congo	-0.8852	18.1337	RUN46	TAG10	87838	47.1055	51.2042
Raphia	*laurentii*	R198	Kamga Mogue 42 (WAG)	Democratic Republic of Congo	-0.60673	18.2468	RUN46	TAG14	86786	46.5246	51.1853
Raphia	*matombe*	R206	Lautenschläger 1095 (JACQ)	Angola	-7.94817	15.83894	RUN46	TAG16	59006	31.6789	34.7279
Raphia	*matombe*	R134	19392103 (BR)	Cultivated, Meise Botanical Garden, Belgium	NA	NA	RUN41	TAG16	76250	39.5975	44.2225
Raphia	*matombe*	R181	Kamga Mogue 37 (WAG)	Democratic Republic of Congo	-5.73485	14.2162	RUN46	TAG8	55883	29.9284	31.68
Raphia	*matombe*	R183	Kamga Mogue 38 (WAG)	Democratic Republic of Congo	-5.65308	14.3181	RUN46	TAG9	66676	35.6465	38.4587
Raphia	*monbuttorum*	R059	Kandem 211 (WAG)	Cameroon	3.885055556	14.39930556	RUN41	TAG9	104568	54.2312	58.009
Raphia	*monbuttorum*	R066	Kamga Mogue 04 (WAG)	Cameroon	3.07684	13.36761	RUN37	TAG78	407023	208.475	245.939
Raphia	*monbuttorum*	R070	Kamga Mogue 13 (WAG)	Cameroon	3.58237	13.14197	RUN33	TAG40	352731	180.143	231.782
Raphia	*monbuttorum*	R173	Kamga Mogue 31 (WAG)	Cameroon	4.18332	13.10538	RUN41	TAG46	98860	50.8202	56.6257
Raphia	*palma-pinus*	R133	Ouatara & Stauffer 14 (G)	Ghana	5.39625	-1.38278	RUN37	TAG5	792856	411.261	507.288
Raphia	*palma-pinus*	TC-S1328	Stauffer 857 (G)	Ivory Coast	7.24598	-5.39625	RUN67	TAG-26	459813	318.875	309.014
Raphia	*regalis*	R055	Couvreur 685 (WAG)	Cameroon	2.49409	10.34844	RUN33	TAG30	184893	94.4991	128.908
Raphia	*regalis*	R056	Couvreur 753 (WAG)	Cameroon	3.4825	13.59469	RUN33	TAG31	210853	107.898	137.475
Raphia	*regalis*	R058	Couvreur 398 (WAG)	Cameroon	3.19962	10.51772	RUN33	TAG33	388670	196.877	269.794
Raphia	*regalis*	R081	Wieringa 8539 (WAG)	Gabon	-2.37125	11.16443	RUN33	TAG46	161079	81.8365	100.151
Raphia	*regalis*	R083	Wieringa 8547 (WAG)	Gabon	-2.3714	11.16399	RUN33	TAG48	150544	76.7195	98.4529
Raphia	*rostrata*	R229	Kamga Mogue 43 (WAG)	Cameroon	2.339	10.6025	RUN67	TAG-30	476709	329.286	313.366
Raphia	*ruwenzorica*	RA_RU	Robyns 4039 (BR) - herbarium sample	Democratic Republic of Congo	NA	NA	N/A	N/A	N/A	N/A	N/A
Raphia	*sese*	R179	Kamga Mogue 36 (WAG)	Democratic Republic of Congo	-5.02068	15.1545	RUN46	TAG7	62972	33.7432	37.0717
Raphia	*sese*	R195	Kamga Mogue 41 (WAG)	Democratic Republic of Congo	-0.8642	18.1458	RUN46	TAG12	88659	47.5337	50.4421
Raphia	*sudanica*	R086	Michon 09 (G)	Benin	7.55826	2.19247	RUN33	TAG49	379755	193.881	231.648
Raphia	*sudanica*	R129	Bayton 70 (K)	Burkina Faso	10.5938888	-5.3069444	RUN37	TAG2	471997	246.119	299.37
Raphia	*sudanica*	R088	Michon 56 (G)	Togo	8.98317	1.49297	RUN33	TAG50	355099	181.913	220.289
Raphia	*taedigera*	RA_TA	Noblick 5015 (K) - (herbarium sample)	Brazil	-1.5666	-48.73333	N/A	N/A	N/A	N/A	N/A
Raphia	*taedigera*	Env0563	MBC 94803 A (MBC)	Cultivated, Montgomery Botanical Garden	N/A	N/A	RUN67	TAG-22	666580	446.883	465.14
Raphia	*textilis*	R209	Lautenschläger 1086 (JACQ)	Angola	-6.74019	16.20117	RUN46	TAG19	50163	26.882	30.0834
Raphia	*textilis*	R192	Kamga Mogue 40 (WAG)	Democratic Republic of Congo	-0.88513	18.1337	RUN46	TAG11	78738	42.0848	46.2868
Raphia	*textilis*	R149	Couvreur 743 (WAG)	Gabon	-1.84988	13.85903	RUN41	TAG29	132014	68.4236	80.0799
Raphia	*textilis*	R151	Couvreur 1075 (WAG)	Gabon	-1.40695	12.5712	RUN41	TAG31	80524	41.722	45.5424
Raphia	*vinifera*	R105	Couvreur 638 (WAG)	Cameroon	6.364476985	10.894598	RUN41	TAG12	87308	45.4189	49.1511
Raphia	*vinifera*	R113	No voucher, Couvreur 638 (WAG)	Cameroon	6.27413	10.51091	RUN37	TAG94	286322	145.81	177.287
Raphia	*vinifera*	R116	No voucher	Cameroon	5.48034	10.05056	RUN37	TAG97	367120	189.172	220.664
Raphia	*zamiana*	R057	Couvreur 427 (WAG)	Cameroon	3.59972	11.2877	RUN33	TAG32	411897	209.629	268.899
Raphia	*zamiana*	R093	Ayole 20 (YA)	Cameroon	3.23685	10.02514	RUN37	TAG79	300001	153.527	177.289
Raphia	*zamiana*	R095	Ayole 32 (YA)	Cameroon	2.80897	10.52734	RUN37	TAG81	420757	215.428	250.534
Raphia	*zamiana*	R230	Kamga Mogue 45 (WAG)	Cameroon	3.137672	9.971397	RUN67	TAG-31	468825	318.682	376.272
Raphia	*zamiana*	R007	Kamga Mogue 17 (WAG)	Gabon	1.59848	11.62294	RUN33	TAG8	217692	110.68	136.444
Raphia	*zamiana*	R009	Kamga Mogue 17 (WAG)	Gabon	-2.25428	11.14284	RUN33	TAG10	228828	116.278	144.853
Raphia	*zamiana*	R154	Couvreur 1122 (WAG)	Gabon	-0.1473	11.726	RUN41	TAG33	56712	29.1983	31.1171

Table A1. Table 1: Herbarium specimen collector and number and herbarium origin, coordinates (in decimal degrees) and country of collection for all samples included in our study. All samples were extracted from silicagel dried leaves, expect when stated otherwise. The last five columns refer to the sequencing identification (TAG and INDEX used) and different sequencing statistics (number of reads mapped to the reference; mean and standard depth, respectively.

498 **Appendix B Supplementary Figures**

Figure A1. IQTREE *Raphia* inferred using 162kb of sequence data. Values for ultrafast bootstrap support are depicted on nodes.

Figure A2. ASTRAL tree of *Raphia* including inferred branch lengths (except terminal branch lengths) and tip labels coloured with species delimitation as inferred with SODA (α = 0.01).

References

1. Dransfield, J.; Uhl, N.W.; Asmussen, C.B.; Baker, W.J.; Harley, M.M.; Lewis, C.E. *Genera Palmarum: The evolution and classification of palms*; Kew Publishing: Kew, 2008.

2. Stauffer, F.W.; Ouattara, D.N.; Roguet, D.; da Giau, S.; Michon, L.; Bakayoko, A.; Ekpe, P. An update to the African palms (Arecaceae) floristic and taxonomic knowledge, with emphasis on the West African region. *Webbia* **2017**, *72*, 1–14. doi:10.1080/00837792.2017.1313381.

3. Cosiaux, A.; Gardiner, L.M.; Stauffer, F.W.; Bachman, S.P.; Sonké, B.; Baker, W.J.; Couvreur, T.L.P. Low extinction risk for an important plant resource: Conservation assessments of continental African palms (Arecaceae/Palmae). *Biological Conservation* **2018**, *221*, 323–333. doi:10.1016/j.biocon.2018.02.025.

4. Moore, H.E. Palms in the tropical forest ecosystems of Africa and South America. In *Tropical forest ecosystems of Africa and South America: a comparative review*; Meggers, B.J.; Ayensu, E.; Duckworth, W.D., Eds.; Smithsonian Institution Press: Washintong DC, 1973; pp. 63–88.

5. Couvreur, T.L.P. Odd man out: Why are there fewer plant species in African rain forests? *Plant Systematics and Evolution* **2015**, *301*, 1299–1313. doi:10.1007/s00606-014-1180-z.

6. Dransfield, J. The palms of Africa and their relationships. In *Modern systematic studies in African botany*; Goldblatt, P.; Lowry, P.P., Eds.; Missouri Botanical Garden Press, 1988; pp. 95–103.

7. Mogue Kamga, S.; Niangadouma, R.; Stauffer, F.W.; Sonké, B.; Couvreur, T.L.P. Two new species of *Raphia* (Palmae/Arecaceae) from Cameroon and Gabon. *PhytoKeys* **2018**, *111*, 17–30. doi:10.3897/phytokeys.111.27175.

8. Urquhart, G.R. Long-term Persistence of *Raphia taedigera* Mart. Swamps in Nicaragua1. *Biotropica* **1999**, *31*, 565–569. doi:10.1111/j.1744-7429.1999.tb00403.x.

9. Urquhart, G.R. Paleoecological evidence of *Raphia* in the Pre-Columbian Neotropics. *Journal of Tropical Ecology* **1997**, *13*, 783–792. doi:10.1017/S0266467400010993.

10. Otedoh, M.O. Systematic Studies in *Raphia* Palms. Doctoral Thesis, University of Reading, UK., 1976.

11. Otedoh, M.O. The African origin of *Raphia taedigera* – Palmae. *J. Niger. Inst. Oil Palm Res* **1977**, *42*, 11–16.

12. Mogue Kamga, S.; Brokamp, G.; Cosiaux, A.; Awono, A.; Fürniss, S.; Barfod, A.S.; Muafor, F.J.; Le Gall, P.; Sonké, B.; Couvreur, T.L.P. Use and cultural significance of *Raphia* palms. *Economic Botany*.

13. Lautenschläger, T.; Neinhuis, C., Eds. *Riquezas naturais de Uíge – uma breve introdução sobre o estado atual, a utilização, a ameaça e a preservação da biodiversidade*, technische universität dresden ed.; 2014.

14. Obahiagbon, F.I. A review of the origin, morphology, cultivation, economic products, health and physiological implications of raphia palm. *African Journal of Food Science* **2009**, *3*, 447–453. 00011.

15. Couvreur, T.L.P.; Fumtim, J. A l'Echelle du Raphia / On Raphia and Man, 2017.

16. Profizi, J.P. Swampy Area Transformations by Exploitation of *Raphia hookeri* (Arecaceae) in Southern Benin (West Africa). *Human Ecology* **1988**, *16*, 87–94.

17. Dargie, G.C.; Lewis, S.L.; Lawson, I.T.; Mitchard, E.T.A.; Page, S.E.; Bocko, Y.E.; Ifo, S.A. Age, extent and carbon storage of the central Congo Basin peatland complex. *Nature* **2017**, *542*, 86–90. doi:10.1038/nature21048.

18. Rainey, H.J.; Iyenguet, F.C.; Malanda, G.A.F.; Madzoké, B.; Santos, D.D.; Stokes, E.J.; Maisels, F.; Strindberg, S. Survey of *Raphia* swamp forest, Republic of Congo, indicates high densities of Critically Endangered western lowland gorillas Gorilla gorilla gorilla. *Oryx* **2010**, *44*, 124–132. doi:10.1017/S003060530999010X.

19. Couvreur, T.L.P.; Helmstetter, A.J.; Koenen, E.J.M.; Bethune, K.; Brandão, R.D.; Little, S.A.; Sauquet, H.; Erkens, R.H.J. Phylogenomics of the Major Tropical Plant Family Annonaceae Using Targeted Enrichment of Nuclear Genes. *Frontiers in Plant Science* **2019**, *9*, 1941. doi:10.3389/fpls.2018.01941.

20. Tuley, P. *The palms of Africa*; The Tendrine Press, Trendrine Press: Zennor, 1995.

21. Otedoh, M.O. A revision of the genus *Raphia* Beauv. (Palmae). *J. Niger. Inst. Oil Palm Res* **1982**, *6*, 145–189. 00011.

22. Beccari, O. Studio monografico del genere "Raphia". *Webbia* **1910**, *3*, 37–130. doi:10.1080/00837792.1910.10669558.

23. Russell, T.A. The *Raphia* Palms of West Africa. *Kew Bulletin* **1965**, *19*, 173–196. doi:10.2307/4108027.

24. Tuley, P.; Russell, T.A. The Raphia palms reviewed. *Nigerian Field* **1966**, *31*, 54–65.

25. Baker, W.J.; Dransfield, J.; Hedderson, T.A. Phylogeny, character evolution, and a new classification of the calamoid palms. *Systematic Botany* **2000**, *25*, 297–322.

26. Heyduk, K.; Trapnell, D.W.; Barrett, C.F.; Leebens-Mack, J. Phylogenomic analyses of species relationships in the genus Sabal (Arecaceae) using targeted sequence capture. *Biological Journal of the Linnean Society* **2016**, *117*, 106–120. doi:10.1111/bij.12551.

27. Ouattara, D.N.; Stauffer, F.W.; Bakayoko, A. Lectotypification de *Raphia sudanica* A. Chev.(Arecaceae, Calamoideae), et commentaires sur la biologie et la conservation de l'espèce. *Adansonia* **2014**, *36*, 53–61. 00002.

28. Liu, X.; Fu, Y.X. Exploring population size changes using SNP frequency spectra. *Nature Genetics* **2015**, *47*, 555–559. doi:10.1038/ng.3254.

29. Tonini, J.; Moore, A.; Stern, D.; Shcheglovitova, M.; Ortí, G. Concatenation and Species Tree Methods Exhibit Statistically Indistinguishable Accuracy under a Range of Simulated Conditions. *PLoS Currents* **2015**, *7*. doi:10.1371/currents.tol.34260cc27551a527b124ec5f6334b6be.

30. Springer, M.S.; Gatesy, J. The gene tree delusion. *Molecular Phylogenetics and Evolution* **2016**, *94*, 1–33. doi:10.1016/j.ympev.2015.07.018.

31. Zhang, C.; Sayyari, E.; Mirarab, S. ASTRAL-III: Increased scalability and impacts of contracting low support branches. Lecture Notes in Computer Science (including subseries Lecture Notes in Artificial Intelligence and Lecture Notes in Bioinformatics), 2017, Vol. 10562 LNBI, pp. 53–75. doi:10.1007/978-3-319-67979-2_4.

32. Mogue Kamga, S.; Sonké, B.; Couvreur, T.L.P. Raphia vinifera (Arecaceae; Calamoideae): Misidentified for far too long. *Biodiversity Data Journal* **2019**, *7*, e37757. doi:10.3897/BDJ.7.e37757.

33. Robyns, W.; Tournay, R. Monocotylées nouvelles ou critiques de la région du Parc National Albert (Congo belge). *Bulletin du Jardin botanique de l'État a Bruxelles* **1955**, *25*, 239–260. doi:10.2307/3667057.

34. Stauffer, F.W.; Ouattara, D.; Stork, A.L. Palmae. In *Tropical African Flowering Plants: Monocotyledons 2*; Lebrun, J.P.; Stork, A.L., Eds.; Conservatoire et Jardin botaniques de la Ville de Genève, Switzerland.: Genève, 2014; Vol. 8, pp. 326–354.

35. Couvreur, T.L.P. Monograph of the syncarpous African genera *Isolona* and *Monodora* (Annonaceae). *Systematic Botany Monographs* **2009**, *87*, 1–150.

36. Kadu, C.a.C.; Schueler, S.; Konrad, H.; Muluvi, G.M.M.; Eyog-Matig, O.; Muchugi, A.; Williams, V.L.; Ramamonjisoa, L.; Kapinga, C.; Foahom, B.; Katsvanga, C.; Hafashimana, D.; Obama, C.; Geburek, T.

Phylogeography of the Afromontane Prunus africana reveals a former migration corridor between East and West African highlands. *Molecular Ecology* **2011**, *20*, 165–178. doi:10.1111/j.1365-294X.2010.04931.x.

37. Dale, I.R. Palms of Kenya. *The Journal of the East Africa and Uganda Natural History Society* **1938**, *23*, 183–186.

38. Russell, T.A. Palmae. In *Flora of West Tropical Africa*, 2 ed.; Hutchinson, J.; Dalziel, J.M.; Hepper, F., Eds.; Crown Agents for Oversea Governments and Administrations: London, 1968; Vol. 3, pp. 159–169.

39. Dransfield, J. Palmae. In *Flora of Tropical East Africa*; Polhill, R., Ed.; A.A. Balkema: Rotterdam, 1986; pp. 1–58.

40. Dransfield, J.; Beentje, H.J. *The palms of Madagascar*; Royal Botanic Gardens and International Palm Society: Kew, 1995. 00134.

41. Baker, W.J.; Smith, G.F. Palmae. In *Plants of Angola*; Figueiredo, E., Ed.; Number 23 in Strelitzia, 2008; pp. 172–173.

42. Faye, A.; Moutsamboté, J.; Couvreur, T.L.P. Palms in southern Republic of Congo. *PALMS* **2016**, *59*, 181–190.

43. Dransfield, J.; Rakotoarinivo, M. The biogeography of Madagascar palms. In *The Biology of Island Floras*, Cambridge University Press ed.; Bramwell, D.; Caujapé-Castells, J., Eds.; 2011; pp. 179–196.

44. Jumelle, H.; Perrier de la Bathié, M. Palmiers. In *Flore de Madagascar et des Comores (plantes vasculaires)*; Humbert, H., Ed.; Impr. officielle. Tananarive: Tananarive, 1945; Vol. 30, pp. 1–185.

45. Mann, G.; Wendland, H.A. On the Palms of Western Tropical Africa. *Transactions of the Linnean Society of London* **1864**, *24*, 421–440. doi:10.1111/j.1096-3642.1863.tb00165.x.

46. Hardy, O.J.; Born, C.; Budde, K.; Daïnou, K.; Dauby, G.; Duminil, J.Ô.; Ewédjé, E.E.B.; Gomez, C.; Heuertz, M.; Koffi, G.K.; Lowe, A.J.; Micheneau, C.; Ndiade-Bourobou, D.; Piñeiro, R.; Poncet, V. Comparative phylogeography of African rain forest trees: A review of genetic signatures of vegetation history in the Guineo-Congolian region. *Comptes Rendus - Geoscience* **2013**, *345*, 284–296. doi:10.1016/j.crte.2013.05.001.

47. Helmstetter, A.J.; Amoussou, B.E.N.; Bethune, K.; Kandem, N.G.; Kakaï, R.G.; Sonké, B.; Couvreur, T.L.P. Phylogenomic data reveal how a climatic inversion and glacial refugia shape patterns of diversity in an African rain forest tree species. *bioRxiv* **2019**, p. 807727. doi:10.1101/807727.

48. Lissambou, B.J.; Couvreur, T.L.P.; Atteke, C.; Stévart, T.; Piñeiro, R.; Dauby, G.; Monthe, F.K.; Ikabanga, D.U.; Sonké, B.; M'batchi, B.; Hardy, O.J. Species delimitation in the genus Greenwayodendron based on morphological and genetic markers reveals new species. *TAXON* **2019**, *68*, 442–454. doi:10.1002/tax.12064.

49. Monthe, F.K.; Duminil, J.; Kasongo Yakusu, E.; Beeckman, H.; Bourland, N.; Doucet, J.L.; Sosef, M.S.M.; Hardy, O.J. The African timber tree Entandrophragma congoense (Pierre ex De Wild.) A.Chev. is morphologically and genetically distinct from Entandrophragma angolense (Welw.) C.DC. *Tree Genetics & Genomes* **2018**, *14*, 66. doi:10.1007/s11295-018-1277-6.

50. Daïnou, K.; Blanc-Jolivet, C.; Degen, B.; Kimani, P.; Ndiade-Bourobou, D.; Donkpegan, A.S.L.; Tosso, F.; Kaymak, E.; Bourland, N.; Doucet, J.L.; Hardy, O.J. Revealing hidden species diversity in closely related species using nuclear SNPs, SSRs and DNA sequences – a case study in the tree genus Milicia. *BMC Evolutionary Biology* **2016**, *16*, 259. doi:10.1186/s12862-016-0831-9.

51. Melo, W.A.; Freitas, C.G.; Bacon, C.D.; Collevatti, R.G. The road to evolutionary success: insights from the demographic history of an Amazonian palm. *Heredity* **2018**, *121*, 183–195. doi:10.1038/s41437-018-0074-1.

52. Loiseau, O.; Olivares, I.; Paris, M.; de La Harpe, M.; Weigand, A.; Koubínová, D.; Rolland, J.; Bacon, C.D.; Balslev, H.; Borchsenius, F.; Cano, A.; Couvreur, T.L.P.; Delnatte, C.; Fardin, F.; Gayot, M.; Mejía, F.; Mota-Machado, T.; Perret, M.; Roncal, J.; Sanin, M.J.; Stauffer, F.; Lexer, C.; Kessler, M.; Salamin, N. Targeted Capture of Hundreds of Nuclear Genes Unravels Phylogenetic Relationships of the Diverse Neotropical Palm Tribe Geonomateae. *Frontiers in Plant Science* **2019**, *10*. doi:10.3389/fpls.2019.00864.

53. Baker, W.J.; Savolainen, V.; Asmussen-Lange, C.B.; Chase, M.W.; Dransfield, J.; Forest, F.; Harley, M.M.; Uhl, N.W.; Wilkinson, M. Complete generic-level phylogenetic analyses of palms (Arecaceae) with comparisons of supertree and supermatrix approaches. *Systematic Biology* **2009**, *58*, 240–256. doi:10.1093/sysbio/syp021.

54. Rohland, N.; Reich, D. Cost-effective, high-throughput DNA sequencing libraries for multiplexed target capture. *Genome Research* **2012**, *22*, 939–946. doi:10.1101/gr.128124.111.

55. Johnson, M.G.; Gardner, E.M.; Liu, Y.; Medina, R.; Goffinet, B.; Shaw, A.J.; Zerega, N.J.C.; Wickett, N.J. HybPiper: Extracting Coding Sequence and Introns for Phylogenetics from High-Throughput Sequencing Reads Using Target Enrichment. *Applications in Plant Sciences* **2016**, *4*, 1600016. doi:10.3732/apps.1600016.

56. Katoh, K.; Standley, D.M. MAFFT multiple sequence alignment software version 7: Improvements in performance and usability. *Molecular Biology and Evolution* **2013**, *30*, 772–780. doi:10.1093/molbev/mst010.

57. Castresana, J. Selection of conserved blocks from multiple alignments for their use in phylogenetic analysis. *Molecular Biology and Evolution* **2000**, *17*, 540–552. doi:10.1093/oxfordjournals.molbev.a026334.

58. Sayyari, E.; Mirarab, S. Fast Coalescent-Based Computation of Local Branch Support from Quartet Frequencies. *Molecular biology and evolution* **2016**, *33*, 1654–1668, [1601.07019]. doi:10.1093/molbev/msw079.

59. Stamatakis, A. RAxML version 8: A tool for phylogenetic analysis and post-analysis of large phylogenies. *Bioinformatics* **2014**, *30*, 1312–1313, [arXiv:10.1093/bioinformatics/btu033]. doi:10.1093/bioinformatics/btu033.

60. Junier, T.; Zdobnov, E.M. The Newick utilities: high-throughput phylogenetic tree processing in the UNIX shell. *Bioinformatics* **2010**, *26*, 1669–1670. doi:10.1093/bioinformatics/btq243.

61. Rabiee, M.; Mirarab, S. SODA: Multi-locus species delimitation using quartet frequencies. *bioRxiv* **2019**, p. 869396. doi:10.1101/869396.

62. Yang, Z. The BPP program for species tree estimation and species delimitation. *Current Zoology* **2015**, *61*, 854–865. doi:10.1093/czoolo/61.5.854.

63. Brown, J.W.; Walker, J.F.; Smith, S.A. Phyx: Phylogenetic tools for unix. *Bioinformatics* **2017**, *33*, 1886–1888. doi:10.1093/bioinformatics/btx063.

64. Nguyen, L.T.; Schmidt, H.A.; Von Haeseler, A.; Minh, B.Q. IQ-TREE: A fast and effective stochastic algorithm for estimating maximum-likelihood phylogenies. *Molecular Biology and Evolution* **2015**, *32*, 268–274. doi:10.1093/molbev/msu300.

65. Kalyaanamoorthy, S.; Minh, B.Q.; Wong, T.K.; Von Haeseler, A.; Jermiin, L.S. ModelFinder: Fast model selection for accurate phylogenetic estimates. *Nature Methods* **2017**, *14*, 587–589. doi:10.1038/nmeth.4285.

66. Lanfear, R.; Calcott, B.; Kainer, D.; Mayer, C.; Stamatakis, A. Selecting optimal partitioning schemes for phylogenomic datasets. *BMC Evolutionary Biology* **2014**, *14*, 82. doi:10.1186/1471-2148-14-82.

67. Hoang, D.T.; Chernomor, O.; Von Haeseler, A.; Minh, B.Q.; Vinh, L.S. UFBoot2: Improving the ultrafast bootstrap approximation. *Molecular Biology and Evolution* **2018**, *35*, 518–522. doi:10.1093/molbev/msx281.

68. Guindon, S.; Dufayard, J.F.; Lefort, V.; Anisimova, M.; Hordijk, W.; Gascuel, O. New algorithms and methods to estimate maximum-likelihood phylogenies: Assessing the performance of PhyML 3.0. *Systematic Biology* **2010**, *59*, 307–321. doi:10.1093/sysbio/syq010.

69. Andermann, T.; Cano, Á.; Zizka, A.; Bacon, C.; Antonelli, A. SECAPR—a bioinformatics pipeline for the rapid and user-friendly processing of targeted enriched Illumina sequences, from raw reads to alignments. *PeerJ* **2018**, *6*, e5175. doi:10.7717/peerj.5175.

70. Li, H.; Durbin, R. Fast and accurate short read alignment with Burrows-Wheeler transform. *Bioinformatics* **2009**, *25*, 1754–1760, [1303.3997]. doi:10.1093/bioinformatics/btp324.

71. McKenna, A.; Hanna, M.; Banks, E.; Sivachenko, A.; Cibulskis, K.; Kernytsky, A.; Garimella, K.; Altshuler, D.; Gabriel, S.; Daly, M.; DePristo, M.A. The Genome Analysis Toolkit: a MapReduce framework for analyzing next-generation DNA sequencing data. *Genome research* **2010**, *20*, 1297–303. doi:10.1101/gr.107524.110.

72. Li, H.; Barrett, J. A statistical framework for SNP calling, mutation discovery, association mapping and population genetical parameter estimation from sequencing data. *Bioinformatics* **2011**, *27*, 2987–2993. doi:10.1093/bioinformatics/btr509.

73. Jombart, T.; Devillard, S.; Balloux, F. Discriminant analysis of principal components: A new method for the analysis of genetically structured populations. *BMC Genetics* **2010**, *11*, 94. doi:10.1186/1471-2156-11-94.

74. Jombart, T. Adegenet: A R package for the multivariate analysis of genetic markers. *Bioinformatics* **2008**, *24*, 1403–1405. doi:10.1093/bioinformatics/btn129.