Effects of dead conspecifics, hunger states, and seasons on the foraging behavior of
the purple urchin *Heliocidaris crassispina*

Dominic Franco C. Belleza[1], Yuuki Kawabata[1], Tatsuki Toda[2], Gregory N. Nishihara[3*]

[1]Graduate School of Fisheries and Environmental Science, Nagasaki University,
Nagasaki, Japan, 852-8521.

[2]Soka University, Tokyo, Japan, 192-8577.

[3]Organization for Marine Science and Technology, Institute for East China Sea
Research, Nagasaki University, Nagasaki, Japan, 851-2213.

*Corresponding author: G. N. Nishihara (greg@nagasaki-u.ac.jp)

Running title: Urchin foraging behavior modification

DFC Belleza ORCiD: https://orcid.org/0000-0002-0141-7495
Y Kawabata ORCiD: https://orcid.org/0000-0001-8267-5199
T Toda ORCiD: none
GN Nishihara ORCiD: https://orcid.org/0000-0003-2543-1301
15

16   Version date: March 10, 2020
17

ABSTRACT

The top-down component of trophic cascades exerts a powerful effect between

predator and prey relationships in an ecosystem. In aquatic environments, the

signals associated with predators and predation are used by prey as a cue to avoid

encountering predators when foraging for food. These cues are powerful enough to

control prey populations and indirectly protect primary producers. We evaluated

the effects of cues associated with predation on the purple urchin, *Heliocidaris*

crassispina and also examined the how starvation and season influenced the

effects. Using time-lapse photography, we conducted a series of manipulative and

in situ behavior experiments to determine foraging behavior patterns which

demonstrate behavior modification. The results suggest that starved urchins were

less sensitive to predation cues when compared to normally fed urchins. Field

experiments indicated that a proportion of fed urchins fled when exposed to a

predation cue (presence of a dead urchin), whereas all starved urchins remained

regardless of the cue, which complemented the results from the laboratory. Sea

urchin activity and feeding rates were lower in winter-spring than in summer-

autumn. We suggest that starvation has a large influence over the decision-making

process of sea urchins, and their response is affected by season due to metabolic

control. In general, the hunger response overrides predator avoidance behaviors

and exposes prey species to higher risks of predation.

Keywords: top-down control, trophic cascade, behavior modification

1. INTRODUCTION

Predation and resource availability control food webs in the ecosystem (Nielsen &

Navarrete 2004, Lynam et al. 2017). These interacting forces, together with

variabilities in environmental stress depend on the regulating effect (i.e., energy

allocation, expenditure, and transfer) they exert on the community of producers

and consumers (Menge & Sutherland 1987). Hairston et al. (1960) hypothesized

that populations of herbivores and the level of herbivory were generally controlled

by predation rather than by food supply (i.e., "green world" hypothesis) and

therefore the collapse of predator populations increased the likelihood of

herbivore domination (Estes & Palmisano 1974).

In temperate regions, macroalgal forests are an important coastal ecosystem

(Steneck et al. 2002, Smale et al. 2010, Langlois et al. 2011). Within the canopy, the

diversity of fish and invertebrate are high and they rely on the canopy for food and

refuge (Lowry & Pearse 1973, Holbrook et al. 199, Kamimura & Shoji 2009). The

sea urchin is a keystone species in marine forests, because they can overwhelm net

benthic primary production (Tuya et al. 2004). In ecosystems where apex predator

populations are intact, urchin populations are maintained through predation

(Tegner & Levin 1983, Pearse & Hines 1987, Sala & Zabala 1996, Sievers &

Nebelsick 2018). When predation pressure is removed, the urchin population leads

to overgrazing, eventually converting them into barrens. In general, barren areas

are characterized by low diversity and habitat complexity (Mangialajo et al. 2008).

Large canopy forming macroalgae are replaced by grazing resistant turf forming

macroalgae (Wright et al. 2005). However, as grazing pressure surpasses the

thresholds of the remaining primary producers, the community state eventually

transitions into a species-poor stable state (Steneck et al. 2002, Filbee-Dexter &

Scheibling 2014) that is easily maintained by a few urchins (Tuya et al. 2004,

Bonaviri et al. 2011). However, sustained human intervention or the recovery of

predator populations can revert barrens into a macroalgal-dominated state

(Blamey et al. 2013, Steneck et al. 2013).

Attempts to revert barrens into seaweed forests are not uncommon. Methods

include manual removal or destruction of urchins and other herbivores to

encourage natural recruitment of juvenile seaweeds (Yotsui & Maesako 1993,

Watanuki et al. 2010, Nanri et al. 2011), small-to-medium scale transplantation of

fertile seaweed thalli and mass dispersal of viable spores (Hernandez-Carmona et

al. 2000, Yoon et al. 2013, Ogata et al. 2016), and installment of artificial reefs

(Watanuki & Yamamoto 1990, Westermeier et al. 2013). Experimental evidence

has shown that human intervention may succeed and promote seaweed forest

recovery (Ling et al. 2010). However, maintaining restored algal forests is difficult

due to the uncontrolled recruitment of urchin that eventually achieve population

levels to establish dense feeding fronts (Lauzon-Guay & Scheibling 2007, Ling &

Johnson 2009).

Harnessing the effect of natural predators on prey to indirectly maintain the

population of primary producers may be a more practical solution (Schmitz et al.

2004). The direct reduction in the population of herbivores through consumption

is called density-mediated indirect interaction (DMII) while the modification of

prey behavior is called trait-mediated indirect interaction (TMII) (Schmitz et al.

2004). These interactions were observed in a variety of terrestrial and aquatic

ecosystems (Shurin et al. 2002). In the aquatic ecosystem, the effects of top-down

trophic cascade seem to be more prominent than in terrestrial ecosystems (i.e.,

marine benthos > marine plankton > terrestrial food web) (Strong 1992, Halaj &

Wise 2001, Shurin et al. 2002). The non-lethal effect of TMII may be comparable in

magnitude to that of DMII, because behavior change has population-wide effects,

whereas direct predation only affects the individual (Peacor & Werner 2001).

Historically, algal forests composed of large brown algae created dense expansive

belts around the coastline around Japan and supported a large diversity of

economically important fish and invertebrates (Uki et al. 1986, Kamimura & Shoji

2009). Presently, seaweed forests in Japan are undergoing a catastrophic decline

("isoyake") and the remaining seaweed forests are at high risk (Okuda 2008,

Haraguchi & Sekida 2008, Fujita 2010). The loss of seaweed forests has led to the

decline in coastal fisheries production (Kiyomoto et al. 2013). Efforts to revert the

decline in seaweed forests has produced numerous guidelines and technology,

however success is limited (Terawaki et al. 2003, Fujita 2010, Kuwahara et al. 2010,

Fujita 2015).

We focused on determining the impact of a non-lethal perceived threat on the

foraging behavior of the purple urchin *Heliocidaris crassispina* Agassiz.

Experimental studies have shown urchins to have complex foraging behaviors

(Vanderklift & Kendrick 2005, Kriegisch et al 2019) and that negative responses

from manipulative experiments ranged from strong (Campbell et al. 2001, Hagen et

al. 2002) to weak (Harding & Scheibling 2015). Here we use dead conspecifics as a

deterrent (Campbell et al 2001, Morishita & Barreto 2011) to explore the effects of

other factors such as the prey species' nutritional condition and season on the

urchin's decision-making process. The individual's nutritional condition (i.e. satiety

vs. starvation) has been known to modulate an individual's perception of risk

(Clark 1994) while the season is associated with reproductive phenology

(Agatsuma et al. 2000, Yatsuya & Nakahara 2004a). In this study we examine prey

behavior towards predation signals to better understand the role of predators in

indirectly maintaining the integrity of the seaweed bed ecosystem.

The following questions were addressed in this study: 1) How does *H. crassispina*

modify its foraging behavior when they encounter odor cues perceived to be a

threat and a non-threat? 2) How does urchin nutritional state influence their

tolerance to predation risk? and 3) How does season affect urchin response to

predation risk?

2. MATERIALS AND METHODS

2.1. Collection and maintenance of urchins and algae

Purple urchins (*Heliocidaris crassispina* Agassiz, 1864) were collected from the

coastal waters of Kashiyama Town, Nagasaki Prefecture, Japan. Urchins were

brought to the Institute for East China Sea Research, Nagasaki University,

approximately 3.7 km south of the collection site. Urchins were placed inside an

outdoor one-ton tank with a constant flow of sand-filtered seawater and aeration.

A Tidbit v2 (Onset Corp.) temperature logger monitored the ambient water

temperature. Urchins were fed *ad libitum* with an assortment of fresh algae

collected from Shihondo, Omura Bay, Nagasaki Prefecture, Japan. The feeding

experiments used *Sargassum patens* C. Agardh and were also collected from Omura

Bay. Stock *S. patens* were kept in a separate outdoor tank which received water

overflowing from the urchin stock tank. Urchins were acclimatized to laboratory

conditions for one week prior to the experiments. Experiments involving Summer-

Autumn and Winter-Spring seasons are hereafter referred to as S-A and W-S,

respectively.

Urchins were starved by placing selected individuals in a separate container with

no food for one week prior to experiments. This allowed for standardization of

their nutritional condition and to elicit a stronger hunger response prior to

experiments (Scheibling & Anthony 2001).

2.2.   Laboratory experiment 1: Urchin grazing rate

To test the hypothesis that the ambient *H. crassispina* grazing rate was influenced

by temperature and season, a flow-through rectangular tank (70 x 112 x 12 cm)

was prepared. Ten numbered containers (2.96 L) were set in the tank, separated

into two treatments. A feeding treatment that included urchins and algae and a

control which contained only algae. A continuous water supply (11 L min^{-1}) was

provided by an overhead perforated polyvinyl chloride (PVC) frame. A Tidbit v2

(Onset Corp.) temperature logger recorded ambient water temperature.

A total of 24 trials were conducted for both S-A (July-November 2018) and W-S

(February-April 2019). There were 4 control treatments and 6 feeding treatments

for each trial. Whole *S. patens* thalli were removed of epiphytes and other debris

and cut into portions. The cut portions were dried with paper towels and weighed

to the nearest 0.1 g to obtain initial fresh weight. Urchins were weighed to the

nearest 0.1 g while their horizontal test width was measured using a firm-joint

outside-caliper and a Vernier caliper to the nearest 0.01 cm. Cuttings and urchins

were haphazardly assigned to containers. A mesh-net frame was placed over the

tank to cover all of the containers to prevent urchins from escaping the container.

After 24 hours, the remaining uneaten algae were collected, dried with paper

towels and re-weighed to obtain final fresh-weight. Urchin feeding rate was the

difference between the final and initial weight with units g algae day^{-1}.

2.3. Laboratory experiment 2: Effect of positive and negative chemosensory cues

on urchin foraging behavior

The experiment was designed to test the hypothesis that urchins will modify

foraging behavior when exposed to cues coming from dead conspecifics compared

to controls (no odor cues).

The experiment used a flow-through tank similar to that of the previous

experiment. However, water was supplied at a steady rate of 2.5 L min^{-1} through a

hose. The hose was placed so that water flowed through the floor, in the middle of

the tank. Water exited the tank through a 6 cm diameter hole in the tank wall,

located 3 cm above the tank floor. Five concentric rings, 5 cm apart were marked

on the tank floor around the water supply. The outermost ring defined the edges of

the region of interest (ROI) where urchin behavior was recorded with a time-lapse

camera (GoPro, Hero 4). At the center of the ring and above the hose, a perforated

PVC cap was placed.

The experiment was started by placing the test urchin 10 cm from the center. The

camera was mounted 40 cm above the ROI and the field of view (FOV) included the

entire tank. The images were recorded every 30 sec and experiment was conducted

for 1 hr. Four treatments were defined, 1) a control (no dead urchin or algae; no

chemosensory cue) 2) an algae treatment (positive chemosensory cue), 3) a dead

urchin treatment (negative chemosensory cue), and 4) a dead urchin and algae

interaction treatment (combined chemosensory cues). To expose the test urchin to

the treatment effect, algae were attached to the top of the PVC cap with clips and

the dead urchin was placed within a mesh bag below the cap. Therefore, water

flowing through the hose and through the cap ensured that chemosensory cues

from the treatment would be dispersed outwards across the tank. The experiment

was conducted on both urchin nutritional conditions (i.e., starved and fed) and in

W-S and S-A.

An opaque plastic sheet covered the entire experimental apparatus to remove all

ambient light. However, below the sheet, a red LED lamp (ISL-150X150, CCS Inc.)

provided enough light to record images while minimizing light disturbance of the

urchins (Flukes et al. 2012). After every trial, the test urchins were removed, and

the experiment chamber was rinsed thoroughly with freshwater and seawater to

eliminate chemical cues from the previous experiment.

Experiments were conducted in the S-A (August-November 2018) and W-S (April-

May 2019). 111 individuals were used for the S-A experiment and 88 individuals

were used for the W-S experiment.

For each trial, the time-lapse images were concatenated into an mpeg-4 video

using FFmpeg (FFmpeg Developers 2018) at a frame rate of 10 frames per second

(fps). Videos were analyzed with Tracker (ver.5.0.6; Brown 2018) to determine the

movement pattern of the urchins. Each video frame (i.e., image) was counted as

one event of a particular behavior (Table 1). The x and y coordinates of the sea

urchin was analyzed to determine the time spent by an urchin performing a

particular behavior (minutes) as well as sea urchin movement speed (cm min^{-1}).

2.4. Cue dispersal rate

The chemical cue plume was visualized and quantified using a Fluorescein tracer

solution as a proxy. A 3 mm diameter hose was attached to the tank floor so that

the tracer was injected below the PVC cap and perpendicular to the water flow. The

50 ml of tracer was injected at a rate of 1.6 ml sec^{-1}. The tracer concentration was

2% Fluorescein. A one-hour video recorded the dispersal of tracer. Three trials

were conducted per treatment including control. The time for the tracer to reach

the 10 and 20 cm ring was recorded and analyzed to determine if there were any

1   differences among treatments.

2.5. Light measurement

3   The spatial homogeneity of the red light provided by the LED lamp was also

4   assessed. Light was measured using five light loggers (HOBO MX2202 Temp/Light,

5   Onset Corp.) that were placed on each ring to form a line. After the initial

6   measurement, the line was rotated 30 degrees, for a total of four times. At every

7   rotation, light was measured for one hour.

2.6. Field experiment: Effect of positive and negative chemosensory cues on

urchin foraging behavior in the field

To examine the effects of chemosensory cues by food and dead conspecifics on sea

urchin behavior under natural conditions, we prepared a site that was located at a

depth of 4-5 m in a barren rocky area adjacent to natural stands of *Sargassum*

macrocarpum in Arikawa Bay (32.988014 °N, 129.118638 °E), Nakadorijima

Island, Nagasaki, Japan. A 2 m^2 plot of flat rocky substrate was selected. For each

experimental trial, a 2 m tall slotted angle-bar tripod frame with an approximately

1 m^2 plan area was deployed. A time-lapse camera (TLC200 PRO, Brinno, Taiwan)

enclosed in a custom acrylic housing was secured to the top of the frame. Images

were taken every 30 s for a total of 3 hours and stored as a video with a frame rate

of 15 fps. The experiment was during slack tide, when the tidal current was

negligible. A velocity logger (Compact-EM, Alec Electronics Co.) and a water level

logger (HOBO U20-001, Onset Corp.) were deployed 1 m from the experimental

plot to record hydrodynamic conditions during the experiment.

A weighted plastic cage was placed in the middle of the 1 m^2 experimental plot.

Drift algae (i.e. *Sargassum* spp., *Dictyopteris* spp.) common during the experiment

period was collected and clipped outside of the plastic cage. Test sea urchins were

placed in direct contact with the algae until they attached. This was considered as

the control. In a separate treatment, a dead urchin was added inside the cage to

determine whether urchins would be repelled by a dead conspecific. The

experiments were conducted on both urchin nutritional conditions.

Video was analyzed with Tracker (ver. 5.0.6) (Brown 2018) to track the urchins,

however the tripod attracted small fish, which occluded the field of view.

Additionally, during a number of days, the area experienced relatively high waves,

which vibrated the tripod that resulted in poor quality images. Thus, only the initial

and final position (stay or flee) of the urchin was noted after the 3 hr experiment

period.

2.7. Data analyses

All statistical analyses were done using R version 3.6.1 (R Development Core Team

2019) and all Bayesian inference was done with Stan (Stan Development Core

Team 2019) through the brms (Bürkner 2017) and rstanarm packages (Goodrich

et al. 2018). Four Markov chains generated 2000 samples per chain. All posterior

distributions were examined visually to evaluate the convergence of the Markov

chains and posterior predictive ability.

The urchin behavior in the laboratory experiments were analyzed with a hurdle-

negative binomial model (Eq. 1),

	$y = (1 - \pi)\text{NegBin}(0, \theta) + \pi\text{NegBin}(\mu, \theta)$	(1)

$$\mu = x\beta$$

$$log \frac{\boldsymbol{\pi}}{1 - \boldsymbol{\pi}} = \boldsymbol{x\alpha}$$

where, y is a vector of observations, and in this case the number of occurrences

for a behavior during the 1-hour observation period. π is a vector of probabilities

for non-zero values, α and β are vectors of coefficients for a model including all

treatment interactions. x is a matrix of factors that include all treatment

interactions. The number of occurrences is assumed to follow a negative binomial

distribution, with a vector of locations μ and a scale θ. The main treatments are

the presence or absence of algae and the dead urchin, the nutritional state of the

test urchin, season (i.e., S-A or W-S), and the type of behavior excluding the "none"

behavior (Fletcher et al. 2005) (Table 1).

In the case of the sea urchin speed and time spent per behavior, a hurdle-gamma

model was applied. The structure of the model is similar to Eq. 1, however rather

than a negative binomial distribution a gamma distribution is assumed. In this case

$y = (1 - \pi)\,\Gamma\,(0, \theta) + \pi\,\Gamma\,(\mu, \theta)$. For more details on the merits of the hurdle

model, see Ridout et al. 1998.

The β coefficients of the all hurdle models were given weakly informative

Student's t-distributions as prior distributions, with 3 degrees of freedom, a

location of 0, and a scale of 1. The α coefficients were given logistic distributions

as priors with a location of 0 and a scale of 1. The prior for θ was a Γ

distribution with a shape and scale of 0.01.

The field experiments were analyzed with a simpler binomial generalized linear

model with a random intercept for the tidal state (Eq. 2).

	$$y = Binom(n, \pi)$$ $$log \frac{\pi}{1 - \pi} = x\gamma + \delta$$	(2)

9   In Eq. 2, the y is the vector of observations, n is the vector of total trials, and π

10   is the vector of probabilities. x is the matrix of treatments, which in this case is a

11   linear combination of the nutritional state of the test urchin (i.e., starved or fed)

1 and the presence or absence of the dead urchin. δ is a random intercept for tidal

2 state. The vector of coefficients is γ. The prior distributions for the coefficients and

3 random intercept was a Student's t-distribution with 3 degrees of freedom, a

4 location of 0, and a scale of 5.

3. RESULTS

3.1. Laboratory experiment 1: Urchin grazing rate

7 The results revealed differences in feeding rates between seasons. Sea urchins had

8 higher mean feeding rates in the S-A (1.3 ± 0.1 g algae urchin^{-1} day^{-1}) (Table 2: A).

9 Conversely, sea urchin feeding rates decreased by about 38% during the W-S (0.8 ±

10 0.1 g algae urchin^{-1} day^{-1}). There was a 23% difference between seasons in the

11 mean maximum water temperature of the experiment conditions (Figure 1).

3.2. Laboratory experiment 2: Effect of positive and negative chemosensory cues

on urchin foraging behavior

3.2.1. Frequency of behaviors

15 The time-lapse experiment showed that the counts of the four behavior types

varied widely across sea urchin condition, season, and treatment (Supp. Figure 1).

The occurrence of the immobile behavior "none" occurred more in W-S (starved:

15 urchins, fed: 21 urchins) than in S-A (starved: 16 urchins, fed: 15 urchins). The

occurrence of this behavior in W-S represents 40.9% of the sea urchins used in that

season while this behavior represented about 27.9% of the total sea urchins used

in the S-A.

Sea urchins were generally more active in S-A than in W-S season. For interaction

events, this behavior was more frequent in S-A and in starved sea urchins,

occurring in 6% and 5% of all experiment trials, respectively. For the W-S,

interaction occurred only in 2% of trials while only occurring in 4% of fed urchins.

The model predictions on the effect of the dead urchin on interaction events can be

observed when comparing algae only (Figure 2C, G) and combined chemosensory

cues treatment (Figure 2D, H). Predictions show that when a dead urchin was

present together with the algae, fed urchins had decreased interaction counts of up

to 64.8% and 72.3% during S-A and W-S, respectively. For starved urchins, they had

a 14% increase and a 49% decrease in interaction counts for S-A and W-S,

respectively. In W-S, starved urchins also had a higher proportion of immobile

individuals. At the same time, the presence of the dead urchin also increased the

number of events outside of the ROI across both seasons for fed (44.8% and 39.2%

for S-A and W-S, respectively) and starved (93.2% and 54.1% for S-A and W-S,

respectively). Overall, fed urchins seem to be sensitive to the chemical cues from

dead urchins while starved urchins were more tolerant during the summer but

were also less active in winter.

3.2.2. Time spent per behavior

The time-lapse experiment demonstrated the ability of the treatments, condition

and the season to influence the sea urchin's allocated time performing a specific

behavior (Supp. Figure 2). Overall, 34% of sea urchins spent the entire 1-hr

experiment period immobile. Of all the immobile urchins, 42% were immobile

during the winter-spring season while 27.9% represented the summer-spring

season experiments.

The model predictions on the effects of a dead urchin chemosensory cues on time

spent per behavior are shown in Figure 3. The predictions show that compared to

algae treatment (Figure 3C, G), the presence of a dead urchin (Figure 3D, H)

reduced mean interaction times for fed urchins by about 5.57 and 4.07 minutes

during S-A and W-S, respectively. For starved urchins, there was a mean increase by

about 1.7 min and a decrease of about 5.14 min for S-A and W-S, respectively. The

time spent outside the ROI also increased across both seasons for fed (10.2 and 5.2

minutes for S-A and W-S, respectively) and starved (5.7 and 5.07 minutes for S-A

and W-S, respectively) urchins. The decrease in time spent for interaction by fed

urchins and the relative increase in the proportion of time spent outside indicates

that urchins were repelled by the presence of the dead urchin together with the

algal food. Conversely, the 11.4% increase in time spent by the starved urchins

interacting with the algae despite the presence of the dead urchin indicates that

hunger was able to increase its tolerance to signals of predation.

3.2.3. Urchin movement speed

The time-lapse experiment revealed urchin movement speeds varied across the

treatments depending on their condition and season. It was noted that even when

the urchins were exhibiting the behavior "none", minute movement speeds were

recorded as urchins were shuffling in place to within the 3cm limit (Table 1). The

urchins exhibited greater speeds when they were interacting with the center of the

ROI in the presence of a dead urchin while overall, fed urchins had higher

movement speeds relative to the starved urchins (Supp Figure 3).

The model predictions on urchin speed (cm min^{-1}) indicates that urchins tend to

move at a different pace depending on the presence or absence of a dead urchin

(Figure 4D, H) than with algae only (Figure 4C, G). When a dead urchin was

present, interaction speeds for fed urchins typically decreased by 1.48 and 0.95 cm

min^{-2} during S-A and W-S, respectively. The opposite was found for starved urchins

because their speeds while interacting with the algae increased by 1.25 and 5.13

cm min^{-1} for S-A and W-S, respectively when a dead urchin was present. However,

when outside the ROI, fed urchins had increased speeds by up to 2.39 cm min^{-2}

during S-A, while showing decreased speeds by up to 1.39 cm min^{-2} during W-S.

Starved urchins still show the same pattern of high speeds outside the ROI by up to

9.18 and 5.26 cm min^{-2} for S-A and W-S, respectively. Examining urchin speeds

show that signals of predation may cause considerable stress to *H. crassispina* as

indicated by the relatively high movement speeds even when outside the ROI.

3.2.4. Cue dispersal rate

The results from the experiment on the rate of spread of the Fluorescein tracer dye showed high variation among the trials (Table 2: B2). The time it took for the tracer to reach the 10 cm mark was modelled since it represented the area where the urchin would first encounter the chemosensory cues coming from the center of the ROI. The control, with nothing beneath and above the treatment container, took the least amount of time at 13.5±5.4 s (mean ± standard deviation). The treatments could be ranked from those that took the least amount of time to the greatest amount of time and resulted in an order of control, algae effect, dead urchin and algae interaction effect, and dead urchin effect. The high standard deviations for all treatments was due to a low sample size (3 trials per treatment). However, high variations between trials among the algae, dead urchin and dead urchin and algae treatments suggested that the variation was associated with the size of the dead urchin or the density of the algae used as treatment for the experiment.

3.2.5. Light measurement

The results of the light experiment showed that the highest light intensity was at

the center of the experiment chamber, directly over the treatment container, at 3.0

± 0.1 lux (Table 2: B1). The light intensity slowly decreased outwards from the ROI,

with the 10 cm area receiving approximately 2.1 ± 0.3 lux, and the outer edge

receiving only 0.4 ± 0.3 lux. Despite the uneven light distribution across the

experiment chamber, the camera was able to record the sea urchin behavior in

enough detail to allow later analysis using the tracking software. More importantly,

sea urchin activity was not discernibly affected by the light intensity throughout

the experiment.

3.3.   Field experiment: Effect of positive and negative chemosensory cues on

urchin foraging behavior in the field

The field experiment showed that sea urchin condition produced discrete

responses between starved and fed urchins to the presence of dead conspecifics

adjacent to an available food source. Of the 20 sea urchins used in the experiment,

all 10 starved sea urchins (100% of the starved condition) stayed and remained in

contact with the treatment cage. For the fed sea urchins, only 3 stayed (30%) while

7 fled from the treatment cage (70%). Of the 7 sea urchins which fled, 4 sea urchins

(40%) were from the treatment which contained the dead urchin. The binomial

model predictions suggested a strong link between urchin condition and outcome

of behavior (Figure 5).

3.4.   Field experiment: Ambient environment conditions

The data recorded by the temperature, current speed, and depth loggers indicate

the specific environmental conditions during field experiments (Supp. Figure 4). Of

the 20 trials conducted, 11 experiments were conducted during low slack tide

while 9 were conducted during high slack tide. In general, mean temperatures and

mean current speeds were higher during low tide relative to high tides.

4.   DISCUSSION

4.1.   Factors affecting sea urchin behavior patterns

Time-lapse photography revealed high variations in the frequency of behavior

occurrences and movement patterns among individual *Heliocidaris crassispina*

across the different treatments. This lack of distinct directionality was also

reported for other urchin species (Flukes et al. 2012, Kriegisch et al. 2019). Despite

these observations, we found discrete behavior patterns between starved and fed

H. crassispina, which suggests that the tolerance to a perceived risk increased when

sea urchins were deprived of food. For starved urchins, there was a greater

proportion of urchin interaction events and increased interaction time with the

seaweed despite the presence of a dead urchin. Fed urchins exhibited predator

avoidance behaviors, observed as decreased interaction and increased occurrences

of behaviors spent outside the ROI when a dead urchin was placed together with

the seaweed.

Studies on predator-prey relationships highlight the "Hobson's choice" (i.e., face

the risk of predation or starve) dilemma all prey species face upon venturing out

from the safety of their refuge when they forage for food (Clark 1994). Ultimately,

the decisions prey species make lean towards optimizing the trade-off to their

advantage by minimizing risk while maximizing benefits. However, intrinsic (i.e.,

reproductive condition, and hunger level) and extrinsic (i.e., temperature, light, and

salinity) factors also play an important role in affecting decision-making processes

for aquatic organisms. For example, the effects of starvation in urchins not only

impacted their energy reserves but also their gut and gonad indices (Lawrence

1970). A study on the effects of starvation on *H. crassispina* and *Hemicentrotus*

pulcherrimus showed that gut clearance was achieved in 3 days for *H. crassispina*

and 6 days for *H. pulcherrimus*. Additionally, *H. pulcherrimus* survived a maximum

of 49 days without food albeit negatively impacting gut and gonad indices (Kaneko

et al. 1981 in Agatsuma et al 2013). In our study, the one-week starvation period

induced a hunger response which appeared to override predator avoidance

behaviors. The starved urchins, which had an energy deficit, were willing to accept

greater risks by feeding longer and more frequently in the presence of a dead

urchin to increase energy reserves, hence supporting the asset protection principle

(Clark 1994). It should be noted that the microcosm experiment utilized a

relatively small chamber which may have allowed the saturation of sea urchin

effluents faster and thus increase the urchin responses artificially. Furthermore,

the manipulation of sea urchin condition by starvation as done in our study does

not fully mimic conditions in the field. Recall that sea urchins are generalist algal

feeders (Vadas 1977) and have flexible dietary preferences and are omnivorous

(Rodriguez-Barreras et al. 2015). *H. crassispina* in seaweed bed habitats generally

had higher gonad indices and were larger in size compared to urchins collected

from a habitat dominated by *Corallina* spp. (Yatsuya & Nakahara 2004a). In the

barrens, they were more cryptic and switched to feeding on a mixed diet composed

of drift *Sargassum* and calcareous algae (Yatsuya & Nakahara 2004b). Other signs

of food limitation manifested themselves as differences in morphometric changes

such as smaller than average test size and longer jaw lengths to maximize grazing

efficiency (Pedersen & Johnson 2008).

A recent study utilizing starved and satiated *Strongylocentrotus droebachiensis*

showed that both sea urchin groups did not react adversely to the presence of a live

nearby predator (*Cancer borealis*) (Harding & Scheibling 2015). Their findings

indicate that the olfactory cues coming from live predatory crabs did not reduce

urchin foraging behavior in the laboratory and in the field. In contrast, studies on

chemical alarm cues showed dead conspecifics and chemically-labelled predators

had a strong adverse effect on urchin behavior (Campbell et al. 2001, Morishita &

Barreto 2011). Specifically, the urchins distinctly avoided waters conditioned with

gut, coelomic, and gonad homogenates, which were the materials most likely to be

exposed when a predator breaks an urchin's test (Campbell et al. 2001).

In our study, instead of completely avoiding the source of the chemical cues coming

from the dead urchin treatment, some urchins actually approached the dead urchin

treatment. In small prey species of fish, this behavior is known as predator

"inspection", and had distinct importance for prey species because this functions as

a learning tool to enable naive prey to associate predators with danger (Magurran

& Girling 1986). For urchins, since olfaction occurs when odor molecules reach

receptors in their tube-feet, predator inspection may need to occur at close range

as odor molecules increase in concentration. The next time they encounter familiar

chemical cues relating to risk of predation, they may better assess the motivation

of the predator and the relative risk of an impending predation event (Clark 1994).

Furthermore, for urchins living in urchin barrens, it may be possible that these

urchins had reached large population sizes due to the absence of their natural

predators. The absence of predators meant that it was likely that they were naive

and had little chance in encountering chemical cues relating to predation.

When comparing sea urchin behavior patterns across seasons, we found that there

1 was a discrete pattern observed between the S-A and W-S experiments. In general,

2 sea urchins were more active and had higher movement speeds during S-A

3 compared to sea urchins used in W-S. The greater decrease in the interaction

4 frequency and interaction time in W-S for starved urchins was attributed to lesser

5 urchins interacting with the treatment as well as more urchins moving outside.

6 Interestingly, urchin speed was predicted to be highest in W-S when starved

7 individuals were exposed to dead urchins together with food. This was likely to be

8 an evasive behavior in response to the scent of the dead urchin since the

9 proportion of outside behaviors and movement speeds also increased. At the same

10 time, when only food was present, starved urchins interacted with the algae

11 longer and moved slower, indicating a stronger intent to feed, compared to fed

12 urchins (Figure 3). Their level of activity was also reflected in their feeding rates as

13 urchins in August had the highest average feeding rates while urchins in February

14 had the lowest rates. This was a similar pattern found among cold-water urchin

15 species where temperature was one of the main drivers of metabolic activity

16 (Agatsuma et al. 2000, Brockinton & Clarke 2001).

17 Studies on the reproductive biology of *H. crassispina* showed that this species had a

distinct seasonal cycle in terms of gonadal development and maturation. In

Nagasaki, a study on the reproductive patterns of *H. crassispina* (Yamasaki &

Kiyomoto 1993 in Agatsuma et al 2013) has found that this species spawns during

the months of July to August while recovery period was from September to January.

The rest of the following months were dedicated to growth and maturation of the

gonads. This pattern was also similar with studies elsewhere in Japan (Kyoto:

Yatsuya & Nakahara 2004a, Oga Peninsula: Feng et al. 2019) and in Korea (Yoo et al.

1982). In Hong Kong, a 7 to 8 month spawning period was recorded. This relatively

long spawning period was represented by two distinct spawning events in May-

June and September-October (Urriago et al. 2016). After every spawning event,

urchins experienced an abrupt decrease in gonad indices as well as lipid and fatty-

acid profiles (Martinez-Pita et al. 2010, Diaz de Vivar et al. 2019). This lipid and

nutrient deficient state indicated that the urchins were in a low nutritional

condition (Lawrence 1970). Urchins compensated by increasing their feeding rates

beginning from the end of summer until next spring, coinciding with winter

macroalgal blooms (Kaehler & Kennish 1996). Increasing feeding rates from

summer ensured the accumulation of energy to support gonadal growth and

maturation as reflected from the biochemical composition and other intrinsic

gonad properties (Rocha et al. 2019). Hence, the rise in summer metabolic activity

in urchins was only partially explained by temperature but was likely

predominantly driven by feeding, growth and reproduction (Brockington & Clarke

2001).

4.2. Field experiment

The results provide evidence of urchin condition affecting the strength of behavior

modification in the field. Compared to fed sea urchins, all starved urchins stayed

regardless of the presence or absence of a dead urchin. The result of our

experiment was in contrast with the field experiment using live cancrid predators

where only a 6% flee response rate was recorded (Harding & Scheibling 2015).

Few studies have previously investigated effects of predation risk cues on prey

species in the field because of the inherent difficulty of controlling for local flow

conditions. The data recorded from the field shows high variability in flow speeds

as well as temperature between low and high tides. A laboratory study simulating

the flow of chemical odor plumes in turbulent conditions suggests that the success

of odor-guided navigation was greatly dependent on dilution and degree of shear-

induced mixing of odor signals (Webster & Weissberg 2001). This was particularly

true for small benthic invertebrates because sampling the water for odor molecules

occurs at a relatively faster rate (Zimmer & Butman 2000) but at a lower height

relative to the substrate (Smee & Weissberg 2006). Furthermore, organisms

attempting to orient themselves relative to the direction of the odor-plume would

find it challenging because odor dispersal occurs as intermittent odor packets

interspersed with clean water (Finelli et al. 1999). In the present study, the sea

urchins would have had no problem detecting the odor from the dead urchin and

seaweed because they were placed in direct contact with the treatment cage at the

start of the experiment, unlike in the laboratory experiment. Although

concentrations of urchin effluent were not tested, it may be plausible that the

immediate area surrounding the treatment cage would have been saturated with

the dead urchin effluent. The fleeing response of some of the fed sea urchins

appear to be a behavior related to minimizing predation risk in lieu of feeding

opportunity. However, with only a short 3-hr experiment period, we were not able

to observe how the starved urchins would behave once they have adequately fed

on the seaweed or how long the dead urchin effluents remained effective. Hence,

for future studies, we propose a longer observation time for urchin behaviors and

identification of the components responsible for urchin alarm response and their

maximum length of efficacy.

5. CONCLUSIONS

The behavioral responses of *Heliocidaris crassispina* in our experiments appear to

encompass all classifications in the prey behavior types proposed by Fraser &

Huntingford (1986). Our results showed that *H. crassispina* foraging behavior was

flexible and was able to assess and adjust accordingly to the presence of chemical

cues associated with predation. To some extent, our results support the idea that

the type of season and phenology appear to modulate urchin behavior and foraging

activity (Luttberg et al. 2003). Our experiments also demonstrate that the presence

of a dead urchin does not prevent live urchins from interacting with the seaweed

but instead decreases the interaction frequency, length of interaction time and

increases movement speeds, indicating escape behaviors. Furthermore, starved

urchins seemed to be more insensitive and indifferent to predation cues. The 100%

stay response from starved urchins despite the presence of a dead urchin in the

field experiment further reinforced our hypothesis. Our findings suggest that

animal condition was a key determinant in its decision-making process and that

their level of hunger may override behaviors associated with predator avoidance.

This puts them at a disadvantage as starved animals feed more boldly, exposing

themselves further to the dangers of predation.

6. ACKNOWLEDGEMENTS

We thank Yukio Inoue, Koichi Osaki and Kenjiro Hinode for assistance in the

collection of seaweed samples. We also thank Mikiya Hidaka and Takeshi Urae for

assistance in the collection of sea urchins. This study was supported by the Grant-

in-Aid for Scientific Research (C-#40508321) from the Japan Society for the

Promotion of Science (JSPS) and by the Japanese Ministry of Education, Culture,

Sports, Science and Technology (MEXT).

7. REFERENCES

Agatsuma Y, Nakata A, Matsuyama K (2000) Seasonal foraging activity of the sea

urchin *Strongylocentrotus nudus* on coralline flats in Oshoro Bay in south-

western Hokkaido, Japan. Fish Sci 66: 198-203.

Agatsuma Y (2013) Chapter 30: *Hemicentrotus pulcherrimus*, *Pseudocentrotus*

depressus, and *Heliocidaris crassispina*. In Lawrence, J.M. (2013) (Ed.), Sea

urchins: Biology and Ecology, Third Edition. Developments in Aquaculture

and Fisheries Science, vol 38, Elsevier, United Kingdom, p 461-472.

Blamey LK, Plaganyi EE, Branch GM (2013) Modeling a regime shift in a kelp forest

ecosystem caused by a lobster range expansion. Bull Mar Sci 89(1): 347-

375.

Brockinton S, Clarke A (2001) The relative influence of temperature and food on

the metabolism of a marine invertebrate. J Exp Mar Biol Ecol 258: 87-99.

Clark CW (1994) Antipredator behavior and the asset-protection principle.

Behavioral Ecology 5(2): 159-170.

Bonaviri C, Ferenandez TV, Fanelli G, Badalamenti F, Gianguzza P (2011) Leading

role of the sea urchin *Arbacia lixula* in maintaining the barren state in

southwestern Mediterranean. Mar Biol 158: 2505-2513.

Bürkner PC (2017) brms: An R package for Bayesian multilevel models using stan. J

Stat Softw 80(1): doi: 10.18637/jss.v080.i01.

Campbell AC, Coppard S, D'Abreo C, Tudor-Thomas R (2001) Escape and

aggregation responses of three echinoderms to conspecific stimuli. Biol Bull

201: 175-185.

Diaz de Vivar ME, Zarate EV, Rubilar T, Epherra L, Avaro MG, Sewell MA (2019)

Lipid and fatty acid profiles of gametes and spawned gonads of *Arbacia*

dufresnii (Echinodermata: Echinoidea): Sexual differences in the lipids of

nutritive phagocytes. Mar Biol 166(96): doi.org/10.1007/s00227-019-

3544-y.

Estes JA, Palmisano JF (1974) Sea otters: Their role in structuring nearshore

communities. Science 185(4156): 1058-1060.

Feng WP, Nakabayashi N, Narita K, Inomata E, Aoki MN, Agatsuma Y (2019)

Reproduction and population structure of the sea urchin *Heliocidaris*

crassispina in its newly extended range: The Oga Peninsula in the Sea of

Japan, northeastern Japan. PLoS ONE 14(1):

doi.org/10.1371/journal.pone.0209858.

FFmpeg Developers (2018) FFmpeg version 4.1.1. [software]. Available from

http://www.ffmpeg.org/.

Filbee-Dexter K, Scheibling RE (2014) Sea urchin barrens as alternative stable

states of collapsed kelp ecosystems. Mar Ecol Prog Ser 495: 1-25.

Finelli CM, Pentcheff ND, Zimmer-Faust RK, Wethey DS (1999) Odor transport in

turbulent flow: constraints on animal navigation. Limnol Oceanogr 44(4):

1056-1071.

Fletcher D, Mackenzie D, Villouta E (2005) Modelling skewed data with many

zeros: A simple approach combining ordinary and logistic regression.

Environ Ecol Stat 12: 45-54.

Flukes EB, Johnson CR, Ling SD (2012) Forming sea urchin barrens from the inside

out: an alternative pattern of overgrazing. Mar Ecol Prog Ser 464: 179-194.

Fraser DF, Huntingford FA (1986) Feeding and avoiding predation hazard: the

behavioral response of the prey. Ethology 73: 56-68.

Fujita D (2010) Current status and problems of isoyake in Japan. Bull Fish Res

Agency 32: 33-42.

Fujita D (2015) Toward the revision of isoyake taisaku guideline. Fish Eng 51(3):

259-262.

Goodrich B, Gabry J, Ali I, Brilleman S (2018) rstanarm: Bayesian applied

regression modeling via Stan. http://mc-stan.org/.

Hagen NT, Andersen A, Stabell OB (2002) Alarm responses of the green sea urchin,

Strongylocentrotus droebachiensis, induced by chemically labelled

durophagous predators and simulated acts of predation. Mar Biol 140: 365-

374.

Hairston NG, Smith FE, Slobodkin LB (1960) Community structure, population

control and competition. Am Nat 94(879): 421-425.

Halaj J, Wise DH (2001) Terrestrial and trophic cascades: How much do they

trickle? Am Nat 157(3): 262-281.

Haraguchi H, Sekida S (2008) Recent changes in the distribution of Sargassum

species in Kochi, Japan. Kuroshio Sci 2(1): 41-46.

Harding APC, Scheibling RE (2015) Feed or flee: Effect of a predation risk cue on

sea urchin foraging activity. J Exp Mar Biol Ecol 466: 59-69.

Hernandez-Carmona G, Garcia O, Robledo D, Foster M (2000) Restoration

techniques for *Macrocystis pyrifera* (Phaeophyceae) populations at the

southern limit of their distribution in Mexico. Bot Mar 43: 273-284.

Holbrook SJ, Carr MH, Schmitt RJ, Coyer JA (1990) Effect of giant kelp on local

abundance of reef fishes: The importance of ontogenetic resource

requirements. Bull Mar Sci 47(1): 104-114.

Kaehler S, Kennish R (1996) Summer and winter comparisons in the nutritional

value of the marine macroalgae from Hong Kong. Bot Mar 39: 11-17.

Kamimura Y, Shoji J (2009) Seasonal changes in the fish assemblage in a mixed

vegetation area of seagrass and macroalgae in the central Seto Inland Sea.

Aquacult Sci 57(2): 233-241.

Kaneko I, Ikeda Y, Ozaki H (1981) Biometrical relationships between body weight

and organ weights in freshly sampled and starved sea urchins. Bull Jap Soc

Fish 47: 539-537.

Kiyomoto S, Tagawa M, Nakamura Y, Horii T, Watanabe S, Tozawa T, Yatsuya K,

Yoshimura T, Tamaki, A (2013) Decrease of abalone resources with

disappearance of macroalgal beds around the Ojika Islands, Nagasaki,

Southwestern Japan. J Shellfish Res 32(1): 51-58.

Kriegisch N, Reeves SE, Flukes EB, Johnson CR, Ling SD (2019) Drift-kelp

suppresses foraging movement of overgrazing urchins. Oecol.

https://doi.org/10.1007/s00442-019-04445-6

Kuwahara H, Hashimoto O, Sato A, Fujita D (2010) Introduction of isoyake recovery

guideline (Fisheries Agency, Japan). Bull Fish Res Agency 32: 51-60.

Langlois TJ, Radford BT, Van Niel KP, Meeuwig JJ, Pearce AF, Rousseaux CSG,

Kendrick GA, Harvey ES (2011) Consistent abundance distributions of

marine fishes in an old, climatically buffered, infertile seascape. Glob Ecol

Biogeogr 21(9): 886-897. http://dx.doi.org/10.1111/j.1466-

8238.2011.00734.x.

Lauzon-Guay JS, Scheibling RE (2007) Behaviour of sea urchin *Strongylocentrotus*

droebachiensis grazing fronts: food mediated aggregation and density-

dependent facilitation. Mar Ecol Prog Ser 329: 191-204.

Lawrence JM (1970) The effects of starvation on the lipid and carbohydrate levels

of the gut of the tropical sea urchin *Echinometra mathaei* (de Blainville). Pac

Sci, 24: 487-489.

Ling SD, Johnson CR (2009) Population dynamics of an economically important

range-extender: kelp beds versus sea urchin barrens. Mar Ecol Prog Ser 374:

113-125.

Ling SD, Ibbott S, Sanderson JC (2010) Recovery of canopy-forming macroalgae

following removal of the enigmatic grazing sea urchin *Heliocidaris*

erythrogramma. J Exp Mar Biol Ecol 395: 135-146.

Lowry LF, Pearse JS (1973) Abalones and sea urchins in an area in habited by sea

otters. Mar Biol 23: 213-219.

Luttberg B, Rowe L, Mangel M (2003) Prey state and experimental design affect

relative size of trait- and density-mediated indirect effects. Ecology 84(5):

1140 -1150.

Lynam CP, Llope M, Möllmann C, Helaouët P, Bayliss-Brown GA, Stenseth NC (2017)

Interaction between top-down and bottom-up control in marine food webs.

Proc Nat Acad Sci 114(8): 1952-1957.

Magurran AE, Girling SL (1986) Predator model recognition and response

habituation in shoaling minnows. Anim Behav 34: 510-518.

Mangialajo L, Chiantore M, Cattaneo-Vietti R (2008) Loss of fucoid algae along a

gradient of urbanisation, and structure of benthic assemblages. Mar Ecol

Prog Ser 358: 63-74.

Martinez-Pita I, Garcia FJ, Pita ML (2010) The effects of seasonality on gonad fatty

acids of the sea urchins *Paracentrotus lividus* and *Arbacia lixula*

(Echinodermata: Echinoidea). J Shellfish Res 29(2): 517-525.

Menge BA, Sutherland JP (1987) Community regulation: Variation in disturbance,

competition, and predation in relation to environmental stress and

recruitment. Am Nat 130(5): 730-757.

Morishita VM, Barreto RE (2011) Black sea urchins evaluate predation risk using

chemical signals from a predator and injured con- and heterospecific prey.

Mar Ecol Prog Ser 435: 173-181.

Nanri K, Nakajima Y, Yatzuya K, Kiyomoto S, Andou W, Yoshimura T (2011) Some

approaches for the recovery from barren grounds in Shin-Mie, Nagasaki

Prefecture. Fish Eng 48(1): 59-64.

Nielsen KJ, Navarrete SA (2004) Mesoscale regulation comes from the bottom-up:

intertidal interactions between consumers and upwelling. Ecol Lett 7: 31-

41.

Ogata R, Takagi K, Mishiku A, Fujita D (2016) A trial of restoration of a Sargassum

forest by transplanting thalli with a suspended net in Uchiura Bay, Shizuoka

Prefecture. Fish Eng 52(3): 177-184.

Okuda K (2008) Coastal environment and seaweed-bed ecology in Japan. Kuroshio

Sci 2(1): 15-20.

Peacor SD, Werner EE (2001) The contribution of trait-mediated indirect effects to

the net effects of a predator. Proc Nat Acad Sci, 98(7): 3904-3908.

Pearse JS, Hines AH (1987) Long-term population dynamics of sea urchins in a

central   California Kelp forest: rare recruitment and rapid decline. Mar Ecol

Prog Ser, 39: 275-283.

Pedersen HG, Johnson CR (2008) Growth and age structure of sea urchins

(*Heliocidaris erythrogramma*) in complex barrens and native macroalgal

beds in eastern Tasmania. ICES J Mar Sci 65(1): 1-11.

R Development Core Team 2019. A language and environment for statistical

computing. R Foundation for Statistical Computing, Vienna, Austria. http:

//www.R-project.org.

Ridout M, Demetrio CGB, Hinde J (1998) Models for count data with many zeros.

Proc 19th Intl Biom Conf, Cape Town, South Africa.

Rocha F, Baiao LF, Moutinho S, Reis B, Oliveira A, Arenas F, Maia MRG, Fonseca AJM,

Pintado M, Valente LMP (2019) The effect of sex, season and gametogenic

cycle on gonad yield, biochemical composition and quality traits of

Paracentrotus lividus along the North Atlantic coast of Portugal. Sci Rep

9(2994). doi.org/10.1038/s41598-019-39912-w.

Rodriguez-Barreras R, Cuevas E, Cabanillas-Teran N, Sabat AM (2015) Potential

omnivory in the sea urchin *Diadema antillarum*? Reg Stud Mar Sci 2: 11-18.

Sala E, Zabala M (1996) Fish predation and the structure of the sea urchin

Paracentrotus lividus populations in the NW Mediterranean. Mar Ecol Prog

Ser 140: 71-81.

Scheibling RE, Anthony SX (2001) Feeding, growth and reproduction of sea urchins

(*Strongylocentrotus droebachiensis*) on single and mixed diets of kelp

(*Laminaria* sp.) and the invasive alga *Codium fragile* ssp. *tomentosoides*. Mar

Biol 139: 139-146.

Schmitz OJ, Krivan V, Ovadia O (2004) Trophic cascades: The primacy of trait-

mediated indirect interactions. Ecol Lett 7: 153-163.

Shurin JB, Borer ET, Seabloom EW, Anderson K, Blanchette CA, Broitman B, Cooper

SD, Halpern BS (2002) A cross-ecosystem comparison of the strength of

trophic cascades. Ecol Lett 5: 785-791.

Sievers D, Nebelsick JH (2018) Fish predation on a Mediterranean echinoid:

Identification and preservation potential. Palaios 33: 47-54.

Smale DA, Kendrick GA, Wernberg T (2010) Assemblage turnover and taxonomic

sufficiency of subtidal macroalgae at multiple spatial scales. J Exp Mar Biol

Ecol 384: 76-86.

Smee DL, Weissberg MJ (2006) Clamming up: environmental forces diminish the

perceptive ability of bivalve prey. Ecology 87(6): 1587-1598.

Stan Development Core Team (2019) RStan: The R interface to Stan. http://mc-

stan.org/.

Steneck RS, Graham MH, Bourque BJ, Corbett D, Erlandson JM, Estes JA, Tegner, MJ

(2002) Kelp forest ecosystems: Biodiversity, stability, resilience and future.

Environ Conserv 29(4): 436-459.

Steneck RS, Leland A, McNaught DC, Vavrinec J (2013) Ecosystem flips, locks and

feedbacks: The lasting effects of fisheries on Maine's kelp forest ecosystem.

Bull Mar Sci 89(1): 31-55.

Strong DR (1992) Are trophic cascades all wet? Differentiation and donor-control

in speciose ecosystems. Ecology 73(3): 747-754.

Tegner MJ, Levin LA (1983) Spiny lobsters and sea urchins: Analysis of predator-

prey interaction. J Exp Mar Biol Ecol 73: 125-150.

Terawaki T, Yoshikawa K, Yoshida G, Ushimura M, Iseki K (2003) Ecology and

restoration techniques for Sargassum beds in the Seto Inland Sea, Japan.

Mar Poll Bull 47: 198-201.

Tuya F, Boyra A, Sanchez-Jerez P, Barbera C, Haroun R (2004) Can one species

determine the structure of the benthic community on a temperate rocky

reef? The case of the long-spined sea-urchin *Diadema antillarum*

(Echinodermata: Echinoidea) in the Eastern Atlantic. Hydrobiol, 519: 211-

214.

Uki N, Sugiura M, Watanabe T (1986) Dietary value of seaweeds occurring on the

Pacific Coast of Tohoku for growth of the abalone *Haliotis discus hannai*. Bull

Jap Soc Sci Fish 52(2): 257-266.

Urriago JD, Wong JCY, Dumont CP, Qiu JW (2016) Reproduction of the short-spined

sea urchin *Heliocidaris crassispina* (Echinodermata: Echinoidea) in Hong

Kong with a subtropical climate. Reg Stud Mar 8: 445-453.

Vadas RL (1977) Preferential feeding: an optimization strategy in sea urchins. Ecol

Monogr 47: 337-371.

Vanderklift MA, Kendrick GA (2005) Contrasting influence of sea urchins on

attached and drift macroalgae. Mar Ecol Prog Ser 299: 101-110

Watanuki A, Yamamoto H (1990) Settlement of seaweeds on coastal structures.

Hydrobiol 204/205: 275-280.

Watanuki A, Aota T, Otsuka E, Kawai T, Iwahashi Y, Kuwahara H, Fujita D (2010)

Restoration of kelp beds on an urchin barren: Removal of sea urchins by

citizen divers in southwestern Hokkaido. Bull Fish Res Agency 32: 83-87.

Webster DR, Weissberg MJ (2001) Chemosensory guidance cues in a turbulent

chemical odor plume. Limnol Oceanogr 46(5): 1034-1047.

Westermeier R, Murua P, Patiño DJ, Muñoz L, Atero C, Müller DG (2013)

Repopulation techniques for *Macrocystis integrifolia* (Phaeophyceae:

Laminariales) in Atacama, Chile. J Appl Phycol 26(1): 511-518. DOI

10.1007/s10811-013-0069-5

Wright JT, Dworjanyn SA, Rogers CN, Steinberg PD, Williamson JE, Poore AGB

(2005) Density-dependent sea urchin grazing: Differential removal of

species, changes in community composition and alternative community

states. Mar Ecol Prog Ser 298: 143-156.

Yamasaki M, Kiyomoto S (1993) Reproductive cycle of the sea urchin *Anthocidaris*

crassispina from Hirado Island, Nagasaki Prefecture. Bull Seikai Nat Fish Res

Inst 71: 33-40.

Yatsuya K, Nakahara H (2004a) Density, growth and reproduction of the sea urchin

Anthocidaris crassispina (A. Agassiz) in two different adjacent habitats, the

Sargassum area and Corallina area. Fish Sci 70: 233-240.

Yatsuya K, Nakahara H (2004b) Diet and stable isotope ratios of gut contents and

gonad of the sea urchin *Anthocidaris crassispina* (A. Agassiz) in two different

adjacent habitats, the *Sargassum* area and *Corallina* area. Fish Sci 70: 285-

292.

Yoo SK, Hur SB, Ryu SY (1982) Growth and spawning of the sea urchin *Anthocidaris*

crassispina (A. Agassiz). Bull Kor Fish Soc 15(4): 345-358.

Yoon JT, Sun SM, Chung G (2013) Sargassum bed restoration by transplanting of

germlings grown under protective mesh cages. J Appl Phycol 26(1): 505-

509. DOI 10.1007/s10811-013-0058-8.

Yotsui T, Maesako N (1993) Restoration experiments of *Eisenia bicyclis* beds on

barren grounds at Tsushima Islands. Suisanzoshoku 41(1): 67-70.

Zimmer RK, Butman CA (2000) Chemical signaling processes in the marine

environment. Biol Bull 198: 168-1187.

1   8. TABLES

2   Table 1. List of sea urchin behavior defined in this study.

Behavior type	Definition
None	Any immobile behavior within the region of interest. The urchin does not move 3 cm away from its starting point.
Movement	The urchin is freely moving inside the region of interest.
Interaction	The urchin makes contact with the center of the region of interest which may or may not contain seaweed or dead urchin.
Outside	When the urchin is outside the region of interest.

1 Table 2. Results of the Bayesian linear models on A) Experiment 1: feeding rate and

2 season, B) Experiment 2: temperature conditions, B1) light distribution and B2)

3 cue dispersal time to reach the 10 cm mark. The table shows the estimates, mean

4 and the lower and upper limits of the 95% confidence interval.

Estimates	Mean	2.5%	97.5%
A.) Lab experiment 1: Feeding rate (g algae day^{-1})			
Summer-Autumn	1.3	1.2	1.5
Winter-Spring	0.8	0.6	1.0
B.) Lab experiment 2: Temperature conditions (°C)			
Stock tank	18.6	18.6	18.7
Control	18.7	18.4	19.0
Algae	19.2	18.9	19.5
Dead urchin	18.8	18.4	19.3
Algae + dead urchin	19.1	18.7	19.5
B1.) Light distribution (Lux)			
Center	3.0	2.8	3.3

Middle	2.1	2.0	2.3
Outer	0.5	0.3	0.70

B2.) Cue dispersal time (sec)

Control	13.5	2.0	25.3
Algae	17.3	5.0	30.2
Dead urchin	22.6	9.8	35.3
Algae + dead urchin	19.5	6.4	31.8

Fig.1. Mean temperature difference between seasons in the feeding rate

experiment.

Fig. 2. Model predictions on mean count of urchin behaviors. The left and right

columns are represented by S-A and W-S, respectively. Rows are represented by the

treatments. Beginning from the top are control (A, E), dead urchin (B, F), algae (C,

G), and dead urchin + algae (D, H). On the x-axis, behaviors are abbreviated as M

(movement), I (Interaction), and (Outside).

Fig. 3. Model prediction on mean urchin time spent per behavior (min). The left

and right columns are represented by the S-A and W-S, respectively. Rows are

represented by the treatments. Beginning from the top are control (A, E), dead

urchin (B, F), algae (C, G), and dead urchin + algae (D, H). On the x-axis, behaviors

are abbreviated as M (movement), I (Interaction), and (Outside).

Fig. 4. Model prediction on mean urchin speed (cm min^{-1}). The left and right

columns are represented by the S-A and W-S, respectively. Rows are represented by

the treatments. Beginning from the top are control (A, E), dead urchin (B, F), algae

(C, G), and dead urchin + algae (D, H). On the x-axis, behaviors are abbreviated as M

(movement), I (Interaction), and (Outside).

Fig. 5. Binomial model predictions on urchin behavior outcome in the field.

Starvation makes urchins less likely to flee regardless of the presence or absence of

a dead urchin.

Figure 1

Figure 2

Figure 3

Figure 4

Figure 5