1

# KCNQ GENE FAMILY MEMBERS ACT AS BOTH TUMOR SUPPRESSORS AND ONCOGENES IN GASTROINTESTINAL CANCERS

*David Shorthouse[1], Eric Rahrmann[2], Cassandra Kosmidou[1], Benedict Greenwood[1], Michael Hall[1], Ginny Devonshire[2], Richard Gilbertson[2], Rebecca C. Fitzgerald[1], Benjamin A Hall[1*]*

[1]MRC Cancer Unit,

University of Cambridge,

Hutchison/MRC Research Centre,

Box 197,

Cambridge Biomedical Campus,

Cambridge,

CB2 0XZ

[2]Cancer Research UK Cambridge Institute

University of Cambridge

Li Ka Shing Centre

Robinson Way

Cambridge CB2 0RE

*to whom correspondence should be addressed.

Email: bh418@mrc-cu.cam.ac.uk

Short title: KCNQ genes in gastrointestinal cancer

## SUMMARY

We present evidence that KCNQ genes are drivers and suppressors of gastrointestinal (GI) cancer in humans. The KCNQ family of genes encode for subunits of a potassium channel complex involved in membrane polarisation and little is known about their role in cancer. We use human cancer data and a multidisciplinary computational-based approach including structural modelling and simulation, coupled with *in vitro* experiments to show that KCNQ1 is a tumor suppressor, and KCNQ3 and KCNQ5 are oncogenic across human GI cancers. We link the expression of KCNQ genes to WNT signalling, EMT, and survival and propose that mutation/copy number alteration of KCNQ genes can significantly alter patient prognosis in GI cancers.

(110 words)

## KEYWORDS

KCNQ, Ion Channels, Gastrointestinal Cancer, Epithelial to Mesenchymal Transformation (EMT), WNT signalling.

42    **GRAPHICAL ABSTRACT**

**SIGNIFICANCE**

Our results show that the ion channel family KCNQ contribute to progression of

gastrointestinal cancer. KCNQ genes are highly mutated and detected mutations

select for inactivation of KCNQ1, but increased activity of KCNQ3. Analysis of gene

expression profiles uncovers KCNQ expression correlating with the WNT pathway

and EMT. Moreover, clinical data correlates with expression data showing KCNQ1

has hallmarks of a tumor suppressor, whilst KCNQ3 and KCNQ5 are oncogenic.

These findings implicate KCNQ genes in control/mediation of the WNT pathway in

cancer, and potentially as prognostic markers or therapeutic targets.

(90 words)

## INTRODUCTION

The KCNQ family of ion channels are a set of evolutionarily related genes encoding for proteins involved in potassium transport (Abbott, 2014; Robbins, 2001; Wang and Li, 2016). KCNQ channels typically repolarise the plasma membrane of a cell after depolarisation through other channels. KCNQ are therefore involved in wide ranging biological functions such as cardiac action potentials (Nerbonne and Kass, 2005; Robbins, 2001), hearing sensitivity (Kharkovets et al., 2002; Kubisch et al., 1999), neural excitability (Brown and Passmore, 2009; Wang and Li, 2016), and ionic homeostasis in the gastrointestinal tract (Ohya et al., 2015; Warth et al., 2002).

Diseases resulting from loss or gain-of-function (LOF and GOF respectively) mutations in the KCNQ family are wide ranging, and include epilepsy (Allen et al., 2014; Millichap et al., 2016; Rogawski, 2000), long and short QT syndrome (Morita et al., 2008), deafness (Kubisch et al., 1999), and more recently, Autism-like disorders (Sands et al., 2019). Neurological disorders caused by KCNQ mutations are an active area of research, and specific mutations in many individuals have been characterised using electrophysiological methods - many mutations therefore have prior functional characterisation as LOF of GOF (Miceli et al., 2008, 2015; Panaghie and Abbott, 2007; Sands et al., 2019). Whilst some of the mechanisms by which these diseases occur have been partially elucidated, there are complications in that the KCNQ family of proteins interact heavily with the KCNE family, to form complex heteromeric combinations of varying subunits that are not completely understood. In particular, KCNQ1 interacts with all known KCNE ancillary proteins in varying tissues, but is otherwise homotetrameric (Abbott, 2014). KCNQ2, 3, 4, and 5 however, can interact with each other and the KCNE family to theoretically form hundreds of combinations of channels, and the impact of specific mutations on any

one member will have an unknown influence on the greater function of the pool of

channels (Howard et al., 2007; Yu et al., 2013).

Previous work has further highlighted the potential roles of individual ion channels

across a broad range of cancers, including TRPA1 in breast cancer (Takahashi et

al., 2018), HERG channels in leukaemia (Pillozzi et al., 2002), and osmotic

regulatory machinery in several cancers (Haas and Sontheimer, 2010; Pedersen et

al., 2013). The complex interplay between ionic gradients and cellular phenotypes

has been studied at a high level before, including in cancer (Pardo and Stühmer,

2013; Pedersen and Stock, 2013; Shorthouse et al., 2018), but the functional

contributions of the KCNQ family to the phenotype of non-excitable tissue is not fully

known.

There is preliminary evidence to suggest that KCNQ1 plays a tumor suppressive role

in the stomach and colon (Rapetti-Mauss et al., 2017; Than et al., 2014), and

hepatocellular carcinoma (Fan et al., 2018), and that KCNQ3 potentially plays a role

in oesophageal adenocarcinoma (Frankell et al., 2019). It is of particular interest to

study the roles of ion channels in the gastrointestinal tract due to the critical nature of

the ionic homeostasis that is required for their varied physiological functions. Cells in

the gastrointestinal tract are often involved at the interface between the tissue and

the luminal environment, and as such membrane transport is critical to their

homeostatic function. The gradients electrochemical ion channels generate are used

in protective mucus production (Tarran, 2004), acid secretion (Grahammer et al.,

2001), and immune recognition (Feske et al., 2015), among other processes.

Additionally, there is some evidence that activity of membrane transporters may play

roles in key cancer pathways, such as a reported interaction between KCNQ1 and

beta catenin (Rapetti-Mauss et al., 2017).

With the wide availability of publicly available databases such as The Cancer

Genome Atlas (TCGA) (https://portal.gdc.cancer.gov/) and the International Cancer

Genome Consortium (ICGC) (https://icgc.org/) it is possible to analyse samples from

large populations of cancer patients in terms of gene expression, mutations, copy

number variation, and gene methylation. Whilst in many cases, mutations have been

previously characterised as LOF or GOF, in cases where specific data is not

available to categorise the mutation, structures of the protein become critically

important in studying their potential effects. Where structures are not available for a

particular gene, homology modelling is a useful method for generating consistently

reliable predictive structures and allow the study and simulation of the spatial context

of individual mutations (Šali and Blundell, 1993; Šali et al., 1995).

In  this study, we aimed to investigate the role of the KCNQ family in gastrointestinal

cancer through study of highly annotated clinical data sets with DNA and RNA

sequencing data paired with structural modelling of KCNQ proteins.

## RESULTS

### KCNQ genes are highly genetically altered in GI cancers

We studied the mutations in the KCNQ and related KCNE family across all cancers

using the TCGA database. We calculated the missense mutational frequency of all

KCNQ genes within each TCGA cohort, and grouped them into previously defined

(Hoadley et al., 2018) subgroups (**Figure 1 A**). Ranking subgroups by percentage of

patients that have a mutation in a KCNQ/E gene shows enrichment for the core GI

subgroup when compared to other subgroups, a higher overall mutational burden is

seen only in the melanoma cohort, which has a significantly higher baseline

mutational burden (Chalmers et al., 2017). Comparison of the rates of synonymous

to nonsynonymous mutations within the core GI cohort potentially indicates the

presence of mutational selection, there are significantly more nonsynonymous

mutations when compared to synonymous.

We further studied all genetic alterations in the KCNQ and related KCNE family

across GI cancers using the publicly available TCGA and our own Oesophageal

Adenocarcinoma data which has detailed clinical annotation (OCCAMS). The GI

cancers studied were: Oesophageal Squamous Cell Carcinoma (ESCC, n = 103),;

Oesophageal Adenocarcinoma in two cohorts: the TCGA (EAC-TCGA, n = 93)(Kim

et al., 2017) and our own data (EAC-OCCAMS part of ICGC, n = 378); Stomach

Adenocarcinoma (STAD-TCGA, n = 426) (Bass et al., 2014), and Colorectal

Adenocarcinoma (COADREAD-TCGA, n = 594)(Muzny et al., 2012). We found that

31% of patients with GI cancers (n = 1594) had genetic alterations in the KCNQ/E

families of genes. 30-40% of patients in all subtypes contain a mutation or copy

number alteration (CAN), except for COADREAD-TCGA, in which 26% of patients

are altered for KCNQ/E. The ratio of amplifications to mutation and deletion are also

conserved between cohorts aside from ESCC-TCGA, which mostly contains

amplification events (**Figure 1 B**). Comparing the copy number profiles, KCNQ1,

KCNE1 and KCNE2 appear mostly to be deleted overall, whereas KCNQ2, KCNQ3,

and KCNE3 are generally amplified. In terms of SNVs we find that KCNQ2, KCNQ3,

and KCNQ5 are heavily mutated (**Figure 1 C**). Interestingly, 175 (11%) of all patients

have a mutation/copy number change in KCNQ3, the most altered member of the

family. The alterations are generally distributed across tissue types (**Figure 1 D**), and

there is no observed correlation between mutations in the KCNQ/E family and

cancer stage where annotated (**Figure S1**).

Co-occurrence analysis with common GI cancer drivers (CCND1, CDKN2A,

CTNNB1, ERBB2, KRAS, PTEN, and TP53) using DISCOVER (Canisius et al.,

2016) show that many members of the KCNQ family are mutually exclusive with

common drivers (**Table S1**), in particular we find that KCNQ2 and KCNQ5 are

statistically significantly ($q <= 0.1$) mutually exclusive with CCND1 and CDKN2A, and

KCNQ2 and KCNQ3 are mutually exclusive with both KRAS and TP53, indicating

that these mutations tend to occur separately, potentially revealing that co-

occurrence of these mutations is unfavourable to overall cell survival (Campbell,

2017).

RNA expression comparison between cancer and normal samples reveals that

KCNQ3 is generally upregulated in GI cancers compared to normal tissue, and

KCNQ1 shows a range of altered expression, but generally has its expression

reduced (**Figure S2**).

Analysis of methylation data also indicates that the CPG promoters of the KCNQ

family are often differently methylated when compared with their normal counterparts

(**Figure S3**), KCNQ5 appears hypermethylated, consistent with a decrease in gene

expression. We also find that KCNQ1 and KCNQ3 show methylation patterns

opposite to their general change in gene expression, with a hypermethylation of CPG

promoter of KCNQ3 observed, despite an increase in RNA expression, and a

general decrease in the methylation of CPG promoter in KCNQ1. This indicates the

gene expression is likely altered through means other than methylation in the case of

KCNQ1 and KCNQ3 such as mutation of the promoter region.

<u>Mutations in the KCNQ family show evidence of selection</u>

In order to study mutations in the KCNQ family across gastrointestinal cancers we

chose to look in depth at the COSMIC database of somatic mutations in cancers,

selecting for mutations occurring in any tissue along the gastrointestinal tract. The

COSMIC database aggregates mutations from multiple sources, including TCGA and

ICGC, and so is a comprehensive database for mutations across human cancers.

We performed a MUSCLE alignment of protein sequences for members of the

KCNQ family to allow alignment of sequences into conserved domains, primarily

transmembrane helices.

We find that there are regions of increased mutational frequency in different genes

within the KCNQ family (**Figure 2 A**). In particular, KCNQ1 shows increased

frequency of mutations in the S2-S3 helical region, and the S5-pore-S6 region of the

protein. KCNQ3 shows a clear region of increased mutational burden within the S4

voltage sensing helix, and a second region in the Calmodulin-binding HB-HC helices.

There are regions of high mutational burden also present within the S6 helix for

KCNQ2, and the S4 helix for KCNQ5 potentially indicating mutational selection.

Fathmm scores calculated for KCNQ1-5 by COSMIC for mutations observed across

GI cancers show that 65% of all mutations observed are predicted with high

confidence to be pathogenic (**Table S2**). We additionally calculated the cancer-

associated CScape scores (Rogers et al., 2017) for all possible mis-sense mutations

in KCNQ1 and KCNQ3, the two genes showing highest relative mutational

clustering. We find evidence that particular structural regions, with good overlap with

those regions showing an increased mutational burden in GI cancer, have a high

proportion of predicted pathogenic mutations. (**Figure S4**). Additionally there is a

significantly larger number of missense vs nonsense mutations within the domains of

all KCNQ genes (**Figure S5**).

To statistically explore regions of mutational clustering within the sequence of each

member of the KCNQ family, we applied the NMC method (Ye et al., 2010) to look

for potential clusters of mutations occurring more often than expected by chance

within the 1D sequence (**Figure 2 B**). We additionally overlay a mutational signature-

based observed vs expected mutation ratio applied across a sliding window of the

protein sequence, similar to the dn/ds method previously applied to mutational

selection in cancer (Martincorena et al., 2017). We find that for KCNQ1, there is a

clear significant cluster of mutations within the S2-S3 linker region (cluster 1.1), and

within the S6 helix (cluster 1.2). KCNQ3 showed a significant region of mutational

selection within the S4 voltage sensing helix, particularly arginine residues 227, 230,

and 236 (cluster 3.1) KCNQ5 shows a major region of mutational clustering within

the S4 helix, and a structurally and functionally undefined region at residue 680-700,.

For KCNQ2, we identified a region of increased mutational frequency around the S6

helix, and no regions of mutational selection within KCNQ4 (**Figure S6**).

Finally, we performed a literature search for functional characterisation of observed

mutations. Many of the mutations observed in the COSMIC database are functionally

characterised mutations in other KCNQ-related disorders (**Table 1**).

Overall we find mutations in KCNQ1 appear to cluster primarily in the S2/S3 and S6

regions and mutations to KCNQ3 and 5 show similar clusters within the S4 helix.

A) Mutational distribution for KCNQ family proteins

B)

<u>Mutations to KCNQ genes are clustered in space</u>

The structural assembly of the KCNQ/KCNE family is complex – KCNQ1 is known to

assemble into homotetramers, which are modulated by members of the KCNE gene

family, whereas KCNQ2, 3, 4, and 5 have varying stoichiometries, and are also

modulated by KCNE gene products (**Figure 3 A**). We performed a bayesian

inference analysis to study the gene expression correlations between members of

the KCNQ/KCNE family in different gasto-intestinal tissues. In particular for the

COADREAD RNAseq dataset **(Figure 3 B)** we find two clusters of genes that

positively correlate with each other, this effect is observed to a lesser extent in other

tissue specific analysis (**Figure S7**). The two clusters that show negative correlation

to each other contain KCNQ1, KCNE2, and KCNE3 (Set 1), and KCNQ3, KCNQ4,

KCNQ5, KCNE1, and KCNE4 (Set 2). KCNQ1 and KCNE2/KCNE3 complexes have

been observed in vivo wild-type gastrointestinal tissue, and are essential for the

homeostatic control of proton and potassium gradients (Grahammer et al., 2001;

Heitzmann et al., 2004). This analysis potentially indicates an antagonistic

relationship between the two clusters.

In order to further demonstrate functional effects of mutations on the structure of

KCNQ genes, we chose to structurally model members of the KCNQ family. KCNQ

proteins contain 6 transmembrane helices (shown schematically in **Figure 3 C**,

shown structurally in **Figure 3 D,** full structure shown in **Figure S8**). Helices S1, S2,

S3, and S4 (red) make up a voltage sensor domain, whose conformation changes in

response to voltage shifts across the membrane control the gating of channel

through positively charged arginine residues in the S4 helix. The S5, pore, and S6

domain (yellow) contain the gating components of the channel, additionally there are

three helices HA, HB, and HC (blue), which make up a structure to which calmodulin

binds and modulated the channel.

Homology models of the human proteins of each member were generated from the

cryo-em structure of *Xenopus laevis* KCNQ1 5VMS. Models exhibit high nativity

scores (**Table S3**), and were validated as viable by 100ns of equilibrium atomistic

level molecular dynamics (**Figure S8**).

Mapping COSMIC mutations onto the resultant structures and colouring by

frequency reveals structural regions with a high frequency of mutations. KCNQ3 in

particular shows a high number of mutations clustered in the S4 helix as expected

(**Figure 3 E**). Other members show some mutational clustering (**Figure S9**).

To explore the spatial clustering of mutations quantitatively within the predicted

structures we calculated clusters of colocalized mutations within each structure.

Mutations within a 12 Å cutoff we grouped together, we find that each structure

contains mutational clusters that give more functional information than 1D analysis

alone. For KCNQ1 we find two distinctive spatial clusters, those in the S2/S3 linker

domain, and a large interlinked cluster of residues around the S6 helix (**Figure 3 F**).

Mutations in cluster 1.1 are clearly part of the previously defined phosphatidylinositol

(PIP) binding site, essential to channel function (Zaydman and Cui, 2014). Mutations

in cluster 1.2 converge in the centre of the protein and appear to cluster around the

pore restriction (**Figure 3 G**). For KCNQ3, we find that the previously defined

mutational cluster in the S4 helix, denoted cluster 3.1, makes up a large region of

spatially clustered mutations within the S4/S5 helices of the assembled tetramer

(**Figure 3 H**). We additionally identify a further region of mutational clustering within

the HA/HB helices of the protein, which are spatially close within the full tetrameric

protein, but not within the 1D structure (cluster 3.2).

Overall, clustered mutations in KCNQ1 appear to mainly influence cofactor binding

(cluster 1.1) and the pore of the protein (cluster 1.2), whereas KCNQ3 mutational

clusters are focussed in the activation domain (cluster 3.1).

<u>Mutations to KCNQ genes in GI cancers are functionally relevant</u>

Due to the spatial clustering of mutations in the S6 gating helix and the S4 voltage

sensing helix in KCNQ1 and KCNQ3 respectively we sought to investigate the

functional effects of these mutations on the activity of the assembled channels.

For mutations in the KCNQ1 S6 helix, we generated additional homology models for

each of the mutations found to be clustered in that region (Cluster 1.2 - F339L,

L342F, P343L, and P343S), and an additional nearby mutation that is the most

frequently observed KCNQ1 mutation across GI cancers (A329T). Analysis of the

pore domain of the protein (**Figure 4 A**), shows that all mutations except F339L are

predicted to occlude the pore, reducing or eliminating its ability to gate potassium

ions, even when a single subunit is mutated, as may be the case when a patient has

a KCNQ1 mutation in only a single allele. We conclude that mutations in cluster 1.2

for KCNQ1 are likely loss-of-function.

The S4 domain of KCNQ channels is particularly important due to its involvement in

channel gating. It is of special interest because it is a "mutational hotspot" in KCNQ2

mutation induced epilepsies, and strong prior evidence exists for mutations in S4

controlling channel activity (DeMarco, 2012; Miceli et al., 2008, 2015; Sands et al.,

2019). The positively charged arginines within S4 are the principal drivers of the

proteins response to membrane polarity. We find a significant number of mutations in

KCNQ2, KCNQ3, and KCNQ5 are within arginines in S4, with KCNQ3 containing

mutations in every one of the 5 arginines (**Figure 4 C**). The S4 helix of all KCNQ

channels includes 4 (KCNQ1) or 5 (KCNQ2-5) positively charged arginine residues

essential for gating and activation of a potassium current, similarly to other

potassium channels (Aggarwal and MacKinnon, 1996; Jentsch, 2000). We refer to

these arginines by convention in ascending order from N to C terminus as R1, R2,

R4, R5, and R6. The position generally referred to as R3 is replaced with a

glutamine in the KCNQ family and is therefore not considered. We find that R1, R2,

and R4 are highly mutated in KCNQ3, and R4 and R5 are mutated in KCNQ2, whilst

KCNQ5 shows 9 total mutations in R5 only.

We chose to analyse individual mutations in the helix using a previously developed

tool utilizing molecular dynamics, Sidekick (Hall et al., 2014) (**Figure S10**), that

simulates the effects of single mutations on the properties of an alpha helix. We

surmised that single helical positioning in the membrane can act as a proxy for the

forces the helix exerts on the full protein upon mutation, without costly simulations of

the entire mutant protein. We chose to first replicate an experimentally determined

alanine scanning experiment on the KCNQ1 S4 helix (Panaghie and Abbott, 2007),

and a tryptophan scanning experiment of the KCNQ2 S4 helix, before simulating

mutations in the KCNQ S4 helices from both cancer and epilepsy, some of which

have previously reported experimental characterization.

Alanine scanning experiments (Panaghie and Abbott, 2007) revealed that mutations

of the R2 and R4 positions of KCNQ1 to alanine resulted in constitutively active, or

more readily activated currents, whereas mutations to alanine at R1, R6, or D242

(referred to as D6) impaired the activation of KCNQ1. Calculating helical tilt within a

DPPC membrane reveals mutations to alanine at R2 and R4 decrease the tilt angle

of S4, and has no effect (R1) or an increased angle for R6 and D6 mutations. A

decrease in tilt angle within Sidekick analysis therefore correlates with increased

activity of the channel (**Figure 4 D**). We additionally replicated a glutamine scanning

experiment performed on the KCNQ2 S4 helix (Miceli et al., 2008) (**Figure S11**). We

find that sidekick correctly identifies a decrease in tilt angle for constitutively active

mutant R2Q but not for R1Q, which has a reduced activation threshold. We also find

no change in tilt angle for mutants with a reduced activation threshold (R4Q, R5Q,

R6Q, R7Q).

We simulated all arginine mutations in KCNQ3 including 3 that occur within our

dataset, and additional mutations that are recently validated as gain of function

causes of autism related disorders (Sands et al., 2019). Simulations agree with

previous data, indicating a decrease in helical tilt angles for R2H and R2C, known

gain-of-function mutations, and predicting that similar properties occur with

unvalidated mutations R236C, and R236H (**Figure 4 E**). Overall we find that 14/17

(82%) of all mutations within the KCNQ3 S4 helix, and by extension involved in

mutational cluster 3.1 are either previously validated gain of function, or predicted by

our simulations to be gain of function (**Table 2**).

Finally we sought to examine the large number of mutations in the R5 residue of

KCNQ2 and KCNQ5, simulations do not show any change in helical tilt upon

mutation to H or C (**Table 2, Figure S11**), but looking at the arginines in the context

of the full protein in molecular dynamics simulations shows that hydrogen bond

occupancy for residues involved in stabilizing the active form of the protein are

different. Residues thought essential for stabilizing the active form of the channel

(Sands et al., 2019; Soldovieri et al., 2019) are R4, R5, R6, an aspartic acid in helix

S2 (D1), and two glutamic acids within helix S3 (E1, and E2) (**Figure 4 F**). We find

the KCNQ5 bonding network differs in occupancy compared to KCNQ2-4. R5 is

principally involved in a hydrogen bonding interaction with D1 in KCNQ2-4, with

occupancies ranging from 40% to 62% (**Figure S12**), but KCNQ5 R5 shows only

16% occupancy in bonding with D1, and instead is found to have an 81% occupancy

with E2 (**Figure 4 G**). Disruption of this hydrogen bonding network by mutation is

likely to have functional consequences on the activity of the protein, particularly as

the primary mutation to KCNQ5 R5 is to a cysteine, which could potentially form a

disulphide bond with the unpaired cysteine R203 situated ~7Å from R5 in our model.

Overall we find evidence that many mutations observed in the KCNQ in GI cancer

are of functional consequence. In particular, we find that mutations in Cluster 1.2 of

KCNQ1 are mostly predicted to be loss-of-function, whilst those in Cluster 3.1 of

KCNQ3 are mostly gain-of-function. We additionally find predicted functional

consequences for many other mutations observed in the KCNQ family. This

reinforces prior data showing that KCNQ1 appears tumor suppressive, and can be

mutationally inactivated in cancers, whilst KCNQ3 appears to primarily accumulate

gain-of-function mutations.

<u>KCNQ gene expression correlates with WNT signalling and EMT in GI cancer</u>

To interrogate the functional and signalling properties of KCNQ genes in cancer, and

in particular to identify potential interactions with known cancer-associated pathways,

we looked at the RNAseq data associated with both TCGA and the OCCAMS

cohorts. We chose to study KCNQ1, KCNQ3, and KCNQ5, as they show the

strongest mutational clustering. Firstly, genes from each of 10 major cancer-

associated pathways were defined based on previously published analysis of the

TCGA (Sanchez-Vega et al., 2018). (**Table S4**). Given KCNQ genes are altered in

expression and copy number within GI cancer, we infer that patients with high and

low expression of each KCNQ gene may show differences in the activation of

cancer-associated pathways they interact with.

We trained a machine learning model using Gradient Boosted Decision Forests

(GBDF) to learn the difference between high and low expressors of each KCNQ

gene when given only the genes in each pathway. A more accurate model (better

discrimination between high and low expressors) indicates greater association of a

particular pathway to the query gene.

We tested our method on the canonical cancer pathway genes CTNNB1, PTEN,

KRAS, and NOTCH1 by generating 20 classification models, and calculating the

average scores for each pathway. All classifications result in the correct pathway

(WNT for CTNNB1, PI3K for PTEN, RTK-RAS for KRAS, and NOTCH for NOTCH1)

being classified in the top 2 pathways, as expected (**Figure S13**), for this analysis we

excluded the query gene from the pathway definition so as not to pre-bias the model.

Running the model with KCNQ1, KCNQ3 and KCNQ5 as the query genes across all

GI cancers (**Figure 5 A**), we find that the WNT pathway scores the highest for

average Matthews Correlation Coefficient (MCC). We also find that accuracy scores

are generally higher for KCNQ1 and KCNQ3, potentially indicating that they are

better classifiers and thus relatively more associated with their respective pathways.

This result reinforces previous evidence that associates the expression of KCNQ1 in

colorectal cell lines with a direct physical interaction with beta catenin (Rapetti-

Mauss et al., 2017), and potentially expands the interaction to involve other KCNQ

genes.

One advantage of using GBDFs for classification is the ability to interpret models and

analyse which features contribute most to a score, in our case indicating how much

influence each gene in each pathway had on the MCC score. We generated average

Shaply Additive Explanation (SHAP) scores for the WNT pathway for (Lundberg and

Lee, 2017; Lundberg et al., 2018) for each model trained on high and low KCNQ

expressors (**Table S5**). Genes with a high SHAP score (>0.1) are extremely

influential in the classification and therefore most associated with the query gene.

The highest ranked WNT genes that correlate with KCNQ1 expression are

transcription factors (TCF7, TLE3, RNF43) (**Figure 5 B**), interestingly TCF7 has

been shown to be responsive to potassium in a tumor setting in immune cell

populations (Vodnala et al., 2019). KCNQ3 classifications were mostly driven by

membrane bound members of the WNT pathway and ligands (FZDs and WNTs).

Overall however, all regions of the WNT pathway scored highly for each gene,

indicating involvement of the entire pathway, rather than a small subset of individual

members.

To further explore the previous suggestions of an association of the KCNQ family

with the WNT pathway, we generated clustered heatmaps of all WNT genes for the

top and bottom 25 expressers of KCNQ1, KCNQ3, and KCNQ5 (**Figure 5 C**).

Heatmaps show patient-type clustering as predicted, and genes associate strongly

into co-regulated groups that correlate with KCNQ expression, showing that KCNQ

expression is strongly linked to activity of genes within the WNT pathway. We further

performed GSEA analysis of the top 25 and bottom 25 overall expressers for each

KCNQ and identify WNT signalling as a hallmark pathway significantly associated

with KCNQ3 and KCNQ5 (FDR q value <= 0.1)(**Figure S14**).

Because of the high association of KCNQ with WNT through our analysis, and

previous work linking KCNQ1 to beta-catenin (Rapetti-Mauss et al., 2017), we chose

to study the link between KCNQ genes and epithelial-to-mesenchymal transition

(EMT). We trained a further GBDF on a set of genes previously implicated in

RNAseq analysis of EMT (Gibbons and Creighton, 2018). Feature extraction of the

models shows a strong correlation between KCNQ1 and E-cadherin and SNAI2

(**Figure 5 D**), and of KCNQ3 with fibronectin (FN1). Additionally, we identify KCNQ1

and KCNQ5 as being within the top ~20% and ~10% respectively of all genes for

correlation with an EMT signature in the bulk RNAseq data (**Figure 5 E**).

To further unpick the relationship between KCNQ1/3/5, and EMT we looked at the

association of a previously defined EMT-score generated from RNAseq (Creighton

and Gibbons, 2013) with the expression of members of the KCNQ family. The EMT

score is significantly correlated with the RNA expression levels for KCNQ1, KCNQ3,

and KCNQ5 (**Figure 5 F**). KCNQ1 shows a negative correlation with EMT score,

indicating that higher KCNQ1 expression is associated with a more epithelial

phenotype (pearson rho = -0.34, p <0.0001). Conversely, KCNQ3 and KCNQ5 show

a positive correlation, indicating that high expression is associated with a more

mesenchymal phenotype (pearson rho KCNQ3 = 0.29, KCNQ5 = 0.39, p <0.0001),

for comparison, the pearson rho for the classical EMT transcription factor SNAI1 is

0.49 (p <0.0001). This reinforces the previous mutational data implicating KCNQ1

with a tumor suppressive role, and KCNQ3/5 with an oncogenic-like role. GSEA

analysis finds a high correlation between EMT signatures in KCNQ1 and

KCNQ3/KCNQ5 high and low expressors (FDR q value <= 0.05) (**Figure 5 G**).

As a validation of the RNAseq data we find significant disruption in the expression of

WNT proteins from analysis of The Cancer Protein Atlas (TCPA) data (Li et al.,

2017) when protein expression data is matched with TCGA RNAseq for the same

patients (**Figure S15**), in particular we find levels of beta-catenin and alpha-catenin

are commonly significantly altered in patients that are high vs low for KCNQ RNA

(corrected p <= 0.1). We also find a statistically significant difference in the

expression of EMT proteins between high and low RNA expressors of KCNQ genes

(**Figure S16**), in particular, E-cadherin, claudin-7, are generally higher in high

expressors of KCNQ1 and low expressors of KCNQ3/5, and fibronectin is generally

found lower in high KCNQ1 patients and low KCNQ3/5 patients. Interestingly we also

find a significant difference in both RNA and protein levels of BRCA2 for expressors

of KCNQ3, potentially indicating a difference in DNA damage response.

Additional validation was performed by analysis of oesophageal adenocarcinoma

organoid systems (Li et al., 2018). Whilst RNA levels of KCNQ3 and 5 in each

system are extremely low, KCNQ1 is expressed at different levels in the different

organoid systems. We find a strong correlation between the expression level of

KCNQ1 and activation of the WNT pathway, similar to that seen in the patient

RNAseq data (**Figure S17**).

■ SNAI1: r = 0.49

KCNQ gene expression correlates strongly with clinical prognosis

Kaplan Meier analysis of overall survival (**Figure 6 A**) and disease-free survival

(**Figure S18**) for TCGA datasets shows that many members of the KCNQ family

have a significant correlation with prognosis and recurrence. Across all GI cancers

KCNQ1 expression positively correlates with survival (p<0.0001), confirming

previous work suggesting that it has tumor suppressive properties. When broken

down into constitutive cancers however, we find statistical significance for KCNQ1

only in the STAD dataset (p<0.001) (**Figure S19**), potentially indicating a stomach

specificity. Though previous work has shown tumor suppressive properties for

KCNQ1 in colorectal tissue (Rapetti-Mauss et al., 2017), we do not find a statistically

significant correlation (**Figure S19**). We additionally find a statistically significant

correlation with poorer overall survival and the high expression of KCNQ3, and

KCNQ5 over all GI patients, indicating an oncogene-like effect associated with their

expression. Tissue breakdowns suggest that KCNQ3 expression is negatively

correlated with overall survival in all subtypes except ESCC, where there is no

correlation, though statistical significance (p < 0.05) is not reached in all cases

(**Figure S19**). We also studied the impact of KCNQ3 on survival in EAC in the

combined TCGA-EAC/OCCAMS dataset, and find a statistically significant

correlation with survival (p=0.017).

To study the inter-gene relationship between KCNQ1, KCNQ3, KCNQ5, and survival

we separated patients into groups that were high and low for every pair of

combinations of the three genes. Comparing KCNQ1 and KCNQ3 (**Figure 6 B**) we

find the patients high for KCNQ1, and low for KCNQ3 have the best prognosis. The

worst prognosis is found for patients with low KCNQ1 and high KCNQ3, whilst

patients high or low for both have an intermediate survival. The graded change in

survival for these 4 sets of patients indicates that KCNQ1 and KCNQ3 may be

independent of each other, as alterations in each individual gene combines with the

effect of the other one survival. The same is also true when KCNQ1 and KCNQ5 are

compared. The same is not true for comparison of patients with differing expression

of KCNQ3 and KCNQ5 (**Figure 6 C**). Patients who are low expressers of both

KCNQ3 and KCNQ5 have the best prognosis, whilst those with an increased

expression of either, or both genes have a similarly worse prognosis. This indicates

that KCNQ3 and KCNQ5 are mutually redundant, in that an alteration in a single one

of them is enough to cause a significant prognostic change, and alteration of the

second has no further impact on the phenotype. Taken together, this provides

evidence that KCNQ1 can be treated as independently influencing the cell from

KCNQ3 and KCNQ5.

We performed differential expression analysis between the best and worse

prognosis patients in each case presented in Figure 6C for oesophageal

adenocarcinoma. Classifying genes as differently expressed with a corrected p > 0.1,

and log2 fold change > ±1.5 (**Figure 6 D**). We find that some commonly deregulated

genes are chemokines, proton-coupled membrane transporters, and some

transcription factors, including VHLL (**Table S6**). Additionally, we find a clear

alteration in the expression levels of WNT transcription factors TCF4, TCF7, SNAI1,

and SNAI2, shown for KCNQ1/KCNQ3 best and worst prognoses in (**Figure 6 E)**.

Finally we combine all of our findings into a single descriptive model. We generated

a discrete, executable systems biology model of KCNQ activity in GI cancers using

the BioModelAnalyzer. (**Figure S20**). The model describes the effects of mutations

and expression changes in the KCNQ family based on the previously presented

data, and predicts the impact of alteration on 5 cellular phenotypes, Proliferation,

Cell Cycle Arrest, Apoptosis, Cell-Cell Adhesion, and EMT.

Overall our results point to a model where WNT signalling is mediated negatively by

KCNQ1, and positively by KCNQ3/KCNQ5 (**Figure 6 F**).

**DISCUSSION**

There is emerging evidence that ion channels play a role in many, and potentially all

cancers. Therapeutics against voltage gated potassium channels have been shown

in vivo to improve prognosis for glioblastoma and breast cancer (Arcangeli and

Becchetti, 2017; Pointer et al., 2017; Wang et al., 2015), and there are studies

implicating specific sodium (Brisson et al., 2011; Roger et al., 2015) and calcium

channels (Phan et al., 2017; Qu et al., 2014) in cancers. We present strong evidence

that the KCNQ family of genes also play a significant and functional role in human

gastrointestinal cancers. Through integration of data at the patient, cell, and protein

structural levels we show that KCNQ genes and protein products contribute to

cancer phenotype, and provide a selective advantage to cancers. We show that a

large number of patients with gastrointestinal cancers show genetic alterations in a

member of the KCNQ family, and that the expression levels of these genes correlate

strongly with survival. We find that mutations in the KCNQ family of genes are

clustered within KCNQ1, KCNQ3, and KCNQ5, and that there is evidence of

mutational selection. Furthermore, we prove that many of these mutations have

functional effects on the protein products, through alteration of the gating properties

of the channel, or through occlusion of the pore region. We go on to analyse

RNAseq and RPPA level data, using novel machine learning techniques, to show

that KCNQ gene expression highly correlates with the WNT pathway, and with EMT,

indicating a potential role for the gene family in cancer signalling and transformation.

Finally, we generate a systems biology model to sum our findings into a descriptive

model, and show that patients with different expression levels of KCNQ genes have

varying prognosis.

By studying data from varying levels simultaneously (patient level, cellular level, and

protein level) we find consistently that KCNQ1 exhibits properties of a tumor

suppressor – it is often deleted or lost in cancer, mutations are generally inactivating,

and it is negatively correlated with WNT signalling and EMT. Opposingly, we find that

other KCNQ genes (mainly KCNQ3 and KCNQ5) show hallmarks of onocogenic

properties, they are often amplified in cancers, and their mutations are synonymous

with gain-of-function mutations. Additionally, KCNQ3 and KCNQ5 are positively

correlated with WNT signalling and the EMT pathway, thus genes within the same

family, with very similar chemical activity have opposing influences on cancer

phenotype. The correlation with survival is of particular interest for potential patient

stratification, and may additionally represent a clinical window for reduction of certain

cancer phenotypes, especially given the recent developments of highly specific and

potent KCNQ subtype specific drugs (Manville and Abbott, 2019). Caution must be

taken when considering therapeutic applications of KCNQ involvement in cancer,

due to the extreme importance of KCNQ1 in cardiac activity, nevertheless

compounds specific to KCNQ3 or KCNQ5 may have a therapeutic window, as is

thought to be the base with hERG inhibitors (Fukushiro-Lopes et al., 2018).

That every level of data (genetic alteration/clinical, mutational/structural, gene

expression/pathway) consistently agrees that KCNQ1 appears tumor suppressive,

whilst KCNQ3 and 5 have oncogenic properties is a strong indicator that the results

are clinically relevant.

Whilst we confirm previous work that KCNQ1 may be involved in WNT signalling

(Rapetti-Mauss et al., 2017), and expand to show that KCNQ3 and KCNQ5 play an

opposing role, further work is necessary to elucidate exactly how ion channels can

drive these pathways, through either membrane polarity, physical protein-protein

interactions, or another currently undiscovered mechanism.

A further result of this finding is the implication that patients with congenital

mutations in KCNQ genes may be more at risk or protected against particular GI

cancers, this is especially poignant in the light of recent results suggesting that

KCNQ channels may be acted upon by particular dietary vitamins such as vitamin B6

(Reid et al., 2015).

In summary, we identify a strong role for KCNQ1, KCNQ3, and KCNQ5 in the

progression of human gastrointestinal cancers. We find that KCNQ1 is tumor

suppressive, whereas KCNQ2, KCNQ3, and KCNQ5 have oncogenic properties. We

confirm our findings across multiple levels of data, and evidence is consistent at the

patient, cellular, and protein/gene levels. We also find a role for KCNQ genes in the

WNT pathway and EMT, expanding upon previous work that found an interaction

between KCNQ1 and beta-catenin. This work highlights the emerging functional role

of the KCNQ family in human gastrointestinal cancers.

**ACKNOWLEDGEMENTS**

We acknowledge the help and support of Benjamin Marshal on the Bayesian

Inference analysis. We are grateful to the OCCAMS consortium for providing the

data for the oesophageal analysis linked with clinical outcome data, and Dr Xiaodun

Li for advice with the organoid data. We acknowledge the services of Cancer

Research UK Cambridge Institute Genomics Core for performing RNAseq. This work

has been supported by the Royal Society (URF to BH grant no. UF130039,

studentship to VK), Medical Research Council (Grant-in-Aid to the MRC Cancer unit

grant number MC_UU_12022/9 and NIRG to BH grant number MR/S000216/1), and

the Harrison Watson Fund at Clare College, Cambridge (M.W.J.H.).

## AUTHOR CONTRIBUTIONS

DS and BH conceived the study, DS curated the data and performed all analysis.

MH wrote code for DN/DS style mutation frequency, ER and CK performed cell

culture and prepared RNA for sequencing, GD supported dataset processing and

analysis. RCF provided the oesophageal data and expertise on human GI cancer.

DS, BH, and RCF cowrote the manuscript. All authors were responsible for editing of

the manuscript.

## DECLARATION OF INTERESTS

678 The authors declare no competing interests.

**FIGURE LEGENDS**

Figure 1:

KCNQ genes are highly altered in GI cancer.

A) Mutations in any member of the KCNQ/KCNE family across all cancers in The Cancer Genome Atlas (TCGA)

B) Percentage of patients with an alteration in a KCNQ gene for 5 cohorts: Oesophageal Adenocarcinoma (EAC – OCCAMS and TCGA), Oesophageal Squamous Cell Carcinoma (ESCC), Stomach Adenocarcinoma (STAD), and Colorectal Adenocarcinoma (COADREAD).

C) Percentage of patients across all cohorts with an alteration in each KCNQ/KCNE gene.

D) Oncoprint showing alterations in KCNQ/KCNE genes per patient.

Figure 2:

Mutations in KCNQ genes cluster in cancer

A) Mutational Frequency for all KCNQ genes after sequence alignment

B) Mutational clustering for KCNQ1, line represents Observed vs Expected mutation ratio

C) Mutational clustering for KCNQ3, line represents Observed vs Expected

mutation ratio

D) Mutational clustering for KCNQ5, line represents Observed vs Expected

mutation ratio

Figure 3:

Mutations in KCNQ genes cluster in space

A) Schematic of the assembly of KCNQ/KCNE genes.

B) Bayesian inference analysis of the RNA expression levels of KCNQ/KCNE

genes in the colorectal (COADREAD) dataset.

C) Schematic of the KCNQ canonical structure.

D) Render of the KCNQ1 structure.

E) Render of the KCNQ3 structure with mutational frequency overlaid, darker red

regions represent a higher mutational frequency.

F) Spatial clustering plot for KCNQ1 (top) and KCNQ3 (bottom), arcs represent

residues that are within 12Å of each other, size of each circle represents the

number of mutations occurring at that location in the structure. Labelled are

mutational clusters identified in figure 2 B.

G) Structure of KCNQ1 with mutational clusters highlighted in red (cluster 1.1),

and blue (cluster 1.2).

H) Structure of KCNQ3 with mutational clusters highlighted in pink (cluster 3.1),

and orange (cluster 3.2).

Figure 4:

Mutations in KCNQ1 and KCNQ3 in GI cancers impact protein function.

A) Render of the pore region of KCNQ1. Residues mutated in cluster 1.2 are

729            rendered in blue. Mutations that constrict the pore region when modelled are

730            starred.

731        B) HOLE analysis of the pore region of KCNQ1 wt (black) WT mutations in

732            cluster 1.2 (coloured according to Figure 4, A).

733        C) Mutational frequency from the COSMIC GI cancer dataset for the S4 region of

734            the KCNQ family. Residues involved in gating (R1, R2, Q3, R4, R5, R6) are

735            highlighted (top).

736        D) Helical tilt angle for alanine scanning mutagenesis for the KCNQ1 S4 helix.

737        E) Helical tilt angle for mutational analysis of KCNQ3 S4 helix.

738        F) Residues involved in stabilizing the active conformation of KCNQ2 through

739            hydrogen bonding.

740        G) Hydrogen bonding networks generated from molecular dynamics trajectories

741            for KCNQ3 (left) and KCNQ5 (right). Percentages represent total bond

742            occupancy over 100 ns equilibrium molecular dynamics.

744    Figure 5:

745    KCNQ1, KCNQ3, and KCNQ5 gene expression correlates with WNT signalling and

746    EMT

747        A) Gradient Boosted Decision Forest (GBDF) based pathway analysis results for

748            comparing patients that are high and low for KCNQ1 (grey), KCNQ3 (blue)

749            and KCNQ5 (orange). Pathways are ranked by average Matthews Correlation

750            Coefficient.

B) Feature extraction schematic for WNT pathway classification. Genes with a

high Shapely Additive Explanation (SHAP score > 0.1) are counted for each

model

C) WNT pathway RNA expression clustering between patients high (dark) and

low (light) for KCNQ1 (left), KCNQ3 (middle), and KCNQ5 (right).

D) Feature extraction for GBDF model of the EMT pathway when classifying

patients high and low for KCNQ1 (grey), KCNQ3 (blue), and KCNQ5 (orange).

E) Ranking metric of KCNQ genes for the EMT pathway. KCNQ genes are

ranked against all genes for ability to cluster EMT pathway genes between

high and low expressing patients.

F) EMT score for KCNQ1 (left), KCNQ3 (middle), KCNQ5 (right). Red line

represents EMT score for EMT associated transcription factor SNAI1 (r =

0.49, p < 0.01).

G) GSEA hallmark: EMT analysis for high vs low expressors of KCNQ1 (top),

KCNQ3 (middle), and KCNQ5 (bottom).

Figure 6:

KCNQ expression correlates with clinical prognosis:

A) Survival analysis for KCNQ1 (left), KCNQ3 (middle), and KCNQ5 (right) high

vs low expressers.

B) Survival analysis for patients with different co-expressions of KCNQ1 vs

KCNQ3 (left), and KCNQ1 vs KCNQ5 (right). Schematics indicate up and

downregulation of KCNQ1 (grey), KCNQ3 (blue), and KCNQ5 (orange).

C) Survival analysis for patients with different co-expressions of KCNQ3 vs

KCNQ5. Schematics indicate up and downregulation of KCNQ3 (blue), and

KCNQ5 (orange).

D) Venn diagram of overlapping genes from differential gene expression analysis

of patients in 6 cohorts divided into 3 sets. Shown are differentially expressed

(corrected $p < 0.05$, $\log_2$ fold change > 1.5) genes for comparison of KCNQ1

high/KCNQ3 low patients vs KCNQ1 low/KCNQ3 high patients (blue), KCNQ1

high/KCNQ5 low patients vs KCNQ1 high/KCNQ5 low patients (pink), and

KCNQ3 low/KCNQ5 low patients vs KCNQ3 high/KCNQ5 high patients (blue).

Grey region represents genes differentially expressed in all cases.

E) RNA expression of WNT associated transcription factors between patients

with high KCNQ1/low KCNQ3 (red), high KCNQ1/high KCNQ3 (blue), low

KCNQ1/low KCNQ3 (pink), and low KCNQ1/high KCNQ3 (orange).

F) Schematic of systems biology model representing influence of KCNQ1 and

KCNQ3/5 on cellular phenotype.

803 **TABLES**

| Gene | Mutation | No. Obs. | Specific Effect | Functional Effect | Clinical Impact | References |
|------|----------|----------|-----------------|-------------------|-----------------|------------|
| KCNQ1 | K422fs | 6 | Does not traffic | LOF | Long QT Syndrome | Harmer et al 2014 |
| KCNQ1 | F339L | 3 | Reduced Currents | LOF | Romano-Ward Syndrome | Thomas et al (2005) |
| KCNQ1 | K422R | 2 | Reduced Currents | LOF | Long QT Syndrome | Veerman et al (2013) |
| KCNQ1 | P343S | 2 | Reduced Currents | LOF | Romano-Ward Syndrome | Zehelein et al (2004) |
| KCNQ1 | R293H | 2 | Loss of Predicted salt bridge to KCNE1 | - | - | Xu et al (2013) |
| KCNQ1 | R192C | 2 | Loss of PIP2 binding site | LOF | Long QT Syndrome | Zaydman et al (2013), Eckey et al (2014) |
| KCNQ1 | R192H | 2 | Loss of PIP2 binding site | LOF | Long QT Syndrome | Zaydman et al (2013), Eckey et al (2014) |
| KCNQ2 | R210H | 4 | Increased Activation threshold | LOF | Benign Familial Neonatal Convulsions | Neverisky et al (2017) |
| KCNQ2 | A294V | 4 | Reduced currents | LOF | Epilepsy of Infancy with Migrating Focal Seizures | Duan et al (2018) |
| KCNQ2 | E119D | 2 | Negative Shift in Activation Threshold | LOF | Benign Familial Neonatal Convulsions | Wuttke et al (2008) |
| KCNQ2 | G301S | 2 | Loss of Retigabine Binding Site | - | - | Kalappa et al (2015) |
| KCNQ2 | R581Q | 2 | - | - | Benign Familial Neonatal Convulsions | Singh et al (2003) |
| KCNQ3 | K533fs | 9 | Loss of PIP2 binding site | - | - | Choveau et al (2018) |
| KCNQ3 | R230C | 5 | Increased current amplitude, increased activation | GOF | Neurodevelopmental Disability | Sands et al (2019) |
| KCNQ3 | R230H | 5 | Increased current amplitude, increased activation | GOF | Neurodevelopmental Disability | Sands et al (2019) |
| KCNQ3 | K366N | 4 | Loss of PIP2 binding site | - | - | Choveau et al (2018) |
| KCNQ3 | R227Q | 3 | Reduced Activation Threshold | GOF | Neurodevelopmental Disability | Sands et al (2019) |

805  Table 1: KCNQ mutations in GI Cosmic patients. Shown are mutations with known

806  functional or clinical consequences

| Gene | Mutation | Equivalent Name | No. Obs. | Specific Effect | Functional Effect | Sidekick Result | Reference |
|---|---|---|---|---|---|---|---|
| KCNQ1 | R228A | R1A | - | Impaired activation | LOF | No change | Panaghie et al (2007) |
| KCNQ1 | R231A | R2A | - | Constitutively open | GOF | Decrease tilt | Panaghie et al (2007) |
| KCNQ1 | R237A | R4A | - | Accumulates in the open state | GOF | Decrease tilt | Panaghie et al (2007) |
| KCNQ1 | D242A | D6A | - | More difficult to activate | LOF | Increase tilt | Panaghie et al (2007) |
| KCNQ1 | R243A | R6A | - | Impaired activation | LOF | Increase tilt | Panaghie et al (2007) |
| KCNQ2 | R201H | R2H | - | Reduced Activation Threshold | GOF | Decrease tilt | Miceli et al (2015) |
| KCNQ2 | R201C | R2C | - | Constitutively Active | GOF | Decrease tilt | Miceli et al (2015) |
| KCNQ2 | I205V | - | - | Loss of function | LOF | No change | Niday et al (2017) |
| KCNQ2 | R207W | R4W | - | Increased Activation Threshold | LOF | Decrease tilt | Dedek et al (2001) |
| KCNQ2 | R210H | R5H | 4 | - | - | No change | Rikee database[1] |
| KCNQ2 | R213W | R6W | - | Increased Activation Threshold | LOF | Increase tilt | Miceli et al (2013) |
| KCNQ2 | R198Q | R1Q | - | Reduced Activation Threshold | GOF | No change | Miceli et al (2008) |
| KCNQ2 | R201Q | R2Q | - | Constitutively Active | GOF | Decrease tilt | Miceli et al (2008) |
| KCNQ2 | R207Q | R4Q | 1 | Increased Activation Threshold | LOF | No change | Miceli et al (2008) |
| KCNQ2 | R210Q | R5Q | - | Increased Activation Threshold | LOF | No change | Miceli et al (2008) |
| KCNQ2 | R213Q | R6Q | - | Increased Activation Threshold | LOF | No change | Miceli et al (2008) |
| KCNQ2 | R214Q | R7Q | - | Increased Activation Threshold | LOF | No change | Miceli et al (2008) |
| KCNQ3 | R227Q | R1Q | 3 | Reduced Activation Threshold | GOF | No change | Sands et al (2019) |
| KCNQ3 | R230H | R2H | 4 | Reduced Activation Threshold | GOF | Decrease tilt | Sands et al (2019) |
| KCNQ3 | R230C | R2C | 1 | Reduced Activation Threshold | GOF | Decrease tilt | Sands et al (2019) |
| KCNQ3 | R236C | R4C | 2 | - | - | Decrease tilt | - |
| KCNQ3 | R236H | R4H | 4 | - | - | Decrease tilt | - |
| KCNQ3 | R239W | R5W | 2 | - | - | No change | - |
| KCNQ3 | R242W | R6W | 1 | - | - | No change | - |
| KCNQ4 | R207H | R2H | 2 | - | - | Decrease tilt | - |
| KCNQ4 | R207C | R2C | 1 | - | - | Decrease tilt | - |
| KCNQ4 | R213C | R4C | 1 | - | - | Decrease tilt | - |

---

[1] Rikee database at https://rikee.org

| KCNQ4 | R216H | R5H | - | Reduced Activation Threshold | GOF | Decrease tilt | Panaghie et al (2007) |
|-------|-------|-----|---|------------------------------|-----|---------------|-----------------------|
| KCNQ5 | R244H | R5H | 3 | - | - | No change | - |
| KCNQ5 | R244C | R5C | 6 | - | - | No change | - |

808    Table 2: Results of Molecular Dynamics Simulations on single S4 helices from the

809    KCNQ family.

## METHODS

<u>TCGA Data</u>

TCGA level 3 data was downloaded using Firebrowse (RNAseq), cBioportal (CNAs,

mutation and clinical data)(Gao et al., 2013), or TCGA wanderer (Methylation

data)(Díez-Villanueva et al., 2015).

<u>COSMIC Data</u>

COSMIC data(Forbes et al., 2017; Tate et al., 2019) was downloaded from

cancer.sanger.ac.uk. We subset mutations into those only found in gastrointestinal

tissue, defined as those where the primary site is in one of the following categories:

"large_intestine", "small_intestine", "gastrointestinal_tract_(site_indeterminate)",

"oesophagus", "stomach".

<u>Oncoprints</u>

Oncoprints were produced using the R library Oncoprint. Copy number alterations

were determined as follows – relative copy number for each gene was defined as:

$$relative\ copy\ number = \ log_2(\frac{Total\ Copy\ Number\ of\ Gene}{Total\ ploidy\ of\ sample})$$

Genes were defined as deleted if total copy number == 0 OR relative copy number <

-1, genes were defined as amplified if relative copy number > 1.

<u>Mutual Exclusivity/Cooperativity</u>

Mutual exclusivity and cooperativity for KCNQ genes was calculated using the

DISCOVER library in R(Canisius et al., 2016).

<u>Sequence alignments</u>

Sequence alignment was performed using the MUSCLE algorithm(Edgar, 2004),

implemented in python through Biopython(Cock et al., 2009), using default settings.

FASTA files for KCNQ sequences were obtained from uniprot using codes P51787

(KCNQ1), O43526 (KCNQ2), O43525 (KCNQ3), P56696 (KCNQ4), and Q9NR82

(KCNQ5).

Expected vs Observed mutational distribution

For calculating the expected vs observed mutational distribution, exon data was

downloaded from the Ensembl Biomart (ensembl.org/biomart). Ensembl 96,

hg38.p12 was selected, and data downloaded for chromosomes 1-22, X, and Y. We

used bedtools(Quinlan and Hall, 2010) to sort data and overlapping exons were

merged. To sort data we used the following command:

tail -n +2 human_exon_bed_1-Y.txt | cut -f 1,2,3 | bedtools sort -i stdin > human_exon_bed_file_1-Y_sorted.bed

Merging was performed using:

bedtools merge -i human_exon_bed_file_1-Y_sorted.bed > human_exon_bed_file_1-Y_merged.bed

The R library Deconstructsigs(Rosenthal et al., 2016) was used to generate the

mutational signature for all COSMIC mutations in KCNQ genes within GI tissue for

these exons. The mutational spectrum was normalised using the mutational

signature to generate the expected relative mutation rate for each possible missense

mutation. This was then multiplied by the total number of mutations in each gene to

get the expected distribution of events along the gene. The observed and expected

mutational frequency ratio was average over a sliding window of 50 bases.

1D Mutational Clustering

Mutational clustering was calculated using the NMC clustering method from the R

library IPAC(Ye et al., 2010) applied to sequence alone. All mutations to each KCNQ

gene in the COSMIC database for GI cancers were considered. The top 5 mutational

clusters ranked by adjusted P-value were plotted.

Fathmm/CScape score calculation

To generate fathmm(Shihab et al., 2013) and CScape(Rogers et al., 2017) scores for

each mutation we downloaded the fathmm and cscape databases

(fathmm.biocompute.org.uk, cscape.biocompute.org.uk). We constructed a query for

each database of every possible mutation in each KCNQ gene of interest using the

python scripts available with each database and pyensembl using GRCH37 and the

canonical transcript.

<u>Bayesian inference analysis</u>

RSEM normalised RNAseq data was used to calculate the association of KCNQ/E

genes with each other. The covariance matrix for KCNQ genes was calculated and

converted to correlation matrix. Permutation tests (N = 5000) were performed to

calculate a p-value testing the null hypothesis that the corresponding pair of

variables has 0 correlation. P-values of less than 0.05 were deemed a significant co-

association.

<u>Homology modelling</u>

Homology modelling was performed using the template structure 5VMS from the

protein data bank(Sun and MacKinnon, 2017). Modeller was used to generate 5

models, and the model with the highest GA341 (nativity) scores, and lowest DOPE

(total energy of the structure) scores was selected. Alignments used to generate the

models are available in the SI. Single point mutations were induced in the models

using the mutate_model protocol in the modeller tool as described in Feyfant et al

(Feyfant et al., 2007), and using the mutate_model.py script available on the

modeller website (https://salilab.org/modeller/wiki/Mutate%20model).

<u>Pore analysis</u>

Pore analysis was performed using the algorithm HOLE (Smart et al., 1996). Pore

profile was visualised using Visual Molecular Dynamics (VMD) (Humphrey et al.,

1996).

<u>Molecular dynamics simulations</u>

Molecular dynamics was performed using Gromacs version 2018.1(Abraham et al.,

2015).

For simulations of homology models in AT we used the charmm36 forcefield(Lee et

al., 2014). In each case the protein was placed in a 15 x 15 x 15nm box with roughly

650 DPPC lipid molecules. Setup was performed in the same manner as systems in

the memprotMD pipeline (Stansfeld et al., 2015). The system was converted to

MARTINI coarse-grained structures (CG-MD) with an elastic network in the martiniv2

forcefield(Monticelli et al., 2008) and self-assembled by running a 1000ns molecular

dynamics simulation at 323k to allow the formation of the bilayer around the protein.

The final frame of the CG-MD simulation was converted back to atomistic detail

using the CG2AT method (Stansfeld and Sansom, 2011). The AT system was

neutralised with counterions, and additional ions added up to a total NACL

concentration of 0.05 mol/litre. The system was minimized using the steepest

descents algorithm until maximum force Fmax of the system converged.

Equilibration was performed using NVT followed by NPT ensembles for 100 ps each

with the protein backbone restrained. We used the Verlet cutoff scheme with PME

electrostatics, and treated the box as periodic in the X, Y, and Z planes. Simulations

were run for 200ns of unrestrained molecular dynamics. Root mean square deviation

(RMSD) was calculated for structures using the g_rmsdist command in GROMACS.

CG simulations of single helices were performed as described previously (Hall et al.,

2014). Models of single helices were generated and converted to MARTINI coarse

grained structures. Helices were then inserted into POPC bilayers and simulated for

short (100ns) simulations for 100 repeats of each sequence.

H-bond analysis

H-bond occupancy for molecular dynamics simulations was calculated using the

VMD HBonds plugin.

Gradient Boosted Decision Forests

Gradient boosted decision forests were implemented using Xgboost in the python

package scikit-learn(Chen and Guestrin, 2016). RNAseq data from TCGA and

OCCAMS was combined. For training the models, a target gene is selected, and the

top and bottom 50 expressors selected. A model was then trained to classify the 100

patients into high and low expressors based only on genes from a query pathways.

Pathways and genes were generated from Sanchez-Vega et al (Sanchez-Vega et

al., 2018). We trained 20 separate models for each pathway, in each case a random

20% (10) patients were removed from the dataset and used for testing. The model
was run using the command:

Model = XGBClassifier(max_depth = 3, n_estimators = 100, learning_rate = 0.1, colsample_bytree = 0.3)

The model was fitted using the logloss evaluation metric and stopped using the early
stopping method 40 epochs prior to a loss of accuracy. The method was tested on
known genes from established pathways (KRAS in the KRAS pathway, CTNNB1 in
the WNT pathway, and P53 in the P53 pathway). In these cases the query gene was
removed from the respective pathway in order to unbias the model towards them.
Models were evaluated and interpreted by generating the SHAP score using the
SHAP implementation in python (Lundberg and Lee, 2017).

<u>Gene ranking</u>

For ranking individual genes against all other genes in the dataset for correlation with
a gene set, a target gene is selected, and the top and bottom 50 expressors from the
RNAseq dataset were selected. The RNAseq data is then subset by genes in a
geneset (for EMT this subset consists of: CDH1, DSP, TJP1, VIM, CDH2, FOXC2,
SNAI1, SNAI2, TWIST1, GSC, FN1, ITBG6, MMP2, MMP3, MMP9, and SOX10 – as
defined by Chae et al (Chae et al., 2018), and the optimally ordered linkage is
calculated for all patients using the hierarchical clustering method from scipy. The
command used to generate the linkage is:

leave_order = scipy.cluster.hierarchy.linkage(gene_expression_matrix, optimal_ordering = True)

Where gene_expression_matrix is a matrix of gene expression values for a pathway
and the top and bottom 50 patients for expression of a query gene. The order of
leaves in the tree is calculated and converted to binary format (where 0 is low
expressors and high expressors). We infer that genes highly associated with a
geneset query will cluster well, and so the binary string will contain contiguous
strings of 1s and 0s. To calculate the "orderedness" of the binary string we calculate
two metrics:

1) The longest continuous string of 1s or 0s in the binary string
2) The number of times a 1 is flipped to a 0 or vice versa

We calculate these two scores for every gene in the dataset for the query pathway.

We plot the resultant values against one another to get the degree to which a set of

query genes correlates with every gene in the dataset.

Gene Set Enrichment Analysis

Gene set enrichment was performed using the GSEA desktop application

(Subramanian et al., 2005). We subset patients into the 25 highest and lowest

expressors for each of KCNQ1, KCNQ3, and KCNQ5. GSEA was run for 1000

permutations using the geneperm permuation method for the hallmarks and

oncogenic signatures gene sets (h.all.v7.0 and c6.all.v7.0 respectively) and ranked

by Ttest.

EMT score

EMT score was calculated following the protocol defined by Chae et al (Chae et al.,

2018). RNAseq expression data for each sample in the TCGA and OCCAMS dataset

was z-score normalised and genes corresponding to epithelial and mesenchymal

phenotypes were extracted. We classed CDH1, DSP, and TJP1 as epithelial

markers, and VIM, CDH2, FOXC2, SNAI1, SNAI2, TWIST1, GSC, FN1, ITBG6,

MMP2, MMP3, MMP9, and SOX10 as mesenchymal. The EMT score for each

patient sample was calculated by subtracting the mean z-score of epithelial markers

from the mean for mesenchymal markers. For comparison we calculated the

correlation for SNAI2, a known driver of EMT. For correlation of EMT scores against

gene expression, we calculated the spearmans correlation for the TPM expression of

a query gene against the EMT score.

TCPA analysis

For analysis of data from The Cancer Protein Atlas (TCPA) (Li et al., 2013)we

downloaded RPPA data from the TCPA portal (tcpaportal.org). We matched samples

from the TCPA with their respected TCGA ID, and took only patients where both

RNAseq and protein level data was available. Patients were then split into the

highest and lowest 25 expressors of a query gene in the RNA expression data, and

protein level data compared. High and low expression cohorts were compared using

a students T test, and multiple test corrected using the Bonferroni method.

Survival analysis

Survival analysis was performed using the R library Survival. We used the clinical

data associated with the TCGA and OCCAMS, which includes overall survival and

recurrence free survival data. Survival times were converted to days, and survival

curves generated on the top and bottom 25% expressors of each query gene.

<u>Differential expression analysis</u>

Differential expression analysis was performed using the R library Deseq2 (Love et

al., 2014).

<u>Systems biology</u>

Systems biology models were constructed with the BioModelAnalyzer tool (Chuang

et al., 2015). Nodes were assigned to genes or interest, and relationships inferred

through data analysis and through studying the literature (see **Table S7** for

interactions in the models)

**REFERENCES**

Abbott, G.W. (2014). Biology of the KCNQ1 Potassium Channel. New J. Sci.

Abraham, M.J., Murtola, T., Schulz, R., Páll, S., Smith, J.C., Hess, B., and Lindahl,

E. (2015). GROMACS: High performance molecular simulations through multi-level

parallelism from laptops to supercomputers. SoftwareX *1–2*, 19–25.

Aggarwal, S.K., and MacKinnon, R. (1996). Contribution of the S4 Segment to

Gating Charge in the Shaker K+ Channel. Neuron *16*, 1169–1177.

Allen, N.M., Mannion, M., Conroy, J., Lynch, S.A., Shahwan, A., Lynch, B., and King,

M.D. (2014). The variable phenotypes of KCNQ-related epilepsy. Epilepsia.

Arcangeli, A., and Becchetti, A. (2017). hERG Channels: From Antitargets to Novel

Targets for Cancer Therapy. Clin. Cancer Res. *23*, 3–5.

Bass, A.J., Thorsson, V., Shmulevich, I., Reynolds, S.M., Miller, M., Bernard, B., Hinoue, T., Laird, P.W., Curtis, C., Shen, H., et al. (2014). Comprehensive molecular characterization of gastric adenocarcinoma. Nature *513*, 202–209.

Brisson, L., Gillet, L., Calaghan, S., Besson, P., Le Guennec, J.-Y., Roger, S., and Gore, J. (2011). Na(V)1.5 enhances breast cancer cell invasiveness by increasing NHE1-dependent H(+) efflux in caveolae. Oncogene *30*, 2070–2076.

Brown, D.A., and Passmore, G.M. (2009). Neural KCNQ (Kv7) channels. Br. J. Pharmacol.

Campbell, P.J. (2017). Cliques and Schisms of Cancer Genes. Cancer Cell *32*, 129–130.

Canisius, S., Martens, J.W.M.M., and Wessels, L.F.A.A. (2016). A novel independence test for somatic alterations in cancer shows that biology drives mutual exclusivity but chance explains most co-occurrence. Genome Biol. *17*, 261.

Chae, Y.K., Chang, S., Ko, T., Anker, J., Agte, S., Iams, W., Choi, W.M., Lee, K., and Cruz, M. (2018). Epithelial-mesenchymal transition (EMT) signature is inversely associated with T-cell infiltration in non-small cell lung cancer (NSCLC). Sci. Rep.

Chalmers, Z.R., Connelly, C.F., Fabrizio, D., Gay, L., Ali, S.M., Ennis, R., Schrock, A., Campbell, B., Shlien, A., Chmielecki, J., et al. (2017). Analysis of 100,000 human cancer genomes reveals the landscape of tumor mutational burden. Genome Med. *9*, 34.

Chen, T., and Guestrin, C. (2016). XGBoost. In Proceedings of the 22nd ACM SIGKDD International Conference on Knowledge Discovery and Data Mining - KDD

'16, (New York, New York, USA: ACM Press), pp. 785–794.

Chuang, R., Hall, B.A., Benque, D., Cook, B., Ishtiaq, S., Piterman, N., Taylor, A.,

Vardi, M., Koschmieder, S., Gottgens, B., et al. (2015). Drug Target Optimization in

Chronic Myeloid Leukemia Using Innovative Computational Platform. Sci. Rep. *5*,

8190.

Cock, P.J.A., Antao, T., Chang, J.T., Chapman, B.A., Cox, C.J., Dalke, A., Friedberg,

I., Hamelryck, T., Kauff, F., Wilczynski, B., et al. (2009). Biopython: freely available

Python tools for computational molecular biology and bioinformatics. Bioinformatics

*25*, 1422–1423.

Creighton, C., and Gibbons, D. (2013). Pan-cancer survey of epithelial–

mesenchymal transition markers across The Cancer Genome Atlas. Dev. Dyn.

DeMarco, M.L. (2012). Three-Dimensional Structure of Glycolipids in Biological

Membranes. Biochemistry *51*, 5725–5732.

Díez-Villanueva, A., Mallona, I., and Peinado, M.A. (2015). Wanderer, an interactive

viewer to explore DNA methylation and gene expression data in human cancer.

Epigenetics and Chromatin.

Edgar, R.C. (2004). MUSCLE: Multiple sequence alignment with high accuracy and

high throughput. Nucleic Acids Res.

Fan, H., Zhang, M., and Liu, W. (2018). Hypermethylated KCNQ1 acts as a tumor

suppressor in hepatocellular carcinoma. Biochem. Biophys. Res. Commun. *503*,

3100–3107.

Feske, S., Wulff, H., and Skolnik, E.Y. (2015). Ion channels in innate and adaptive

immunity. Annu. Rev. Immunol. *33*, 291–353.

Feyfant, E., Sali, A., and Fiser, A. (2007). Modeling mutations in protein structures.

Protein Sci.

Forbes, S.A., Beare, D., Boutselakis, H., Bamford, S., Bindal, N., Tate, J., Cole,

C.G., Ward, S., Dawson, E., Ponting, L., et al. (2017). COSMIC: Somatic cancer

genetics at high-resolution. Nucleic Acids Res.

Frankell, A.M., Jammula, S., Li, X., Contino, G., Killcoyne, S., Abbas, S., Perner, J.,

Bower, L., Devonshire, G., Ococks, E., et al. (2019). The landscape of selection in

551 esophageal adenocarcinomas defines genomic biomarkers for the clinic. Nat.

Genet. *51*, 506–516.

Fukushiro-Lopes, D.F., Hegel, A.D., Rao, V., Wyatt, D., Baker, A., Breuer, E.-K.,

Osipo, C., Zartman, J.J., Burnette, M., Kaja, S., et al. (2018). Preclinical study of a

Kv11.1 potassium channel activator as antineoplastic approach for breast cancer.

Oncotarget *9*, 3321–3337.

Gao, J., Aksoy, B.A., Dogrusoz, U., Dresdner, G., Gross, B., Sumer, S.O., Sun, Y.,

Jacobsen, A., Sinha, R., Larsson, E., et al. (2013). Integrative analysis of complex

cancer genomics and clinical profiles using the cBioPortal. Sci. Signal.

Gibbons, D.L., and Creighton, C.J. (2018). Pan-cancer survey of epithelial-

mesenchymal transition markers across the Cancer Genome Atlas. Dev. Dyn. *247*,

555–564.

Grahammer, F., Herling, A.W., Lang, H.J., Schmitt-Gräff, A., Wittekindt, O.H.,

Nitschke, R., Bleich, M., Barhanin, J., and Warth, R. (2001). The cardiac K+ channel

KCNQ1 is essential for gastric acid secretion. Gastroenterology *120*, 1363–1371.

Haas, B.R., and Sontheimer, H. (2010). Inhibition of the sodium-potassium-chloride

cotransporter isoform-1 reduces glioma invasion. Cancer Res.

Hall, B.A., Halim, K.B.A., Buyan, A., Emmanouil, B., and Sansom, M.S.P. (2014).

Sidekick for Membrane Simulations: Automated Ensemble Molecular Dynamics

Simulations of Transmembrane Helices. J. Chem. Theory Comput. *10*, 2165–2175.

Heitzmann, D., Grahammer, F., von Hahn, T., Schmitt-Gräff, A., Romeo, E.,

Nitschke, R., Gerlach, U., Lang, H.J., Verrey, F., Barhanin, J., et al. (2004).

Heteromeric KCNE2/KCNQ1 potassium channels in the luminal membrane of gastric

parietal cells. J. Physiol.

Hoadley, K.A., Yau, C., Hinoue, T., Wolf, D.M., Lazar, A.J., Drill, E., Shen, R., Taylor,

A.M., Cherniack, A.D., Thorsson, V., et al. (2018). Cell-of-Origin Patterns Dominate

the Molecular Classification of 10,000 Tumors from 33 Types of Cancer. Cell *173*,

291-304.e6.

Howard, R.J., Clark, K.A., Holton, J.M., Minor, D.L., and Jr. (2007). Structural insight

into KCNQ (Kv7) channel assembly and channelopathy. Neuron *53*, 663–675.

Humphrey, W., Dalke, A., and Schulten, K. (1996). VMD: Visual molecular dynamics.

J. Mol. Graph. *14*, 33–38, 27–28.

Jentsch, T.J. (2000). Neuronal KCNQ potassium channels: Physislogy and role in

disease. Nat. Rev. Neurosci.

Kharkovets, T., Hardelin, J.-P., Safieddine, S., Schweizer, M., El-Amraoui, A., Petit,

C., and Jentsch, T.J. (2002). KCNQ4, a K+ channel mutated in a form of dominant

deafness, is expressed in the inner ear and the central auditory pathway. Proc. Natl.

Acad. Sci.

Kim, J., Bowlby, R., Mungall, A.J., Robertson, A.G., Odze, R.D., Cherniack, A.D.,

Shih, J., Pedamallu, C.S., Cibulskis, C., Dunford, A., et al. (2017). Integrated

genomic characterization of oesophageal carcinoma. Nature *541*, 169–174.

Kubisch, C., Schroeder, B.C., Friedrich, T., Lütjohann, B., El-Amraoui, A., Marlin, S.,

Petit, C., and Jentsch, T.J. (1999). KCNQ4, a novel potassium channel expressed in

sensory outer hair cells, is mutated in dominant deafness. Cell.

Lee, S., Tran, A., Allsopp, M., Lim, J.B., Hénin, J., and Klauda, J.B. (2014).

CHARMM36 united atom chain model for lipids and surfactants. J. Phys. Chem. B.

Li, J., Lu, Y., Akbani, R., Ju, Z., Roebuck, P.L., Liu, W., Yang, J.Y., Broom, B.M.,

Verhaak, R.G.W., Kane, D.W., et al. (2013). TCPA: A resource for cancer functional

proteomics data. Nat. Methods.

Li, J., Akbani, R., Zhao, W., Lu, Y., Weinstein, J.N., Mills, G.B., and Liang, H. (2017).

Explore, Visualize, and Analyze Functional Cancer Proteomic Data Using the Cancer

Proteome Atlas. Cancer Res. *77*, e51–e54.

Li, X., Francies, H.E., Secrier, M., Perner, J., Miremadi, A., Galeano-Dalmau, N.,

Barendt, W.J., Letchford, L., Leyden, G.M., Goffin, E.K., et al. (2018). Organoid

cultures recapitulate esophageal adenocarcinoma heterogeneity providing a model

for clonality studies and precision therapeutics. Nat. Commun. *9*, 2983.

Love, M.I., Huber, W., and Anders, S. (2014). DESeq2.

Lundberg, S.M., and Lee, S.-I.I. (2017). A unified approach to interpreting model

predictions. In Advances in Neural Information Processing Systems, pp. 4765–4774.

Lundberg, S.M., Erion, G.G., and Lee, S.-I. (2018). Consistent Individualized Feature

Attribution for Tree Ensembles.

Manville, R.W., and Abbott, G.W. (2019). In silico re-engineering of a

neurotransmitter to activate KCNQ potassium channels in an isoform-specific manner. Commun. Biol. *2*, 401.

Martincorena, I., Raine, K.M., Gerstung, M., Dawson, K.J., Haase, K., Van Loo, P., Davies, H., Stratton, M.R., and Campbell, P.J. (2017). Universal Patterns of Selection in Cancer and Somatic Tissues. Cell *171*, 1029-1041.e21.

Miceli, F., Soldovieri, M.V., Hernandez, C.C., Shapiro, M.S., Annunziato, L., and Taglialatela, M. (2008). Gating consequences of charge neutralization of arginine residues in the S4 segment of Kv7.2, an epilepsy-linked K+ channel subunit. Biophys. J.

Miceli, F., Soldovieri, M.V., Ambrosino, P., De Maria, M., Migliore, M., Migliore, R., and Taglialatela, M. (2015). Early-onset epileptic encephalopathy caused by gain-of-function mutations in the voltage sensor of Kv7.2 and Kv7.3 potassium channel subunits. J. Neurosci. *35*, 3782–3793.

Millichap, J.J., Park, K.L., Tsuchida, T., Ben-Zeev, B., Carmant, L., Flamini, R., Joshi, N., Levisohn, P.M., Marsh, E., Nangia, S., et al. (2016). *KCNQ2* encephalopathy. Neurol. Genet. *2*, e96.

Monticelli, L., Kandasamy, S.K., Periole, X., Larson, R.G., Tieleman, D.P., and Marrink, S.-J. (2008). The MARTINI Coarse-Grained Force Field: Extension to Proteins. J. Chem. Theory Comput. *4*, 819–834.

Morita, H., Wu, J., and Zipes, D.P. (2008). The QT syndromes: long and short. Lancet *372*, 750–763.

Muzny, D.M., Bainbridge, M.N., Chang, K., Dinh, H.H., Drummond, J.A., Fowler, G., Kovar, C.L., Lewis, L.R., Morgan, M.B., Newsham, I.F., et al. (2012). Comprehensive

molecular characterization of human colon and rectal cancer. Nature *487*, 330–337.

Nerbonne, J.M., and Kass, R.S. (2005). Molecular Physiology of Cardiac Repolarization. Physiol. Rev.

Ohya, S., Asakura, K., Muraki, K., Watanabe, M., and Imaizumi, Y. (2015). Molecular and functional characterization of ERG, KCNQ, and KCNE subtypes in rat stomach smooth muscle. Am. J. Physiol. Liver Physiol.

Panaghie, G., and Abbott, G.W. (2007). The role of S4 charges in voltage-dependent and voltage-independent KCNQ1 potassium channel complexes. J. Gen. Physiol. *129*, 121–133.

Pardo, L.A., and Stühmer, W. (2013). The roles of K+ channels in cancer. Nat. Rev. Cancer *14*, 39–48.

Pedersen, S.F., and Stock, C. (2013). Ion channels and transporters in cancer: pathophysiology, regulation, and clinical potential. Cancer Res. *73*, 1658–1661.

Pedersen, S.F., Hoffmann, E.K., and Novak, I. (2013). Cell volume regulation in epithelial physiology and cancer. Front. Physiol. *4*, 233.

Phan, N.N., Wang, C.-Y., Chen, C.-F., Sun, Z., Lai, M.-D., and Lin, Y.-C. (2017). Voltage-gated calcium channels: Novel targets for cancer therapy. Oncol. Lett. *14*, 2059–2074.

Pillozzi, S., Brizzi, M., Balzi, M., Crociani, O., Cherubini, A., Guasti, L., Bartolozzi, B., Becchetti, A., Wanke, E., Bernabei, P., et al. (2002). HERG potassium channels are constitutively expressed in primary human acute myeloid leukemias and regulate cell proliferation of normal and leukemic hemopoietic progenitors. Leukemia *16*, 1791–1798.

Pointer, K.B., Clark, P.A., Eliceiri, K.W., Salamat, M.S., Robertson, G.A., and Kuo,

J.S. (2017). Administration of Non-Torsadogenic human Ether-à-go-go-Related

Gene Inhibitors Is Associated with Better Survival for High hERG–Expressing

Glioblastoma Patients. Clin. Cancer Res. *23*, 73–80.

Qu, Z., Yao, W., Yao, R., Liu, X., Yu, K., and Hartzell, C. (2014). The Ca(2+) -

activated Cl(-) channel, ANO1 (TMEM16A), is a double-edged sword in cell

proliferation and tumorigenesis. Cancer Med. *3*, 453–461.

Quinlan, A.R., and Hall, I.M. (2010). BEDTools: A flexible suite of utilities for

comparing genomic features. Bioinformatics.

Rapetti-Mauss, R., Bustos, V., Thomas, W., McBryan, J., Harvey, H., Lajczak, N.,

Madden, S.F., Pellissier, B., Borgese, F., Soriani, O., et al. (2017). Bidirectional

KCNQ1:β-catenin interaction drives colorectal cancer cell differentiation. Proc. Natl.

Acad. Sci. *114*, 4159–4164.

Reid, E.S., Williams, H., Stabej, P.L.Q., James, C., Ocaka, L., Bacchelli, C., Footitt,

E.J., Boyd, S., Cleary, M.A., Mills, P.B., et al. (2015). Seizures Due to a KCNQ2

Mutation: Treatment with Vitamin B6. In JIMD Reports, (Wiley-Blackwell), pp. 79–84.

Robbins, J. (2001). KCNQ potassium channels: Physiology, pathophysiology, and

pharmacology. Pharmacol. Ther.

Rogawski, M.A. (2000). KCNQ2/KCNQ3 K+channels and the molecular

pathogenesis of epilepsy: Implications for therapy. Trends Neurosci.

Roger, S., Gillet, L., Le Guennec, J.-Y., and Besson, P. (2015). Voltage-gated

sodium channels and cancer: is excitability their primary role? Front. Pharmacol. *6*,

152.

Rogers, M.F., Shihab, H.A., Gaunt, T.R., and Campbell, C. (2017). CScape: A tool

for predicting oncogenic single-point mutations in the cancer genome. Sci. Rep.

Rosenthal, R., McGranahan, N., Herrero, J., Taylor, B.S., and Swanton, C. (2016).

deconstructSigs: Delineating mutational processes in single tumors distinguishes

DNA repair deficiencies and patterns of carcinoma evolution. Genome Biol.

Šali, A., and Blundell, T.L. (1993). Comparative Protein Modelling by Satisfaction of

Spatial Restraints. J. Mol. Biol. *234*, 779–815.

Šali, A., Potterton, L., Yuan, F., van Vlijmen, H., and Karplus, M. (1995). Evaluation

of comparative protein modeling by MODELLER. Proteins Struct. Funct. Bioinforma.

Sanchez-Vega, F., Mina, M., Armenia, J., Chatila, W.K., Luna, A., La, K.C.,

Dimitriadoy, S., Liu, D.L., Kantheti, H.S., Saghafinia, S., et al. (2018). Oncogenic

Signaling Pathways in The Cancer Genome Atlas. Cell.

Sands, T.T., Miceli, F., Lesca, G., Beck, A.E., Sadleir, L.G., Arrington, D.K.,

Schönewolf‑Greulich, B., Moutton, S., Lauritano, A., Nappi, P., et al. (2019). Autism

and developmental disability caused by *KCNQ3* gain‑of‑function variants. Ann.

Neurol. *86*, 181–192.

Shihab, H.A., Gough, J., Cooper, D.N., Stenson, P.D., Barker, G.L.A., Edwards, K.J.,

Day, I.N.M., and Gaunt, T.R. (2013). Predicting the Functional, Molecular, and

Phenotypic Consequences of Amino Acid Substitutions using Hidden Markov

Models. Hum. Mutat.

Shorthouse, D., Riedel, A., Kerr, E., Pedro, L., Bihary, D., Samarajiwa, S., Martins,

C.P.C.P., Shields, J., and Hall, B.A.B.A. (2018). Exploring the role of stromal

osmoregulation in cancer and disease using executable modelling. Nat. Commun. *9*,

3011.

Smart, O.S., Neduvelil, J.G., Wang, X., Wallace, B.A., and Sansom, M.S.P. (1996).

HOLE: A program for the analysis of the pore dimensions of ion channel structural

models. J. Mol. Graph.

Soldovieri, M.V., Ambrosino, P., Mosca, I., Miceli, F., Franco, C., Canzoniero, L.M.T.,

Kline-Fath, B., Cooper, E.C., Venkatesan, C., and Taglialatela, M. (2019). Epileptic

Encephalopathy In A Patient With A Novel Variant In The Kv7.2 S2 Transmembrane

Segment: Clinical, Genetic, and Functional Features. Int. J. Mol. Sci. *20*.

Stansfeld, P.J., and Sansom, M.S.P.P. (2011). From Coarse Grained to Atomistic: A

Serial Multiscale Approach to Membrane Protein Simulations. J. Chem. Theory

Comput. *7*, 1157–1166.

Stansfeld, P.J., Goose, J.E., Caffrey, M., Carpenter, E.P., Parker, J.L., Newstead, S.,

and Sansom, M.S.P. (2015). MemProtMD: Automated Insertion of Membrane

Protein Structures into Explicit Lipid Membranes. Structure.

Subramanian, A., Tamayo, P., Mootha, V.K., Mukherjee, S., Ebert, B.L., Gillette,

M.A., Paulovich, A., Pomeroy, S.L., Golub, T.R., Lander, E.S., et al. (2005). Gene

set enrichment analysis: A knowledge-based approach for interpreting genome-wide

expression profiles. Proc. Natl. Acad. Sci. U. S. A.

Sun, J., and MacKinnon, R. (2017). Cryo-EM Structure of a KCNQ1/CaM Complex

Reveals Insights into Congenital Long QT Syndrome. Cell.

Takahashi, N., Chen, H.Y., Harris, I.S., Stover, D.G., Selfors, L.M., Bronson, R.T.,

Deraedt, T., Cichowski, K., Welm, A.L., Mori, Y., et al. (2018). Cancer Cells Co-opt

the Neuronal Redox-Sensing Channel TRPA1 to Promote Oxidative-Stress

Tolerance. Cancer Cell.

Tarran, R. (2004). Regulation of Airway Surface Liquid Volume and Mucus Transport

by Active Ion Transport. Proc. Am. Thorac. Soc. *1*, 42–46.

Tate, J.G., Bamford, S., Jubb, H.C., Sondka, Z., Beare, D.M., Bindal, N.,

Boutselakis, H., Cole, C.G., Creatore, C., Dawson, E., et al. (2019). COSMIC: The

Catalogue Of Somatic Mutations In Cancer. Nucleic Acids Res.

Than, B.L.N., Goos, J.A.C.M., Sarver, A.L., O'Sullivan, M.G., Rod, A., Starr, T.K.,

Fijneman, R.J.A., Meijer, G.A., Zhao, L., Zhang, Y., et al. (2014). The role of KCNQ1

in mouse and human gastrointestinal cancers. Oncogene.

Vodnala, S.K., Eil, R., Kishton, R.J., Sukumar, M., Yamamoto, T.N., Ha, N.-H., Lee,

P.-H., Shin, M., Patel, S.J., Yu, Z., et al. (2019). T cell stemness and dysfunction in

tumors are triggered by a common mechanism. Science.

Wang, J.J., and Li, Y. (2016). KCNQ potassium channels in sensory system and

neural circuits. Acta Pharmacol. Sin.

Wang, Y., Wang, L., Yin, C., An, B., Hao, Y., Wei, T., Li, L., and Song, G. (2015).

Arsenic trioxide inhibits breast cancer cell growth via microRNA-328/hERG pathway

in MCF-7 cells. Mol. Med. Rep. *12*, 1233–1238.

Warth, R., Garcia Alzamora, M., Kim, J., Zdebik, A., Nitschke, R., Bleich, M.,

Gerlach, U., Barhanin, J., and Kim, S. (2002). The role of KCNQ1/KCNE1 K+

channels in intestine and pancreas: lessons from the KCNE1 knockout mouse.

Pflügers Arch. - Eur. J. Physiol. *443*, 822–828.

Ye, J., Pavlicek, A., Lunney, E.A., Rejto, P.A., and Teng, C.H. (2010). Statistical

method on nonrandom clustering with application to somatic mutations in cancer.

BMC Bioinformatics.

Yu, H., Lin, Z., Mattmann, M.E., Zou, B., Terrenoire, C., Zhang, H., Wu, M.,

McManus, O.B., Kass, R.S., Lindsley, C.W., et al. (2013). Dynamic subunit

stoichiometry confers a progressive continuum of pharmacological sensitivity by

KCNQ potassium channels. Proc. Natl. Acad. Sci.

Zaydman, M.A., and Cui, J. (2014). PIP2 regulation of KCNQ channels: biophysical

and molecular mechanisms for lipid modulation of voltage-dependent gating. Front.

Physiol. *5*, 195.

1268