Establishment of *Etv5* gene knockout mice as a recipient model for

spermatogonial stem cell transplantation

Xianyu Zhang[1,2], Xin Zhao[2], Guoling Li[2], Mao Zhang[2], Pingping Xing[2], Zicong Li[2],

Huaqiang Yang[2], Bin Chen[1*], & Zhenfang Wu[2*]

[1] College of Animal Science and Technology, Hunan Agricultural University,

Changsha, Hunan, 410128, P.R. China.

[2] National Engineering Research Center for Breeding Swine Industry, College of

Animal Science, South China Agricultural University, Guangzhou, 510642, P.R.

China.

Correspondence: Bin Chen, Email: chenbin7586@126.com; and Zhenfang Wu, Email:

wzfemail@163.com

Abstract: Transplantation of allogeneic spermatogonial stem cells (SSCs) into the

testes could generate heterologous sperm and thus produce offspring by natural

breeding. This has a significant impact on the conservation of economically valuable

and endangered animals, as well as to the understanding of SSCs function. In this

study, we report the generation of Ets-variant gene5 (*Etv5*) knockout (KO) mice using

the CRISPR/Cas9 system. The testicular weight of $Etv5^{-/-}$ mice was significantly

lower compared to wild-type (WT) mice ($P<0.01$) at 3 weeks of age. In addition, the

germ cell layer of the seminiferous tubules gradually receded with age in $Etv5^{-/-}$ mice.

At 12 weeks of age, the tubules of $Etv5^{-/-}$ mice lacked germ cell layers

(Sertoli-cell-only syndrome) and there was a lack of sperm in the epididymis. Based

on the above observations, we determined that $Etv5^{-/-}$ mice were suitable for SSCs

transplantation. We transplanted allogeneic SSCs with or without enhanced green

fluorescent protein (EGFP) into 3 week (immature) or 7 week (mature) $Etv5^{-/-}$ mice

via efferent duct injection. Spermatogenesis was observed in all immature testes but

not in mature adult testes two months post-transplantation. In addition, we measured

the expression of heterologous gene *Etv5* and *EGFP* in recipient testicular tissue and

epididymal sperm by PCR and confirmed that sperm originated from the transplanted

donor cells. Our study provides an appropriate animal model for investigating

spermatogonial stem cell function and an alternative for mammalian breeding.

Keyword: *Etv*5, SSCs, Gene knockout, Transplantation, Spermatogenesis

Introduction

Spermatogonial stem cells (SSCs) are an important component of male germ cells. They form spermatocytes through mitosis and then produce sperm after two consecutive meiosis cycles. SSCs are located on the basement membrane of seminiferous tubules and reside in a very complex micro-environment. Although they account for only 0.02 %-0.035 % of the total germ cells in the testis, they are essential for maintaining a stable number of germ cells. This is via constant self-renewal and differentiation throughout the reproductive cycle [1]. When allogeneic SSCs are transplanted into SSCs-deficient testes, they can repair damaged tissue via spermatogenesis and regeneration [2]. Based on the biological function of SSCs, allogeneic germ cells when transplanted into the recipient's testicles have the capability to undergo spermatogenesis, produce donor-derived sperm, and generate offspring via natural conception. Using this procedure, the conservation of economically valuable and endangered animals could be achieved. In addition, this model will be significant for understanding the function of SSCs.

At present, successful SSCs-deficient animal models have been generated through natural mutations, busulan intraperitoneal or testicular injections [3, 4]. However, natural mutants are difficult to obtain, and Busan ablation is difficult to perform and has a high mortality rate [5, 6], which significantly limits its use. With the advent of gene-editing tools such as the clustered regularly interspaced short palindromic repeat (CRISPR)/CRISPR-associated 9 (Cas9) artificial nuclease system,

animal models with specific mutations could be efficiently generated. Ets-variant

gene5 *(Etv5)*, a member of the PEA3 family, is an important transcription factor

encoded by E-twenty-six (ETS) genes. It is essential for self-renewal, proliferation,

and differentiation of SSCs [7]. Tyagi *et al*. [8] observed that *Etv5* KO mice generated

by homologous recombination had a degenerative loss of SSCs in the seminiferous

tubules, had Sertoli cell-only syndrome and were infertile. Regrettably, they did not

perform follow-up studies to determine whether the microenvironment of this model

could support the re-spermatogenesis of donor SSCs, nor did they investigate suitable

transplantation techniques and transplantation times. In this study, we generated *Etv5*

gene KO mouse models using the CRISPR/Cas9 system, and then measured relevant

changes in germ cells. In addition, we performed allogeneic SSCs transplantation into

Etv5$^{-/-}$ mice testicles. Our study provides a gene-editing method to generate receptor

models for SSCs transplantation. In addition, it may provide possible avenues for the

conservation of endangered species as well as understanding spermatogonial stem cell

function.

Materials and methods

Ethics statement

All animal experiments were performed in accordance with the guidelines of the

National Institutes of Health's Guide for the Care and Use of Laboratory Animals.

Our study was approved by the IACUCs at South China Agricultural University.

Cell culture

Testes were harvested from 6 to 8 days postpartum C57BL/6 male pups, and then

were digested using a two-step enzymatic digestion protocol. Briefly, testes were

washed in DPBS with 2% penicillin-streptomycin (Gibco) and then the tunica

albuginea and convoluted epididymis were removed. Afterward, the seminiferous

tubules were digested in 5 mLs of DPBS solution I consisting of 1 mg/mL collagenase

type IV (Gibco, 17104019) and 20 U/mL DNase I (Takara, 2212) at 37 °C for 10 min

with intermittent agitation every 5 min. Samples were then centrifuged at 300 g for 5

min and the supernatants discarded. The precipitates were washed in DPBS and then

incubated in 5 mLs of solution II consisting of 20 U/mL DNase I and 0.25%

trypsin/EDTA at 37 °C for 5 min until cells were completely dispersed. Digestion was

terminated using 10% fetal bovine serum in DMEM/F12 medium (Gibco). The

samples containing single cells and clumps were filtered through a nylon cell strainer

with 40 μm pore size (B&D). Single-cell suspensions were collected in the filtrate and

then centrifuged. Supernatants were discarded and the cell pellet was washed three

times with DPBS, and then resuspended in complete medium. The number and

viability of the resulting dissociated single cells had a density of 1.06×10^7 cells/mL

with viability greater than 99%.

Lentivirus infection

Cells were infected with LPP-EGFP-Lv156-400 lentivirus expressing enhanced

green fluorescent protein (EGFP) into the genome of donor mice SSCs prior to

transplantation. Briefly, 30 μL lentivirus packaging vector (1×10^8 TU/mL) was added

into 1 mL of cell suspension and then seeded into 12-well plates. The cells were

incubated at 37 °C and 5% CO_2 atmosphere for up to 12 h. Subsequently, cells were

washed once in DPBS and then resuspended gently in DPBS using a Pasteur pipette.

Cell suspensions were then transferred into a new 15 mL centrifuge tube and

centrifuged. Supernatants were discarded and cells were re-suspended in 300 μL of

Hanks' Balanced Salt Solution (HBSS, Gibco, 14170112). The cell suspension was

then transferred to a designated transplantation suite for transplantation.

Embryo injections for KO mice production

gRNA sequences are shown in Fig. 1A. The mouse (C57BL/6) *Etv5* gene

(GenBank accession number: NM_023794.2; Ensembl*:* ENSMUSG00000013089) is

located on chromosome 16, with thirteen identifiable exons, with an ATG start codon

on exon 2 and a TAA stop codon on exon 13. The introns on both sides of exon 1 to 5

were selected as the target sites for CRISPR/Cas9-mediated genome editing. A pair of

gRNA targeting vectors (pRP[CRISPR]-hCas9-U6) was constructed and confirmed

by sequencing: gRNA1 (matching the forward strand of the gene):

GAACGGCCATTGTCGGTGGCAGG, and gRNA2 (matching the forward strand of

the gene): CTTCTATGCTAATAACGGGTGGG were selected. These two targeting

vectors were used for embryo injections, together with Cas9 mRNA. gRNA generated

by *in vitro* transcription was then co-injected into fertilized eggs for the generation of

KO mice.

Western blot and PCR analysis

CRISPR-induced mutation assays. Female and male heterozygous edited (*Etv5$^{+/-}$*)

mice were bred to generate homozygous edited (*Etv5$^{-/-}$*) offspring. DNA was extracted

from the toes of pups from the same litter at 2 days post-birth for PCR analysis and

sequencing. The following two primers were used for genotype identification by PCR;

Mouse ETV5 Primer-Forward: 5'-CAACTGGTGCCCTTCCCAGTCT-3', Mouse

ETV5 Primer-Reverse: 5'-GCCGCTCTTAAACCTGTTCATTCG-3'. Wild-type (WT)

mice were designated as the control group and $Etv5^{-/-}$ mice as the experimental group.

Polyclonal anti-ERM/ETV5 antibody targeting amino acids 244 of the human

ERM/ETV5 was used to measure ETV5 protein expression in $Etv5^{-/-}$ male and WT

mice by western blotting. The anti-β-actin monoclonal antibody was used as the

loading control. $Etv5$ expression in male mice was measured by gene and protein

expression levels.

Monitoring body and testicular weight

Animal care and use protocols were approved by our institutional review board.

Animals were provided sufficient food and water and were housed in appropriate

conditions to safeguard their welfare. Bodyweight of 5 male littermates during the

development period from 4 postnatal days to adulthood were measured and recorded.

Testes weights were determined at weekly intervals from 1 to 12 weeks of age.

H.E. **staining and immunohistochemistry**

Whole testes were fixed in 4% paraformaldehyde and then paraffin sectioned for

hematoxylin-eosin (H.E.) staining and immunohistochemistry. For H.E. staining,

paraffin sections were counterstained with hematoxylin and eosin and then observed

under an electron microscope for comparing spermatogonium between $Etv5^{-/-}$ mice

and WT littermates. Immunohistochemistry was performed after antigen retrieval with

EDTA (pH9.0),   3% hydrogen peroxide was used for blocking endogenous

peroxidase, and 3% BSA was used to block non-specific binding. Afterward, sections

were incubated with specific antibodies. Anti-promyelocytic leukemia zinc-finger

(PLZF) mouse antibody (sc-28319), a marker of undifferentiated spermatogonia, was

used at 1:100 dilution. HRP labeled goat anti-mouse IgG diluted at 1:200 was used as

the secondary antibody. The sections were then washed and developed using DAB

color rendering and nuclear staining. Hematoxylin staining of the nucleus was in blue

and brownish yellow color indicated positive expression of DAB.

Real time-PCR analysis

Total RNA samples were extracted using Eastep Super Total RNA Extraction

Kit (LS1040, Promega) according to the manufacturer's instructions. Total RNA was

then converted to cDNA using PrimeScript RT reagent Kit with gDNA Eraser

(Perfect Real, Takara). mRNA expression levels of *Ccl9*, *Prim2*, and *Cyp17a1* genes

were then measured by quantitative PCR using PowerUp™ SYBR™ Green Master

Mix (A25742). *β-actin* served as an internal control. Each tested gene from control

and experimental samples were performed in triplicate. Relative gene expression was

calculated using the $2^{-\Delta\Delta CT}$ method. Primer sequences used for RT-qPCR are shown in

Supplementary Table 1.

Measurement of Testosterone levels

Blood was collected from mice via the orbital vein and then centrifuged for serum

collection. Testosterone levels in *Etv5$^{-/-}$* male and WT control littermates at 12 weeks of

age were determined using the mouse testosterone ELISA kit. Brief, 50 µL of standard

or sample were added to each well, with blank wells left empty. Then 50 μL of

HRP-conjugate was added to each well except for the blank well. Afterward, 50 μL of

antibody was added to each well. The wells were mixed and then incubated for 1 h at

37°C. Contents from each well were then aspirated and washed 3 times. The assay plate

was blotted dry and 50 μL of substrate A and 50 μL of substrate B was added to each

well and incubated for 15 min at 37°C in the dark. After incubation, 50 μL of stop

solution was added to each well, and the optical density (OD) of each well was

recorded within 10 min using a microplate reader set to 450 nm.

Semen collection and analysis

Mice were euthanized by cervical dislocation. The unilateral epididymis was

then surgically removed, cut into pieces and immersed in 1 mL of semen diluter. After

incubation at room temperature, 10 μL sperm diluent was transferred to a

hemocytometer to determine the number and vitality of sperm. The average of 3 data

records from different mice in each group was then determined. The sperm smear test

was used to compare the sperm concentrations between $Etv5^{-/-}$ male and WT mice.

Briefly, approximately 1 mL of semen diluter (mentioned above) was centrifuged at

2000 rpm for 10 min. The resulting precipitate was re-suspended with 500 μL semen

diluter followed by the addition of trypan blue (1:2 in volume). Slides were sealed

using glycerol gelatin after 10 μL of the sample was placed onto glass slides.

Transplantation

Mice with homozygous mutations of $Etv5$ gene were used as recipients for

transplantation. Mice were anesthetized intraperitoneally (M2910, 0.2 mL/10g body

weight) and then were placed dorsally under a stereomicroscope for the

transplantation procedure. Briefly, the cuticular layer at the midline of the abdomen,

not far from the genitals, was lifted using small forceps. Then a transverse 0.3 cm

incision was made followed by an incision to the peritoneum and abdominal muscle

layer until the peritoneal cavity was visible. The lateral fat pad attached to the testis

was gently pulled until the testis was exteriorized. The efferent duct that connected

the testis to the epididymis was identified and fat tissues around the duct were gently

removed. The SSCs suspension was then carefully transferred to a 100 µL volume

microinjection syringe. The syringe was then connected into a capillary-glass tube

with an inner diameter of 40-50 µm at the tip. The cell suspension was gently forced

into the seminiferous tubules of the testis via the efferent duct by applying pressure to

the syringe. The injection pipette was held parallel to the ordinate axis of the efferent

duct. The injection rate and cell suspension flow rate were controlled manually by

monitoring the movement of the cell suspension in the tubules. Approximately 5-10

µL of the donor cell suspension was injected into each recipient testis. The other testis

was not surgically manipulated and served as a control. After transplantation, the

testis was placed back into the peritoneal cavity. The incision was closed with one

stitch of a 7.0 absorbable suture from the inside out, with the abdominal muscle layer

first, followed by the peritoneum layer, and finally the cuticular layer. Mice were then

returned back to their cages. Mice were monitored for distress and assessed regularly

until testis samples were collected.

Identification of Allogeneic SSCs

Testes of recipient mice were harvested 2 months after transplantation to

determine the functional recovery of spermatogenesis. Recipient testes were fixed in

4% paraformaldehyde and embedded in paraffin wax for histomorphology and

immunohistochemistry analysis. Heterogenous spermatogenesis was observed under a

fluorescence stereomicroscope using EGFP-positive sperm obtained from the

epididymis of recipient mice. In addition, sperm smear slides were generated.

Expression of exogenous genes in the testes and spermatozoa, i.e., *Etv5* and *EGFP*

were measured by PCR (primer sequences are shown in Supplementary Table 1).

Statistical Analysis

PASW Statistics 21 (IBM SPSS, Chicago, IL, USA) was used to determine

statistical significance and standard deviation. Bodyweight, testes weight, gene

expression, testosterone levels, and cell counts were analyzed using the student *t*-test.

Differences were considered significant at $P<0.05$ (*) and $P<0.01$(**).

Results

Generation of *Etv5* KO mice using CRISPR/Cas9

We designed two gRNAs targeting the introns on both sides of exon 1 (gRNA1)

to exon 5 (gRNA2) to delete about 5610 bp of the *Etv5* gene to generate *Etv5*$^{-/-}$ mice

by embryo injection of Cas9 mRNA and gRNAs(Fig. 1A). The genotypes of the

founder and their offspring were identified using primers F/R (Fig. 1A). DNA gel

electrophoresis results demonstrated that homozygous KO mice generated a 710bp

band, while heterozygotes had two bands, 710bp, and 6326bp, and wild type mice had

a single band of 6326bp (Fig. 1B). PCR and DNA sequencing demonstrated that

homozygous KO mice had a 5610 bp deletion (Fig. 1C). In addition, western blot

analysis confirmed a lack of ETV5 expression in homozygous $Etv5^{-/-}$ KO mice (Fig.

1D). Interestingly, $Etv5^{-/-}$ mice at 3 weeks of age were obviously thinner compared to

WT mice, and this was more pronounced at 12 weeks of age (Sup-Fig. 1A). The body

weights of $Etv5^{-/-}$ mice were markedly lower at each point from 14 postnatal days to

84 days ($P < 0.05$) compared to WT controls, with the magnitude of bodyweight

decline between 34.3% to 45.7% (Sup-Fig. 1B).

Lack of SSCs in the seminiferous tubules of *Etv5* KO mice

We measured the weight of testes in $Etv5^{-/-}$ mice from day 4 to 12 weeks of age

and compared them to WT mice of the same age. The weights were almost similar in

mice aged 4 days to 1 week. However, the testes of $Etv5^{-/-}$ mice were much smaller

compared to WT mice at 3 weeks of age (reduced by as much as 43.1%). The

difference became more evident with age, with up to a 67.4% reduction at 12 weeks

of age (Fig. 2A and B). H.E. staining demonstrated a remarkable decrease in the

number and class of testicular cells in the seminiferous tubules of $Etv5^{-/-}$ mice at 3

weeks of age. Testicular cells were absent in the tubules of $Etv5^{-/-}$ mice and

Sertoli-cell-only syndrome was observed at 12 weeks of age (Fig. 2C). Additionally,

we measured the protein expression levels of PLZF, which is a marker for SSCs. We

observed that PLZF was still expressed in KO mice in some seminiferous tubules at 3

weeks of age. However, by 12 weeks of age, SSCs in $Etv5^{-/-}$ had completely

disappeared, while they were present in WT mice (Fig. 2D). In addition, the number

of premeiotic germ cells, meiotic germ cells, and round spermatids gradually declined

in $Etv5^{-/-}$ mice, and by 12 weeks of age, were undetectable (Fig. 2E). Additionally, the

percentage of vacuolation observed in seminiferous tubules both in 3 and 12 week old

$Etv5^{-/-}$ mice were higher compared to WT control mice, reaching 100% at 12 weeks of

age (Fig. 2F). There were no differences in the number of seminiferous tubules

between $Etv5^{-/-}$ and WT mice at 3 weeks of age ($P>0.05$), however, there was a

significant difference at 12 weeks of age ($P<0.01$) (Fig. 2G).

We next examined sperm production in KO mice at different ages. The results

were consistent with H.E. staining and immunohistochemistry. At the time of sexual

maturity, there were numerous motile sperm in WT mice compared to age-matched

$Etv5^{-/-}$ mice, whose sperm numbers were extremely low (Table 1, Sup-Fig. 2A). At 12

weeks of age, numerous motile sperm were observed in the epididymis WT mice,

however, there was a lack of spermatozoa in age-matched $Etv5^{-/-}$ mice (Table 1,

Sup-Fig. 2C). In addition, we found that the $Etv5$ target gene $Ccl9$, spermatid-specific

gene $Prim2$ and interstitial gland-specific gene $Cyp17a1$ expression levels were

significantly decreased ($P<0.01$, $P<0.01$ and $P<0.05$ respectively) in $Etv5^{-/-}$ mice

(Sup-Fig. 3A, B and C). We also tested testosterone levels in WT and $Etv5^{-/-}$

littermates at 6, 8 and 12 weeks. There was a significant reduction of testosterone

levels in $Etv5^{-/-}$ mice, and at 12 weeks of age reached a maximum difference

compared to WT mice ($P<0.01$) (Sup-Fig. 3D).

Transplantation of allogeneic SSCs through the efferent duct

In order to determine whether the $Etv5^{-/-}$ mouse model could serve as a

''surrogate'', we transplanted allogeneic SSCs (expressing or not expressing EGFP)

through the efferent duct into immature (3 weeks of age) and mature (7 weeks of age)

mice (Fig. 3A). Two months post-transplantation, complete spermatogenesis was

observed in all post-transplanted immature testes with spermatozoa observed in the

epididymides (approximately $9.58 \pm 5.05 \times 10^4$) (Table 2, Fig. 3B), however, no

spermatozoa were detected in adult testes (Table 2, Sup-Fig. 4A). The proportion of

tubules with spermatogenesis was $29.51\% \pm 0.08$ and epididymides with spermatozoa

were as high as 77.78% (7/9) (Table 2). All germ cell layers could be observed in

these tubules by H.E. staining (Fig3. C) and spermatogonia were positive for PLZF

protein expression (Fig. 3D) in KO mice after SSCs transplantation. To determine the

origin of the sperm, the expression of heterologous genes *Etv5* and *EGFP* were

measured in recipient epididymal sperm by PCR. *Etv5* and *EGFP* expression were not

detected in KO mice, however, they were expressed in posttransplant *Etv5*$^{-/-}$ mice

spermatozoa (Fig. 3E). This indicated that the sperm were derived from donor cells.

Discussion

The mitogen-activated protein kinase 1 (Map2k1) signaling pathway is vital for

SSC self-renewal and is activated by fibroblast growth factor 2 (Fgf2). Activated

Map2k1 induces the expression of ETV5 which in turn up-regulates the expression of

Bcl6b that is critical for SSC self-renewal [9, 10]. Chen et al. [11] demonstrated the

presence of multiple germ cell layers in the tubules of 4 week old ERM$^{-/-}$ mice, with

gradual loss of spermatogonia. However, by 10 weeks of age, most tubules were

devoid of any germ cells, with only the presence of Sertoli cells. In this study, we

successfully generated *Etv5* KO mice using CRISPR/Cas9. These KO mice gradually

lost SSCs and only retained Sertoli cells, indicating that they were suitable models for

SSCs transplantation. This is because the loss of endogenous SSCs facilitates the

transplantation of exogenous SSCs [12].

Knockout of the *Etv5* gene affected the differentiation of SSCs and the formation

of sperm. Based on our results, only a small region of the tubules retained multilayer

germ cells, which included premeiotic germ cells, meiotic germ cells, and round

spermatids. This suggested that differentiated germ cells were lost in the majority of

tubules. The testicular sizes of 3-week-old *Etv5$^{-/-}$* mice were slightly smaller

compared to age-matched WT controls. In addition, 3-week-old Etv5-/- mice had

partial germ cell loss in the seminiferous tubules. By 12 weeks of age, the testicular

sizes of 3-week-old *Etv5$^{-/-}$* mice were much smaller compared to WT controls, with

germ cell layers lost in the tubules. These results suggest that testicular size may

reflect the extent of germ cell loss. In addition, several previous studies have

demonstrated that the maintenance of spermatogenesis was associated with

testosterone levels. Testosterone, secreted by Leydig cells in the interstitial space of

the testis, is a sex hormone responsible for maintaining spermatogenesis in the testis

[13]. The lack of testosterone in males leads to infertility. This is because

spermatogenesis does not progress beyond meiosis. In the absence of testosterone,

germ cells, even if they underwent meiosis, would detach from Sertoli cells and

undergo apoptosis [14]. Lack of testosterone in males leads to two major reproductive

disorders. First, the integrity of the blood-testis barrier (BTB) is compromised, which

exposes germ cells to autoimmune attack and cytotoxicity [15, 16]. This can be

explained by the classical testosterone signaling pathway: Testosterone induces gene

expression that alters the function of Sertoli cells. Second, defects in cell adhesion not

only make round sperm cells separate from Sertoli cells before they convert into long

sperm cells but also prevent mature sperm from being released by being engulfed by

Sertoli cells. This can be explained by the nonclassical kinase activation pathway [17,

18], i.e., kinase activation especially Erk and Src kinases. Testosterone functions in

promoting adhesion between germ cells and Sertoli cells, maturation, and release,

which are crucial to maintaining spermatogenesis. However, the direct link between

Etv5 and testosterone levels have not been demonstrated. Our results demonstrated

that continuous loss of germ cells was observed with a decrease in testosterone levels

in *Etv5*$^{-/-}$ mice.

Interestingly, 3-week-old *Etv5*$^{-/-}$ mice were thinner compared to WT mice. At 12

weeks of age, this trend was even more obvious. The reduced body weights in *Etv5*$^{-/-}$

mice indicated that ETV5 had an influence on overall growth. *Etv5* mRNA has been

detected in a variety of tissues, including the heart, lungs, thymus, lymphocytes,

kidneys, and skeletal muscles [19, 20]. The widespread expression of ETV5 during

development may be crucial for growth. In a previous study that investigated the

viability of *Etv5*$^{-/-}$ mice, it was observed that *Etv5* mRNA was abundantly expressed

in the brain, lungs, and colon, but was the most abundantly expressed in the testes

[21].

We next investigated whether our *Etv5*^{-/-} mouse model would support sperm

generation after allogeneic SSC transplantation. Although complete spermatogenesis

was visible in all post-transplantation immature *Etv5*^{-/-} mouse testes, the number and

sperm motility in the epididymes were not ideal. Possible reasons could be attributed

to the following: Immunologic rejection. It is well known that the testis is one of the

immune-privileged organs. Leydig and Sertoli cells have immunosuppressive activity

to inhibit the proliferation of lymphocytes [22], and testicular macrophages also play

an immunosuppressive role [23]. These immune-privileged conditions in the testis

may still make the transplant of donor SSCs differentiate to a certain extent even in

immunocompetent recipients. This is because the blood-testis barrier protects

allogeneic sperm from an attack from the immune system [24]. A previous study

reported that the percentage of tubules with spermatogenesis in recipient testes

increased by as much as 48.1% compared to 9.4% in control mice after treatment with

monoclonal anti-CD4 antibody [25]. Complete spermatogenesis takes approximately

34.5 days from type a- single (As) spermatogonia to mature spermatozoa in WT mice

[26]. Colonization and establishment of two-dimensional colonies are usually

complete within 2 weeks and post-2 weeks respectively after transplantation. Meioses

usually initiates within the second month, with several spermatids being observed

after 2 months post-transplantation. The degree of germ cell differentiation will

continue to increase, with normal spermatogenesis being observed 3 months

post-transplantation [27]. In addition, it takes at least 10 days for sperm to travel from

the testes to the epididymides to mature [28]. With regard to the environment in the

epididymis, a previous study [29] reported that flagellar motion is crucial for the reproductive function of sperm. Flagellar movement depends on several factors, including intracellular pH, sperm membrane, flagellar elements, and external factors such as matrix availability, ionic signals, and physical elements in the sperm microenvironment. It is not clear what factors reduce the motility of sperm in transplant recipients.

In summary, we successfully generated $Etv5^{-/-}$ mice suitable for SSCs transplantation and verified that donor spermatogonial stem cells transplanted into recipient testes could restore heterogenous spermatogenesis and produced donor-derived sperm. However, spermatogenesis and production of donor-derived sperm were not optimal and may have been due to factors, such as immune rejection, sperm recovery time, etc., resulting in unsatisfactory sperm quantity and vitality. Additional studies are required to investigate factors that contribute to sperm quality and vitality after SSC transplantation.

Acknowledgment

This work was supported by the national science and technology major project for the breeding of new transgenic organisms (2016ZX08006002) and the national natural science foundation of china (31772555).

Author Contributions

Conceived and designed the experiments: Xianyu Zhang, Huaqiang Yang, Zhenfang Wu.

Performed the experiments: Mao Zhang, Xianyu Zhang, Xin Zhao, Pingping Xing.

Analyzed the data: Xianyu Zhang, Guoling, Li.

Contributedreagents/materials/analysis tools: Xianyu Zhang, Zicong Li.

Wrote the paper: Xianyu Zhang, Guoling, Li, Huaqiang Yang.

Legend

Fig. 1. Generation of *Etv5* KO mice by CRISPR/Cas9. (A) Schematic diagram

depicting the strategy for generation of knockout *Etv5* mice, which includes gRNA1

and gRNA2 targeting sites. Primer-F and primer-R used to determine the genotype of

the mice. (B) Genotype determined by PCR for newborn male mice littermates. WT*,*

Etv5$^{+/-}$ and *Etv5*$^{-/-}$ represent the wild type, heterozygote, and homozygote, respectively.

Blue asterisks indicate complete sequence and red asterisks indicate remaining

sequence after deleting. (C) Sequencing the PCR products of *Etv5*$^{-/-}$ mice and WT

controls in figure B. (D) ETV5 protein expression in WT and *Etv5*$^{-/-}$ mice measured

by western blotting.

Fig. 2. Absence of SSCs in the seminiferous tubules of *Etv5* KO mice. (A) The

phenotype of testes of WT and *Etv5*$^{-/-}$ mice at 3 weeks and 12 weeks of age. (B)

Weight of Testes in *Etv5*$^{-/-}$ mice compared to WT littermates at periodic points during

development. For all time points, n=4 per group for both WT and *Etv5*$^{-/-}$ mice. (C and

D) Representative figures for H.E. staining and immunohistochemistry of testes in

WT and *Etv5*$^{-/-}$ mice littermates at 3 weeks and 12 weeks. Bars=50 μm. Red arrows

indicate spermatogonial stem cells with brown positive staining. Pentacles reveal

vacuolation in some seminiferous tubules in *Etv5*$^{-/-}$ mice. (E) The number of testicular

cells were mainly composed of premeiotic germ cells, meiotic germ cells, round

spermatids, and elongated spermatids in seminiferous tubules, (F) the percentage of

vacuolation of seminiferous tubules, and (G) number of seminiferous tubules from

WT and $Etv5^{-/-}$ mice at 3 weeks and 12 weeks of age at different time points, n=3 per

group for both WT and $Etv5^{-/-}$ mice. A total of 9 slices per group was prepared and

data from each slice was collected at least from 3 different regions. ** represents

$p<0.01$.

Fig. 3. Transplantation of allogeneic SSCs through the efferent duct. (A)

Transplantation of allogeneic SSCs through the efferent duct. (B) The epididymis of

3-week-old recipient mice was observed using a stereomicroscope 2 months

post-transplantation. NT and AT represent no transplantation and after transplantation.

AT-1 and AT-2 represent allogeneic SSCs with and without $EGFP$ gene expression,

respectively. The insets represent higher magnification of cells. Blue arrows indicate

sperm under visible light (left) and fluorescence (right). (C and D) H.E. staining and

PLZF immunohistochemistry of testes for NT and AT. Bars=50 μm. Red arrows

indicate spermatogonial stem cells with brown positive staining. (E) Heterogeneous

expression of $Etv5$ and $EGFP$ were measured to determine the origin of epididymal

sperm. M1 and M2 represent testis and epididyme transplanted with donor cells on the

same side, and M3 represents donor cells.

Supplementary fig. 1. Body weighs in KO mice. (A) The phenotype of WT and

$Etv5^{-/-}$ mice at 3 weeks and 12 weeks of age. Arrows indicate $Etv5$ gene KO and WT

mice. (B) Bodyweight of WT and $Etv5^{-/-}$ mice at periodic timepoints during

development from 4 postnatal days to adulthood. For all time points, n=5 for both WT

and $Etv5^{-/-}$ mice. Data presented as mean±*SD*. *p<0.05, **p<0.01.

Supplementary fig. 2. Sperm density in the epididymis at postpartum 6 to

12-week WT and $Etv5^{-/-}$ mice. Arrows indicate sperm and the insert represents the

higher magnification of sperm. For all time points, n=3 per group, Bars=50 μm.

Supplementary fig. 3. Changes in gene expression related to spermatogenesis and

testosterone levels. mRNA expression levels of *Etv5* target gene *Ccl9* (A),

spermatids-specific gene *Prim2* (B) and interstitial gland-specific gene *Cyp17a1* (C)

in postnatal week 12 testes of WT and $Etv5^{-/-}$ mice littermates. WT values were set as

100% in the histogram. Testosterone levels of WT and $Etv5^{-/-}$ littermates at 6, 8 and

12 weeks (D). n=3 per group, *p<0.05, **p<0.01. Data presented as mean±*SD*.

Supplementary fig. 4. Transplantation of allogeneic SSCs into mature mice. (A)

The epididymis of 7-week-old recipient mice was observed using a stereomicroscope

2 months post-transplantation. Semen containing epididymal tissue (left) and removal

of epididymal tissue (right). NT and AT represent no transplantation and

post-transplantation. AT-1 and AT-2 represent allogeneic SSCs with or without

EGFP gene expression, respectively. The insert depicts the higher magnification of

cells. (B and C) H.E. staining and PLZF immunohistochemistry of the testes for NT

and AT. Bars=50 μm.

References:

1. Hess RA, Cooke PS, Hofmann MC, et al. Mechanistic insights into the regulation

of the spermatogonial stem cell niche. Cell cycle. 2006; 5(11): 1164-70.

2. Yoshida S. Elucidating the identity and behavior of spermatogenic stem cells in

the mouse testis. Reproduction. 2012; 144(3): 293-302.

3. Ohta H, Tohda A, Nishimune Y. Proliferation and differentiation of

spermatogonial stem cells in the w/wv mutant mouse testis. Biology of reproduction.

2003; 69(6): 1815-21.

4. Qin Y, Liu L, He Y, et al. Testicular injection of busulfan for recipient

preparation in transplantation of spermatogonial stem cells in mice. Reproduction,

fertility and development. 2016; 28(12): 1916-25.

5. Qin Y, Liu L, He Y, et al. Testicular busulfan injection in mice to prepare

recipients for spermatogonial stem cell transplantation is safe and non-toxic. PloS one.

2016; 11(2): e148388.

6. Lin Z, Bao J, Kong Q, et al. Effective production of recipient male pigs for

spermatogonial stem cell transplantation by intratesticular injection with busulfan.

Theriogenology. 2017; 89: 365-73.

7. Jamsai D, Clark BJ, Smith SJ, et al. A missense mutation in the transcription

factor ETV5 leads to sterility, increased embryonic and perinatal death, postnatal

growth restriction, renal asymmetry and polydactyly in the mouse. PloS one. 2013;

8(10): e77311.

8. Tyagi G, Carnes K, Morrow C, et al. Loss of Etv5 decreases proliferation and Ret

levels in neonatal mouse testicular germ cells and causes an abnormal first wave of

spermatogenesis1. Biology of reproduction. 2009; 81(2): 258-66.

9. Hofmann M C. Gdnf signaling pathways within the mammalian spermatogonial

stem cell niche. Molecular and cellular endocrinology. 2008; 288(1-2): 95-103.

10. Ishii K, Kanatsu-Shinohara M, Toyokuni S, et al. FGF2 mediates mouse

spermatogonial stem cell self-renewal via upregulation of Etv5 and Bcl6b through

MAP2K1 activation. Development. 2012; 139(10): 1734-43.

11. Chen C, Ouyang W, Grigura V, et al. ERM is required for transcriptional control

of the spermatogonial stem cell niche. Nature. 2005; 436(7053): 1030-4.

12. Kubota H, Brinster RL. Spermatogonial stem cells. Biology of reproduction.

2018; 99(1): 52-74.

13. McLachlan RI, O'Donnell L, Meachem SJ, et al. Identification of specific sites of

hormonal regulation in spermatogenesis in rats, monkeys, and man. Recent progress

in hormone research. 2002; 57(1): 149-79.

14. Walker WH. Testosterone signaling and the regulation of spermatogenesis.

Spermatogenesis. 2011; 1(2): 116-20.

15. Meng J, Greenlee AR, Taub CJ, et al. Sertoli cell-specific deletion of the

androgen receptor compromises testicular immune privilege in mice. Biology of

reproduction. 2011; 85(2): 254-60.

16. Willems A, Batlouni SR, Esnal A, et al. Selective ablation of the androgen

receptor in mouse sertoli cells affects sertoli cell maturation, barrier formation and

cytoskeletal development. PLoS one. 2010; 5(11): e14168.

17. Holdcraft RW, Braun RE. Androgen receptor function is required in Sertoli cells

for the terminal differentiation of haploid spermatids. Development. 2004; 131(2):

459-67.

18. O'Donnell L, McLachlan RI, Wreford NG, et al. Testosterone withdrawal

promotes stage-specific detachment of round spermatids from the rat seminiferous

epithelium. Biology of reproduction. 1996; 55(4): 895-901.

19. Liu Y, Jiang H, Crawford HC, et al. Role for ETS domain transcription factors

Pea3/Erm in mouse lung development. Developmental biology. 2003; 261(1): 10-24.

20. T'Sas F, Brenner C, Mauen S, et al. Expression of the Ets transcription factor Erm

is regulated through a conventional PKC signaling pathway in the Molt4

lymphoblastic cell line. FEBS letters. 2005; 579(1): 66-70.

21. Schlesser HN, Simon L, Hofmann M, et al. Effects of ETV5 (ets variant gene 5)

on testis and body growth, time course of spermatogonial stem cell loss, and fertility

in mice. Biology of reproduction. 2008; 78(3): 483-9.

22. Wyatt CR, Law L, Magnuson JA, et al. Suppression of lymphocyte proliferation

by proteins secreted by cultured Sertoli cells. Journal of reproductive immunology.

1988; 14(1): 27-40.

23. Pöllänen P, Maddocks S. Macrophages, lymphocytes and MHC II antigen in the

ram and the rat testis. Reproduction. 1988; 82(2): 437-45.

24. Qu N, Naito M, Li J, et al. Xenogeneic and endogenous spermatogenesis

following transplantation of rat germ cells into testes of immunocompetent mice.

Reproduction, fertility and development. 2012; 24(2): 337-43.

25. Kanatsu-Shinohara M, Ogonuki N, Inoue K, et al. Allogeneic offspring produced

by male germ line stem cell transplantation into infertile mouse testis. Biology of

reproduction. 2003; 68(1): 167-73.

26. Oakberg EF. Duration of spermatogenesis in the mouse and timing of stages of

the cycle of the seminiferous epithelium. American Journal of Anatomy. 1956; 99(3):

507-16.

27. Nagano M, Avarbock MR, Brinster RL. Pattern and kinetics of mouse donor

spermatogonial stem cell colonization in recipient testes. Biology of reproduction.

1999; 60(6): 1429-36.

28. Kanatsu-Shinohara M, Morimoto H, Shinohara T. Fertility of male germline stem

cells following spermatogonial transplantation in infertile mouse models. Biology of

reproduction. 2016; 94(5): 112-1.

29. Goossens E, De Block G, Tournaye H. Computer-assisted motility analysis of

spermatozoa obtained after spermatogonial stem cell transplantation in the mouse.

Fertility and sterility. 2008; 90(4): 1411-6.

A

Etv5 KO mice

B

C

WT

Deletion (5610bp)

Etv5⁻ᐟ⁻

D

Fig.1

Fig.2

Fig.3

Table 1. Number and motility of sperm in one side epididymis between WT and *Etv5⁻/⁻* mice

Age (w)	Sperm		Motility	
	WT	*Etv5⁻/⁻*	WT	*Etv5⁻/⁻*
6	$3.73\pm0.49\times10^{6}$ [c]	$6.25\pm0.40\times10^{3}$ [e]	$68.92\%\pm2.51$ [c]	$20.01\%\pm2.08$ [e]
8	$6.13\pm0.33\times10^{6}$ [b]	$2.88\pm0.25\times10^{4}$ [d]	$69.33\%\pm0.47$ [b]	$47.63\%\pm0.75$ [d]
12	$7.50\pm0.32\times10^{6}$ [a]	ND	$88.01\%\pm1.50$ [a]	ND

ND: No sperm can be detected; WT: wild type

Different letters (a, b, c, d and e) within a column represent significance (P < 0.01).

Table 1

Table 2. Development of allogeneic spermatogenesis in *Etv5⁻ᐟ⁻* recipient mice

Recipient mice		Spermatogenesis after 2 months transplantation							
		No transplantation			After transplantation				
Age (w)	number	Testes	Tubles	Epididymides	Testes	Tubles	epididymides	Sperm	Motility
3	9	ND	ND	ND	9/9(100%)	29.51 ±0.08	7/9(77.78%)	9.58±5.05×10⁴	0
7	9	ND	ND	ND	ND	ND	ND	ND	ND

ND: No sperm can be detected.